

OPREA ADRIAN, SÎRBU CULIȚĂ, ELIÁŠ PAVOL jun., FERUS PETER

*J. Plant Develop.*  
19(2012): 141-156

## NEW DATA ADDITION TO THE ROMANIAN ALIEN FLORA

OPREA ADRIAN<sup>1</sup>, SÎRBU CULIȚĂ<sup>2</sup>, ELIÁŠ PAVOL jun.<sup>3</sup>, FERUS PETER<sup>4</sup>

**Abstract:** A number of seven plant taxa are presented in this paper, some of them being now for the first time reported in Romania (e. g. *Cenchrus longispinus*, *Panicum miliaceum* subsp. *ruderales*; *Panicum miliaceum* subsp. *agricolum*; *Robinia neomexicana*), while other taxa are mentioned for the first time in the flora of some historical provinces (e. g. *Oenothera parviflora* L. in Oltenia, or. *Setaria faberi* R. A. W. Herrm. in Moldavia); other taxa are newly identified in the flora of various counties (e. g. *Reynoutria × bohémica* in Bihor, Sălaj, Cluj, Caraș-Severin, Vâlcea, Prahova, and Bacău counties; *Reynoutria sachalinensis* in Caraș-Severin County).

**Keywords:** alien plant species, flora, new records, Romania

### Introduction

Biological invasions of alien species are large-scale phenomena of widespread importance and represent one of the major current threats to economic value, biological diversity and function of invaded ecosystems [HULME, 2007; LAMBDON & al. 2008].

Identification and reporting of alien species entering on a certain area, and assessing their invasive character, are the first important steps in developing national or regional strategies to prevent negative effects of biological invasions.

According to LAMBDON & al. (2008) and PYŠEK & al. (2009), a constant increase in the number of neophyte species was noticed in the whole Europe, especially in the last two centuries; on average, 6.2 new alien plant species are naturalized in Europe in each year.

Similar to the situation on the European level, research conducted in recent years also showed a continuous enrichment with neophytes of the Romania's flora. According to our estimations, through the contribution of many authors (see SÎRBU & OPREA, 2011 and NEGREAN, 2011, for extensive reference lists), a number of 47 new alien plant species were registered in the Romania's flora, after the year of 2000. This suggests that further floristic investigations are required for a better knowledge of the alien flora.

Our paper is a new contribution in this regard, including: i) newly registered taxa in Romania's flora, and, ii) new chorological contributions of some species previously reported in the literature.

<sup>1</sup> "Anastase Fătu" Botanical Garden, "Alexandru Ioan Cuza" University, 7-9 Dumbrava Roșie St., 700487, Iași – Romania

<sup>2</sup> University of Agricultural Sciences and Veterinary Medicine Iași, Faculty of Agriculture, 3, Mihail Sadoveanu Alley, Iași – Romania, e-mail: culita69@yahoo.com

<sup>3</sup> Slovak Agricultural University, Faculty of Agrifood Resources, Tr. A. Hlinku 2, 949 76, Nitra – Slovakia

<sup>4</sup> Mlynský Arboretum, Slovak Academy of Sciences, Vieska nad Zitavou 178, 95152, Slepčany – Slovakia

### Materials and methods

The floristic and chorological data in this paper are based on lately field investigations, conducted in different regions and localities of Romania. For each identified species, there are given information concerning their general distribution all over the world, previously reported occurrence, as well as their current distribution in Romania's flora.

The herbarium vouchers are deposited in the general herbarium of the University of Agricultural Sciences and Veterinary Medicine of Iași (IASI). The species nomenclature is given according to *Flora Europaea* [TUTIN & al. 1964-1980, 1993]. Terminology and definitions recommended by RICHARDSON & al. (2000) and PYŠEK & al. (2004) were used for establishing the status of alien plants.

### Results and discussion

During our field surveys in the last six years, there have been identified other new vascular plants in the spontaneous and/or sub-spontaneous flora of Romania, as *Reynoutria* × *bohemica* Chrtek et Chrtková, *Reynoutria sachalinensis* (F. Schmidt) Nakai., *Panicum miliaceum* L. subsp. *agricolum* Scholz et Mikoláš, *Panicum miliaceum* L. subsp. *ruderales* (Kitag.) Tzvelev, *Robinia neomexicana* Gray, *Cenchrus longispinus* (Hack.) Fernald, *Oenothera parviflora* L., and *Setaria faberi* R. A. W. Herrm.

Among these alien taxa, two are xenophytes, being introduced accidentally (e. g. *Cenchrus longispinus* and *Setaria faberi*), while the others were intentionally introduced by humans for different uses (mainly as ornamental plants) and, subsequently, escaped from cultivation in the wild (= hemerophyte species).

Having in mind the number of localities in which these plants were identified in Romania, the following taxa show an invasive tendency, as they are: *Cenchrus longispinus*, *Reynoutria* × *bohemica*, *Reynoutria sachalinensis* etc.

All of these taxa are integrated into anthropogenic habitats. Some of these species, as: *Reynoutria* × *bohemica*, *Reynoutria sachalinensis* also penetrate into the natural habitats, as: riverbanks, river meadows, grasslands etc.

#### a) Newly registered taxa in Romania's flora

##### *Cenchrus longispinus* (Hack.) Fernald, *Rhodora* **45**: 388 (1943)

This species of Poaceae family is native in the United States of America and in southern parts of Canada, in Mexico, Central America and the West Indies [VERLOOVE & SÁNCHEZ-GULLÓN, 2012]. It is also naturalised in Australia, South Africa, the Mediterranean area of Europe, Asia in Middle East (Israel) and northern Africa (Morocco) [VERLOOVE & SÁNCHEZ-GULLÓN, 2012].

According to some authors [cited in VERLOOVE & SÁNCHEZ-GULLÓN, 2012] it seems that the naturalization of this species in Europe has begun on the Adriatic Sea coasts (in Italy), where it is known at least from the year of 1933, today being a fully naturalized species on a large scale. Still, the presence of this species in the European flora has been neglected until recently, being confused in literature, as well as in herbaria collections, with *Cenchrus spinifex* Cav. (*C. incertus* M. A. Curtis; *C. pauciflorus* Benth.) [VERLOOVE & SÁNCHEZ-GULLÓN, 2012].

Nowadays, in Europe, *C. longispinus* is spread in Italy, France, Greece, Croatia, Hungary and Ukraine [VERLOOVE & SÁNCHEZ-GULLÓN, 2012].

In Romania, *C. longispinus* has been identified by us on the Black Sea coasts, on sandy beaches along the seashore, at Mamaia resort (Constanța County) (leg. Oprea, 2007 & 2009); further, it has been identified in central railway station of Galați town, among the rails (leg. Sîrbu, Oprea, Eliáš, Ferus, 2011), being erroneously identified and published as *C. incertus* M. A. Curtis [SÎRBU & al. 2011].

Revision of our herbarium material collected in Romania led us to a correctly identification of this plant species, being a newly registered one in the Romanian vascular flora.

According to data in the taxonomical references on this species [CHASE, 1920; FERNALD, 1943; VERLOOVE & SÁNCHEZ-GULLÓN, 2012; WARD, 2010], *C. longispinus* differs by *C. spinifex* Cav., as it is stated in the next identification key:

- Involucres with 30-50 spines, the outer ones numerous, shorter and slender (bristles-like), patents to reflexed; the inner spines longer and stout, terete, not or hardly flattened at base, at most 1 mm wide; spikelets of 5-6 mm long; plants always annuals .....  
 ..... *C. longispinus*
- Involucres with c. 20-30 spines usually, shorter, very stout; the outer ones fewer to almost lacking, bristle-like, reflexed; the inner ones stout, more or less conical, distinctly flattened, their bases up to 3 mm wide; spikelets of 5.8 mm long; plants annual or more often paucennial ..... *C. spinifex*

In addition, in *Jepson Manual of Vascular Plants of California* [BALDWIN & al. 2012] there are showed other diagenomas, in order to help to discriminate between these two species (sheats strongly keeled at *C. longispinus* (Hack.) Fernald vs. sheats compressed, but not strongly keeled at *C. spinifex* Cav.).

Till now, in Romania's flora, the species *Cenchrus incertus* M. A. Curtis (Syn. *C. carolinianus* Walter; *C. pauciflorus* Benth.) has been reported only, as being present along the Black Sea shorelines at Vama Veche [CIOCÂRLAN, 1991], Constanța harbour [CIOCÂRLAN & al. 2004], Jurilovca at Doloșman Cape (Tulcea County) [OȚEL, 1995 & 2006; DOROFTEI & al. 2011], Măcin (Tulcea County) [CIOCÂRLAN & al. 2004], Jijila (Tulcea County) [OPREA, 2005] (Fig. 1, a-d).

In *Flora Europaea* [CLAYTON, in TUTIN & al. 1980, p. 264] there are given three species of *Cenchrus*, namely: 1. *Cenchrus ciliaris* L. - originated in Africa and South-western Asia, distributed in Sicilia and Isole Lipari (Italy), only; 2. *C. incertus* M. A. Curtis – originated in tropical and warm-temperate regions of America's and naturalized in the centre of Mediterranean region (Italy, Corsica, Azores?, France?, Spain?); 3. *C. longispinus* (Hack.) Fernald – of North and Central American origins, naturalized in Southern Europe.

It is not excluded the possibility that other herbarium material collected in Romania to show the same confusion between the two species and therefore is need to be reviewed other herbarium collections.

***Panicum miliaceum* L. Sp. Pl. ed. I: 58 (1753)**

(Proso) Millet is a species originating in China and Central Asia. It is an annual herbaceous, therophyte plant species, flowering in June-August.

#### NEW DATA ADDITION TO THE ROMANIAN ADVENTIVE FLORA

It has been cultivated extensively in the past, before the advent of maize (*Zea mays* L.) in Romania (XVII century); at those times people prepared a kind of traditionally food made from the grains of (proso) millet, boiled in water. Nowadays, the (proso) millet is cultivated as a fodder plant, only. PRODAN (1935) cited this plant as a sub-spontaneous one in Danube Delta's flora. It is sporadically met in many localities in Romania's flora, being described also a plant association with this species as a dominant one, namely Ass. *Erigeron canadensis*-*Panicetum miliacei* Ștefan 1993 [ȘTEFAN, 1993].

As a result of field surveys made in the last few years, there were identified two subspecies of *Panicum miliaceum*, one of these being a newly registered infrataxa in Romania's flora, namely:

***Panicum miliaceum* L. subsp. *agricolum* Scholz et Mikoláš, *Thaiszia*, 1: 33-36 (1991)**

This wild subspecies of the (Proso) Millet has been described from Austria (Carinthia) [SCHOLZ & MIKOLÁŠ, 1991].

According to SCHOLZ & MIKOLÁŠ (1991), this subspecies differs of subsp. *ruderalis* by relatively dense, nodding panicle, persistent glumes at maturity (glumes not disarticulating from the pedicel); from subsp. *miliaceum* (a taxa only in cultivation) it differs by fruits easily falling off at maturity, and lighter and thinner rippen cariopses.

It is distributed in the wild in Europe, as: Italy (identified early in 1842, in Herbarium Firenze, cf. SCHOLZ & MIKOLÁŠ, 1991), Austria [SCHOLZ & MIKOLÁŠ, 1991], Czech Republic [SCHOLZ & MIKOLÁŠ, 1991], Slovakia [SCHOLZ & MIKOLÁŠ, 1991], Slovenia [MELZER, cited by SCHOLZ & MIKOLÁŠ, 1991], Hungary [KOVÁCS & SZABÓ, cited by SCHOLZ & MIKOLÁŠ, 1991; TERPÓ-POMOGYI, 1976], France [SCHOLZ & MIKOLÁŠ, 1991].

In Asia it was found in Afghanistan [SCHOLZ & MIKOLÁŠ, 1991].

This infrataxa seems not to grow in South and North America (cf. SCHOLZ & MIKOLÁŠ, 1991).

This subspecies was identified in the next localities, in the Eastern part of Romania:

- Galați-West toward the village of Barboși (Galați County), in ruderal places (leg. Sîrbu & Oprea, 01.08.2011);
- the village of Cudalbi (Galați County), in ruderal places (leg. Sîrbu & Oprea, 01.08.2011);
- the neighborhood of Borzești (Onești town, Bacău County), on fallow grounds (leg. Sîrbu, 09.07.2012) (Fig. 2);
- the village of Crișan (Tulcea County), in ruderal places (leg. Sîrbu & Oprea, 15.09.2011).

The genetic samples, analyzed on individuals from Slovakia, relieved the next results:  $2n=36$  [SCHOLZ & MIKOLÁŠ, 1991].

*Flora Europaea* counts only the species of (Proso) Millet (*Panicum miliaceum* L.) [CLAYTON, in TUTIN & al. 1980, p. 261].

***Robinia neomexicana* Gray, *Pl. Nov. Thurb.* 314 (1854) var. *neomexicana***

(Syn.: *Robinia luxurians* (Dieck) Schneid. ex Tarouca et Schneid., *R. neomexicana* var. *luxurians* Dieck, *R. neomexicana* var. *subvelutina* (Rydb.) Kearney et Peebles, *R. rusbyi* Wooton et Standl.)

New Mexican Locust, New Mexico, Southwest, Desert, Pink, or Rose Locust is a tree, up to 10 m high, often shrub; shoots glandular-hairy, with stipular-subulate spines;

leaves impari-pinnate up to 20 cm long, with (13-) 15-21 oblong-elliptical leaflets, 2-3.5 cm long, rounded or gradually acute at the apex, silky hairy on the underside, and pubescent, non-glandular rachis; flowers pink-whitish, ca 20 mm long, in multiflowers, pendent racemes, with peduncle and rachis glandulous-hairy; legumes of 6-10 cm long, glandular setaceous; V-VII.

Origins: Centre of the Northern America and Northern Mexico. Introduced in Europa.

Cultivated in Romania: the Arboretum of Gurahonț (Arad County) [ZANOSCHI & al. 2006], and the Botanic Garden “Anastasiu Fătu” of Iași [DUMITRIU-TĂTĂRANU, 1960].

This taxa has been identified as a subsontaneous one, along the railways, in the western parts of Iași city, in the neighborhoods of “Canta” and “Alexandru cel Bun” [leg. Sîrbu & Oprea, 2012] (Fig. 3, a-c).

*Flora Europaea* does not mention this plant species in Europe [BALL, in TUTIN & al. 1968, p. 106].

#### b) Alien species identified in new localities of Romania

*Panicum miliaceum* L. subsp. *ruderales* (Kitag.) Tzvelev, *Zlaki S. S. S. R.* (1976) (Syn. *P. miliaceum* L. var. *ruderales* Kitag., *P. spontaneum* Lyssov ex Zhuk., nom. illegit.)

This wild subspecies of the (Proso) Millet or Weed-Broomcorm Millet has been described from China (Manshuria), as var. *ruderales*, in 1937 [KITAGAWA, 1937]. It has structures developed for an effective natural fruit dispersal at maturity, as well as the brittle spikelets (cariopses are falling off together with the glumes), in contrast to the grains of the subsp. *miliaceum* (under cultivation in Romania, only), which are not readily falling off at their maturity [LYSSOV, 1975; BOUGH et al. 1986].

In Europe, this subspecies is distributed in the wild, in: Italy (the earliest record in Europe, in 1888) [LIPPERT, 1984], Austria [MELZER, 1982, 1983, 1984, 1987, 1988; SCHOLZ, 1983], Germany [LIPPERT, 1984], Hungary [TERPÓ-POMOGYI, 1976], Ukraine [MOSYAKIN, 1991], Czech and Slovakia [JEHLÍK, 1986]; also, probably exist in the wild flora of Poland [FREY & al. 1981], Latvia [TABAKA & al. 1988], and Lithuania [GUDZHINSKAS, in SCHOLZ & MIKOLAŠ, 1991].

On other continents, *Panicum miliaceum* L. subsp. *ruderales* is distributed in: U. S. A. and Canada [CARPENTER & HOPEN, 1985; CAVERS & BOUGH, 1985; MOORE & CAVERS, 1985; BOUGH & al. 1986; COLOSI & al. 1988], and various Asian countries [TZVELEV, 1976].

This infrataxa has been identified also in the wild in Romania, since the early 1978-1980, when Sakamoto and Kobayashi made two field investigations in here [SAKAMOTO & KOBAYASHI, 1982a, 1982b]. These two Japanese authors cited this subspecies growing along the road from Iași toward the Experimental Farm of Agronomical Institute.

We also identified this infrataxa in other localities, in the Eastern part of Romania, as:

- the village of Lețcani (Iași County), in maize fields (leg. Sîrbu, 15.09.2007);
- between the villages of Ursoaia and Horlești (Iași County), in maize fields (leg. Oprea, 24.10.2012);
- the village of Horlești, in maize fields (leg. Eliáš, Ferus, Sîrbu, 16.08.2011);

**NEW DATA ADDITION TO THE ROMANIAN ADVENTIVE FLORA**

– the village of Tăcuta (Vaslui County), in maize fields (leg. Sîrbu & Oprea, 02.08.2009) (Fig. 4).

The genetic samples, analyzed on individuals from Slovakia, relieved next results:  $2n=36$ , and  $2n=c. 36$  [SCHOLZ & MIKOLÁŠ, 1991].

These three subspecies of *Panicum miliaceum* L. identified in Romania's flora could be discriminated by the next traits (Tab. 1):

**Tab. 1.** Infrataxa determination of *Panicum miliaceum* L. (cf. Scholz & Mikoláš 1991):

Specifications	subsp. <i>ruderales</i>	subsp. <i>agricolum</i>	subsp. <i>miliaceum</i>
panicle	erect, with branches erect-patent, lax	relative densely, nodding	densely, nodding
glumes	falling off	persistent	persistent
fruits	falling off	falling off, smaller	persistent, greater

*Oenothera parviflora* L. *Syst. Nat.* ed. 10, 2: 998 (1759)

(Syn.: *Oenothera cruciata* Nutt. ex G. Don)

Small-flowered evening-primrose is a herbaceous biennial or short-lived perennial species, up to 2 m high, originated in North America [RAVEN, in TUTIN & al. 1968], introduced in Europe early in 1871 (in England) [MIHULKA & PYŠEK, 2001]. It is an adventive plant in ex-Czechoslovakia, France, Germany, Netherland, Hungary, Italy, Norway, Poland [RAVEN, in TUTIN & al. 1968], Great Britany (rare nowadays) [ROSTAŃSKI, 1982; SELL & MURRELL, 2009], R. of Moldova [BURAC & MITITELU, 1995]. On other continents, it is spread in Asia, Southern Africa and New Zealand [HOCH & WAGNER, 2007].

It grows in open habitats, disturbed grounds [WEAKLEY, 2007], along the roads, from the sea level to ca 1000 m alt. (in China) [HOCH & WAGNER, 2007].

In Romania's wild flora, *O. parviflora* was previously reported in some localities from Danube Delta (Tulcea County), Moldavia, Maramureş, and Transylvania [different authors, cited by SÎRBU & OPREA, 2011].

However, as it was shown in some references of the romanian literature [e.g. SÎRBU & OPREA, 2011a], the name of *O. parviflora* is an misapplied synonyme for *O. muricata*. This is a reason for uncertainty of the presence and chorology of these two species in Romania.

It is a newly identified species for the historical province of Oltenie, in the villages of Stolniceni and Marcea (Vâlcea County) [leg. Sîrbu & Oprea, 2012] (Fig. 5).

A reliable identification key for all species of *Oenothera* in European flora is presented in a paper of ROSTAŃSKI (1982), while an identification key for all alien species of *Oenothera* in Romania's flora is given in SÎRBU & OPREA (2011a).

*O. parviflora* is a diploid plant species ( $2n=14$ ) [CIOCÂRLAN, 2009].

*Setaria faberi* R. A. W. Herrm. *Beitr. Biol. Pflanzen*, 10: 51 (1910) (as “*faberii*”)

The giant foxtail (Chinese foxtail, Chinese millet, giant bristlegrass, nodding foxtail) is a neophyte grass, originated in East Asia. It is distributed as an alien plant in Northern and Central America, Central Europe and Russia [HITCHCOCK, 1950; ZHENG-YI & RAVEN, 1994; DARBYSHIRE, 2003; NURSE & al. 2009].

*S. faberi* entered accidentally into European flora in the beginning of the last century, being mentioned till now from Czech Republic (1961) [PYŠEK & al. 2002], Belgium (1977) [VERLOOVE, 2006], Austria (1981) [ESSL & RABITSCH, 2002], Ukraine [MOSYAKIN & YAVORSKA, 2002], Greece and Creta [BERGMEIER, in GREUTER & RAUS, 2007], European part of Russia (1985), Lithuania (1988), Germany and Sweden [NOBANIS], France, Italy, Portugal, Slovenia, Spain (naturalized), Azore Islands, Great Britany, and Slovakia (as casual) [DAISIE], etc.

In Romania, it has been mentioned for the first time by COSTEA (1996), from the harbour of Constanța and along the railway between Medgidia and Constanța (leg. 1993-1995). Further on, it has been identified, also, in Brăila county [CIOCÂRLAN, 2000]. Giving its resemblance to the species of *S. viridis* [HITCHCOCK, 1950; ZHENG-YI & RAVEN, 1994; NURSE & al. 2009; CIOCÂRLAN, 2009], it is possible that its distribution in Romania's flora is much larger actually, but remained unnoticed in the recent times (*S. faberi* has scabrid leaves on both parts, with long hairs on the above part; *S. viridis* is scabrid on the upper part of the leaves, only [CIOCÂRLAN, 2009]).

*S. faberi* is an annual, polyploid species ( $2n=36$ ) [NURSE & al. 2009].

It is a newly identified species for the historical province of Moldavia (being identified in the railway station of Bacău town [leg. Sîrbu, Oprea, Ferus, 2012]) (Fig. 6), as well as for Buzău County (it has been identified in the railway station of Buzău town [leg. Sîrbu, Oprea, Ferus, 2012]).

*Flora Europaea* does not mention this plant species [CLAYTON, in TUTIN & al. 1980, p. 263-264].

*S. faberi* is a tetraploid species ( $2n=36$ ) [CIOCÂRLAN, 2009].

***Reynoutria* × *bohemica*** Chrték et Chrtková, *Čas. Nár. Muz. Praha, Ser. nat.*, **152**: 120 (1983)

It is a perennial species, of hybrid origin, between *R. japonica* and *R. sachalinensis* [WEBB, in TUTIN & al. 1964], relatively newly described as a taxa, from Central Bohemia (Czech Republic) by CHRTEK & CHRTKOVÁ (1983). Nowadays, this taxa is one of the most invasive alien plant in Europe [PYŠEK & al. 2002; MANDÁK & al. 2004; SÎRBU & OPREA, 2008, 2011b], as well as in North America [ZIKA & JACOBSON, 2003; BARNEY & al. 2006; FNA].

*R. × bohemica* is known for some time in Romania, having already been identified in dozens of localities in the historical provinces of Transylvania, Muntenia, Moldavia, and Oltenia [SÎRBU & OPREA, 2011a], being designed as a recogniton species (together with *R. japonica*) for the synanthropic perennial vegetation of humid habitats (*Galio-Urticetea*), of type "*Fallopia japonica* agg. DC." [KOVÁCS, 2004, 2006; FENESI, 2004].

This species is largely distributed as invasive in Romania's flora, as it has been stated in many papers [In: SÎRBU & OPREA, 2011a].

It is a newly identified taxon in the following counties of Romania:

- Bihor County: Ștei, Petrileni, Sudrișiu, Beiuș, Răbăgani, Băile Felix, and Vașcău (leg. Sîrbu & Oprea, 2012);
- Sălaj County: Jibou and Gârbou (leg. Sîrbu & Oprea, 2012);
- Cluj County: Frășinet, Valea Ierei, Cerc, Caps, Buru (leg. Sîrbu & Oprea, 2012), Dej [leg. Sîrbu, 2012];
- Caraș-Severin County: Oțelu Roșu and Rusca Montană (leg. Sîrbu & Oprea, 2012);

#### NEW DATA ADDITION TO THE ROMANIAN ADVENTIVE FLORA

- Vâlcea County: Râmnicu Vâlcea, Stolniceni (Fig. 7), and Munteni (leg. Sîrbu & Oprea, 2012);
- Prahova County: Comarnic, Valea Sarului, Sinaia, Azuga, and Bușteni (leg. Sîrbu, 2012);
- Vrancea County: West of Focșani (at the crossroad towards Odobești, Câmpineanca and Vidra) (leg. Sîrbu, Oprea & Ferus, 2012);
- Bacău County: Căiuți (leg. Sîrbu, 2012).

***Reynoutria sachalinensis*** (F. Schmidt) Nakai, in Mori, *Enum. Pl. Cor.* **135** (1922)  
(Syn.: *Polygonum sachalinense* F. Schmidt; *Fallopia sachalinensis* (F. Schmidt) Ronse Decr.)

Giant Knotweed or Sakhalin Knotweed is a native species to East Asia (Sakhalin Islands, Kurile, Hokkaido, Honshu, Ullung-do, Korea) [MANDÁK & al. 2004; WITTENBERG, 2005; PYŠEK, 2006; ALBERTERNST & BÖHMER, 2006], similar to *R. japonica* var. *japonica* in many aspects, but is more vigorous (4-5 m height), with leaves much larger (up to 43 cm long and 27 cm wide), cordate at base, with multicells hairs on the lower side [BAILEY & al. 1996; ALBERTERNST & BÖHMER, 2006; BARNEY & al. 2006, etc.].

It is nowadays a naturalised or invasive alien species in most of Europe [WITTENBERG, 2005; MANDÁK & al. 2004; ALBERTERNST & BÖHMER, 2006], as well as in North America [ZIKA & JACOBSON, 2003; WESTON & al. 2005]

In Romania, this species is less known, being introduced early in 1901 (at Herăstrău, in Bucharest) [GRINȚESCU, in SĂVULESCU, 1952]. It was reported, till now, as an alien plant, in some localities from Transylvania [ȚOPA, 1947; OROIAN, 1998, cited by KOVÁCS, 2006; SĂMĂRGHIȚAN, 2000, 2005], South-West of Romania (in Mehedinți County) [MATACĂ, 2005], and North-East of Romania (Iași County) [SÎRBU & OPREA, 2011a].

*R. sachalinensis* is a newly identified species in the following county:

- Caraș-Severin: the villages of Rusca Montană, Rușchița (and along the road between these two villages) (Fig. 8, a-c), and Vama Marga (along the railways) [leg. Sîrbu & Oprea 2012].

*Flora Europaea* mention this species as naturalized from gardens as *R. japonica*, but much less frequently [WEBB, in TUTIN & al. 1964, 1993].

A reliable identification key for the species of *Reynoutria* from the Romania's flora is given in SÎRBU & OPREA (2011a).

#### Conclusions

All the taxa presented in this paper are alien plants (neophytes, xenophytes and hemerophytes). Some of them are new for the Romania's flora (as *Cenchrus longispinus* (Hack.) Fernald; *Panicum miliaceum* L. subsp. *agricolum* Scholz et Mikoláš; *Robinia neomexicana* Gray), while the others (*Panicum miliaceum* L. subsp. *ruderales* (Kitag.) Tzvelev; *Oenothera parviflora* L.; *Setaria faberi* R. A. W. Herrm.; *Reynoutria* × *bohemica* Chrték et Chrtková; *Reynoutria sachalinensis* (F. Schmidt) Nakai) have been identified in other regions and counties of Romania.

Some of them have an invasive character in Romania (e. g. *Reynoutria* × *bohemica* Chrték et Chrtková; *Reynoutria sachalinensis* (F. Schmidt) Nakai; *Cenchrus longispinus* (Hack.) Fernald), while the others could become invasive in near future (*Panicum*



*miliaceum* L. subsp. *runderale* (Kitag.) Tzvelev; *Panicum miliaceum* L. subsp. *agricolum* Scholz et Mikoláš; *Robinia neomexicana* Gray).

### Acknowledgements

This work was supported by CNCISIS-UEFISCDI Romania, grant PN II - IDEI\_1227, contract no. 1110 / 2009 and by ANCS-UEFISCDI Romania, grant PN II - Capacități, contract no. 474 / 07.03.2010.

### References

1. ALBERTERNST B. & BÖHMER H. J. 2006. In: NOBANIS - Invasive Alien Species Fact Sheet - *Fallopia japonica* - from: Online Database of the North European and Baltic Network on Invasive Alien Species-NOBANIS. www.nobanis.org. Date of access: 09/08/2007.
2. BALDWIN B. G., GOLDMAN D. H., KEIL D. J., PATTERSON R. W., ROSATTI TH. J. & WILKEN D. H. 2012. *The Jepson Manual: Vascular Plants of California*. 2<sup>nd</sup> ed. Berkeley, Los Angeles, London, University of California Press, 1568 pp.
3. BAILEY J. P., CHILD L. E. & CONOLLY A. P. 1996. A survey of the distribution of *Fallopia x bohemica* (Chrtek & Chrtková) J. Bailey (*Polygonaceae*) in the British Isles. *Watsonia*. **21**: 187-198.
4. BARNEY J. N., THARAYIL N., DITOMMASO A. & BHOWMIK P. C. 2006. The Biology of Invasive Alien Plants in Canada. 5. *Polygonum cuspidatum* Sieb. & Zucc. [= *Fallopia japonica* (Houtt.) Ronse Decr.]. *Canadian J. Pl. Sci.*: 887-905.
5. BOUGH M., COLOSI J. C. & CAVERS P. B. 1986. The major weedy biotypes of proso millet (*Panicum miliaceum*) in Canada. *Canad. J. Bot.* **64**: 1188-1198.
6. BURAC T. & MITITELU D. 1995. Flora vasculară din lunca Prutului (Republica Moldova). *Bul. Grăd. Bot. Iași*. **5**: 231-239.
7. CARPENTER J. L. & HOPEN H. J. 1985. A comparison of the biology of wild and cultivated proso millet (*Panicum miliaceum*): *Weed Sci.* **33**: 1188-1198.
8. CAVERS P. B. & BOUGH M. A. 1985. Proso millet (*Panicum miliaceum* L.): A crop and a weed. In: White J. (ed.), Studies on plant demography. A Festschrift for John Harper, p. 143/155. London/New York.
9. CHASE A. 1920. The North American species of *Cenchrus*. *Contr. U. S. Nat. Herb.* **22**: 45-77.
10. CHRTEK J. & CHRTEKOVÁ A. 1983. *Reynoutria x bohemica*, nový kříženec z čeledi rdesnovitých. *Čas. Nár. Muz. Praha, Ser. nat.* **152**: 120.
11. CIOCĂRLAN V. 1991. *Cenchrus incertus* M. A. Curtis în flora României. *Stud. cerc. Biol., Ser. biol. veget.* **43**(1-2): 7-8.
12. CIOCĂRLAN V. (coord.), BERCA M., CHIRILĂ C., COSTE I. & POPESCU GH. 2004. *Flora segetală a României*. București: Edit. Ceres, 351 pp.
13. CIOCĂRLAN V. 2000, 2009. *Flora Ilustrată a României. Pteridophyta et Spermatophyta*. București: Edit. Ceres, 1138 pp., 1141 pp.
14. COLOSI J. C., CAVERS P. B. & BOUGH M. A. 1988. Dormancy and survival in buried seeds of proso millet (*Panicum miliaceum*). *Canad. J. Bot.* **66**: 161-168.
15. COSTEA M. 1996. The recording of some new adventive taxa for Romania in the harbor of Constanța. *Rev. Roum. Biol., ser. Biol. végét.* **41**(2): 91-96.
16. DARBYSHIRE S. J. 2003. *Inventory of Canadian Agricultural Weeds*. Agriculture and Agri-Food Canada, Research Branch, Ottawa, Ontario, 396 pp.
17. DOROFTEI M., OPREA Ad., ȘTEFAN N. & SÂRBU I. 2011. Vascular wild flora of Danube Delta Biosphere Reserve. *Sci. Annals of Danube Delta Institute*, Tulcea. **17**: 15-52.
18. DUMITRIU-TĂTĂRANU I. 1960. *Arbori și arbuști forestieri și ornamentali cultivați în R. P. R.* Edit. Agro-Silvică, București, p. 624-629.
19. ESSL F. & RABITSCH W. (eds.). 2002. *Neobiota in Österreich*. Umweltbundesamt GmbH, Wien, 432 pp.
20. FENESI A. 2004. Két invázív növényfaj (*Reynoutria japonica* és *R. x bohemica*) sarjtelepszintű viselkedése ártéri élőhelyein. *Kolozsvári Biológus Napok / 5<sup>th</sup> Biology Days Cluj*.
21. FERNALD M. L. 1943. Virginian botanizing under restrictions. Part I. Field-studies of 1942 and 1943. *Rhodora*. **45**(538): 357-413.
22. FREY L., MIZIANTY M. & MIREK Z. 1981. Chromosome numbers of Polish Vascular Plants. *Fragm. Flor. Geobot.* **27**: 581-590.

#### NEW DATA ADDITION TO THE ROMANIAN ADVENTIVE FLORA


23. GREUTER W. & RAUS Th. (ed.). 2006. Med-Checklist Notulae, 24. *Willdenowia*. **36**: 719-730.
24. HITCHCOCK A. S. 1950. *Manual of the grasses of the United States*. 2<sup>nd</sup> edition, revised by Chase A. Washington: United States Government Printing Office, 1051 pp.
25. HOCH P. C. & WAGNER W. L. 2007. *Oenothera* L. In: *Flora of China*. **13**: 423-426.
26. HULME P. E. 2007. Biological invasions in Europe: drivers, pressures, states, impacts and responses. In: Hester R. E. & Harrison R. M. (Eds.). *Biodiversity under threat*. Royal Society of Chemistry, Cambridge, pp. 55-80.
27. JEHLÍK V. 1986. Nový návrh karanténních plevelů Československa. *Proceed. Czechoslovak Plant Protec. Conf. Brno*, September, p. 2-5.
28. KITAGAWA M. 1937. Contributio ad cognitionem florum Manchuricae, X. *Bot. Mag.*, Tokyo. **51**: 150-157.
29. KOVÁCS J. A. 2004. Syntaxonomical checklist of the plant communities of Szeklerland (Eastern Transylvania). *Szombathely, Kanitzia*. **12**: 75-150.
30. KOVÁCS J. A. 2006. Distribution of invasive alien plant species stands in Eastern Transylvania. *Szombathely, Kanitzia*. **14**: 109-136.
31. LAMBDON P. W., PYŠEK P., BAŞNOU C., HEJDA M., ARIANOUTSOU M., ESSL F., JAROŠÍ V., PERGL J., WINTER M., ANASTASIU P., ANDRIOPOULOS P., BAZOS I., BRUNDU G., CELESTI-GRAPOW L., CHASSOT Ph., DELIPETROU P., JOSEFSSON M., KARK S., KLOTZ S., KOKKORIS Y., KÜHN I., MARCHANTE H., PERGLOVÁ I., PINO J., VILÁ M., ZIKO A., ROY D. & HULME P. E. 2008. Alien flora of Europe: species diversity, temporal trends, geographical patterns and research needs. *Preslia*. **80**: 101-149.
32. LIPPERT W. 1984. Beiträge zu kritischen oder wenig beachteten Arten und Artengruppen der Bayerischen Flora. *Ber. Bayer. Bot. Ges.* **55**: 63-65.
33. LYSSOV V. N. 1975. Proso-Panicum L. In: Krotov A. S. (ed.). *Flora of cultivated plants, III, groat crops (buckwheat, millet, rice)*, p. 119-236. Leningrad.
34. MANDÁK B., PYŠEK P. & BÍMOVÁ K. 2004. History of the invasion and distribution of *Reynoutria* taxa in the Czech Republic: a hybrid spreading faster than its parents. *Preslia*. **76**: 15-64.
35. MATACĂ S. Şt. 2005. *Parcul Natural Porțile de Fier. Floră, vegetație și protecția naturii*. Craiova: Edit. Universitaria, 550 pp.
36. MELZER H. 1982. Neues zur Flora von Steiermark, XXIV. *Mitt. Naturw. ver. Steiermark*. **112**: 131-139.
37. MELZER H. 1983. Floristisch Neues aus Kärnten. *Carinthia*, II, **173/93**: 151-165.
38. MELZER H. 1984. Neues und Kritisches über Kärntner Blütenpflanzen. *Carinthia*, II, **174/94**: 189-203.
39. MELZER H. 1987. Beiträge zur Kärntner Flora. *Carinthia*, II, **177/97**: 237-248.
40. MELZER H. 1988. Über *Chorispora tenella*, einen südosteuropäisch-asiatischen Kreuzblütler, *Viola cucullata*, das Amerikanische Veilchen, und andere Pflanzenfunde in Kärnten. *Carinthia*, II, **178/98**: 561-566.
41. MIHULKA S. & PYŠEK P. 2001. Invasion history of *Oenothera* congeners in Europe: a comparative study of spreading rates in the last 200 years. *J. Biogeography*. **28**: 597-609.
42. MOORE D. R. J. & CAVERS P. B. 1985. A comparison of seedling vigour in crop and weed biotypes of proso millet (*Panicum miliaceum*). *Canad. J. Bot.* **63**: 1659-1663.
43. MOSYAKIN S. L. 1991. Novi vidomosti pro poshirennykh adventivnykh vidiv rodini Poaceae v m. Kievi. *Ukr. Bot. Zh.* **48**(3): 45-48.
44. MOSYAKIN S. L. & YAVORSKA O. G. 2002. The nonnative flora of the Kiev (Kyiv) urban area, Ukraine: a checklist and brief analysis. *Urban Habitats*. **1**: 45-65.
45. NEGREAN G. 2011. Addenda to "Flora Romaniaae", volumes 1-12. Newly published plants, nomenclature, taxonomy, chorology and commentaries (Part 1). *Kanitzia*. **18**: 89-194.
46. NURSE R. E., DARBYSHIRE S. J., BERTIN C. & DITOMMASO A. 2009. The biology of Canadian weeds. 141. *Setaria faberi* Herrm. *Can. J. Plant Sci.* **89**: 379-404.
47. OPREA Ad. 2005. *Lista critică a plantelor vasculare din România (Checklist of vascular plants of Romania)*. Edit. Univ. "Alexandru Ioan Cuza", Iași, 668 pp.
48. OȚEL V. 1995. Evaluarea și protecția genofondului sălbatic din perimetrul R. B. D. D. Inventariere floră R. B. D. D. / Nucleu Project "Danube Delta". Danube Delta Research Institute of Tulcea. Internal Report.
49. OȚEL V. 2006. Cercetări privind redelimitarea zonelor funcționale ale R. B. D. D. Identificarea și distribuția speciilor și habitatelor de interes comunitar, inventariate în perioada precedentă, în zonele de protecție integrală din R. B. D. D. / Nucleu Project "Danube Delta". Danube Delta Research Institute of Tulcea. Internal Report.
50. PRODAN I. 1935. *Conspectul florei Dobrogei: p. a Ia*. *Bul. Acad. Înalte Stud. Agron. Cluj*, V(1)/1934/ 170 pp.
51. PYŠEK P., SÁDLO J. & MANDÁK B. 2002. Catalogue of alien plants of the Czech Republic. *Preslia*. **74**: 97-186.

52. PYŠEK P., RICHARDSON D. M., REJMANEK M., WEBSTER G. L., WILLIAMSON M. & KIRSCHNER J. 2004. Alien plants in checklists and floras: towards better communication between taxonomists and ecologists. *Taxon*. **51**(1): 131-143.
53. PYŠEK P. 2006. *Fallopia japonica* Factsheet. <http://www.europe-aliens.org/pdf/>
54. PYŠEK P., LAMBDON P. W., ARIANOUTSOU M., KÜHN I., PINO J. & WINTER M. 2009. Alien vascular plants of Europe. In: DAISIE, *Handbook of alien species in Europe*. Springer Verlag, Dordrecht, pp. 43-61.
55. RICHARDSON D. M., PYŠEK P., REJMANEK M., BARBOUR M., PANETTA F. D. & WEST C. J. 2000. Naturalization and invasion of alien plants: concepts and definitions. *Diversity and Distrib.* **6**: 93-107.
56. ROSTAŃSKI K. 1982. The species of *Oenothera* L. in Britain. *Watsonia*. **14**: 1-34.
57. SAKAMOTO S. & KOBAYASHI H. 1982a. Variation and geographical distribution of cultivated plants, their relatives and weeds native to Turkey, Greece and Romania. In: Tani Y. (ed.). *Preliminary report of comparative studies on agrico-pastoral peoples in Southwestern Eurasia*, II, 1980, p. 41-104, Kyoto.
58. SAKAMOTO S. & KOBAYASHI H. 1982b. Variation and distribution of some cultivated *Triticeae* and associated weeds collected in Romania. *Notulae Botanicae Horti Agrobotanici, Cluj-Napoca*. **12**: 3-14.
59. SĂMĂRGHITAN M. 2000. Conspectul plantelor vasculare de pe Valea Gurghiului. *Marisia*. **26**(4): 43-144.
60. SĂMĂRGHITAN M. 2005. *Flora și vegetația Văii Gurghiului*. Târgu Mureș: Edit. University Press, 519 pp.
61. SĂVULESCU Tr. (coord.). 1952-1976. *Flora R. P. R. – R. S. R. I-XIII*. București: Edit. Acad. Române.
62. SCHOLZ H. 1983. Die Unkraut-Hirse (*Panicum miliaceum* subsp. *ruderales*) – neue Tatsachen und Befunde. *Pl. Syst. Evol.* **143**: 233-244.
63. SCHOLZ H. & MIKOLÁŠ V. 1991. The weedy representatives of Proso Millet (*Panicum miliaceum*, Poaceae) in Central Europe. *Thaïsia*. **1**: 31-41.
64. SELL P. D. & MURRELL G. 2009. *Flora of Great Britain and Ireland. Mimosaceae-Lentibulariaceae*. **3**, 2nd ed. Cambridge: Cambridge University Press, 595 pp.
65. SÎRBU C. & OPREA Ad. 2008. Two alien species in the spreading process in Romania: *Reynoutria* × *bohemica* Chrték & Chrtková and *Grindelia squarrosa* (Pursh) Dunal. *Cerc. Agr. Mold.* **41**(2/134): 41-50.
66. SÎRBU C. & OPREA Ad. 2011a. *Plante adventive în flora României*. Iași: Edit. „Ion Ionescu de la Brad”, 733 pp.
67. SÎRBU C., OPREA Ad., ELIÁȘ P. jun. & FERUS P. 2011b. New contribution to the study of alien flora in Romania. *J. Plant Develop.* **18**: 121-134.
68. ȘTEFAN N. 1993. Deux nouvelles associations de mauvaises herbes. *Analele Ști. Univ. “Al. I. Cuza” Iași, s. II, a. Biol. veget.* **39**: 77-80.
69. TABAKA L., GAVRILOVA G. & TATARE I. 1988. *Flora sosudistykh rasteny*. Latvyskoy S. S. R. Riga.
70. TERPÓ-POMOGYI M. 1976. Néhány egyszikű gyomnövény térhódítása Magyarországon. Különlen. *Kétesz. Egyet. Közlo.* **40**: 515-527.
71. TUTIN T. G., HEYWOOD V. H., BURGESS N. A., MOORE D. M., VALENTINE D. H., WALTERS S. M. & WEBB D. A. (eds.) (1964-1980). *Flora Europaea*, **1-5**. 1<sup>st</sup> ed., Cambridge: Cambridge University Press, 464 pp., 455 pp., 370 pp., 523 pp., 467 pp.
72. TUTIN T. G., BURGESS N. A., CHATER A. O., EDMONSON J. R., HEYWOOD V. H., MOORE D. M., VALENTINE D. H., WALTERS S. M. & WEBB D. A. (eds.) (1993). *Flora Europaea*, **1** (*Psilotaceae* to *Platanaceae*). 2<sup>nd</sup> ed., Cambridge: Cambridge University Press, 581 pp.
73. TZVELEV N. N. 1976. *Zlaki SSSR*. Leningrad.
74. ȚOPA Em. 1947. Contribuțiuni noi la cunoașterea ergasiophytophytelor din Cluj. *Bul. Grăd. Bot. Muz. Bot. Cluj*. **27**(3-4): 181-188.
75. VERLOOVE F. 2006. Catalogue of neophytes in Belgium (1800-2005). *Scripta Botanica Belgica*. **39**: 1- 89.
76. VERLOOVE F. & SÁNCHEZ-GULLÓN E. 2012. A taxonomic revision of non-native *Cenchrus* s.str. (Paniceae, Poaceae) in the Mediterranean area. *Willdenowia-Annals of the Botanic Garden and Botanical Museum Berlin-Dahlem*. **42**(1): 67-75.
77. WARD D. B. 2010. Keys to the flora of Florida – 26. *Cenchrus* (Gramineae). *Phytologia*. **92**(3): 442-450.
78. WEAKLEY A. S. 2007. *Flora of the Carolinas, Virginia, Georgia, and surrounding areas*. University of North Carolina Herbarium (NCU), North Carolina Botanical Garden; University of North Carolina at Chapel Hill, 836 pp.
79. WESTON L. A., BARNEY J. N. & DITOMMASO A. 2005. A review of the biology and ecology of three invasive perennials in New York State: Japanese knotweed (*Polygonum cuspidatum*), Mugwort (*Artemisia vulgaris*) and Pale Swallow-wort (*Vincetoxicum rossicum*). *Plant and Soil*. **277**(1-2): 53-69.
80. WITTENBERG R. (ed.). 2005. *An inventory of alien species and their threat to biodiversity and economy in Switzerland*. CABI Bioscience Switzerland Centre report to the Swiss Agency for Environment, Forests and Landscape, 416 pp.

#### NEW DATA ADDITION TO THE ROMANIAN ADVENTIVE FLORA

---

81. ZANOSCHI V., SÂRBU I. & TONIUC A. 2006. *Flora lemnoasă spontană și cultivată din România*. Edit. Univ. "Alexandru Ioan Cuza" Iași. **3**, p. 80-88.
82. ZHENG-YI W. & RAVEN P. H. (eds.). 1994. *Flora of China* (English edition. <http://www.efloras.org/>).
83. ZIKA P. F. & JACOBSON A. L. 2003. An overlooked hybrid Japanese knotweed (*Polygonum cuspidatum* × *sachalinense*; *Polygonaceae*) in North America. *Rhodora*. **105**: 143-152.
84. \*\*\*DAISIE – online Database of Delivering Alien Invasive Species Inventories for Europe). <http://www.europe-aliens.org/>
85. \*\*\*FNA (*Flora of North America Editorial Committee, eds. 1993+.* *Flora of North America North of Mexico*). 12+ vols. New York and Oxford. 1, 1993; 2, 1993; 3, 1997; 4, 2003; 5, 2005; 19, 2006; 20, 2006; 21, 2006. 22, 2000; 23, 2002; 25, 2003; 26, 2002.
86. \*\*\*NOBANIS – online Database of the North European and Baltic Network on Invasive Alien Species ([www.nobanis.org](http://www.nobanis.org). date of access: 09/08/2007).



**Fig. 1.** *Cenchrus longispinus* (Hack.) Fernald at Mamaia (Constanța County), on sandy beaches  
a. general habitus; b-c. panicle-like inflorescences; d. burs



**Fig. 2.** *Panicum miliaceum* L. subsp. *agricolum* Scholz et Mikoláš at Borzești (Bacău County),  
on fallow grounds

NEW DATA ADDITION TO THE ROMANIAN ADVENTIVE FLORA



**Fig. 3.** *Robinia neomexicana* Gray var. *neomexicana* in Iași city  
a. general habitus; b. shoots; c. racemes



**Fig. 4.** *Panicum miliaceum* L. subsp. *runderale* (Kitag.) Tzvelev at Tăcuta village (Vaslui County),  
in maize crops



Fig. 5. *Oenothera parviflora* L. at Stolniceni village (Vâlcea County)



Fig. 6. *Setaria faberi* R. A. W. Herrm. in railway station of Bacău town



**Fig. 7.** *Reynoutria* × *bohemica* Chrtek et Chrtková at Stolniceni village (Vâlcea County)



**Fig. 8.** *Reynoutria sachalinensis* (F. Schmidt) Nakai between Rusca Montană and Rușchița villages (Caraș Severin County)  
a. general habitus; b-c. leaves