

Neutrosophic Sets and Systems

Volume 9

Article 10

1-1-2015

Refined Literal Indeterminacy and the Multiplication Law of Sub-Indeterminacies

Florentin Smarandache

Follow this and additional works at: https://digitalrepository.unm.edu/nss_journal

Recommended Citation

Smarandache, Florentin. "Refined Literal Indeterminacy and the Multiplication Law of Sub-Indeterminacies." *Neutrosophic Sets and Systems* 9, 1 (2019). https://digitalrepository.unm.edu/nss_journal/vol9/iss1/10

This Article is brought to you for free and open access by UNM Digital Repository. It has been accepted for inclusion in *Neutrosophic Sets and Systems* by an authorized editor of UNM Digital Repository. For more information, please contact amywinter@unm.edu.



Refined Literal Indeterminacy and the Multiplication Law of Sub-Indeterminacies

Florentin Smarandache¹

¹ University of New Mexico, 705 Gurley Ave., Gallup, NM 87301, USA. E-mail: smarand@unm.edu

Abstract. In this paper, we make a short history about: the neutrosophic set, neutrosophic numerical components and neutrosophic literal components, neutrosophic numbers, neutrosophic intervals, neutrosophic hypercomplex numbers of dimension n , and elementary neutrosophic algebraic structures. Afterwards, their generalizations to refined neutrosophic set, respectively refined neutrosophic numerical and literal components, then refined neutrosophic numbers and refined neutrosophic algebraic structures. The aim of this paper is to construct examples of

splitting the literal indeterminacy (I) into literal sub-indeterminacies (I_1, I_2, \dots, I_r), and to define a multiplication law of these literal sub-indeterminacies in order to be able to build refined I -neutrosophic algebraic structures. Also, examples of splitting the numerical indeterminacy (\mathfrak{I}) into numerical sub-indeterminacies, and examples of splitting neutrosophic numerical components into neutrosophic numerical sub-components are given.

Keywords: neutrosophic set, elementary neutrosophic algebraic structures, neutrosophic numerical components, neutrosophic literal components, neutrosophic numbers, refined neutrosophic set, refined elementary neutrosophic algebraic structures, refined neutrosophic numerical components, refined neutrosophic literal components, refined neutrosophic numbers, literal indeterminacy, literal sub-indeterminacies, I -neutrosophic algebraic structures.

1 Introduction

Neutrosophic Set was introduced in 1995 by Florentin Smarandache, who coined the words "neutrosophy" and its derivative „neutrosophic". The first published work on neutrosophics was in 1998 see [3].

There exist two types of neutrosophic components: numerical and literal.

2 Neutrosophic Numerical Components

Of course, the *neutrosophic numerical components* (t, i, f) are crisp numbers, intervals, or in general subsets of the unitary standard or nonstandard unit interval.

Let \mathcal{U} be a universe of discourse, and M a set included in \mathcal{U} . A generic element x from \mathcal{U} belongs to the set M in the following way: $x(t, i, f) \in M$, meaning that x 's degree of membership/truth with respect to the set M is t , x 's degree of indeterminacy with respect to the set M is i , and x 's degree of non-membership/falsehood with respect to the set M is f , where t, i, f are independent standard subsets of the interval $[0, 1]$, or non-standard subsets of the non-standard interval $]^{-0}, 1^{+}[$ in the case when one needs to make distinctions between *absolute and relative* truth, indeterminacy, or falsehood.

Many papers and books have been published for the cases when t, i, f were single values (crisp numbers), or

t, i, f were intervals.

3 Neutrosophic Literal Components

In 2003, W. B. Vasantha Kandasamy and Florentin Smarandache [4] introduced the *literal indeterminacy* " I ", such that $I^2 = I$ (whence $I^n = I$ for $n \geq 1$, n integer). They extended this to *neutrosophic numbers* of the form: $a + bI$, where a, b are real or complex numbers, and

$$(a_1 + b_1I) + (a_2 + b_2I) = (a_1 + a_2) + (b_1 + b_2)I \quad (1)$$

$$(a_1 + b_1I)(a_2 + b_2I) = (a_1a_2) + (a_1b_2 + a_2b_1 + b_1b_2)I \quad (2)$$

and developed many I -neutrosophic algebraic structures based on sets formed of neutrosophic numbers.

Working with imprecisions, Vasantha Kandasamy & Smarandache have proposed (approximated) I^2 by I , yet different approaches may be investigated by the interested researchers where $I^2 \neq I$ (in accordance with their believe and with the practice), and thus a new field would arise in the neutrosophic theory.

The neutrosophic number $N = a + bI$ can be interpreted as: " a " represents the determinate part of number N , while " bI " the indeterminate part of number N .

For example, $\sqrt{7} = 2.6457\dots$ that is irrational has infinitely many decimals. We cannot work with this exact number in our real life, we need to approximate it. Hence, we

may write it as $2 + I$ with $I \in (0.6, 0.7)$, or as $2.6 + 3I$ with $I \in (0.01, 0.02)$, or $2.64 + 2I$ with $I \in (0.002, 0.004)$, etc. depending on the problem to be solved and on the needed accuracy.

Jun Ye [9] applied the neutrosophic numbers to decision making in 2014.

4 Neutrosophic Intervals

We now for the first time extend the neutrosophic number to (open, closed, or half-open half-closed) neutrosophic interval. A *neutrosophic interval* A is an (open, closed, or half-open half-closed) interval that has some indeterminacy in one of its extremes, i.e. it has the form $A = [a, b] \cup \{cI\}$, or $A = \{cI\} \cup [a, b]$, where $[a, b]$ is the determinate part of the neutrosophic interval A , and I is the indeterminate part of it (while a, b, c are real numbers, and \cup means union). (Herein I is an interval.)

We may even have neutrosophic intervals with double indeterminacy (or refined indeterminacy): one to the left (I_1), and one to the right (I_2):

$$A = \{c_1 I_1\} \cup [a, b] \cup \{c_2 I_2\}. \tag{3}$$

A classical real interval that has a neutrosophic number as one of its extremes becomes a neutrosophic interval. For example: $[0, \sqrt{7}]$ can be represented as $[0, 2] \cup I$ with $I = (2.0, 2.7)$, or $[0, 2] \cup \{10I\}$ with $I = (0.20, 0.27)$, or $[0, 2.6] \cup \{10I\}$ with $I = (0.26, 0.27)$, or $[0, 2.64] \cup \{10I\}$ with $I = (0.264, 0.265)$, etc. in the same way depending on the problem to be solved and on the needed accuracy.

We gave examples of closed neutrosophic intervals, but the open and half-open half-closed neutrosophic intervals are similar.

5 Notations

In order to make distinctions between the numerical and literal neutrosophic components, we start denoting the *numerical indeterminacy* by lower case letter “ i ” (whence consequently similar notations for *numerical truth* “ t ”, and for *numerical falsehood* “ f ”), and *literal indeterminacy* by upper case letter “ I ” (whence consequently similar notations for *literal truth* “ T ”, and for *literal falsehood* “ F ”).

6 Refined Neutrosophic Components

In 2013, F. Smarandache [3] introduced the refined neutrosophic components in the following way: the neutrosophic numerical components t, i, f can be refined (split) into respectively the following refined neutrosophic numerical sub-components:

$$\langle t_1, t_2, \dots, t_p; i_1, i_2, \dots, i_r; f_1, f_2, \dots, f_s \rangle, \tag{4}$$

where p, r, s are integers ≥ 1 and $\max\{p, r, s\} \geq 2$, meaning that at least one of p, r, s is ≥ 2 ; and t_j represents types of numeral truths, i_k represents types of numeral indeterminacies, and f_l represents types of numeral falsehoods, for $j = 1, 2, \dots, p; k = 1, 2, \dots, r; l = 1, 2, \dots, s$.

t_j, i_k, f_l are called numerical sub-components, or respectively *numerical sub-truths*, *numerical sub-indeterminacies*, and *numerical sub-falsehoods*.

Similarly, the neutrosophic literal components T, I, F can be refined (split) into respectively the following neutrosophic literal sub-components:

$$\langle T_1, T_2, \dots, T_p; I_1, I_2, \dots, I_r; F_1, F_2, \dots, F_s \rangle, \tag{5}$$

where p, r, s are integers ≥ 1 too, and $\max\{p, r, s\} \geq 2$, meaning that at least one of p, r, s is ≥ 2 ; and similarly T_j represent types of literal truths, I_k represent types of literal indeterminacies, and F_l represent types of literal falsehoods, for $j = 1, 2, \dots, p; k = 1, 2, \dots, r; l = 1, 2, \dots, s$.

T_j, I_k, F_l are called literal sub-components, or respectively *literal sub-truths*, *literal sub-indeterminacies*, and *literal sub-falsehoods*.

Let consider a *simple example of refined numerical components*.

Suppose that a country C is composed of two districts D_1 and D_2 , and a candidate John Doe competes for the position of president of this country C . Per whole country, $NL(\text{Joe Doe}) = (0.6, 0.1, 0.3)$, meaning that 60% of people voted for him, 10% of people were indeterminate or neutral – i.e. didn’t vote, or gave a black vote, or a blank vote –, and 30% of people voted against him, where NL means the neutrosophic logic values.

But a political analyst does some research to find out what happened to each district separately. So, he does a refinement and he gets:

$$\langle 0.40 \quad 0.20 \quad 0.08 \quad 0.02 \quad 0.05 \quad 0.25 \rangle \tag{6}$$

$$\langle t_1 \quad t_2; i_1 \quad i_2; f_1 \quad f_2 \rangle$$

which means that 40% of people that voted for Joe Doe were from district D_1 , and 20% of people that voted for Joe Doe were from district D_2 ; similarly, 8% from D_1 and 2% from D_2 were indeterminate (neutral), and 5% from D_1 and 25% from D_2 were against Joe Doe.

It is possible, in the same example, to refine (split) it in a different way, considering another criterion, namely: what percentage of people did not vote (i_1), what percentage of people gave a blank vote – cutting all candidates on the ballot – (i_2), and what percentage of people gave a blank vote – not selecting any candidate on the ballot (i_3). Thus, the numerical indeterminacy (i) is refined into i_1, i_2 , and i_3 :

$$\langle 0.60 \quad 0.05 \quad 0.04 \quad 0.01 \quad 0.30 \rangle \tag{7}$$

$$\langle t; i_1 \quad i_2 \quad i_3; f \rangle$$

7 Refined Neutrosophic Numbers

In 2015, F. Smarandache [6] introduced the *refined literal indeterminacy (I)*, which was split (refined) as I_1, I_2, \dots, I_r , with $r \geq 2$, where I_k , for $k = 1, 2, \dots, r$ represent types of literal sub-indeterminacies. A refined neutrosophic number has the general form:

$$N_r = a + b_1I_1 + b_2I_2 + \dots + b_rI_r, \tag{8}$$

where a, b_1, b_2, \dots, b_r are real numbers, and in this case N_r is called a *refined neutrosophic real number*; and if at least one of a, b_1, b_2, \dots, b_r is a complex number (i.e. of the form $\alpha + \beta\sqrt{-1}$, with $\beta \neq 0$, and α, β real numbers), then N_r is called a *refined neutrosophic complex number*.

An example of refined neutrosophic number, with three types of indeterminacies resulted from the cubic root (I_1), from Euler's constant e (I_2), and from number π (I_3):

$$N_3 = -6 + \sqrt[3]{59} - 2e + 11\pi \tag{9}$$

Roughly

$$N_3 = -6 + (3 + I_1) - 2(2 + I_2) + 11(3 + I_3) \\ = (-6 + 3 - 4 + 33) + I_1 - 2I_2 + 11I_3 = 26 + I_1 - 2I_2 + 11I_3$$

where $I_1 \in (0.8, 0.9)$, $I_2 \in (0.7, 0.8)$, and $I_3 \in (0.1, 0.2)$, since $\sqrt[3]{59} = 3.8929\dots$, $e = 2.7182\dots$, $\pi = 3.1415\dots$

Of course, other 3-valued refined neutrosophic number representations of N_3 could be done depending on accuracy.

Then F. Smarandache [6] defined the *refined I-neutrosophic algebraic structures* in 2015 as algebraic structures based on sets of refined neutrosophic numbers.

Soon after this definition, Dr. Adesina Agboola wrote a paper on refined *I-neutrosophic algebraic structures* [7].

They were called "*I-neutrosophic*" because the refinement is done with respect to the literal indeterminacy (I), in order to distinguish them from the refined (t, i, f)-neutrosophic algebraic structures, where "*(t, i, f)-neutrosophic*" is referred to as refinement of the neutrosophic numerical components t, i, f .

Said Broumi and F. Smarandache published a paper [8] on refined neutrosophic numerical components in 2014.

8 Neutrosophic Hypercomplex Numbers of Dimension n

The *Hypercomplex Number of Dimension n* (or *n-Complex Number*) was defined by S. Olariu [10] as a number of the form:

$$u = x_0 + h_1x_1 + h_2x_2 + \dots + h_{n-1}x_{n-1} \tag{10}$$

where $n \geq 2$, and the variables $x_0, x_1, x_2, \dots, x_{n-1}$ are real numbers, while h_1, h_2, \dots, h_{n-1} are the complex units, $h_0 = 1$, and they are multiplied as follows:

$$h_jh_k = h_{j+k} \text{ if } 0 \leq j+k \leq n-1, \text{ and } h_jh_k = h_{j+k-n} \text{ if } n \leq j+k \leq 2n-2. \tag{11}$$

We think that the above (11) complex unit multiplication formulas can be written in a simpler way as:

$$h_jh_k = h_{j+k \pmod n} \tag{12}$$

where *mod n* means *modulo n*.

For example, if $n = 5$, then $h_3h_4 = h_{3+4 \pmod 5} = h_{7 \pmod 5} = h_2$.

Even more, formula (12) allows us to multiply many complex units at once, as follows:

$$h_{j_1}h_{j_2}\dots h_{j_p} = h_{j_1+j_2+\dots+j_p \pmod n}, \text{ for } p \geq 1. \tag{13}$$

We now define for the first time the *Neutrosophic Hypercomplex Number of Dimension n* (or *Neutrosophic n-Complex Number*), which is a number of the form:

$$u+vI, \tag{14}$$

where u and v are n -complex numbers and $I =$ indeterminacy.

We also introduce now the *Refined Neutrosophic Hypercomplex Number of Dimension n* (or *Refined Neutrosophic n-Complex Number*) as a number of the form:

$$u+v_1I_1+v_2I_2+\dots+v_rI_r, \tag{15}$$

where u, v_1, v_2, \dots, v_r are n -complex numbers, and I_1, I_2, \dots, I_r are sub-indeterminacies, for $r \geq 2$.

Combining these, we may define a *Hybrid Neutrosophic Hypercomplex Number* (or *Hybrid Neutrosophic n-Complex Number*), which is a number of the form $u+vI$, where either u or v is a n -complex number while the other one is different (may be an m -complex number, with $m \neq n$, or a real number, or another type of number).

And a *Hybrid Refined Neutrosophic Hypercomplex Number* (or *Hybrid Refined Neutrosophic n-Complex Number*), which is a number of the form $u+v_1I_1+v_2I_2+\dots+v_rI_r$, where at least one of u, v_1, v_2, \dots, v_r is a n -complex number, while the others are different (may be m -complex numbers, with $m \neq n$, and/or a real numbers, and/or other types of numbers).

9 Neutrosophic Graphs

We now introduce for the first time the general definition of a *neutrosophic graph* [12], which is a (directed or undirected) graph that has some indeterminacy with respect to its edges, or with respect to its vertexes (nodes), or with respect to both (edges and vertexes simultaneously). We have four main categories of neutrosophic graphs:

- 1) The *(t, i, f)-Edge Neutrosophic Graph*.

In such a graph, the connection between two vertexes A and B , represented by edge AB :



has the neutrosophic value of (t, i, f) .

- 2) *I-Edge Neutrosophic Graph*.

This one was introduced in 2003 in the book "*Fuzzy Cognitive Maps and Neutrosophic Cognitive Maps*", by Dr. Vasantha Kandasamy and F. Smarandache, that used a different approach for the edge:



which can be just $I =$ literal indeterminacy of the edge, with $I^2 = I$ (as in *I-Neutrosophic algebraic structures*). Therefore, simply we say that the connection between vertex A and vertex B is indeterminate.

- 3) *Orientation-Edge Neutrosophic Graph*.

At least one edge, let's say AB , has an unknown orientation (i.e. we do not know if it is from A to B , or from B to A).

4) *I-Vertex Neutrosophic Graph.*

Or at least one literal indeterminate vertex, meaning we do not know what this vertex represents.

5) *(t, i, f)-Vertex Neutrosophic Graph.*

We can also have at least one neutrosophic vertex, for example vertex *A* only partially belongs to the graph (*t*), indeterminate appurtenance to the graph (*i*), does not partially belong to the graph (*f*), we can say $A(t, i, f)$.

And combinations of any two, three, four, or five of the above five possibilities of neutrosophic graphs.

If (*t, i, f*) or the literal *I* are refined, we can get corresponding *refined neutrosophic graphs*.

10 Example of Refined Indeterminacy and Multiplication Law of Sub-Indeterminacies

Discussing the development of Refined *I-Neutrosophic Structures* with Dr. W.B. Vasantha Kandasamy, Dr. A.A.A. Agboola, Mumtaz Ali, and Said Broumi, a question has arisen: if *I* is refined into I_1, I_2, \dots, I_r , with $r \geq 2$, how to define (or compute) $I_j * I_k$, for $j \neq k$?

We need to design a Sub-Indeterminacy * Law Table.

Of course, this depends on the way one defines the algebraic binary multiplication law * on the set:

$$\{N_r = a + b_1I_1 + b_2I_2 + \dots + b_rI_r | a, b_1, b_2, \dots, b_r \in M\}, \tag{16}$$

where *M* can be \mathbb{R} (the set of real numbers), or \mathbb{C} (the set of complex numbers).

We present the below example.

But, first, let's present several (possible) interconnections between logic, set, and algebra.


operators	<i>Logic</i>	<i>Set</i>	<i>Algebra</i>
	Disjunction (or) \vee	Union \cup	Addition $+$
	Conjunction (and) \wedge	Intersection \cap	Multiplication \cdot
	Negation \neg	Complement \complement	Subtraction $-$
	Implication \rightarrow	Inclusion \subseteq	Subtraction, Addition $-, +$
	Equivalence \leftrightarrow	Identity \equiv	Equality $=$

Table 1: Interconnections between logic, set, and algebra.


In general, if a Venn Diagram has *n* sets, with $n \geq 1$, the number of disjoint parts formed is 2^n . Then, if one combines the 2^n parts either by none, or by one, or by 2, ..., or by 2^n , one gets:

$$C_{2^n}^0 + C_{2^n}^1 + C_{2^n}^2 + \dots + C_{2^n}^{2^n} = (1 + 1)^{2^n} = 2^{2^n}. \tag{17}$$

Hence, for $n = 2$, the Venn diagram, with literal truth



(*T*), and literal falsehood (*F*), will make $2^2 = 4$ disjoint parts, where the whole rectangle represents the whole uni-




Venn Diagram for $n = 2$.

verse of discourse (*U*).

Then, combining the four disjoint parts by none, by one, by two, by three, and by four, one gets

$$C_4^0 + C_4^1 + C_4^2 + C_4^3 + C_4^4 = (1 + 1)^4 = 2^4 = 16 = 2^{2^2}. \tag{18}$$

For $n = 3$, one has $2^3 = 8$ disjoint parts,




Venn Diagram for $n = 3$.


and combining them by none, by one, by two, and so on, by eight, one gets $2^8 = 256$, or $2^{2^3} = 256$.

For the case when $n = 2 = \{T, F\}$ one can make up to 16 sub-indeterminacies, such as:


$$I_1 = C = \text{contradiction} = \text{True and False} = T \wedge F$$




$I_2 = Y = \text{uncertainty} = \text{True or False} = T \vee F$




$I_3 = S = \text{unsureness} = \text{either True or False} = T \underline{\vee} F$




$I_4 = H = \text{nihilness} = \text{neither True nor False} = \neg T \wedge \neg F$



$I_5 = V = \text{vagueness} = \text{not True or not False} = \neg T \vee \neg F$



$I_6 = E = \text{emptiness} = \text{neither True nor not True} = \neg T \wedge \neg(\neg T) = \neg T \wedge T$



Let's consider the literal indeterminacy (I) refined into

only six literal sub-indeterminacies as above.

The binary multiplication law

$$*: \{I_1, I_2, I_3, I_4, I_5, I_6\}^2 \rightarrow \{I_1, I_2, I_3, I_4, I_5, I_6\} \quad (19)$$

defined as:

$I_j * I_k$ = intersections of their Venn diagram representations;

or $I_j * I_k$ = application of \wedge operator, i.e. $I_j \wedge I_k$.

We make the following:

*	I_1	I_2	I_3	I_4	I_5	I_6
I_1	I_1	I_1	I_6	I_6	I_6	I_6
I_2	I_1	I_2	I_3	I_6	I_3	I_6
I_3	I_6	I_3	I_3	I_6	I_3	I_6
I_4	I_6	I_6	I_6	I_4	I_4	I_6
I_5	I_6	I_3	I_3	I_4	I_5	I_6
I_6	I_6	I_6	I_6	I_6	I_6	I_6

Table 2: Sub-Indeterminacies Multiplication Law

11 Remark on the Variety of Sub-Indeterminacies Diagrams

One can construct in various ways the diagrams that represent the sub-indeterminacies and similarly one can define in many ways the $*$ algebraic multiplication law, $I_j * I_k$, depending on the problem or application to solve.

What we constructed above is just an example, not a general procedure.

Let's present below several calculations, so the reader gets familiar:

$$I_1 * I_2 = (\text{shaded area of } I_1) \cap (\text{shaded area of } I_2) = \text{shaded area of } I_1,$$

$$\text{or } I_1 * I_2 = (T \wedge F) \wedge (T \vee F) = T \wedge F = I_1.$$

$$I_3 * I_4 = (\text{shaded area of } I_3) \cap (\text{shaded area of } I_4) = \text{empty set} = I_6,$$

$$\text{or } I_3 * I_4 = (T \underline{\vee} F) \wedge (\neg T \wedge \neg F) = [T \wedge (\neg T \wedge \neg F)] \underline{\vee} [F \wedge (\neg T \wedge \neg F)] = (T \wedge \neg T \wedge \neg F) \underline{\vee} (F \wedge \neg T \wedge \neg F) = (\text{impossible}) \underline{\vee} (\text{impossible})$$

because of $T \wedge \neg T$ in the first pair of parentheses and because of $F \wedge \neg F$ in the second pair of parentheses = (impossible) = I_6 .

$$I_5 * I_5 = (\text{shaded area of } I_5) \cap (\text{shaded area of } I_5) = (\text{shaded area of } I_5) = I_5,$$

$$\text{or } I_5 * I_5 = (\neg T \vee \neg F) \wedge (\neg T \vee \neg F) = \neg T \vee \neg F = I_5.$$

Now we are able to build refined I -neutrosophic algebraic structures on the set

$$S_6 = \{a_0 + a_1 I_1 + a_2 I_2 + \dots + a_6 I_6, \text{ for } a_0, a_1, a_2, \dots, a_6 \in \mathbb{R}\}, \quad (20)$$

by defining the addition of refined I -neutrosophic numbers:

$$(a_0 + a_1 I_1 + a_2 I_2 + \dots + a_6 I_6) + (b_0 + b_1 I_1 + b_2 I_2 + \dots + b_6 I_6) = (a_0 + b_0) + (a_1 + b_1) I_1 + (a_2 + b_2) I_2 + \dots + (a_6 + b_6) I_6 \in S_6. \quad (21)$$

And the multiplication of refined neutrosophic numbers:

$$\begin{aligned} & (a_0 + a_1I_1 + a_2I_2 + \dots + a_6I_6) \cdot (b_0 + b_1I_1 + b_2I_2 + \\ & \dots + b_6I_6) = a_0b_0 + (a_0b_1 + a_1b_0)I_1 + (a_0b_2 + \\ & a_2b_0)I_2 + \dots + (a_0b_6 + a_6b_0)I_6 + \\ & + \sum_{j,k=1}^6 a_jb_k(I_j * I_k) = a_0b_0 + \sum_{k=1}^6 (a_0b_k + \\ & a_kb_0)I_k + \sum_{j,k=1}^6 a_jb_k(I_j * I_k) \in S_6, \end{aligned} \quad (22)$$

where the coefficients (scalars) $a_m \cdot b_n$, for $m = 0, 1, 2, \dots, 6$ and $n = 0, 1, 2, \dots, 6$, are multiplied as any real numbers, while $I_j * I_k$ are calculated according to the previous Sub-Indeterminacies Multiplication Law (Table 2).

Clearly, both operators (addition and multiplication of refined neutrosophic numbers) are well-defined on the set S_6 .

References

- [1] L. A. Zadeh, *Fuzzy Sets*, Inform. and Control, 8 (1965) 338-353.
- [2] K. T. Atanassov, *Intuitionistic Fuzzy Set*. Fuzzy Sets and Systems, 20(1) (1986) 87-96.
- [3] Florentin Smarandache, *Neutrosophy. Neutrosophic Probability, Set, and Logic*, Amer. Res. Press, Rehoboth, USA, 105 p., 1998.
- [4] W. B. Vasantha Kandasamy, Florentin Smarandache, *Fuzzy Cognitive Maps and Neutrosophic Cognitive Maps*, Xiquan, Phoenix, 211 p., 2003.
- [5] Florentin Smarandache, *n-Valued Refined Neutrosophic Logic and Its Applications in Physics*, Progress in Physics, 143-146, Vol. 4, 2013.
- [6] Florentin Smarandache, *(t,i,f)-Neutrosophic Structures and I-Neutrosophic Structures*, Neutrosophic Sets and Systems, 3-10, Vol. 8, 2015.
- [7] A.A.A. Agboola, *On Refined Neutrosophic Algebraic Structures*, mss., 2015.
- [8] S. Broumi, F. Smarandache, *Neutrosophic Refined Similarity Measure Based on Cosine Function*, *Neutrosophic Sets and Systems*, 42-48, Vol. 6, 2014.
- [9] Jun Ye, *Multiple-Attribute Group Decision-Making Method under a Neutrosophic Number Environment*, Journal of Intelligent Systems, DOI: 10.1515/jisys-2014-0149.
- [10] S. Olariu, *Complex Numbers in n Dimensions*, Elsevier Publication, 2002.
- [11] F. Smarandache, *The Neutrosophic Triplet Group and its Application to Physics*, seminar Universidad Nacional de Quilmes, Department of Science and Technology, Bernal, Buenos Aires, Argentina, 02 June 2014.
- [12] F. Smarandache, *Types of Neutrosophic Graphs and neutrosophic Algebraic Structures together with their Applications in Technology*, seminar, Universitatea Transilvania din Brasov, Facultatea de Design de Produs si Mediu, Brasov, Romania, 06 June 2015.

Received: June 2, 2015. Accepted: August 12, 2015.