



Corrigendum to

“Temperature response of denitrification and anammox reveals the adaptation of microbial communities to in situ temperatures in permeable marine sediments that span 50° in latitude” published in *Biogeosciences*, 11, 309–320, 2014

A. Canon^{1,*}, J. E. Kostka^{1,**}, T. M. Gihring¹, M. Huettel¹, J. E. E. van Beusekom^{2,***}, H. Gao^{3,****}, G. Lavik³, and M. M. M. Kuypers³

¹Earth, Ocean, and Atmos. Science Dept, Florida State University, Tallahassee, Florida, USA

²Alfred Wegener Institute for Polar and Marine Research, List, Germany

³Max Planck Institute for Marine Microbiology, Bremen, Germany

* now at: St. Johns River Water Management District, Palatka, Florida, USA

** now at: Schools of Biology and Earth and Atmospheric Science, Georgia Institute of Technology, Atlanta, Georgia, USA

*** now at: Institute of Coastal Research, Helmholtz Zentrum Geesthacht, Geesthacht, Germany

**** now at: School of Ocean and Earth Sciences, Tongji University, Shanghai, China

Correspondence to: J. E. Kostka (joel.kostka@biology.gatech.edu)

In the paper “Temperature response of denitrification and anammox reveals the adaptation of microbial communities to in situ temperatures in permeable marine sediments that span 50° in latitude” by A. Canon et al. (*Biogeosciences*, 11, 309–320, 2014), the following error occurred: the first row for both denitrification and anammox should read “ T_{opt} (°C)”. The 3rd row under anammox labeled “ T_{opt} (kJ mol⁻¹)” should instead read “ E_a (kJ mol⁻¹)”.

Table 3. Optimum temperatures (T_{opt}), apparent activation energy (E_a), and ratio of rates at 5 °C relative to T_{opt} for anammox and denitrification in permeable sediments. The range of the T_{opt} peak where rates did not fall below 90 % of the optimum rate is noted in parentheses. Values of Q_{10} were calculated between T_{opt} and $T_{\text{opt}}-10$ °C.

		SGI Gulf	SGI Bay	Sylt Summer	Sylt Winter	Ymerbukta
Denitrification	T_{opt} (°C)	36 (35–37)	35	34 (26–35)	26 (24–32)	21 (17–24)
	Rate at 5 °C relative to T_{opt}	1.4 %	0.4 %	17.1 %	11.2 %	34.4 %
	E_a (kJ mol ⁻¹)	102.7 ± 2.1	123.5 ± 4.0	51.7 ± 3.8	65.0 ± 4.0	57.8 ± 4.4
	Q_{10}	3.8	5.0	2.0	2.5	2.3
Anammox	T_{opt} (°C)	N.A.	N.A.	26 (22–29)	N.D.	9
	Rate at 5 °C relative to T_{opt}	N.A.	N.A.	34.8 %	N.D.	35 %
	E_a (kJ mol ⁻¹)	N.A.	N.A.	35.0 ± 4.9	N.D.	N.D.
	Q_{10}	N.A.	N.A.	1.6	N.D.	N.D.

N.A., no detectable activity; N.D., not determined.