

Physical injury stimulates aerobic methane emissions from terrestrial plants

Z.-P. Wang¹, J. Gullledge^{2,3}, J.-Q. Zheng¹, W. Liu¹, L.-H. Li¹, and X.-G. Han¹

¹State Key Laboratory of Vegetation and Environmental Change, Institute of Botany, Chinese Academy of Sciences, Nanxincun 20, Xiangshan, Beijing 100093, China

²Department of Zoology and Physiology, University of Wyoming, Laramie, WY, USA

³Pew Center on Global Climate Change, 2101 Wilson Blvd., Arlington, Virginia, USA

Received: 21 November 2008 – Published in Biogeosciences Discuss.: 29 January 2009

Revised: 7 April 2009 – Accepted: 7 April 2009 – Published: 17 April 2009

Abstract. Physical injury is common in terrestrial plants as a result of grazing, harvesting, trampling, and extreme weather events. Previous studies demonstrated enhanced emission of non-microbial CH₄ under aerobic conditions from plant tissues when they were exposed to increasing UV radiation and temperature. Since physical injury is also a form of environmental stress, we sought to determine whether it would also affect CH₄ emissions from plants. Physical injury (cutting) stimulated CH₄ emission from fresh twigs of *Artemisia* species under aerobic conditions. More cutting resulted in more CH₄ emissions. Hypoxia also enhanced CH₄ emission from both uncut and cut *Artemisia frigida* twigs. Physical injury typically results in cell wall degradation, which may either stimulate formation of reactive oxygen species (ROS) or decrease scavenging of them. Increased ROS activity might explain increased CH₄ emission in response to physical injury and other forms of stress. There were significant differences in CH₄ emissions among 10 species of *Artemisia*, with some species emitting no detectable CH₄ under any circumstances. Consequently, CH₄ emissions may be species-dependent and therefore difficult to estimate in nature based on total plant biomass. Our results and those of previous studies suggest that a variety of environmental stresses stimulate CH₄ emission from a wide variety of plant species. Global change processes, including climate change, depletion of stratospheric ozone, increasing ground-level ozone, spread of plant pests, and land-use changes, could cause more stress in plants on a global scale, potentially stimulating more CH₄ emission globally.

1 Introduction

Methane (CH₄) is an important atmospheric trace gas, contributing to global warming and atmospheric redox chemistry. Traditionally, the only known biological source of CH₄ was a limited group of obligately anaerobic prokaryotes called methanogens. However, a recent study (Keppler et al., 2006) reported aerobic CH₄ emission from plants by an unrecognized, non-microbial mechanism, a result that has been controversial (Schiermeier, 2006; Dueck and van der Werf, 2008). The controversy has focused mainly on two scientific aspects: (i) the as yet unidentified mechanism(s) of CH₄ formation in plants, without which the source cannot be confirmed with full confidence; and (ii) how much, if at all, this plant source contributes to the global CH₄ budget.

Some studies (Dueck et al., 2007; Beerling et al., 2008; Kirschbaum and Walcroft, 2008; Nisbet et al., 2009) observed no substantial aerobic CH₄ emission from plants. However, six independent studies (Keppler et al., 2008; McLeod et al., 2008; Vigano et al., 2008; Wang et al., 2008; Brüggemann et al., 2009; Messenger et al., 2009) did detect CH₄ emission from plant tissues/compounds under aerobic conditions in the laboratory. Several studies used isotope signature analysis to confirm that the CH₄ originated directly from plant tissues/compounds rather than from microbial methanogenesis (Keppler et al., 2008; Wang et al., 2008; Brüggemann et al., 2009). Field observations (do Carmo et al., 2006; Crutzen et al., 2006; Sanhueza and Donoso, 2006; Sinha et al., 2007; Cao et al., 2008) and satellite measurements (Frankenberg et al., 2005, 2008; Miller et al., 2007) also provided indirect evidence for the possibility of aerobic CH₄ emissions by plants in the field but did not verify the source. Keppler et al. (2006) initially estimated aerobic CH₄ emission by plants to be in the range of 62–236 Tg CH₄ y⁻¹,



Correspondence to: Z.-P. Wang
(wangzpz5@ibcas.ac.cn)

about 10–40% of the total annual source. Based on a variety of constraints, subsequent analyses yielded lower but potentially still environmentally important source strengths (Houweling et al., 2006; Parsons et al., 2006; Kirschbaum et al., 2006, 2007; Butenhoff and Khalil, 2007; Ferretti et al., 2007).

As one potential constraint on source strength, Dueck and van der Werf (2008) suggested that most species of plants do not actively emit CH₄ in the field. Consistent with this suggestion, Wang et al. (2008) observed CH₄ emission from several upland shrub species but not from a much larger number of herb species from the same grassland ecosystem. Hence, it is possible that some of the negative results in other studies arise from differences in species examined. For example, Kirschbaum and Walcroft (2008) observed no substantial CH₄ emission from *Artemisia absinthium*. Hence, aerobic CH₄ emission might vary across species, even among close relatives. Some negative results could still arise from methodological differences as suggested previously (Wang et al., 2008).

Some studies found that the rate of CH₄ production is strongly affected by environmental variables that can induce physiological stress. CH₄ emission from plants increased linearly with UV radiation and/or temperature (Keppler et al., 2006; McLeod et al., 2008; Vigano et al., 2008). Physical injury as an environmental stress is common in terrestrial plants. For example, machines harvest crops; insects and ruminants graze leaves and twigs; and strong winds break twigs and detach leaves from stems. In relation to physiological stress, McLeod et al. (2008) suggested that reactive oxygen species (ROS) may have a role in CH₄ formation by plants. It is possible therefore that aerobic CH₄ emission from plants may be affected by O₂ stress or any other stress leading to ROS production.

In this study, we examined the effect of physical injury and hypoxia on CH₄ emissions from 10 species of the *Artemisia* genus sampled from the grasslands of Inner Mongolia.

2 Materials and methods

2.1 Site description

Fresh plants were collected from the upland grasslands of the Xilin River basin in August 2008. The climate is semi-arid, temperate, and continental, with a mean annual temperature of ~0.6°C. The coldest monthly mean temperature is -21.4°C in January, and the warmest is 18.5°C in July. The mean annual precipitation is about 350 mm, with a rainy season between mid-June and mid-September. Approximately 10% of precipitation falls as snow. The growing season extends from late April to early October. Detailed descriptions of the Xilin River basin have been published elsewhere (Wang et al., 2005).

2.2 Laboratory incubation

This study examined CH₄ emission in closed-chamber laboratory incubations from fresh twigs of *Artemisia* genus species indigenous to the Xilin River basin. All species examined were xerophytes (arid-adapted plants) from upland habitats with well-drained soils that exhibit net CH₄ consumption from the atmosphere (Wang et al., 2005) and therefore having little if any soil methanogenesis. A previous study (Wang et al., 2008) used isotope signature analysis to confirm that the CH₄ emitted from the shrub *Artemisia frigida* (note that this species was mislabeled as *Achillea frigida* in Wang et al., 2008) was derived directly from plant tissues. We examined the effect of physical injury (simulated by cutting) on CH₄ emission rates from all *Artemisia* species we found in the Xilin River basin (10 species total). All species were analyzed under aerobic conditions (ambient laboratory air) and *A. frigida* was also examined under hypoxic conditions. In order to exclude potentially complicating factors from whole plants, such as soil contamination and logistical difficulties with assaying large amounts of biomass, we restricted this work to detached twigs consisting of leaves and stems/petioles. Laboratory incubations were conducted in the dark at an ambient temperature of 24–26°C. There were triplicate samples for each treatment group described below.

Fresh twigs were sampled early in the morning (6:00–7:00 a.m. local time) preceding each measurement event. Samples were placed in plastic bags and transported to the laboratory immediately after sampling. Total time for harvest and transport to the laboratory was approximately 10 min. The twigs were washed in deionized water and air-dried for about 0.5 h. Four grams of air-dried fresh twigs (5–8 cm length) were either sealed immediately in a gastight serum bottle with a butyl rubber stopper (diameter 20 mm) or cut into 5-mm (moderate cutting) or 1-mm (severe cutting) segments and allowed to vent for 10 min before being sealed in a bottle. To ensure representativeness, we combined twigs from different plants and randomly mixed them in the bottles. Each sample (i.e. one serum bottle) contained five to ten twigs. Both leaves and stems/petioles were cut. To test whether stems and leaves both emitted CH₄, stem ends of detached *A. frigida* were sealed with silicone sealant as a separate treatment group. To establish hypoxic conditions, the bottle was immediately sealed with a butyl rubber stopper and flushed with pure nitrogen (400 ml min⁻¹ for 5 min) from a compressed nitrogen cylinder using “inlet-outlet” needles inserted through the stopper. Parallel blanks were used to test whether the background CH₄ concentrations in the bottles changed in the absence of plant material. The initial CH₄ concentrations were measured immediately after sealing.

In order to examine the relationship between CH₄ emission and respiration, CO₂ release rate was measured in the dark. One gram of fresh twigs (5–8 cm length) of *A. frigida*

was placed into a 120-ml serum bottle ($n=3$). Because the CO_2 accumulation in the bottle was high relative to the detection range, incubations lasted approximately 15 min. After sampling, the bottle was opened to the ambient atmosphere until the next sampling. Samples were incubated in the dark at an ambient temperature of 22–24°C.

2.3 CH_4 and CO_2 flux measurement

CH_4 and CO_2 concentrations were analyzed at various time intervals using a Hewlett-Packard 5890 Series II gas chromatograph equipped with a flame-ionization detector operated at 200°C, a 2-m stainless steel column packed with 13 XMS (60/80 mesh) for CH_4 analysis, and a 2-m stainless steel column packed with Porapak Q (60/80 mesh) for CO_2 analysis. The column oven temperature was 55°C, and the carrier gas was N_2 flowing at 30 ml min^{-1} . A 6-ml gas sample was withdrawn from the 120-ml serum bottle by syringe and immediately replaced by 6 ml of laboratory air (aerobic conditions) or N_2 (hypoxic conditions) to maintain headspace pressure. Certified CH_4 and CO_2 standards (China National Research Center for Certified Reference Materials, Beijing) were used for calibration. At the end of each incubation, biomass was determined as oven-dried weight (60°C for 48 h).

2.4 Statistical analysis

Statistical analysis was performed using the SAS (Statistical Analysis System) program (SAS, 1999). Duncan's multiple range test was employed for mean separation of CH_4 emission rates among treatments at $P < 0.05$. One-way analysis of variance (ANOVA) was used to evaluate statistical difference in the CH_4 emission rates among different injuries or between aerobic and hypoxic conditions.

3 Results

Cutting, a simulated physical injury, stimulated CH_4 emission from fresh twigs of *A. frigida* (Fig. 1). CH_4 concentrations increased linearly with time in all treatments with the exception of the blank, indicating that the twigs constantly emitted CH_4 throughout the ~24-h incubation. The emission rates were higher in cut twigs than uncut twigs ($P < 0.05$, $n=3$), and twigs with severe cutting had higher emission rates than twigs with moderate cutting ($P < 0.05$, $n=3$). Sealing the stem ends of detached twigs did not have a measurable effect on CH_4 emission, suggesting that CH_4 was emitted from leaves, not stems. Under hypoxic conditions, CH_4 concentrations increased linearly with time and cutting increased the rate of CH_4 emission (Fig. 2a). CH_4 emissions were significantly higher ($P < 0.05$, $n=3$) under hypoxic conditions than under aerobic conditions for both cut and uncut twigs (Fig. 2b).




Fig. 1. Effect of physical injury (cutting) on CH_4 emission from fresh twigs of *Artemisia frigida*. (a) CH_4 accumulation over time. (b) CH_4 emission rates showing standard deviations ($n=3$) and statistically significant differences ($P < 0.05$) among treatments indicated by different letters. The CH_4 emission rates were calculated using the CH_4 accumulation over time. Treatments: Blank, Uncut twigs with stem ends sealed with silicone sealant, Uncut twigs with stem ends not sealed, Moderate injury (twigs cut into 5 mm segments), and Severe injury (twigs cut into 1 mm segments).

Dark respiration decreased over time in all treatments but did not cease entirely during incubations (Fig. 3). Respiration rates were about 50% higher in cut than in uncut twigs under aerobic conditions (Fig. 3a). CO_2 production was about 3- to 10-fold slower and decreased more over time under hypoxic than under aerobic conditions (Fig. 3b). In contrast to CH_4 emission, which was increased by cutting under both aerobic and hypoxic conditions, CO_2 emission was unaffected or decreased in response to cutting under hypoxic conditions.

Table 1 lists the CH_4 emission rates under aerobic conditions from fresh twigs of 10 *Artemisia* species sampled from the upland areas of the Xilin River basin. Four species were shrubs and six species were herbs. Only 1 species, *A. frigida*, emitted detectable amounts of CH_4 without cutting. Eight species were further tested for CH_4 emission after cutting, but only five, two shrub species including *A. frigida* and three herb species, emitted CH_4 continuously throughout the 24-h incubation. *A. frigida* exhibited the highest emission rates.




Fig. 2. Effect of hypoxia on CH₄ emission from fresh twigs of *Artemisia frigida*. (a) CH₄ accumulation over time. Initial CH₄ concentrations were approximately 1.9 and 0 ppmv in aerobic and hypoxic conditions, respectively. (b) CH₄ emission rates (mean \pm standard deviation; $n=3$; * indicates statistically significant difference at $P<0.05$ between aerobic and hypoxic conditions). The CH₄ emission rates were calculated using the CH₄ accumulation over time. Treatments: Blank, Uncut twigs, Severe injured twigs in aerobic conditions, Blank in hypoxic conditions, Uncut twigs in hypoxic conditions, and Severe injured twigs in hypoxic conditions (twigs cut into 1 mm segments).

4 Discussion

4.1 Environmental stress stimulates CH₄ emission from plants

Many wetland plants (hydrophytes) transport CH₄ produced by anaerobic soil microorganisms to the atmosphere through their roots and stems (Schimel et al., 1995). Previously, we showed that microbial CH₄ stored in the stems of hydrophytes is eliminated within minutes after cutting, whereas CH₄ derived directly from plant compounds is emitted continuously over many hours; microbial and plant-derived CH₄ were distinguished by their distinct $\delta^{13}\text{C}$ signatures (Wang et al., 2008). Concurring studies have found that neither adsorption-desorption processes nor CH₄ stored within intercellular air spaces are responsible for the observed CH₄ emissions (Kirschbaum et al., 2007; Kirschbaum and Walcroft, 2008; Vigano et al., 2008). In the present study, physi-




Fig. 3. Dark respiration from fresh twigs of *Artemisia frigida*. Treatments are (a) Uncut twigs and cut twigs in aerobic conditions and (b) Uncut twigs and cut twigs in hypoxic conditions. The CO₂ release rate is mean ± 1 standard deviation, $n=3$.

cal injury elevated the rate of CH₄ emission from fresh twigs of *A. frigida* continuously for ~ 24 -h (Fig. 1), indicating that physical injury released CH₄ from plant compounds by an as yet unidentified mechanism.

Hypoxia also induced higher rates of CH₄ emission from *A. frigida* (Fig. 2). Although the initial CH₄ concentration was lower in the hypoxic incubations because of flushing with N₂, we found previously that CH₄ emission rates were indistinguishable in CH₄-free air and in ambient atmospheric CH₄ (Wang et al., 2008). Vigano et al. (2008) also found that emission rates were independent of background CH₄ concentration in an incubation system that continuously replaced the atmosphere. Hence, the significant difference in emission rates under aerobic and hypoxic conditions observed here likely resulted from the difference in O₂ concentrations and not from the difference in initial CH₄ concentrations. Hypoxia induces stress through the accumulation of toxic metabolic products, the lowering of energy charge, and the lack of substrates for respiration (Drew, 1997). Xerophytes, such as the *Artemisia* species examined here, would probably be particularly susceptible to hypoxia stress since they are adapted to well-aerated soils and are not accustomed to hypoxic conditions.

Previous studies have shown that UV radiation and rising temperature independently stimulate CH₄ emission from plants (Keppler et al., 2006; McLeod et al., 2008; Vigano et al., 2008). Keppler et al. (2006) observed an exponential

Table 1. CH₄ emissions from *Artemisia* species sampled from the upland grasslands of the Xilin River basin^a.

Morphotype	Species	Incubated tissue ^b	CH ₄ emission (ng CH ₄ gdw ⁻¹ h ⁻¹)	
			Uncut	Cut
Shrub	<i>Artemisia frigida</i> Willd.	twigs (leaves and stems)	0.38±0.11	4.59±0.45
	<i>Artemisia intramongolica</i> H. C. Fu	twigs (leaves and stems)	ND	1.24±0.04
	<i>Artemisia gmelinii</i> var. <i>vestita</i> Nakai	twigs (leaves and stems)	ND	ND
	<i>Artemisia gmelinii</i> Web. ex Stechm	twigs (leaves and stems)	ND	ND
Herb	<i>Artemisia argyi</i> Levl. et Vant.	twigs (leaf and petiole)	ND	3.25±0.80
	<i>Artemisia eriopoda</i> Bunge	twigs (leaf and petiole)	ND	0.45±0.15
	<i>Artemisia sieversiana</i> Willd.	twigs (leaf and petiole)	ND	0.91±0.22
	<i>Artemisia laciniata</i> Willd.	twigs (leaves and stems)	ND	ND
	<i>Artemisia rubripes</i> Nakai	twigs (leaf and petiole)	ND	NM
	<i>Artemisia scoparia</i> Wald. et kit.	twigs (leaves and stems)	ND	NM

^a Fresh plant tissues were sampled during 17–18 August 2008. CH₄ emissions were measured for uncut (but detached) and cut plant tissues, respectively, incubated in the dark at room temperature of 21–24°C for approximately 24 h with initially ambient CH₄ concentration in air. Emission rate is mean ± standard deviation ($n=3$). ND indicates that no CH₄ emission was detected or was too weak to quantify from linear change in CH₄ concentrations within 24 h, while NM was no measurement.

^b Bracket shows one or more plant organs per twig.

increase in emission rates with rising temperature from 30 to 70°C, which is inconsistent with a microbial or other enzymatic process. Vigano et al. (2008) similarly observed increasing CH₄ emission rates with increasing temperature up to 60°C in the dark. In our study, CH₄ emission increased with increasing physical injury in the dark. Hence, UV radiation, rising temperature, physical injury, and hypoxia all stimulate CH₄ emission from plant tissues, suggesting that environmental stress in general may increase CH₄ emission from plants.

4.2 Mechanistic implications of stress

In this study, physical injury stimulated dark respiration under aerobic conditions (Fig. 3). During physiological stress, respiratory electron transport can give rise to cell-damaging ROS, including hydrogen peroxide (H₂O₂), superoxide ion (O₂⁻), hydroperoxyl radical (HO₂), hydroxyl radical (OH), and singlet oxygen (¹O₂). ROS cleave polymer chains in site-specific reactions (Scandalios, 1993; Fry, 1998; Schweikert et al., 2002; Cheng et al., 2008). Hence, formation of CH₄ from plant tissues could result from ROS attacks that cleave methyl groups from plant lignin or pectin (Keppler et al., 2008; McLeod et al., 2008). In a previous study, we found that CH₄ emitted continuously from *A. frigida* leaves had a carbon isotope signature consistent with methoxyl groups of plant pectin (Wang et al., 2008), and Keppler et al. (2006) found evidence for pectin-derived CH₄ emitted by other plant species. The pool of methoxyl groups in pectin is large in comparison to adsorbed/stored CH₄ and would be available continuously for CH₄ formation (Keppler et al., 2006; Vigano et al., 2008), consistent with the results of this study and previous results.

A variety of environmental stressors, such as radiation, herbicides, pathogens, certain injuries, hypoxia, ground-level ozone, and temperature fluctuations, stimulate the formation of ROS in plants (Scandalios, 1992; Crawford and Brandle, 1996). For example, Chen and Qualls (2003) reported an increase of free radical damage during hypoxia, and Lin et al. (2006) observed a decrease of ROS scavenging in plants under flooding (resulting in hypoxia). Physical injury of certain plant tissues initiates a complex series of reactions that largely involve a self-perpetuating wave of free radical generation (Thompson et al., 1987). Hence, repeated observations in several studies of CH₄ emission from plants being stimulated by rising temperatures, UV irradiation, hypoxia, and physical injury could be explained by physiological stress leading to ROS formation. This mechanism is distinct from but not mutually exclusive with the mechanism of UV-driven cleavage of methyl groups from plant compounds proposed previously (McLeod et al., 2008; Vigano et al., 2008).

4.3 Species differences

In a previous study, we found that several shrubs but no herbs emitted CH₄ derived from plant compounds (Wang et al., 2008). In the present study we again detected no CH₄ emission from uncut herbs (Table 1). However, cutting injuries stimulated CH₄ emission in three out of four herb species that were tested (Table 1). Hence, both shrubs and herbs have the potential to emit CH₄ when injured. Even after injury, however, we still detected no CH₄ emission from two species of shrub and one species of herb, all belonging to the *Artemisia* genus (Table 1). Kirschbaum and Walcroft (2008) observed no substantial CH₄ emission from another *Artemisia* species, *Artemisia absinthium*, in New

Zealand. Thus, CH₄ emissions may be species-dependent, even among closely related species. The differences could reflect variation among leaf structures, UV-screening pigments, UV-photosensitizers, and ROS-scavenging mechanisms (see McLeod et al., 2008), as well as phenology and environmental/experimental conditions.

5 Conclusions

Along with other laboratory studies, our results confirm that a wide variety of plant species and functional types can emit CH₄ under aerobic conditions and in the absence of likely microbial CH₄ sources. However, emission rates vary both among plant species and in different environmental conditions, including hypoxia. Therefore, estimating the contribution of this process to the global CH₄ cycle might not be possible based on simple biomass- and temperature-based algorithms (e.g., Keppler et al., 2006; Butenhoff and Khalil, 2007). Similarly, because CH₄ emission responds to light, Keppler et al. (2006) used average daily sunshine hours to estimate annual global emissions of CH₄ by living plants and leaf litter. However, a wide variety of environmental conditions can stimulate or accelerate CH₄ emission from plants in the dark. Because nighttime accounts for half of a plant's life, the global plant-derived CH₄ emission stimulated by environmental stressors at night could be an important component of aerobic CH₄ emissions.

If stress is a major driver of CH₄ emission from plants, then global change processes, such as climate change, stratospheric ozone depletion, increasing ground-level ozone, spread of plant pests, and land-use changes, could increase stress in plants and cause a global increase in CH₄ emissions. If so, the role of stress in plant CH₄ production in the global CH₄ cycle could be important in a changing world. Future studies are needed that focus on in situ measurements, comparisons between stressed and unstressed plants, dark and light CH₄ emissions, and interspecies differences.

Acknowledgements. This research was supported by a general program from the National Natural Science Foundation of China (30670402), the State Key Basic Research Development Program of China (2007CB106800), and the funding of State Key Laboratory of Vegetation and Environmental Change.

Edited by: G. Wohlfahrt

References

Berling, D. J., Gardiner, T., Leggett, G., McLeod, A., and Quick, W. P.: Missing methane emissions from leaves of terrestrial plants, *Glob. Change Biol.*, 14, 1–6, 2008.

Brüggemann, N., Meier, R., Steigner, D., Zimmer, I., Louis, S., and Schnitzler, J.-P.: Nonmicrobial aerobic methane emission from poplar shoot cultures under low-light conditions, *New Phytol.*, doi:10.1111/j.1469-8137.2009.02797.x, 2009.

Butenhoff, C. L. and Khalil, M. A. K.: Global methane emissions from terrestrial plants, *Environ. Sci. Technol.*, 41, 4032–4037, 2007.

Cao, G. M., Xu, X. L., Long, R. J., Wang, Q. L., Wang, C. T., Du, Y. G., and Zhao, X. Q.: Methane emissions by alpine plant communities in the Qinghai-Tibet Plateau, *Biol. Lett.-UK*, 4, 681–684, 2008.

do Carmo, J. B., Keller, M., Dias, J. D., de Camargo, P. B., and Crill, P.: A source of methane from upland forests in the Brazilian Amazon, *Geophys. Res. Lett.*, 33, L04809, doi:10.1029/2005GL025436, 2006.

Chen, H. J. and Qualls, R. G.: Anaerobic metabolism in the roots of seedlings of the invasive exotic *Lepidium latifolium*, *Environ. Exp. Bot.*, 50, 29–40, 2003.

Cheng, G. P., Duan, X. W., Yang, B., Jiang, Y. M., Lu, W. J., Luo, Y. B., and Jiang, W. B.: Effect of hydroxyl radical on the scission of cellular wall polysaccharides in vitro of banana fruit at various ripening stages, *Acta Physiol. Plant*, 30, 257–263, 2008.

Crawford, R. M. M. and Brandle, R.: Oxygen deprivation stress in a changing environment, *J. Exp. Bot.*, 47, 145–159, 1996.

Crutzen, P. J., Sanhueza, E., and Brenninkmeijer, C. A. M.: Methane production from mixed tropical savanna and forest vegetation in Venezuela, *Atmos. Chem. Phys. Discuss.*, 6, 3093–3097, 2006, <http://www.atmos-chem-phys-discuss.net/6/3093/2006/>.

Drew, M. C.: Oxygen deficiency and root metabolism: injury and acclimation under hypoxia and anoxia, *Ann. Rev. Plant Physiol. Plant Mole. Biol.*, 48, 223–250, 1997.

Dueck, T. A., de Visser, R., Poorter, H., Persijn, S., Gorissen, A., de Visser, W., Schapendonk, A., Verhagen, J., Snel, J., Harren, F. J. M., Ngai, A. K. Y., Verstappen, F., Bouwmeester, H., Voeselek, L. A. C. J., and van der Werf, A.: No evidence for substantial aerobic methane emission by terrestrial plants: a ¹³C-labelling approach, *New Phytol.*, 175, 29–35, 2007.

Dueck, T. and van der Werf, A.: Are plants precursors for methane?, *New Phytol.*, 178, 693–695, 2008.

Ferretti, D. F., Miller, J. B., White, J. W. C., Lassey, K. R., Lowe, D. C., and Etheridge, D. M.: Stable isotopes provide revised global limits of aerobic methane emissions from plants, *Atmos. Chem. Phys.*, 7, 237–241, 2007, <http://www.atmos-chem-phys.net/7/237/2007/>.

Frankenberg, C., Meirink, J. F., van Weele, M., Platt, U., and Wagner, T.: Assessing methane emissions from global space-borne observations, *Science*, 308, 1010–1014, 2005.

Frankenberg, C., Bergamaschi, P., Butz, A., Houweling, S., Meirink, J. F., Notholt, J., Petersen, A. K., Schrijver, H., Warneke, T., and Aben, I.: Tropical methane emissions: A revised view from SCIAMACHY onboard ENVISAT, *Geophys. Res. Lett.*, 35, L15811, doi:10.1029/2008GL034300, 2008.

Fry, S. C.: Oxidative scission of plant cell wall polysaccharides by ascorbate-induced hydroxyl radicals, *Biochem. J.*, 332, 507–515, 1998.

Houweling, S., Röckmann, T., Aben, I., Keppler, F., Krol, M., Meirink, J. F., Dlugokencky, E. J., and Frankenberg, C.: Atmospheric constraints on global emissions of methane from plants, *Geophys. Res. Lett.*, 33, L15821, doi:10.1029/2006GL026162, 2006.

Keppler, F., Hamilton, J. T. G., Braß, M., and Röckmann, T.: Methane emissions from terrestrial plants under aerobic condi-

- tions, *Nature*, 439, 187–191, 2006.
- Kepler, F., Hamilton, J. T. G., McRoberts, W. C., Vigano, I., Braß, M., and Röckmann, T.: Methoxyl groups of plant pectin as a precursor of atmospheric methane: evidence from deuterium labelling studies, *New Phytol.*, 178, 808–814, 2008.
- Kirschbaum, M. U. F., Bruhn, D., Etheridge, D. M., Evans, J. R., Farquhar, G. D., Gifford, R. M., Paul, K. I., and Winters, A. J.: A comment on the quantitative significance of aerobic methane release by plants, *Funct. Plant Biol.*, 33, 521–530, 2006.
- Kirschbaum, M. U. F., Niinemets, U., Bruhn, D., and Winters, A. J.: How important is aerobic methane release by plants?, *Functional Plant Science and Technology*, 1, 138–145, 2007.
- Kirschbaum, M. U. F. and Walcroft, A.: No detectable aerobic methane efflux from plant material, nor from adsorption/desorption processes, *Biogeosciences*, 5, 1551–1558, 2008, <http://www.biogeosciences.net/5/1551/2008/>.
- Lin, K. H., Chao, P. Y., Yang, C. M., Cheng, W. C., Lo, H. F., and Chang, T. R.: The effects of flooding and drought stresses on the antioxidant constituents in sweet potato leaves, *Bot. Stud.*, 47, 417–426, 2006.
- McLeod, A. R., Fry, S. C., Loake, G. J., Messenger, D. J., Reay, D. S., Smith, K. A., and Yun, B. W.: Ultraviolet radiation drives methane emissions from terrestrial plant pectins, *New Phytol.*, 180, 124–132, 2008.
- Messenger, D. J., McLeod, A. R., and Fry, S. C.: The role of ultraviolet radiation, photosensitizers, reactive oxygen species and ester groups in mechanisms of methane formation from pectin, *Plant Cell Environ.*, 32, 1–9, 2009.
- Miller, J. B., Gatti, L. V., d’Amelio, M. T. S., Crotwell, A. M., Dlugokencky, E. J., Bakwin, P., Artaxo, P., and Tans, P. P.: Airborne measurements indicate large methane emissions from the eastern Amazon basin, *Geophys. Res. Lett.*, 34, L10809, doi:10.1029/2006GL029213, 2007.
- Nisbet, R. E. R., Fisher, R., Nimmo, R. H., Bendall, D. S., Crill, P. M., Gallego-Sala, A. V., Hornibrook, E. R. C., López-Juez, E., Lowry, D., Nisbet, P. B. R., Shuckburgh, E. F., Sriskantharajah, S., Howe, C. J., and Nisbet, E. G.: Emission of methane from plants, *Proceedings of the Royal Society B: Biological Sciences*, 276, 1347–1354, 2009.
- Parsons, A. J., Newton, P. C. D., Clark, H., and Kelliher, F. M.: Scaling methane emissions from vegetation, *Trends Ecol. Evol.*, 21, 423–424, 2006.
- Sanhueza, E. and Donoso, L.: Methane emission from tropical savanna *Trachypogon* sp. grasses, *Atmos. Chem. Phys.*, 6, 5315–5319, 2006, <http://www.atmos-chem-phys.net/6/5315/2006/>.
- SAS: SAS/STAT User’s Guide Release 8.0 Cary, North Carolina, USA, 1999.
- Scandalios, J. G.: *Molecular Biology of Free Radical Scavenging Systems*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, 1992.
- Scandalios, J. G.: Oxygen stress and superoxide dismutases, *Plant Physiol.*, 101, 7–12, 1993.
- Schiermeier, Q.: Methane finding baffles scientists, *Nature*, 439, 128–128, 2006.
- Schimel, J. P.: Plant transport and methane production as controls on methane flux from arctic wet meadow tundra, *Biogeochemistry*, 28, 183–200, 1995.
- Schweikert, C., Liskay, A., and Schopfer, P.: Polysaccharide degradation by Fenton reaction- or peroxidase-generated hydroxyl radicals in isolated plant cell walls, *Phytochemistry*, 61, 31–35, 2002.
- Sinha, V., Williams, J., Crutzen, P. J., and Lelieveld, J.: Methane emissions from boreal and tropical forest ecosystems derived from in-situ measurements, *Atmos. Chem. Phys. Discuss.*, 7, 14011–14039, 2007, <http://www.atmos-chem-phys-discuss.net/7/14011/2007/>.
- Thompson, J. E., Legge, R. L., and Barber, R. F.: The role of free radicals in senescence and wounding, *New Phytol.*, 105, 317–344, 1987.
- Vigano, I., van Weelden, H., Holzinger, R., Kepler, F., McLeod, A., and Röckmann, T.: Effect of UV radiation and temperature on the emission of methane from plant biomass and structural components, *Biogeosciences*, 5, 937–947, 2008, <http://www.biogeosciences.net/5/937/2008/>.
- Wang, Z. P., Han, X. G., Li, L. H., Chen, Q. S., Duan, Y., and Cheng, W.: Methane emission from small wetlands and implications for semiarid region budgets, *J. Geophys. Res.*, 110, D13304, doi:10.1029/12004JD005548, 2005.
- Wang, Z. P., Han, X. G., Wang, G. G., Song, Y., and Guldge, J.: Aerobic methane emission from plants in the Inner Mongolia steppe, *Environ. Sci. Technol.*, 42, 62–68, 2008.