

J Vect Borne Dis 41, March & June 2004, pp 37–41

Variations in ornamentation of wings and palpi of *Anopheles (Cellia) subpictus* Grassi collected from northwest India

Jagbir S. Kirti* & Jagdish Kaur

Department of Zoology, Punjabi University, Patiala, India; jskirti@pbi.ernet.in

Key words *Anopheles* – palpi – species – variations – wings

Studies have been conducted on the wings and palpi of *Anopheles (C) subpictus* Grassi to observe variations. As many as 10 types of variations have been observed in wings and palpi of this species. These variations have been described and illustrated in this paper.

An. subpictus Grassi is a dominant species in Haryana and Uttaranchal states. Though it is a non-vector species but recently infected specimens with malarial parasite have been reported from India, Indonesia and Java¹. This species is distributed throughout India, Afghanistan, Borneo, China, Malaysia, Myanmar, Philippines, Sri Lanka, Java and Indonesia. Morphological variations have been recorded in Indian anophelines since 1933. Christophers² first recorded the variations from type form in genus *Anopheles* Meigen. It was followed by Ramakrishna³ and Rehman *et al*⁴, who observed an extra dark band on the palpi of female *An. fluviatilis* James. An extra dark band on the palpi in *An. subpictus* Grassi and *An. pallidus* Theobald was observed by Subramanian & Nagendra⁵. Incomplete development of anal vein in *An. stephensi* Liston was noted by Bhatnagar *et al*⁶. Rajagopal & Chakraborty⁷ recorded variations in the palpi, hind tarsi and wings of *An. annularis* Van der Wulp from Dhanbad. Nagpal & Sharma^{8,9} recorded

*Corresponding author


variation in wings and palpi of *An. vagus* Donitz and variations in palpi of *An. sundaicus* Rodenwaldt. Nagpal¹⁰ reported morphological variations in *An. stephensi* Liston. From the above it becomes crystal clear that no worker recorded variations in palpi and wings of *An. subpictus* Grassi except Subramanian and Nagendra⁵, who reported single variation in palpi of this species. However, the present authors have been able to record 10 types of morphological variations in wings and palpi of *An. subpictus* from field collected specimens. These variations have been described and illustrated in the present paper.

Intensive and extensive collection-cum-survey tours were conducted in different localities falling in the states of Punjab, Haryana, Himachal Pradesh, Uttaranchal, Jammu and Kashmir, Delhi and Union Territory of Chandigarh. A large number of adult mosquitoes were collected with the help of suction tube and torch from different resting places—cattlesheds, human dwellings, mixed dwellings, vegetation, etc. Besides collecting adult specimens, the immature stages such as larvae and pupae were collected with the help of dropper and glass tube from different breeding places, such as paddy fields, stagnant water bodies, water coolers, tree holes, rock pools, rivulets, tyres, unused water boats, gutters and other natural as well as man-made containers. These immature stages were


brought to the laboratory and reared for the emergence of adults. The adult specimens were killed with ethyl acetate vapours. The freshly killed specimens were fixed either on entomological pins or on the tip of paper triangles with mounting glues. The triangles were supported with large entomological pins. The pinned specimens were preserved in the air tight and properly fumigated boxes of insect cabinets. The data on locality, source, altitude, date of collection, etc. were recorded on the field labels. The present species has been identified by following the keys of Puri¹¹; and Wattal & Kalra¹². The comparison was made

with the identified collection of National Institute of Communicable Diseases, Delhi.

As many as 141 specimens of the present species were collected from different localities of northwest India during 1997–2000. In total 10 types of morphological variations have been observed in the speckling, banding pattern and size of band in palpi and ornamentation of wings. Out of these six variations have been recorded in wings and four in palpi. The details of collection date, site of collection, number of collected specimens and morphological variations have been given in Figs. 1 to 12 and Table 1.



Figs. 1–5 : *Anopheles (Cellia) subpictus* Grassi (1: Palpi of type form; 2–5: Variations in palpi; Bdb – Basal dark band; Bpb – Basal pale band; Mdb – Middle dark band; Spb – Subapical pale band; Sdb – Subapical dark band; Apb – Apical pale band)



Figs. 6-12: *Anopheles (Celia) subpictus* Grassi (6: Wings of type form; 7-12: Variations in wings; C - Coasta; CU - Cubitus; CU₁ - Cubitus one; CU₂ - Cubitus two; M - Media; M₁ - Media one; M₂ - Media two; R₁ - Radius one; R₂ - Radius two; R₃ - Radius three; R₄₊₅ - Radius four-plus-five; Rs - Radial sector; SC - Subcosta)

Table 1. Morphological variations in the palpi and wings of *Anopheles (Cellia) subpictus* Grassi

Sl. No.	Date of collection	Site of collection	No. of specimens collected	Description of variation from the type form
<i>Variations in palpi</i>				
1.	1 August 1999	Dehria, Nainital	27	Tips of both palpi dark; one palpus lacking subapical pale band (Fig. 2).
2.	-do-	-do-	27	Middle dark area bearing pale scales in centre all over; basal dark area with pale scales on outer side on one of palpi; one palpus carrying a rounded white spot at basal dark area (Fig. 3).
3.	18 August 1998	Ismilepur, Dehradun	7	Apical and subapical pale areas absent on both palpi (Fig. 4).
4.	1 September 1998	Indira Nagar, Kurukshetra	10	One of the palpi lacking subapical pale band (Fig. 5).
<i>Variations in wings</i>				
5.	1 August 1999	Dehria, Nainital	27	R _s lacking one dark spot; R ₂ and R ₃ pale except dark tips; dark spot at base of M ₁ & M ₂ absent; vein M bearing one dark spot only; Cu ₁ lacking a dark spot at base (Fig. 7).
6.	-do-	-do-	27	Presector, prehumeral and basal dark spots at costa absent; median dark spot of R _s absent (Fig. 8).
7.	-do-	-do-	27	Base of costa largely dark; dark spot of R ₁ below middle dark mark of costa misplaced; R _s lacking both of the dark spots; R ₂ with one dark spot only; M and M ₂ lacking one dark spot each (Fig. 9).
8.	-do-	-do-	27	Costa with basal dark spot missing; R _s lacking one dark spot; R ₁ with an extra small dark spot below middle dark spot of costa; M lacking one dark spot; dark spot at fork of Cu absent (Fig. 10).
9.	2 August 1999	Tilpuri, Udham Singh Nagar	36	R ₁ with additional dark spot; R _s bearing single dark spot only; basal dark spot of vein M ₂ lacking (Fig. 11).
10.	-do-	-do-	36	Costa with large humeral pale spot; R _s lacking one spot; R ₃ totally pale; R ₄₊₅ lacking dark spot at tip; dark spot at base M ₂ , near fork at Cu ₁ and middle of anal vein absent (Fig. 12).

Acknowledgement

Authors are thankful to the Director, Malaria Research Centre, Delhi for providing the necessary facilities and relevant literature.

References

1. Nagpal BN, Sharma VP. *Indian Anophelines*. New Delhi : Oxford & IBH Publishing Co. Pvt. Ltd. 1995; p 1–416.
2. Christophers SR. *The fauna of British India including Ceylon and Burma*, v IV (Diptera : Culicidae). London : Taylor and Francis 1933; p 1–360.
3. Ramakrishna V. Variations in palpai banding of *An. fluviatilis* James (1902). *Bull Natl Soc Ind Mal Mosq Dis* 1954; 2 : 210–1.
4. Rehman J, Singh MY, Singh NN. Note on the study of morphology, prevalence and host preference of an ecotype of *An. fluviatilis* in Nainital Terai (U.P.). *Bull Natl Soc Ind Mal Mosq Dis* 1960; 8 : 137–42.
5. Subramanian R, Nagendra S. Variation in the marking of palpi in certain anophelines. *Bull Natl Soc Ind Mal Mosq Dis* 1955; 3 : 94–5.
6. Bhatnagar VN, Bhatia ML, Krishna KS. On certain morphological abnormalities noted in *An. pallidus* Theobald, 1901 and *An. stephensi* Liston, 1901. *Indian J Malariol* 1958; 12 : 39–2.
7. Rajagopal R, Chakraborty RK. Record of morphological variation in *An. Annularis* Van der Wulp near Dhanbad. *Indian J Malariol* 1960; 14 : 171–4.
8. Nagpal BN, Sharma VP. Morphological variations in a natural population of *Anopheles vagus* Donitz (1902) collected from Andaman Islands. *Indian J Malariol* 1983; 20 : 35–44.
9. Nagpal BN, Sharma VP. Variation in ornamentation of palpi of *An. sudaicus* Rodenwaldt (1982) collected from Andaman Island, India. *Indian J Malariol* 1983; 20 : 85–7.
10. Nagpal BN. Morphological variations in natural populations of *Anopheles stephensi* Liston, 1901 collected from Kutch (Gujarat). *Indian J Malariol* 1990; 27 : 25–35.
11. Puri IM. Synoptic table for the identification of the anopheline mosquitoes of India. *Health Bull* 1954; 10 : 1–57.
12. Wattal BL, Kalra NL. Regionwise pictorial keys to the female Indian *Anopheles*. *Bull Natl Soc Ind Mal Mosq Dis* 1961; 9 : 85–138.