

RESEARCH ARTICLE

Ivelina Nikolova¹
Mariyana Georgieva¹
Lubomir Stoilov¹
Zornica Katerova²
Desisslava Todorova²

Optimization of Neutral Comet Assay for studying DNA double-strand breaks in pea and wheat

Authors' addresses:

¹ Department of Molecular Genetics, Institute of Plant Physiology & Genetics, Bulgarian Academy of Sciences, Sofia, Bulgaria.

² Department of Regulation of Plant Growth and Development, Institute of Plant Physiology & Genetics, Bulgarian Academy of Sciences, Sofia, Bulgaria.

Correspondence:

Mariyana Georgieva
Department of Molecular Genetics, Institute of Plant Physiology & Genetics, Bulgarian Academy of Sciences Acad. G. Bonchev Str., Bldg. 21 1113 Sofia, Bulgaria,
Tel.: +359 2 974 62 28
e-mail: mimy80@abv.bg

Article info:

Received: 25 October 2012

In revised form: 5 July 2013

Accepted: 5 August 2013

ABSTRACT

This study describes an adaptation of the Comet assay under neutral conditions for mono- and dicotyledonous plants pea (*Pisum sativum* L.) and wheat (*Triticum aestivum* L.). Modifications concern lysis and electrophoresis steps, respectively. Electrophoresis was carried out varying the intensity of the electric field. A linear relationship between the percentages of DNA in the tail from control background with alteration of intensity was found. Trypan blue dye exclusion test was used in order to determine the intactness of nuclear membrane of the isolated nuclei from both plant model systems. Assessment was conducted on non-irradiated and irradiated nuclei on a monolayer with three doses of UVC. It was found that the share of intact nuclei (trypan blue negative ones) is about 95% in controls. Gradual dose-related increase of damaged nuclei was observed in both species, reaching statistical significance only at the higher dose applied.

Key words: neutral Comet assay, trypan blue dye exclusion test, UVC.

Introduction

A wide range of methods are presently used for detection of early biological effects of DNA-damaging agents in the environment, irrespective of whether they are toxic, mutagenic, teratogenic and carcinogenic. Plant mutagenicity bioassays have taken their place as screening and biomonitoring tools for detection of DNA damage induction. Most higher-plant tests are based on detection of point mutations, chromosomal rearrangements, DNA fragmentation (Maluszynska & Juchimiuk, 2005; Gady et al., 2009; Uhl et al., 2003). Quantification methods of DNA strand breaks induction, such as Comet assay, have been developed and became increasingly popular during the last 10 years along with other cytogenetic assays in plant biomonitoring. The Comet assay was introduced by Östling & Johanson (1984) under neutral lysis and electrophoresis (pH 9.5) conditions. This technique has been developed in an

empirical way, with two basically different alkaline protocols described in the literature by Singh et al. (1988) and Olive (1989), concerning animal cells. The respective protocol was developed to measure low levels of strand breaks with high sensitivity. The subsequent protocols were optimized to detect a subpopulation of cells with variable sensitivity to drugs or radiation (Singh et al., 1988).

In plants, this technique has been modified and extensively validated over the years. Primary applications of plant Comet assay were used to determinate the effect of ionizing radiation on seed disinfection and preservation of diseases and pathogens (Cerda et al., 1993, 1997; Koppen & Cerda, 1997). Since the radiation doses used in food irradiation extensively damage DNA, a neutral pH was chosen, combined with a low voltage and short electrophoresis time (Cerda et al., 1993). Neutral pH limits the amplitude of DNA unwinding as reported by Singh (1988). Comet assay under these conditions detects mainly

RESEARCH ARTICLE

double strand breaks (DSB). Alkaline conditions (pH 10 or above) were required for unwinding and detection of single strand breaks (SSB). However, as shown in *Nicotiana tabacum* and *Vicia faba*, low sensitivity of the neutral version creates problems in detecting of induced DSB, even after adaptation of the basic assay (Koppen & Angelis, 1998; Koppen *et al.*, 1999). Additionally, neutral comet assay proved to be adequate for estimation of DNaseI-induced DSB in the presence of MnCl₂ (Menke *et al.*, 2000a), although being less sensitive than alkaline lyses/neutral gel electrophoresis (A/N), used to follow repair of damage induced by N-methyl-N-nitrosourea (MNU) and UVC in barley genome (Jovtchev *et al.*, 2001; Armalytè & Žukas, 2003).

Another limitation of the assay in plants was the relatively high value of comet parameter “% DNA in tail” in the controls, that could mask the effect of the treatment (Koppen & Angelis, 1998; Angelis *et al.*, 1999; Menke *et al.* 2000a, 2000b). Moreover, a statistical requirement of this method is a low background level of breaks in untreated cells (i.e. between 10 and 20% DNA in tail) (Collins, 2004; Lovell & Omori, 2008).

The objectives of this study were: (1) to determine and compare the nuclear membrane intactness on non-irradiated and UVC irradiated pea and wheat nuclei by trypan blue exclusion test; (2) to define the parameters for neutral Comet assay by alteration of lysis duration and electrophoresis conditions using nuclei isolated from pea and wheat leaves.

Materials and Methods

Plant growth conditions and isolation of nuclei

Seeds of *Pisum sativum* and *Triticum aestivum* were surface-sterilized by immersion in 0.1% KMnO₄ for 15 min followed by 30 min in distilled water. Germination was performed on wet filter paper and seedlings were subsequently transferred to a container with water and kept in a plant growth chamber at 26/22°C day/night temperature, 16/8 h (day/night) photoperiod, 150 μmol m⁻² s⁻¹ photon flux density and 60% relative air humidity.

Individual leaf samples were removed from the plant and placed in a petri dish with Sørensen buffer (50 mM sodium phosphate, pH 6.8, 0.1 mM ethylenediaminetetraacetic acid (EDTA), 0.5% dimethyl sulfoxide (DMSO) kept on ice. The leaf tissue was gently sliced by a razor blade and the resulting material was repeatedly dipped in the cold Sørensen buffer. The suspension with released nuclei was filtered through a 30

μm disposable filter (Partec, Münster, Germany) to remove most of the debris and centrifuged at 550 g for 5 min at 4°C.

UVC irradiation

UVC irradiation at 254 nm was delivered by BLX 254 UV crosslinker (Life Technologies TM, GIBCO BRL UV Crosslinker). In a precooled, plastic petri dish nuclear suspension was irradiated with doses of 5, 7 and 9 kJ/m² for pea and 14, 16 or 18 kJ/m² for wheat, respectively. After treatment, nuclei were kept on ice and shielded from the ambient light until subsequent use for the viability test.

Measurement of nuclear membrane intactness by Trypan blue dye exclusion test

The nuclear membrane integrity was determined by trypan blue exclusion test. Briefly, nuclei were incubated in 0.4% trypan solution (1:1 vol/vol) for 5 min and scored as membrane damaged if the nuclei were stained blue. As a negative control for the trypan blue exclusion test were used non irradiated nuclei incubated in Sørensen buffer. All samples were examined and counted in duplicates under light microscope at 25x (Zeiss Jenamed-2). The nuclear membrane intactness was expressed as percentage of the number of nuclei excluding trypan blue from the total number of nuclei for each plant (pea and wheat). Statistical analysis was performed by the Kruskal-Wallis test. Subsequently the Duncan test was applied for calculations concerning pairwise comparisons. The level of statistical significance was set at $p < 0.05$. The data obtained are presented as mean with standard error (\pm SEM).

Neutral Comet assay

For the neutral Comet assay, the protocol of Georgieva & Stoilov (2008) with modifications was followed. Microscope slides were coated with 0.5% normal melting agarose and dried at room temperature. 40 μl of the nuclei suspension was mixed with 40 μl of 0.1% low melting agarose, spread on the slide surface and subjected to gel formation for at least 10–15 min on a cooling plate at 4°C. Lysis was carried out in 2.5 M NaCl, 10 mM Na₂EDTA (pH8), 10 mM Tris-HCl (pH8), 1% N-lauroylsarcosine sodium salt, 1% TritonX-100, 10% DMSO for 5 min for pea and 15 min for wheat at 4°C in the dark. Electrophoresis was performed in *ex tempore* prepared TAE buffer (pH8) at 0.5, 1, 2 and 5 V/cm for 10 min for pea nuclei. Lysis of embedded wheat nuclei was performed in the same lysis buffer for 15 min before electrophoresis at 1 V/cm at 4°C for 15 min. The slides were dehydrated in 70% and 96% ethanol for 5 min and dried at room temperature.

RESEARCH ARTICLE

Staining of the comets

The slides were covered with solution of the fluorescent dye acridine orange (10 µg/ml). Visualization of the stained comets was carried out using a fluorescence microscope (Zeiss Jenamed-2) coupled with a digital camera (Samsung Digimax V50). The percentage of DNA in tail was calculated as a measure of DNA damage. Three independent experiments were performed and 50 comets were analyzed per point in each experiment.

Results**Trypan blue dye exclusion test for nuclear membrane intactness after exposure to UVC**

To determine the dose effect of UVC exposure on pea and wheat nuclei, a series of dose-response experiments were performed. Figures 1 and 2 demonstrate the response of the plant nuclei to the various doses.




Figure 1. Nuclei membrane integrity of pea and wheat cells exposed to UVC (*Pisum sativum* – 5, 7 and 9 kJ/m²; *Triticum aestivum* – 14, 16 and 18 kJ/m²) as indicated by staining with Trypan blue. Each result indicates the mean±SE from at least 3 independent experiments.

* significant difference using the Kruskal-Wallis test compared to the control ($p < 0.05$); n.s. no significant differences.

The nuclear membrane intactness determined by Trypan blue dye exclusion test decreased significantly (approximately 20%) in pea nuclei irradiated with 7 kJ/m² UVC radiation as compared to untreated control. The intactness of wheat nuclei after 16 kJ/m² UVC treatment also decreased by approximately 8% compared to control (Figure 1). Significant differences were not observed between intact nuclei in control and nuclei irradiated with the lowest UVC dose. Untreated nuclei displayed a bright-blue color indicating an intact membrane. Dark-blue colored nuclei with membrane corruption and blebbing were also present (Figure 2).




Figure 2. Representative images of Trypan blue stained nuclei isolated from pea leaves. The intact nuclei which provide a negative test are bright and colored in pale blue. Trypan blue-positive nuclei are dark blue as indicated by arrow.

Detection of background level of DNA damage in control samples by neutral Comet assay

For successful accomplishment of the neutral Comet assay a sufficient quantity of intact nuclei is needed. Besides, a neutral Comet assay under different conditions was performed. The methodological changes concerned mainly the lysis duration (data not shown) and electrophoresis conditions – duration and intensity of the electric field. Percentage of DNA in tail as a parameter was most accurate and quantitative indicator of DNA damage since it reflects the total intensity of the tail and the total intensity of the comet being not dependable on the tail length.

RESEARCH ARTICLE




Figure 3. Representative images of control pea nuclei stained with acridine orange isolated after electrophoresis at: a) 0.5 V/cm; b) 5 V/cm.




Figure 4. Values of the parameter % DNA in comet tails under different intensities of the electric field.

Nevertheless, pea nuclei showed high level of DNA migration in control samples at 5 V/cm and 10 min lysis (Figure 3b). Under these experimental conditions DNA fragments migrated out of the head to form a comet tail thus increasing the percentage of DNA in the tail. It is obvious that these conditions were not appropriate to quantify the amount of DNA damage in pea accurately. As shown on the Figure 4, at 5 V/cm electrophoresis conditions the DNA

damage expressed by % DNA in tail was comparatively high and reached 20.44%. At 1 V/cm it was decreased to 11.67%. At 0.5 V/cm and 5 min lysis the amount of damages was reduced to 6.98% for pea (Figure 4). The releasing fragments that diffuse away from the center of the nucleoid create a “halo” with a small compact origin (Figure 3a). The most optimal conditions found to be most suitable for wheat were 15 min lysis and 15 min (1 V/cm) electrophoresis time (Figure 4).

Discussion

DSB are considered to be the most biologically important lesions, producing critical DNA damage, mainly due to the close relationship with chromosomal damage, genome instability, apoptosis etc. (McMillan *et al.*, 2001). Several different experimental protocols have been established for the detection of DSB: ultracentrifugation in neutral sucrose gradients (Iliakis *et al.*, 1991), neutral filter elution (Goutham *et al.*, 2011), pulse-field gel electrophoresis – PFGE (Hong *et al.*, 2010), clamped homogeneous electrical field electrophoresis – CHEF (Blöcher *et al.*, 1989; Blöcher & Kunhi, 1990), static field gel electrophoresis – SFGE (Taucher-Scholz *et al.*, 1995). One of the most sensitive methods is considered to be the single cell gel electrophoresis or the comet assay (Menke *et al.*, 2000a; Collins, 2004).

Both versions of the comet assay protocol under neutral and alkaline conditions can measure a broad range of damage

RESEARCH ARTICLE

as SSB, DSB, alkali labile sites, oxidative DNA base damage, DNA-DNA/DNA-protein/DNA-drug crosslinking (Collins *et al.*, 2008). In all model systems, alkaline version of this analysis is considered to be more sensitive, but SSB are quickly repaired, therefore they are not of significant importance. It is known that DSB are more essential than SSB (Collins *et al.*, 1997) and they can be detected after application of the neutral version.

A number of Comet assay protocols have been developed for different type of studies in plants. Guidelines, recommendations and modifications of the both versions have been already published (Wojewódzka *et al.*, 2002; Christofolletti *et al.* 2009; Lovell & Omori, 2008). Up to now, 3 protocols of the Comet assay are available in plants: A/A (alkaline lysis and electrophoresis), N/N (neutral lysis and electrophoresis) and A/N (alkaline lysis and neutral electrophoresis) protocol. The alkaline variant (A/A) (Klaude *et al.*, 1996; Angelis *et al.*, 1999) has a wider range of SSB and DSB detection. The minimal technical requirements to conduct the alkaline comet assay in pea and wheat have been established (Gichner *et al.*, 2003; Peycheva *et al.*, 2011).

Even though Comet assay in plants is already routinely applied, only few studies have examined DSBs under neutral conditions. Restrictive conditions of the neutral Comet assay in plants can be surmounted by performing the analysis in alkaline lysis and neutral or mildly alkaline electrophoresis (A/N) conditions. In A/N protocol the comet formation is supported by alkaline unwinding of DNA before electrophoresis that allows the expression of free single-stranded DNA fragments in the tail (Menke *et al.*, 2000a). Relevant *in vivo* studies on DNA damage were reported for plants, such as *Allium cepa* L., *Arabidopsis thaliana* L., *Hordeum vulgare* L., *Nicotiana tabacum* L., *Vicia faba* L. However, classical neutral variant (N/N) without the unwinding DNA step was applied only for barley, broad bean and *Arabidopsis* (Koppen & Angelis, 1998; Menke *et al.*, 2000b; Georgieva & Stoilov, 2008; Kozak *et al.*, 2009).

As mentioned above, N/N protocol of the plant comet assay possesses minimal sensitivity (Menke *et al.*, 2000a). For that reason the neutral version of the method needs adaptation for the individual cell types (Wojewydzka *et al.* 2002). Adaptation concerns the main steps of the analysis, including the nuclei isolation as different cytoplasmic components (nucleases and hydrolytic proteins) which influence the background level of induced DSB (Armalytè & Žukas, 2002). The nucleus as an experimental model is more sensitive than the intact cell (Armalytè & Žukas, 2002). One

of the most significant requirements of the analysis is the low level of initial DSB background that could facilitate statistical approaches (Lovell & Omori, 2008). The amount of primary induced DSB is in close relationship with the induction of programmed cell death (Huang *et al.*, 2005; Kuthanova *et al.*, 2008). Studies have shown that plant response after UV irradiation affects nucleic acids, photosynthetic apparatus and membrane lipids (Yao *et al.*, 2012; Kalbina & Strid, 2006; Rao *et al.*, 1996; Hidema *et al.*, 2000; Strid *et al.*, 1990; Chow *et al.*, 1992). Besides typical damages resulting from UV irradiation, direct induction of direct DSB is often discussed. For that reason the quality of the isolated pea and wheat nuclei and the membrane intactness was verified immediately after isolation and after irradiation by applying the Trypan blue dye exclusion test. This blue acid dye contains two azo chromophores group, but it is not capable to enter into the cell wall of plant cells grown in culture (Sini *et al.*, 2012). Previous studies have shown the cell death as result of “classical” ozone symptoms in tobacco cv Bel W3 plants by Trypan blue test (Pasqualini *et al.*, 2003).

Other analytical limitations of the neutral comet assay in plants are connected with migration of DNA under neutral conditions. Often the differences depend on the size of plant genomes and protein content (Armalytè & Žukas, 2002), which impose variation in methodology concerning lysis time, temperature during lysis, and electrophoresis. As shown in previous studies (Singh *et al.*, 1999) we also detected dependence on the length of DNA migration and intensity on the electric field. In addition, these conditions reflect on quantity of DNA damage represented as mean of % DNA in tail of the comets, observed in control samples of pea and wheat nuclei. Similarly to the alkaline comet assay protocol elaborated on *Oreochromis niloticus* (Christofolletti *et al.*, 2009), our results also revealed gradual increase in the intensity of DNA migration due to the effects of electrophoretic conditions on the morphology of comets in control samples. We consider this fact directly related to DNA organization within higher order chromatin structure in the form of ‘matrix attachment sites’ and ‘loops’ (Ventura *et al.*, 2013). The kinetics of the comet tail formation during the alkaline A/A protocol is related to ssDNA fragments, the ends of which are pulled out from the comet head by electric force. In contrast, in N/N protocol the comet tail is formed by extended DNA loops. It is clearly shown that the rate of DNA release is dependent on the topological state of DNA (Afanasieva *et al.*, 2010).

RESEARCH ARTICLE

Conclusion

In this study we found appropriate experimental conditions for neutral Comet assay, specific for pea and wheat by variation in the main steps of the protocol – lysis and electrophoresis. The optimal conditions were specified as: 5 min lysis and electrophoresis at 0.5 V/cm for pea and 15 min lysis and electrophoresis at 1 V/cm for wheat. The amount of nuclei with an intact membrane was decreased after application of high UVC doses in both model plants.

Acknowledgement

This study was supported by the Bulgarian National Science Fund, contract DMU 03/60 and IAEA Research contract 15481. Thanks are due to Maria Krasteva for revising the manuscript linguistically.

References

- Afanasieva K, Zazhytska M, Sivolob A. 2010. Kinetics of comet formation in single-cell gel electrophoresis: loops and fragments. *Electrophoresis*, 31(3): 512-519.
- Angelis KJ, Dusinska M, Collins AR. 1999. Single cell gel electrophoresis: Detection of DNA damage at different levels of sensitivity. *Electrophoresis*, 20: 2133-2138.
- Armalytė J, Žukas K. 2002. Evaluation of UVC-induced DNA damage by SCGE assay and its repair in barley. *Biologija*, 3: 57-59.
- Armalytė J, Žukas K. 2003. Investigation of UVC-induced DNA damage and its repair by SCGE assay in barley. *Ekologia (Vilnius)*, 1: 72-74.
- Blöcher D, Einspinner M, Zajackowski J. 1989. CHEF electrophoresis, a sensitive technique for the determination of DNA double-strand breaks. *Int. J. Radiat. Biol.*, 56(4): 437-448.
- Blöcher D, Kunhi M. 1990. DNA Double-strand Break Analysis by CHEF Electrophoresis. *Int. J. Radiat. Biol.*, 58(1): 23-34.
- Cerda H, Hofsten BV, Johanson KJ. 1993. Identification of irradiated food by microelectrophoresis of DNA from single cells. - In: Leonardi M, Belliardo JJ, Raffi JJ (eds), *Recent Advances of New Methods of Detection of Irradiated Food*, Proceedings of the workshop, Luxembourg, Commission of the European Communities, EUR-14315, p. 401-405.
- Cerda H, Delicée H, Haine H, Rapp H. 1997. The DNA 'comet assay' as a rapid screening technique to control irradiated food. *Mutat. Res.* 375(2): 167-181.
- Chow WS, Strid A, Anderson JM. 1992. Short-term treatment of pea plants with supplementary ultraviolet-B radiation: recovery time-courses of some photosynthetic functions and components. In *Research in Photosynthesis* (ed. N. Murata), 4: 361-364.
- Collins AR, Dobson VL, Dusinska M, Kennedy G, Stetina R. 1997. The Comet assay: what can it really tell us? *Mutation Research*, 375(2): 183-193.
- Collins AR. 2004. The comet assay for DNA damage and repair: principles, applications, and limitations. *Mol. Biotechnol.*, 26: 249-261.
- Collins AR, Oscoz AA, Brunborg G, Gaivao I, Giovannelli L, Kruszewski M, Smith CC, Stetina R. 2008. The comet assay: topical issues. *Mutagenesis*, 23: 143-151.
- Christofolletti CA, David JAO, Fontanetti CS. 2009. Application of the comet assay in erythrocytes of *Oreochromis niloticus* (Pisces): A methodological comparison. *Genet. Mol. Biol.*, 32(1): 155-158.
- Gady ALF, Hermans FWK, Van de Wal MHB, Van Loo EN, Visser RGF, Bachem CWB. 2009. Implementation of two high through-put techniques in a novel application: detecting point mutations in large EMS mutated plant populations. *Plant Methods*, 5: 13.
- Georgieva M, Stoilov L. 2008. Assessment of DNA strand breaks induced by bleomycin in barley by the comet assay. *Environ. Mol. Mutag.*, 49(5): 381-387.
- Gichner T, Patkova Z, Kim JK. 2003. DNA damage measured by the Comet assay in eight agronomic plants. *Biol. Plantarum*, 47(2): 185-188.
- Goutham VH, Kamalesh MD, Guruprasad PK, Vadhiraja MB, Satyamoorthy K, Satish SRB. 2011. A modified fluorimetric neutral filter elution method for analyzing radiation-induced double strand break and repair. *Anal. Biochem.*, 414(2): 287-293.
- Hidema J, Kumagai T, Sutherland BM. 2000. UV radiationsensitive *Norin 1* rice contains defective cyclobutane pyrimidine dimer photolyase. *Plant Cell*, 12: 1569-1578.
- Hong JP, Byun MY, An K, Yang SJ, An G, Kim WT. 2010. OsKu70 is associated with developmental growth and genome stability in rice (*Oryza sativa* L.). *Plant. Physiol.*, 152(1): 374-387.
- Huang X, Halicka HD, Traganos F, Tanaka T, Kurose A, Darzynkiewicz Z. 2005. Cytometric assessment of DNA damage in relation to cell cycle phase and apoptosis. *Cell Prolif.*, 38: 223-243.
- Jovtchev G, Menke M, Schubert I. 2001. The comet assay detects adaptation to MNU-induced DNA damage in barley. *Mutat. Res.*, 493: 95-100.
- Iliakis G, Blöcher D, Metzger L, Pantelias G. 1991. Comparison of DNA double-strand break rejoining as measured by pulsed field gel electrophoresis, neutral sucrose gradient centrifugation and non-unwinding filter elution in irradiated plateau-phase CHO cells. *Int. J. Radiat. Biol.*, 59: 927-939.
- Kalbina I, Strid A. 2006. Supplementary ultraviolet-B irradiation reveals differences in stress responses between *Arabidopsis thaliana* ecotypes. *Plant, Cell and Environment.*, 29: 754-763.
- Klaude M, Eriksson S, Nygren J, Ahnstrom G. 1996. The comet assay: mechanisms and technical considerations. *Mutat. Res.*, 363: 89-96.
- Koppen G, Cerda H. 1997. Identification of low-dose irradiated seeds using the neutral comet assay. *Lebenson-Wiss Technol.*, 30: 452-457.
- Koppen G, Angelis KJ. 1998. Repair of X-ray induced DNA damage measured by the comet assay in roots of *Vicia faba*. *Environ. Mol. Mutag.*, 32: 281-285.
- Koppen G, Toncelli LM, Triest L, Verschaevae L. 1999. The comet assay: A tool to study alteration of DNA integrity in developing plant leaves. *Mech. Ageing. Dev.*, 110: 13-24.

RESEARCH ARTICLE

- Kozak J, West CE, White C, da Costa-Nunes JA, Angelis KJ. 2009. Rapid repair of DNA double strand breaks in *Arabidopsis thaliana* is dependent on proteins involved in chromosome structure maintenance. *DNA Repair*, 8: 413-419.
- Kuthanova A, Opatrny Z, Fischer L. 2008. Is internucleosomal DNA fragmentation an indicator of programmed death in plant cells? *J. Exp. Bot.*, 59(8): 2233-2240.
- Lovell DP, Omori T. 2008. Statistical issues in the use of the comet assay. *Mutagen.*, 23(3): 171-182.
- Maluszynska J, Juchimiuk J. 2005. Plant genotoxicity: A molecular cytogenetic approach in plant bioassays. *Arh. Hig. Rada. Toksikol.*, 56: 177-184.
- McMillan TJ, Tobi S, Mateos S, Lemon C. 2001. The use of DNA double-strand break quantification in radiotherapy. *Int. J. Radiat. Oncol. Biol. Phys.*, 49(2): 373-377.
- Menke M, Angelis KJ, Schubert I. 2000a. Detection of specific DNA lesions by a combination of comet assay and FISH in plants. *Environ. Mol. Mutag.*, 35: 132-138.
- Menke M, Meister A, Schubert I. 2000b. N-Methyl-N-nitrosourea induced DNA damage detected by the comet assay in *Vicia faba* nuclei during all interphase stages is not restricted to chromatid aberration hot spots. *Mutagen.*, 15: 503-506.
- Olive PL. 1989. Cell proliferation as a requirement for development of the contact effect in Chinese hamster V79 spheroids. *Radiat. Res.*, 117: 79-92.
- Östling O, Johanson KJ. 1984. Microelectrophoretic study of radiation-induced DNA damages in individual mammalian cells. *Biochem. Biophys. Res. Commun.*, 123: 291-298.
- Pasqualini S, Piccioni C, Reale L, Ederli L, Della Torre G, Ferranti F. 2003. Ozone-induced cell death in tobacco cultivar Bel W3 plants. The role of programmed cell death in lesion formation. *Plant Physiol.*, 133: 1122-1134.
- Peycheva E, Ivanova E, Ivanov S, Miloshev G. 2011. Improved procedure for Comet assay on active photosynthetic cells from pea plants. *Oxid. Commun.*, 34(2): 320-325.
- Rao MV, Pallyath G, Ormrod DP. 1996. Ultraviolet-B- and ozone-induced biochemical changes in antioxidant enzymes of *Arabidopsis thaliana*. *Plant Physiol.*, 110: 125-136.
- Singh NP, McCoy MT, Tice RR, Schneider EL. 1988. A simple technique for quantitation of low levels of DNA damage in individual cells. *Exp. Cell Res.*, 175: 184-191.
- Singh NP, Stephens RE, Singh H, Lai H. 1999. Visual quantification of DNA double-strand breaks in bacteria. *Mutat. Res.*, 429: 159-168.
- Sini KR, Haribabu Y, Sajith MS, Sreekumar SK. 2012. *In-vitro* cytotoxic activity of *Orthosiphon thymiflorus* Roth. Sleensen leaf extract against Dalton lymphoma ascites Cell Line. *J. Chem. Pharm. Res.*, 4(1): 917-921.
- Strid A, Chow WS, Anderson JM. 1990. Effects of supplementary UV-B radiation on photosynthesis in *Pisum sativum*. *Biochimica et Biophysica Acta*, 1020: 260-268.
- Taucher-Scholz G, Heilmann J, Schneider M, Kraft G. 1995. Detection of heavy-ion-induced DNA double-strand breaks using static-field gel electrophoresis. *Radiation and Environmental Biophysics*, 34(2): 101-106.
- Ventura L, Giovannini A, Savio M, Donà M, Macovei A, Buttafava A, Carbonera D, Balestrazzi A. 2013. Single cell gel electrophoresis (Comet) assay with plants: Research on DNA repair and ecogenotoxicity testing. *Chemosphere*, 92(1): 1-9.
- Uhl M, Plewa M, Majer BJ, Knasmüller S. 2003. Basic principles of genetic toxicology with an emphasis on plant bioassays. In: Maluszynska J, Plewa M, editors. *Bioassays in plant cells for improvement of ecosystem and human health*. Katowice: Wydawnictwo Uniwersytetu Śląskiego, 11-30.
- Wojewódzka M, Buraczewska I, Kruszewski M. 2002. A modified neutral comet assay: Elimination of lysis at high temperature and validation of the assay with anti-single-stranded DNA antibody. *Mutat. Res.*, 518(1): 9-20.
- Yao Y, Danna CH, Ausubel FM, Kovalchuk I. 2012. Perception of volatiles produced by UVC-irradiated plants alters the response to viral infection in naïve neighboring plants. *Plant Signal Behav.*, 7(7): 741-745.