

Interannual variability of the stratospheric wave driving during northern winter

A. J. Haklander^{1,2}, P. C. Siegmund³, and H. M. Kelder^{1,2}

¹Eindhoven University of Technology (TUE), Department of Applied Physics, P.O. Box 513, 5600 MB Eindhoven, The Netherlands

²Royal Netherlands Meteorological Institute (KNMI), Climate and Seismology Department, Climate Observation Division, P.O. Box 201, 3730 AE De Bilt, The Netherlands

³Royal Netherlands Meteorological Institute (KNMI), Climate and Seismology Department, Climate and Chemistry Division, P.O. Box 201, 3730 AE De Bilt, The Netherlands

Received: 6 December 2006 – Published in Atmos. Chem. Phys. Discuss.: 5 January 2007

Revised: 16 March 2007 – Accepted: 8 May 2007 – Published: 16 May 2007

Abstract. The strength of the stratospheric wave driving during northern winter is often quantified by the January–February mean poleward eddy heat flux at 100 hPa, averaged over 40°–80° N (or a similar area and period). Despite the dynamical and chemical relevance of the wave driving, the causes for its variability are still not well understood. In this study, ERA-40 reanalysis data for the period 1979–2002 are used to examine several factors that significantly affect the interannual variability of the wave driving. The total poleward heat flux at 100 hPa is poorly correlated with that in the troposphere, suggesting a decoupling between 100 hPa and the troposphere. However, the individual zonal wave-1 and wave-2 contributions to the wave driving at 100 hPa do exhibit a significant coupling with the troposphere, predominantly their stationary components. The stationary wave-1 contribution to the total wave driving significantly depends on the latitude of the stationary wave-1 source in the troposphere. The results suggest that this dependence is associated with the varying ability of stationary wave-1 activity to enter the tropospheric waveguide at mid-latitudes. The wave driving anomalies are separated into three parts: one part due to anomalies in the zonal correlation coefficient between the eddy temperature and eddy meridional wind, another part due to anomalies in the zonal eddy temperature amplitude, and a third part due to anomalies in the zonal eddy meridional wind amplitude. It is found that year-to-year variability in the zonal correlation coefficient between the eddy temperature and the eddy meridional wind is the most dominant

factor in explaining the year-to-year variability of the poleward eddy heat flux.

1 Introduction

According to the downward-control principle, the stratospheric residual meridional circulation at any level is controlled by the vertically-integrated zonal force due to breaking Rossby and gravity waves above that level (Haynes et al., 1991). The breaking waves deposit westward angular momentum into the relative and planetary angular momentum “reservoirs”, causing a westward acceleration and a poleward displacement of the air (e.g., Andrews et al., 1987). In the low-frequency limit, the westward acceleration is zero and the meridional circulation is referred to as the Brewer-Dobson circulation (e.g., Shepherd, 2000). The term “downward control” can, however, be somewhat misleading, since it is predominantly the wave activity emanating from below that determines the amount of angular momentum deposited aloft.

The net zonal-mean upward flux of wave activity is represented by the upward component F_z of the Eliassen-Palm (E-P) flux, which for quasigeostrophic flow is proportional to the zonal-mean poleward eddy heat flux $[\overline{v^*T^*}]$, where the square brackets denote the zonal average and the asterisk denotes the deviation thereof (e.g. Andrews et al., 1987; Newman and Nash, 2000). In the lower stratosphere, $[\overline{v^*T^*}]$ (and therefore F_z) exhibits a strong, positive correlation with the tendency of total ozone at mid- and high latitudes during northern winter (e.g., Fusco and Salby, 1999; Randel et al.,

Correspondence to: A. J. Haklander
(haklande@knmi.nl)

2002). Also, during late northern winter the midlatitude eddy heat flux in the lower stratosphere is highly and positively correlated with the temperature in early March at high latitudes, and consequently with the strength of the polar vortex (e.g., Newman et al., 2001; Polvani and Waugh, 2004). Both observations can be explained by the wave-induced poleward transport of ozone-rich air from the tropical source and the subsequent adiabatic compression at higher latitudes.

This fundamental link between the poleward eddy heat flux and the dynamics and chemistry of the stratosphere is now well understood and also quantitatively employed as a diagnostic to validate coupled chemistry-climate models (CCMs) (Austin et al., 2003; Eyring et al., 2005). Austin et al. used the $[v^*T^*]$ field at 100 hPa averaged for January–February over 40° – 80° N, hereafter referred to as H_{100} , as a measure of the net upward flux of midwinter wave activity in the lower stratosphere. They evaluated several CCMs with respect to H_{100} , the polar stratospheric temperatures in early spring, and the (almost linear) relationship between both diagnostics. Since almost all of the planetary wave activity crosses the 100 hPa level between 40° – 80° N (Hu and Tung, 2003), H_{100} indeed provides a good measure of the net total wave activity propagating into the stratosphere. Several studies have been performed to analyze trends in H_{100} (or a similar diagnostic) over the last decades, and to predict future trends. Hu and Tung (2003) found a significant downward trend over 1979–2002 for $[v^*T^*]$ at 100 hPa, averaged over 50° – 90° N and January to March. Austin et al. (2003) compared future H_{100} trends in a number of enhanced- CO_2 climate simulations, and found for most models a slightly negative trend over the next few decades, although at best of marginal statistical significance. One of the models showed a highly significant positive trend. For a doubled CO_2 climate, Sigmond et al. (2004) computed a significant increase of the Northern Hemisphere (NH) stratospheric residual circulation during winter, corresponding to an increase of the poleward eddy heat flux. These examples illustrate that future predictions of H_{100} are very uncertain. However, in a recent multi-model study by Butchart et al. (2006), increasing greenhouse gas concentrations were found to yield an overall strengthening of the Brewer-Dobson circulation and the associated wave driving, with the strongest trend in NH winter. Butchart et al. conclude that it remains an important future task to identify the causes of the increase in wave driving. Thus, despite the fact that it is widely acknowledged that the stratospheric wave driving has a large impact on the dynamics and chemistry of the stratosphere, the causes for its trends and variability are still not well understood.

In the present study, H_{100} is used as a measure of the net total midwinter wave activity that propagates from the troposphere into the stratosphere, following Austin et al. (2003), and Eyring et al. (2005). Our analysis puts the emphasis on the interannual variability of H_{100} , in order to obtain a better understanding of the causes of this observed variability, at both the interannual and the decadal timescale. Year-to-year

variations in H_{100} can be attributed to many factors. The strength of the tropospheric wave source is an obvious factor, but also the shape of the source spectrum determines the amount of wave activity that reaches 100 hPa, as is described by the Charney–Drazin criterion (Charney and Drazin, 1961). This criterion states that stationary planetary waves can only propagate upward in a westerly zonal-mean flow that is not too strong. Only the longest waves can propagate through stronger westerlies, which implies that during winter only the longest waves can propagate into the stratosphere. The background zonal-mean flow determines the properties of this low-pass filter, and therefore also affects the interannual variability of H_{100} . The main goal of the present study is to investigate to what extent the year-to-year variability in H_{100} is affected by several factors. The factors that are examined include the strength of the total upward wave activity flux in the troposphere, the shape of the wave activity spectrum, the latitude and height of the wave source, and the refractive properties of the background flow. It is also examined whether the zonal correlation coefficient between the meridional wind and the temperature has a significant effect on the interannual variability of H_{100} . Our analysis is based on ERA-40 reanalysis data for the period of 1979–2002. In Sect. 3, estimated values are often presented along with an error bar. These represent the associated standard deviations.

The structure of the paper is as follows. Section 2 gives a description of the data sources and discusses various ways of decomposing the poleward eddy heat flux. The main results are given in Sect. 3, and a summary and discussion of the results is presented in Sect. 4.

2 Data and method

2.1 Data

We use 6-hourly temperature and horizontal wind fields for 24 years (1979–2002) of ERA-40 reanalysis data (Simmons and Gibson, 2000). Although the ERA-40 dataset starts in 1957, we have only included the satellite era, from 1979 onwards, in our analysis. For this period, the global observation data coverage is considered to be good (Uppala et al., 2005). ERA-40 is produced by the European Centre for Medium-Range Weather Forecasts (ECMWF). The meteorological fields were retrieved for a lat-lon grid of $2.5^\circ \times 2.5^\circ$ between 0° – 90° N, at 23 pressure levels between 1000 and 1 hPa.

2.2 Linear regression analysis

The total poleward eddy heat flux can be decomposed into the sum of several components, such as the stationary and transient wave components. The impact of each component on the interannual variability of the total heat flux can be evaluated by performing a linear regression analysis. This can be understood as follows. It can easily be shown that if

$y \equiv \sum x_i$, then the variance of y equals the sum of the covariances between y and the x_i , i.e., $\text{var}(y) = \sum \text{cov}(x_i, y)$. Thus, $\text{cov}(x_i, y)$ can be interpreted as the contribution of x_i to $\text{var}(y)$. If we define $b_i \equiv \text{cov}(x_i, y) \text{var}^{-1}(y)$, where $\sum b_i = 1$, then b_i is the regression coefficient for the linear least-squares fit given by $\tilde{x}_i = a_i + b_i y$. It should be noted that the correlation coefficient r_i between y and x_i can be large, while the associated value of b_i is small. For this reason, it is useful to consider both b_i and r_i . In the present study, y represents the total heat flux and the x_i represent the various components of the total heat flux (see Sect. 2.3). Hereafter the phrase “a linear regression of v_1 with v_2 is performed” implies that v_1 is the independent and v_2 is the dependent variable.

2.3 Decompositions of the heat flux

The total poleward eddy heat flux, averaged over space and time, can be decomposed by separating the eddy meridional wind and eddy temperature into a stationary and a transient part. If we denote the temporal average over January and February by an overbar, the deviation thereof by a prime, and the cosine-latitude weighted spatial average between 40° – 80° N by angle brackets, we can write H at a certain pressure level as

$$H \equiv \langle \overline{v^* T^*} \rangle = \langle \overline{\bar{v}^* \bar{T}^*} \rangle + \langle \overline{v'^* T'^*} \rangle, \quad (1)$$

where H at 100 hPa was previously denoted by H_{100} . The asterisk denotes the deviation of the zonal mean. The r.h.s. terms of Eq. (1) thus describe the contribution to H by the stationary and transient eddies, respectively.

In addition to the temporal decomposition in Eq. (1), a spatial decomposition can be made by discriminating between the individual zonal wavenumber components of the heat flux. The zonal wave component of the total eddy heat flux is calculated as the product of the wave components of v and T , v_s and T_s (e.g., Newman and Nash, 2000). This yields

$$H = \sum_{s \geq 1} \langle \overline{v_s T_s} \rangle, \quad (2)$$

where the subscript s denotes the zonal wave- s component. Combining Eqs. (1) and (2) yields

$$H = \sum_{s \geq 1} \langle \overline{\bar{v}_s \bar{T}_s} \rangle + \sum_{s \geq 1} \langle \overline{v'_s T'_s} \rangle. \quad (3)$$

The impact of these individual stationary and transient wave components on the interannual variability of the total heat flux will be evaluated by a linear regression analysis in the next section.

A different way of decomposing the total eddy heat flux, is by noting that $[v^* T^*]$ can be expressed as

$$[v^* T^*] = r_{v,T} \sigma_v \sigma_T, \quad (4)$$

where $r_{v,T}$ is the zonal correlation coefficient, and σ_v and σ_T are the zonal standard deviations of v and T . If v or T would consist of only one wave component, then their wave amplitudes would be given by $\sqrt{2}\sigma_v$ and $\sqrt{2}\sigma_T$, respectively, and $r_{v,T}$ would be equal to $\cos \Delta\varphi$, where $\Delta\varphi$ is the phase difference between the v and the T wave patterns. Therefore, $r_{v,T}$ might be considered as the cosine of the ‘effective phase difference’ between v and T for the total spectrum of waves, with $\sqrt{2}\sigma_v$ and $\sqrt{2}\sigma_T$ representing the effective v and T amplitudes, respectively. The larger $r_{v,T}$, the smaller the effective phase difference between the v and T fields, and the more efficiently heat is transported poleward by the waves. To investigate both the qualitative and the quantitative effects of year-to-year variations in $r_{v,T}$, σ_v , and σ_T on the interannual variability of H_{100} , we use the following approach (e.g., Siegmund, 1995). First, we rewrite the average of Eq. (4) over 40° – 80° N and January–February as follows:

$$H \equiv \langle \overline{r_{v,T} \sigma_v \sigma_T} \rangle \equiv \langle \overline{r_{v,T}} \rangle \langle \overline{\sigma_v} \rangle \langle \overline{\sigma_T} \rangle + \tilde{E} \equiv \tilde{H} + \tilde{E}, \quad (5)$$

where \tilde{E} denotes the error that is introduced by neglecting the zonal and temporal cross-correlations between $r_{v,T}$, σ_v , and σ_T . The validity of the approximation $H_{100} \approx \tilde{H}_{100}$ will be discussed in Sect. 4. Dropping the angular brackets and the overbars, denoting the 24-year mean by a superscript m and the deviation thereof by a Δ , $\Delta \tilde{H}$ can be linearized as:

$$\Delta \tilde{H} \equiv \Delta \tilde{H} + \Delta \tilde{E} \equiv \sigma_v^m \sigma_T^m \Delta r_{v,T} + \sigma_T^m r_{v,T}^m \Delta \sigma_v + \sigma_v^m r_{v,T}^m \Delta \sigma_T + \Delta \tilde{E}, \quad (6)$$

where $\Delta \tilde{E}$ denotes the error that arises from the linearization. If Eqs. (5) and (6) are combined, we obtain

$$\Delta H \equiv \sigma_v^m \sigma_T^m \Delta r_{v,T} + \sigma_T^m r_{v,T}^m \Delta \sigma_v + \sigma_v^m r_{v,T}^m \Delta \sigma_T + \Delta \tilde{E} + \Delta \tilde{E}. \quad (7)$$

Equation (7) will be used to analyze the contribution to the interannual variability of H_{100} by the interannual variabilities of $r_{v,T}$, σ_v , and σ_T .

3 Results

3.1 Timeseries of H_{100}

The timeseries of H_{100} (1979–2002) is displayed as the thick solid line with filled circles in Fig. 1. During this 24-year period, H_{100} ranged between 11.2 and 19.2 K m s^{-1} , and the average value of H_{100} was $15.1 \pm 0.5 \text{ K m s}^{-1}$. There is a slightly negative trend in H_{100} , but the trend is not statistically significant (93.2% confidence level). Hu and Tung (2003) did find a highly significant downward trend over 1979–2002 in the vertical EP-flux component north of 50° N for the January–March period, which was suggested to be due to the effect of Arctic ozone depletion during late winter and spring. This effect is likely smaller for H_{100} , which is the average over the January–February period. If we regard H_{100} as the sum of its zonal wave- s components as expressed by Eq. (2), we find that the $s = 1$ –3 components of H_{100} account for more than 90% of H_{100} . The first three wave components




Fig. 1. Poleward eddy heat flux at 100 hPa, averaged over 40°–80° N and January–February, for the years 1979–2002. The sum of all wave contributions H_{100} is shown, as well as the separate wavenumber 1–3 components of H_{100} .

Table 1. Linear regression coefficients \pm their standard error, and correlation coefficients for the linear regression of H_{100} with its total, stationary and transient $s=1-5$ components. The linear regression is performed over 1979–2002.

s	Regr. coeff. (b_i)			Corr. coeff. (r_i)		
	Tot $\pm\sigma$	Stat $\pm\sigma$	Tran $\pm\sigma$	Tot	Stat	Tran
1	0.74 \pm 0.15	0.56 \pm 0.23	0.18 \pm 0.14	0.73	0.47	0.26
2	0.12 \pm 0.16	0.06 \pm 0.16	0.06 \pm 0.08	0.16	0.08	0.15
3	0.09 \pm 0.08	0.02 \pm 0.08	0.08 \pm 0.05	0.24	0.05	0.30
4	0.00 \pm 0.03	-0.02 \pm 0.02	0.02 \pm 0.03	0.01	-0.19	0.13
5	0.00 \pm 0.02	0.00 \pm 0.00	-0.01 \pm 0.02	-0.04	0.23	-0.09
all	1	0.63 \pm 0.16	0.37 \pm 0.16	1	0.65	0.46

of H_{100} are also shown in Fig. 1. Both the variance and the 24-year average decrease with increasing wavenumber. For the majority of the years, variations in H_{100} seem to be dominated by the wave-1 variations, indicated by the line with open circles in Fig. 1. A relatively constant factor is that the wave-1 and wave-2 variations tend to be opposite. The correlation coefficient for the wave-1 and wave-2 components is $r=-0.42$, which is significant at a 95.8% confidence level. The separate $s=1, 2$, and 3 components of H_{100} all exhibit a slightly negative trend over 1979–2002, but the statistical significance of these trends is well below the 95% confidence level.

3.2 Wave contributions to interannual variability of H_{100}

To further examine the interannual variability of H_{100} , we decompose H_{100} into its stationary and transient wave components, as expressed by Eq. (3). Taking H_{100} as the independent variable, the linear regression coefficient b_i for




Fig. 2. Correlation coefficients of H_{100} with $\langle v^*T^* \rangle_{NH}$. Correlation coefficients are calculated over the years 1979–2002.

the regression of H_{100} with its stationary or transient wave component represents the relative contribution by that wave component to the total variance of H_{100} , as discussed in Sect. 2. Table 1 shows the regression coefficients and the corresponding correlation coefficients for the total (stationary plus transient), stationary, and transient wave 1–5 components of H_{100} . Using a Student's t -test, we find for a sample size of 24, that the correlation coefficient r_i is significantly different from zero at a >95%, 99%, or 99.9% confidence level if $|r_i| > 0.40, 0.52$ or 0.63 , respectively. The sum of the regression coefficients for the total wave 1–5 contribution to H_{100} is 0.95 ± 0.05 , with a corresponding correlation coefficient $r_i=0.996$ (not shown). Therefore, wavenumbers 6 and higher can be neglected in the analysis of the interannual variability of H_{100} . In fact, almost all of the interannual variability of H_{100} is due to $s=1-3$, with a combined regression coefficient of $b_i=0.96 \pm 0.06$ and $r_i=0.98$. Only retaining $s=1,2$ yields $b_i=0.86 \pm 0.11$ and $r_i=0.88$. We can thus state that about 85% of the interannual variability of H_{100} can be attributed to its $s=1,2$ components. When analyzing the interannual variability of H_{100} , it is sufficient to only consider wavenumbers $s=1-3$, or perhaps even $s=1,2$. The bottom row of Table 1 clearly shows that the interannual variability of H_{100} is dominated by the stationary waves, with

$b_i=0.63\pm 0.16$ and $r_i=0.65$. However, the correlation coefficient of H_{100} with its transient component is also significant at the 97.6% confidence level.

3.3 Vertical coupling

As described in the introduction, H_{100} is proportional to the total net wave activity flux emanating from below between 40° N and 80° N at the 100 hPa level. Most of the planetary wave activity that constitutes H_{100} has propagated upward from its tropospheric source, and therefore, it is expected that H_{100} has a significant positive correlation coefficient with the poleward eddy heat flux at levels below (and above) 100 hPa. To verify this, we examine the amount of vertical coupling between 100 hPa and other levels. We do this by calculating the vertical profile of the correlation coefficient of H_{100} with $\langle \overline{v^*T^*} \rangle_{NH}$, which we define as the 20° – 90° N average of $\overline{v^*T^*}$, at levels from 1000 hPa to 1 hPa. We compute the correlation coefficients with $\langle \overline{v^*T^*} \rangle_{NH}$, since the wave activity flux contributing to H_{100} may have partially originated from, or may propagate into, latitudes outside the 40° – 80° N band. The correlation coefficient profile is shown in Fig. 2. At 100 hPa, H_{100} is highly correlated with $\langle \overline{v^*T^*} \rangle_{NH}$ ($r=0.95$), suggesting that the 100 hPa wave activity outside the 40° – 80° N area is relatively small. This is confirmed by the fact that H_{100} accounts for about 90% of $\langle \overline{v^*T^*} \rangle_{NH}$ at 100 hPa in the 1979–2002 period. Therefore, to simplify the interpretation of our results, we neglect the 100 hPa upward wave activity flux at latitudes outside the 40° – 80° N window. In Fig. 2, we see that the level of maximum correlation coefficient between H_{100} and $\langle \overline{v^*T^*} \rangle_{NH}$ is found at 70 hPa rather than 100 hPa. Above 70 hPa, the correlation coefficient gradually decreases to marginally significant values near the stratopause ~ 1 hPa. Below 100 hPa, the correlation coefficient falls off quite rapidly, and is not significant ($|r| < 0.40$) below 200 hPa. Thus, there exists a decoupling between variations in the total upward flux of wave activity in the troposphere (below 200 hPa) and variations at 100 hPa. Assuming that the wave sources are predominantly located in the troposphere, this decoupling can be understood as follows. Suppose an amount of wave activity propagates upward in the NH at some tropospheric pressure level p during January–February. A fraction α_p of this wave activity is absorbed below 100 hPa, where α_p lies between 0 and 1. Assuming that no wave activity crosses the equator, the remaining fraction of the wave activity ($1-\alpha_p$) thus reaches the 100-hPa level and contributes to H_{100} . If α_p is less than one and independent of time, then the correlation coefficient between H_{100} and the wave activity flux at pressure level p will be one. On the other hand, if α_p has a large interannual variability, the correlation coefficient will be small.

For a plane and conservative planetary wave, the zonal wavenumber s and the frequency remain constant along its path (or “ray”) in the meridional plane (e.g., Karoly and

Hoskins, 1982). Therefore, it makes sense to repeat the previous analysis for each individual zonal wave component. To examine the amount of vertical coupling as a function of zonal wavenumber (stationary plus transient), we compute the correlation coefficients of the wave- s component of H_{100} with the (same) wave- s component of $\langle \overline{v^*T^*} \rangle_{NH}$ at other levels. We perform this analysis for the individual wavenumbers 1 to 5 and 6+, separately. The results are shown in Fig. 3a. The $s=1$ component of $\langle \overline{v^*T^*} \rangle_{NH}$ is significantly correlated with the $s=1$ component of H_{100} at 500 hPa, and at all levels above 400 hPa. The $s=2$ component of $\langle \overline{v^*T^*} \rangle_{NH}$ exhibits a significant correlation with the $s=2$ component of H_{100} throughout the troposphere and the stratosphere. Note that the wave-2 correlation coefficient reaches a minimum in the mid-troposphere but that the correlation coefficients for $s=2$ are statistically significant at all levels below 1 hPa. For waves with $s > 2$, the correlation coefficients generally only exceed the 95% significance level between 250 hPa and 50 hPa. Thus, the lack of correlation we saw in Fig. 2 between the total upward flux of wave activity in the troposphere and that at 100 hPa is also observed for the $s > 2$ flux components, but not (entirely) for $s=1,2$. This result implies that a significant part of the interannual variability in the $s = 1,2$ components of H_{100} is due to year-to-year variations in the strength of the $s=1,2$ wave source in the troposphere. There is no significant correlation for the $s > 2$ flux components, since the refractive index for those waves exhibits a vertical layer of negative values in the lower stratosphere (Fig. 5b). Therefore, meridional refraction and reflection of the $s > 2$ waves is taking place below 100 hPa, so that H_{100} is dominated by waves 1 and 2.

We have just shown that the separate $s=1,2$ components of H_{100} , to which about 85% of the interannual variability of H_{100} can be attributed, are significantly correlated with the separate $s=1,2$ components of $\langle \overline{v^*T^*} \rangle_{NH}$ in the troposphere. But to what extent can the interannual variability of the total H_{100} be attributed to year-to-year variations in the separate $s=1$ and $s=2$ components (or higher) of $\langle \overline{v^*T^*} \rangle_{NH}$ in the troposphere? To answer this question, we examine the correlation coefficient of H_{100} (i.e., the sum of all wave components) with the separate wave components of $\langle \overline{v^*T^*} \rangle_{NH}$. The results are shown in Fig. 3b, where the wavenumber of the pressure-dependent $\langle \overline{v^*T^*} \rangle_{NH}$ component is given along the horizontal axis. At levels below 200 hPa, the $s=1,2$ components of $\langle \overline{v^*T^*} \rangle_{NH}$ are not significantly correlated with H_{100} . In fact, none of the wave components is. However, in the upper stratosphere, the $s=4$ component of $\langle \overline{v^*T^*} \rangle_{NH}$ exhibits a remarkably strong correlation with H_{100} . It is interesting to compare Figs. 3a and 3b. We observe in Fig. 3a that the wave-4 component of $\langle \overline{v^*T^*} \rangle_{NH}$ in the upper stratosphere is not at all correlated with the wave-4 component of H_{100} . However, Fig. 3b shows that the correlation coefficient with the total H_{100} is highly significant for the wave-4 component of $\langle \overline{v^*T^*} \rangle_{NH}$ in the upper stratosphere ($r=0.58$ at 2 hPa, 99.7% confidence level).




Fig. 3. Correlation coefficients of (a) the zonal wave- s component of H_{100} , and (b) H_{100} , with the zonal wave- s component of $\overline{v^*T^*}_{NH}$. Correlation coefficients are calculated over 1979–2002, and the areas with >95% confidence levels are shaded.

The results suggest that wavenumber 4 is a preferred mode for the breaking of very long planetary waves in the upper stratosphere.

3.4 Correlation patterns in the meridional plane

Thusfar, we have only considered averages over 40° – 80° N and averages over the Northern Hemisphere north of 20° N. In the previous subsection we mentioned that α_p also depends on the latitude at which the waves propagate upward. Therefore, we next examine where the zonal-mean upward wave-activity flux, which is proportional to $\overline{v^*T^*}$, is significantly correlated with H_{100} . The latitude- and pressure-dependent correlation coefficient of H_{100} with $\overline{v^*T^*}$ is shown in Fig. 4a. The highest correlation coefficient is found at 100 hPa and 62.5° N ($r=0.87$). The decoupling between the total upward wave-activity flux at 100 hPa and that in the troposphere (Fig. 2) is also visible in Fig. 4a: H_{100} is not significantly correlated with $\overline{v^*T^*}$ in the lower and middle troposphere. We previously saw that the $s=1,2$ components of H_{100} significantly correlate with the same wave- s components of $\overline{v^*T^*}_{NH}$ at some level in the mid- and lower troposphere. Figure 4b shows that the wave-1 component of H_{100} is significantly correlated with the wave-1 component of $\overline{v^*T^*}$ in the troposphere, between about 40° N and 60° N. Note that the meridionally confined correlation coefficient maximum in the troposphere tilts poleward with height. A possible explanation for this correlation coefficient maxi-

imum would be the presence of a waveguide, through which the wave-1 activity is ducted to 100 hPa (e.g., Karoly and Hoskins, 1982). Since a waveguide can be identified as a ridge in the refractive index field, we verify this by computing the climatological January–February pattern of the refractive index squared for $s=1$, which is shown in Fig. 5a. The wave-1 correlation coefficient maximum is roughly denoted by the dashed line. We see that the refractive index indeed has a ridge in the mid-latitude middle and upper troposphere, which suggests that the tropospheric correlation coefficient maximum in Fig. 4b may be regarded as the signature of this tropospheric wave guide. To illustrate how the upper troposphere and lower stratosphere act as a low-pass filter for planetary wave activity from below, we show the refractive index field for $s=3$ in Fig. 5b. A mid-latitude vertical layer with negative values in the lower stratosphere emerges, of which the vertical extent increases with increasing wavenumber (not shown). For wavenumber 2 (Fig. 4c), a connection with the (lower) troposphere is found that is similar to that of wavenumber 1. A marked difference between Figs. 4c and b is, that the area of maximum correlation coefficients in the troposphere is found at higher latitudes (50 – 70° N) in Fig. 4c. (The distinct and very high maximum of $r=0.82$ at 1000 hPa may not be very meaningful due to the extrapolation below ground.) This correlation coefficient maximum cannot be linked to an $s=2$ waveguide, since the refractive index field in this region is highly variable due to the presence of a zero-wind critical line, which is not sensitive to the zonal




Fig. 4. Correlation coefficient of (a) H_{100} , (b) (c) the $s=1,2$ component of H_{100} , (d) the stationary component of H_{100} , (e) (f) the stationary $s = 1,2$ component of H_{100} (g) the transient component of H_{100} , (h), (i) the transient $s=1,2$ component of H_{100} ; with the same latitude- and pressure-dependent wave component of the zonal-mean eddy heat flux averaged over January–February. Correlation coefficients are calculated over 1979–2002, and only the areas with $>95\%$ confidence levels are shown.

wave number of the refractive index. For wavenumber 2, the latitude of maximum correlation coefficients shifts equatorward with height above 100 hPa. For the wave-3 contribution to the heat flux, the link with the troposphere is absent and the area with significant correlation coefficients is much more confined (not shown). A further decomposition into stationary and transient wave components (Figs. 4d–i) reveals that the link with the (lower) troposphere is only statistically significant for the stationary part of the wave-1 and wave-2 contributions. The tropospheric meridional dipole structure in the stationary wave-1 correlation coefficient map (Fig. 4e) implies that the stationary wave-1 component of H_{100} is sensitive to the latitude of the tropospheric wave-1 source. If the source is located too far south, less wave activity is able to enter the mid-latitude waveguide and contribute to H_{100} . Such a dipole structure is not found in Fig. 4f for $s=2$.

3.5 An alternative analysis of the interannual variability of H_{100}

In Sect. 2.3, we mentioned an alternative way of decomposing H_{100} , by noting that $[v^*T^*]$ equals the product of $r_{v,T}$ with σ_v and σ_T (Eq. 4). Subsequently, we assumed in Eq. (5) that H_{100} can be approximated by \tilde{H}_{100} . To examine the accuracy of this approximation, we first compare the 24-year averages of both \tilde{H}_{100} and H_{100} . This yields $14.1 \pm 0.4 \text{ K m s}^{-1}$ and $15.1 \pm 0.5 \text{ K m s}^{-1}$, respectively with a mean ratio of 0.94 ± 0.02 . To see if \tilde{H}_{100} and H_{100} also have comparable interannual variability, we performed a linear regression of H_{100} with \tilde{H}_{100} . The regression yields a high correlation coefficient ($r=0.85$) and a regression coefficient $b=0.71 \pm 0.09$. We conclude that \tilde{H}_{100} is indeed a useful approximation of H_{100} . We next linearize the deviation of




Fig. 5. Refractive index square $a^2 n_s^2$ in m^2 , based on the 1979–2002 average of the zonal-mean zonal wind during January–February, for (a) $s=1$, and (b) $s=3$. Except for the 150 and 300 m^2 contours, contours are from -100 to $+100$ m^2 , with an interval of 10 m^2 . Negative values are indicated by dark shading, positive values below 20 m^2 by light shading. The 0 , 20 , 40 , and 150 m^2 contours have been labeled, and the location of the $s=1$ correlation coefficient maximum is roughly denoted by the dashed line in (a).

Table 2. Linear regression of H_{100} with $\sigma_v^m \sigma_T^m \Delta r_{v,T}$, $\sigma_T^m r_{v,T}^m \Delta \sigma_v$, and $\sigma_v^m r_{v,T}^m \Delta \sigma_T$ at 100 hPa. The linear regression is performed over 1979–2002. The 1979–2002 averages and standard deviations are for $\langle \overline{r_{v,T}} \rangle$, $\langle \overline{\sigma_v} \rangle$, and $\langle \overline{\sigma_T} \rangle$, respectively.

	Regression coefficient	Corr. coefficient	1979–2002 average	1979–2002 stdev
$\sigma_v^m \sigma_T^m \Delta r_{v,T}$	0.40 ± 0.20	0.39	$r_{v,T}^m = 0.23$	0.04
$\sigma_T^m r_{v,T}^m \Delta \sigma_v$	0.07 ± 0.09	0.17	$\sigma_v^m = 10.4 \text{ m s}^{-1}$	0.8 m s^{-1}
$\sigma_v^m r_{v,T}^m \Delta \sigma_T$	0.23 ± 0.14	0.34	$\sigma_T^m = 5.9 \text{ K}$	0.7 K

\hat{H}_{100} from its 1979–2002 mean as $\Delta \hat{H}_{100}$ (Eq. 6). The error $\Delta \hat{E}_{100}$ that arises from the linearization in Eq. (6) is remarkably small: a linear regression of $\Delta \hat{H}_{100}$ with $\Delta \hat{H}_{100}$ yields $b = 0.98 \pm 0.03$ and $r = 0.99$. The difference between ΔH_{100} and $\Delta \hat{H}_{100}$ in Eq. (7) thus primarily results from $\Delta \hat{E}_{100}$. We can use Eq. (7) to analyze the sensitivity of H_{100} to the year-to-year variations in the effective phase difference $\langle \overline{r_{v,T}} \rangle$, as well as to the interannual variability of the effective amplitudes $\langle \overline{\sigma_v} \rangle$ and $\langle \overline{\sigma_T} \rangle$. The regression and correlation coefficients of the linear regression analysis of H_{100} with the first three terms on the r.h.s. of Eq. (7) at 100 hPa are given in Table 2. The results show that the interannual variability of H_{100} is more sensitive to the $\langle \overline{r_{v,T}} \rangle$ than to $\langle \overline{\sigma_v} \rangle$ and $\langle \overline{\sigma_T} \rangle$. Therefore, a significant part of the year-to-year variability in H_{100} is not determined by variability in the amplitude of the waves but by variability in the efficiency of the

poleward heat transport, as represented by $\langle \overline{r_{v,T}} \rangle$. We also note that the variability in $\langle \overline{\sigma_T} \rangle$ affects H_{100} more strongly than the variability in $\langle \overline{\sigma_v} \rangle$.

4 Summary and discussion

We have studied the interannual variability of the stratospheric wave driving during NH winter, as quantified by H_{100} , being the January–February mean of the 40° – 80° N average of the total poleward heat flux at 100 hPa. For our analysis, we used 24 years (1979–2002) of ERA-40 reanalysis data from ECMWF. The results can be summarized as follows. We have examined the sensitivity of H_{100} to several factors. The first factor is the strength of the total tropospheric wave source. It was found that H_{100} is not

significantly correlated with the total upward wave activity flux below 200 hPa. However, both the individual zonal wave-1 and wave-2 components exhibit significant vertical coupling between 100 hPa and lower (as well as higher) levels. About 85% of the interannual variability of H_{100} can be attributed to its $s=1,2$ components. However, the interannual variability of H_{100} cannot be attributed to either of these individual wave components of the heat flux in the troposphere. Presumably, this is in part due to the statistically significant negative correlation coefficient that we found between the $s=1$ and $s=2$ components of H_{100} . This negative correlation coefficient is also observed on an intraseasonal timescale, in association with the leading mode of variability in the NH winter geopotential field, the Northern Annular Mode (NAM) (Hartmann et al., 2000). During high NAM index periods, with a stronger stratospheric polar vortex, the anomalous $s=1$ component of the heat flux at 100 hPa was negative, and the anomalous $s=2$ component was positive. During low NAM index periods, both anomalies were of opposite sign. Thus, the negative correlation coefficient we found on the interannual timescale is also observed on the shorter timescales. The wave-1 contribution to H_{100} was found to depend on the latitude of the wave-1 source. Particularly, if the tropospheric stationary wave-1 source is located near 30° N instead of near 50° N, significantly less wave activity is able to enter the mid-latitude waveguide and contribute to H_{100} .

Finally, another approach was followed, where the wave driving anomalies were separated into three parts: one part due to anomalies in the zonal correlation coefficient between the eddy temperature and eddy meridional wind, another part due to anomalies in the zonal eddy temperature amplitude, and a third part due to anomalies in the zonal eddy meridional wind amplitude. It was found that year-to-year variability in the zonal correlation coefficient between the eddy temperature and the eddy meridional wind is the most dominant of the three factors.

In the interpretation of our results, the assumption has been that wave activity always propagates upward, so that the source of the wave activity at 100 hPa is situated below 100 hPa. For stationary waves, this is a reasonable assumption. However, transient waves can develop in the stratosphere as a result of purely stationary waves of sufficient amplitude emanating from the troposphere, as demonstrated by Christiansen (1999). Downward propagation of these stratospheric transients might affect H_{100} . However, we expect this to be only a minor influence, since the interannual variability of H_{100} is dominated by stationary waves (Table 1), and transient wave activity is abundantly generated in the troposphere.

In the present study, the 1958–1978 period was omitted from the analysis. However, we have also analyzed the entire 1958–2002 period. The results were very similar, although the statistical significance was generally larger due to the longer period. As a result, Fig. 3a exhibited a significant

correlation coefficient for $s=1$ in the entire free troposphere. A remarkable difference with the 1979–2002 analysis was found for Fig. 3b, in which the level of maximum and significant correlation coefficient with H_{100} was found to increase with increasing wavenumber $s=1-5$. The $s=1$ correlation coefficient decreased to statistically insignificant values of less than 0.3 in the upper stratosphere. In Fig. 4a, a clear equatorward displacement of a statistically significant correlation coefficient maximum was observed above 100 hPa, and in Fig. 4e, the tropospheric dipole structure for $s=1$ in Fig. 4e was more pronounced. Finally, for the 1958–2002 period, a significantly higher refractive index was observed in the mid-latitude stratosphere for a composite with positive H_{100} anomalies exceeding one standard deviation than for a composite of negative H_{100} anomalies exceeding one standard deviation. Such a significant signal could not be obtained for the 1979–2002 period, likely due to the smaller sample size.

One could argue that the results depend on the $40^\circ-80^\circ$ N latitude window that is applied. However, replacing H_{100} with the NH average of $[\overline{v^*T^*}]$ at 100 hPa yields almost identical results. The present study has focused on the interannual variability of the stratospheric wave driving. We note that the factors that dominate the interannual variability may be different from the factors that dominate the trend. These issues will be subject of our further investigation.

Acknowledgements. The authors gratefully acknowledge the constructive suggestions that were made by the editor and three anonymous referees.

Edited by: P. Haynes

References

- Andrews, D. G., Holton, J. R., and Leovy, C. B.: Middle atmosphere dynamics, Academic Press, 489 pp., 1987.
- Austin, J., Shindell, D., Beagley, S. R., Brühl, C., Dameris, M., Manzini, E., Nagashima, T., Newman, P., Pawson, S., Pitari, G., Rozanov, E., Schnadt, C., and Shepherd, T. G.: Uncertainties and assessments of chemistry-climate models of the stratosphere, *Atmos. Chem. Phys.*, 3, 1–27, 2003, <http://www.atmos-chem-phys.net/3/1/2003/>.
- Butchart, N., Scaife, A. A., Bourqui, M., de Grandpre, J., Hare, S. H. E., Kettleborough, J., Langematz, U., Manzini, E., Sassi, F., Shibata, K., Shindell, D., and Sigmond, M.: Simulations of anthropogenic change in the strength of the Brewer-Dobson circulation, *Clim. Dyn.*, 27, 727–741, doi:10.1007/s00382-006-0162-4, 2006.
- Charney, J. G. and Drazin, P. G.: Propagation of planetary-scale disturbances from the lower to the upper atmosphere, *J. Atmos. Sci.*, 18, 83–109, 1961.
- Christiansen, B.: Stratospheric vacillations in a General Circulation Model, *J. Atmos. Sci.*, 56, 1858–1872, 1999.
- Eyring, V., Harris, N. R. P., Rex, M., Shepherd, T. G., Fahey, D. W., Amanatidis, G. T., Austin, J., Chipperfield, M. P., Dameris, M., Forster, P. M. De F., Gettelman, A., Graf, H. F., Nagashima, T., Newman, P. A., Pawson, S., Prather, M. J., Pyle, J. A., Salawitch,

- R. J., Santer, B. D., and Waugh, D. W.: A strategy for process-oriented validation of coupled chemistry-climate models, *Bull. Am. Meteorol. Soc.*, 86, 1117–1133, 2005.
- Fusco, A. C. and Salby, M. L.: Interannual variations of total ozone and their relationship to variations of planetary wave activity, *J. Climate*, 12, 1619–1629, 1999.
- Hartmann, D. J., Wallace, J. M., Limpasuvan, V., Thompson, D. W. J., and Holton, J. R.: Can ozone depletion and global warming interact to produce rapid climate change?, *PNAS*, 97, 1412–1417, 2000.
- Haynes, P. H., Marks, C. J., McIntyre, M. E., Shepherd, T. G., and Shine, K. P.: On the “downward control” of extratropical diabatic circulations by eddy-induced mean zonal forces, *J. Atmos. Sci.*, 48, 651–678, 1991.
- Hu, Y. and Tung, K. K.: Possible ozone-induced long-term changes in planetary wave activity in late winter, *J. Climate*, 16, 3027–3038, 2003.
- Karoly, D. and Hoskins, B. J.: Three-dimensional propagation of planetary waves, *J. Meteorol. Soc. Japan*, 60, 109–123, 1982.
- Newman, P. A. and Nash, E. R.: Quantifying wave driving of the stratosphere, *J. Geophys. Res.*, 105, 12 485–12 497, 2000.
- Newman, P. A., Nash, E. R., and Rosenfield, J. E.: What controls the temperature of the Arctic stratosphere during the spring?, *J. Geophys. Res.*, 106(D17), 19 999–20 010, 2001.
- Polvani, L. M. and Waugh, D. W.: Upward wave activity flux as precursor to extreme stratospheric events and subsequent anomalous surface weather regimes, *J. Climate*, 17, 3548–3554, 2004.
- Randel, W. J., Wu, F., and Stolarski, R.: Changes in column ozone correlated with the stratospheric EP flux, *J. Meteorol. Soc. Japan*, 80, 849–862, 2002.
- Shepherd, T. G.: The middle atmosphere, *J. Atmos. Sol.-Terr. Phys.*, 62, 1587–1601, 2000.
- Siegmund P. C.: The generation of available potential energy: a comparison of results from a general circulation model with observations, *Clim. Dyn.*, 11, 129–140, 1995.
- Sigmond, M., Siegmund, P. C., Manzini, E., and Kelder, H.: A simulation of the separate climate effects of middle atmospheric and tropospheric CO₂ doubling, *J. Climate*, 17(12), 2352–2367, 2004.
- Simmons, A. J. and Gibson, J. K.: The ERA-40 project plan, ERA-40 Proj. Rep. Ser. 1, 63 pp., European Centre for Medium-Range Weather Forecasts, Reading, UK, 2000.
- Uppala, S. M., Kållberg, P. W., Simmons, A. J., Andrae, U., Da Costa Bechtold, V., Fiorino, M., Gibson, J. K., Haseler, J., Hernandez, A., Kelly, G. A., Li, X., Onogi, K., Saarinen, S., Sokka, N., Allan, R. P., Andersson, E., Arpe, K., Balmaseda, M. A., Beljaars, A. C. M., Van De Berg, L., Bidlot, J., Bormann, N., Caires, S., Chevallier, F., Dethof, A., Dragosavac, M., Fisher, M., Fuentes, M., Hagemann, S., Hólm, E., Hoskins, B. J., Isaksen, I., Janssen, P. A. E. M., Jenne, R., McNall, A. P. Y., Mahfouf, J.-F., Morcrette, J.-J., Rayner, N. A., Saunders, R. W., Simon, P., Sterl, A., Trenberth, K. E., Untch, A., Vasiljevic, D., Viterbo, P., and Woollen, J.: The ERA-40 re-analysis, *Quart. J. R. Meteorol. Soc.*, 131, 2961–3012, doi:10.1256/qj.04.176, 2005.