



# Journal of the Serbian Chemical Society

JSCS-info@shd.org.rs • www.shd.org.rs/JSCS

*J. Serb. Chem. Soc.* 78 (4) 477–482 (2013)  
JSCS-4431

UDC 542.9+547.571+547.551:542.913:547.673

*Short communication*

## SHORT COMMUNICATION

## An efficient synthesis and spectroscopic characterization of Schiff bases containing the 9,10-anthracenedione moiety

GHULAM FAREED<sup>1,2\*</sup>, MUHAMMAD ALI VERSIANI<sup>2</sup>, NIGHAT AFZA<sup>1</sup>,  
NAZIA FAREED<sup>2</sup>, MUHAMMAD IRFAN ALI<sup>1</sup> and MAHBOOB ALI KALHORO<sup>1</sup>

<sup>1</sup>Pharmaceutical Research Center, PCSIR Laboratories Complex Karachi, Shahrah-e-Dr. Salim-u-Zaman Siddiqui, Karachi-75280, Sindh, Pakistan and <sup>2</sup>Department of Chemistry, Federal Urdu University of Arts, Science and Technology, Ghulshan-e-Iqbal, Karachi-75300, Sindh, Pakistan

(Received 9 April, revised 28 August 2012)

**Abstract:** A new method was developed for the synthesis of novel Schiff bases containing the anthraquinone moiety using dodecatungstosilicic acid/P<sub>2</sub>O<sub>5</sub> under solvent-free conditions at room temperature. The reaction was completed in 1–3 minutes with excellent yields. This method was found to be very efficient, easy and hazardous-free for the synthesis of azomethines. The development of these types of methodologies in synthetic chemistry may contribute to green chemistry. The structures of synthesized novel Schiff bases were elucidated using <sup>1</sup>H-NMR, <sup>13</sup>C-NMR, LC-MS, FTIR and CHN analysis.

**Keywords:** azomethine; dodecatungstosilicic acid; P<sub>2</sub>O<sub>5</sub>; anthraquinone.

## INTRODUCTION

Schiff bases, a versatile class of organic compounds synthesized by the condensation of primary amines with aromatic carbonyl compounds, bear many pharmaceutical properties, including anticancer, antitumor, antibacterial, antifungal, antioxidant, herbicidal, antiproliferative, antimarial and anticovulsant activities.<sup>1–16</sup> Schiff bases act as ligands in coordination chemistry<sup>17</sup> and are considered a subject of interest due to their industrial and biological applications.<sup>18</sup>

Similarly 9,10-anthracenediones are a very important class of compounds having some remarkable *in vivo* biological activities, such as antibacterial, antifungal, antimarial, antioxidant, antileukemic, hypotensive, mutagenic functions, and analgesic activity.<sup>19–27</sup>

Moreover, in the recent era, the use environmentally benign, facile and low costing method for the synthesis of biologically active compounds has received

\*Corresponding author. E-mail: fareedchm@yahoo.com  
doi: 10.2298/JSC120409092F



great attention of synthetic chemists. Based on green chemistry and facile methodologies, numerous methods have been used for the synthesis of Schiff bases, *e.g.*, catalyzed by  $TiCl_2$ ,<sup>28</sup>  $ZnCl_2$ ,<sup>29</sup>  $Mg(ClO_4)_2$ ,<sup>30</sup>  $P_2O_5/SiO_2$ ,<sup>31</sup>  $Ni(NO_3)_2 \cdot 6H_2O$ ,  $Cu(NO_3)_2 \cdot 6H_2O$ ,  $Fe(NO_3)_2 \cdot 6H_2O$ ,  $Mn(NO_3)_2 \cdot 6H_2O$  and  $Co(NO_3)_2 \cdot 6H_2O$ ,<sup>32</sup> montmorillonite K-10<sup>33</sup> and  $MgSO_4$ -pyridinium *p*-toluenesulfonate (PPTL).<sup>34</sup>


In view of the above facts, results concerning the syntheses of novel Schiff bases using dodecatungstosilicic acid/ $P_2O_5$  as a catalyst are reported herein. The employed method was found to be very efficient and environmentally friendly.

#### EXPERIMENTAL

Melting points were measured in open glass capillaries using a Gallenkamp MF-370 melting point apparatus. The IR spectra were recorded on a Nicolet Avatar 300 DTGS instrument. The  $^1H$ - and  $^{13}C$ -NMR were recorded on Bruker A V-300 instrument operating at 300 and 75 MHz, respectively, using deuterated dimethyl sulfoxide ( $DMSO-d_6$ ) as the solvent and trimethylsilane (TMS) as an internal standard. The mass spectra were recorded on a Finnigan LCQ Advantage Max instrument. The CHN analyses were realized on a Carlo Erba instrument – Mod-1106. The purity of the products were determined by TLC using silica gel 60F<sub>254</sub> precoated cards (0.2 mm thickness) and visualized under UV light (254 and 366 nm) and by iodine vapors. All chemicals were of AR grade. Analytical and spectral data for the synthesized compounds are given in the Supplementary material.

#### *General method for the synthesis of Schiff bases 1–25*

A mixture of 1-aminoanthraquinone (1 mmol), a substituted aromatic aldehyde or ketone (1 mmol) and dodecatungstosilicic acid/ $P_2O_5$  (0.2 g, 1 mol % of 1-aminoanthraquinone/ $P_2O_5$ ) as a catalyst was ground in a mortar with a pestle under solvent-free conditions at room temperature for 1–3 min, Scheme 1. The reaction mixture turned to a pasty material that



Scheme 1. Preparation of Schiff bases.

indicated the completion of the reaction. Crushed ice was added to afford precipitates of the Schiff bases. In order to remove the catalyst, the product was washed several times with ice-cold water. The solid products were obtained in excellent yield.

#### RESULTS AND DISCUSSION


The aim of this work was to develop a new facile method for the preparation of Schiff bases. To the best of our knowledge and a literature search, the synthesis of novel Schiff bases using dodecatungstosilicic acid/P<sub>2</sub>O<sub>5</sub> has not been previously reported.

An efficient methodology was established using an acidic catalyst for the synthesis of novel Schiff bases by the condensation of 1-aminoanthraquinone with aromatic carbonyl compounds under solvent free conditions at room temperature. The reactions were completed in 1–3 minutes and the products were obtained in high yield after simple work-up (Table I). In order to develop optimum conditions for this synthesis, varying amounts of dodecatungstosilicic acid/P<sub>2</sub>O<sub>5</sub> were used using 4-hydroxybenzaldehyde and 1-aminoanthraquinone under solvent-free conditions at room temperature as a model reaction (Table II). The method is very simple and contributes to the hazardous-free synthesis. The catalyst dodecatungstosilicic acid/P<sub>2</sub>O<sub>5</sub> was found to be very efficient compared to other catalysts used for the synthesis of imines (Table III).

TABLE I. Synthesis of Schiff bases using *dodeca*-tungstosilicic acid/P<sub>2</sub>O<sub>5</sub> as a catalyst

Entry	Substrate	t / min	Entry	Substrate	t / min
1	3-Hydroxybenzaldehyde	2	14	2,4-Dihydroxybenzaldehyde	2
2	4-Hydroxybenzaldehyde	1	15	4-(Diethylamino)benzaldehyde	3
3	3-Chlorobenzaldehyde	2	16	α-Bromocinnamaldehyde	3
4	Furfural	1	17	3,5-Dibromo-4-hydroxybenzaldehyde	2
5	4-Hydroxy-3-methoxybenzaldehyde	2	18	4-(Dimethylamino)cinnamaldehyde	3
6	3-Bromo-4-hydroxybenzaldehyde	2	19	2-Ethoxybenzaldehyde	3
7	4-Nitrobenzaldehyde	2	20	4-(Dimethylamino)benzaldehyde	2
8	3-Bromobenzaldehyde	1	21	2-Hydroxyacetophenone	1
9	3-Nitrobenzaldehyde	3	22	3,5-Dimethoxybenzaldehyde	2
10	2-Nitrobenzaldehyde	3	23	4-Methoxycinnamaldehyde	1
11	4-Methoxybenzaldehyde	1	24	4-Ethoxybenzaldehyde	1
12	4-Chlorobenzaldehyde	2	25	4-Methylbenzaldehyde	2
13	4-Chloro-3-nitrobenzaldehyde	1			

Moreover, the structures of the products were elucidated by LC-MS, <sup>1</sup>H-NMR, <sup>13</sup>C-NMR, IR, and CHN analysis. The <sup>1</sup>H-NMR spectra of all the compounds was quite simple and the azomethine group (–N=CH–Ar) of all the synthesized Schiff bases was found in the region of 7.4–8.9 ppm. The aromatic protons ap-



peared as a multiplet in the region of 6.74–7.90 ppm depending on the aromatic substituent. In IR spectrum, –N=C– appears in the region 1511–1545 cm<sup>-1</sup>.

TABLE II. Effect of different amounts of catalyst on the yield of 1-{[(4-hydroxyphenyl)-methylidene]amino}-9,10-anthraquinone

Entry	Catalyst, mol %	Reaction time, min	Yield, %
1	Free	20	50
2	0.5	10	70
3	1	1	87
4	1.5	1	87

TABLE III. Comparison with other catalysts used for the synthesis of Schiff base

Entry	Catalyst	Solvent	Reaction Time	Yield, %	Ref.
1	Mg(ClO <sub>4</sub> ) <sub>2</sub>	1,2-Dichloroethane	8 h	80	30
2	SiO <sub>2</sub> /P <sub>2</sub> O <sub>5</sub>	–	13–26 min	88	31
3	Ni(NO <sub>3</sub> ) <sub>2</sub> ·6H <sub>2</sub> O	MeOH/EtOH/DMF	20–30 min	60–88	32
4 <sup>a</sup>	Montmorillonite K-10	–	1–3 min	80	33
5	ZnCl <sub>2</sub>	1,2-Dichloroethane	8 h	46	30
6	CuSO <sub>4</sub> ·H <sub>2</sub> O	1,2-Dichloroethane	8 h	65	30
7	TiCl <sub>2</sub>	1,2-Dichloroethane	8 h	35	30

<sup>a</sup>Under microwave conditions

#### CONCLUSION

Based of the excellent yields, short reaction time, easy work-up, solvent-free facile and environment greener reaction, it could be concluded that dodecatungstosilic acid/P<sub>2</sub>O<sub>5</sub> is a very useful catalyst for the synthesis of Schiff bases.

#### SUPPLEMENTARY MATERIAL

Analytical and spectral data for the synthesized compounds are available electronically from <http://www.shd.org.rs/JSCS/>, or from the corresponding author on request.

*Acknowledgement:* All authors are grateful to the Pharmaceutical Research Center, PCSIR Laboratories Complex, Karachi, Pakistan, for the financial and technical support of this research work.


#### ИЗВОД

#### ЕФИКАСНА СИНТЕЗА И СПЕКТРОСКОПСКА КАРАКТЕРИЗАЦИЈА ШИФОВИХ БАЗА КОЈЕ САДРЖЕ 9,10-АНТРАЦЕНДИОНСКИ СТРУКТУРНИ ФРАГМЕНТ

GHULAM FAREED<sup>1,2</sup>, MUHAMMAD ALI VERSIANI<sup>2</sup>, NIGHAT AFZA<sup>1</sup>, NAZIA FAREED<sup>2</sup>, MUHAMMAD IRFAN ALI<sup>1</sup>  
и MAHBOOB ALI KALHORO<sup>1</sup>

<sup>1</sup>Pharmaceutical Research Center, PCSIR Laboratories Complex Karachi, Shahrah-e-Dr.Salim-uz-Zaman Siddiqui, Karachi-75280, Sindh, Pakistan и <sup>2</sup>Department of Chemistry, Federal Urdu University of Arts, Science and Technology, Ghulshan-e-Iqbal, Karachi-75300, Sindh, Pakistan

Развијен је нов поступак за синтезу Шифових база које садржи антрахинонски фрагмент, употребом додекаволфрамсилицијумове киселине/P<sub>2</sub>O<sub>5</sub>, у одсуству органских растворача, на собној температури. Реакција је готова у периоду 1–3 min у одличном



приносу. Утврђено је да је поступак ефикаснији, једноставнији и безбеднији. Структуре синтетисаних Шифових база одређене су  $^1\text{H-NMR}$ ,  $^{13}\text{C-NMR}$ , LC-MS, FT-IR и CHN анализама.

(Примљено 9. априла, ревидирано 28. августа 2012)

#### REFERENCES

1. A. Faizul, S. Satendra, L. K. Sukhbir, P. Om, *J. Zhejiang Univ. Sci., B* **8** (2007) 446
2. P. G. More, R. B. Bhalvankar, S. C. Pattar, *J. Indian Chem. Soc.* **78** (2001) 474
3. A. H. El-Masry, H. H. Fahmy, S. H. A. Abdelwahed, *Molecules* **5** (2000) 1429
4. M. A. Baseer, V. D. Jadhav, R. M. Phule, Y. V. Archana, Y. B. Vibhute, *Orient. J. Chem.* **16** (2000) 553
5. S. N. Pandeya, D. Sriram, G. Nath, E. De Clercq, *Farmaco* **54** (1999) 624
6. W. M. Singh, B. C. Dash, *Pesticides* **22** (1988) 33
7. E. M. Hodnett, W. J. Dunn, *J. Med. Chem.* **13** (1970) 768
8. S. B. Desai, P. B. Desai, K. R. Desai, *Heterocycl. Commun.* **7** (2001) 83
9. S. Samadhiya, A. Halve, *Orient. J. Chem.* **17** (2001) 119
10. W. P. Narocka, B. Sztuba, A. Drys, J. Wietrzyk, J. Ietrzyk, J. Kosendiak, A. Opolski, *Pol. J. Chem.* **80** (2006) 279
11. J. P. Kaplan, B. M. Rizon, *J. Med. Chem.* **23** (1980) 702
12. M. Verma, S. N. Pandeya, K. N. Singh, J. P. Stables, *Acta Pharm.* **54** (2004) 49
13. B. Rada, T. L. Leto, *Contrib. Microbiol.* **15** (2008) 164
14. A. Almasirad, R. Hosseini, H. Jalalizadeh, Z. Rahimi-Moghaddam, N. Abeian, M. Jana-frooz, M. Abbaspour, V. Ziaeef, A. Dalvandi, A. Shafiee, *Biol. Pharm. Bull.* **29** (2006) 1180
15. C. J. Clarke, J. W. Eaton, *Clin. Res.* **38** (1990) 300A
16. A. Tsafack, M. Loyevski, P. Ponks, Z. I. Cabantchik, *J. Lab. Clin. Med.* **127** (1996) 575
17. J. C. Pessoa, J. Cavaco, J. Covreia, D. Costa, R. T. Henriques, R. D. Gillard, *Inorg. Chim. Acta* **305** (2000) 7
18. Z. M. Zaki, S. S. Haggag, A. A. Soayed, *Spectrosc. Lett.* **31** (1998) 757
19. N. W. Alcock, D. F. Cook, E. D. McKenzie, J. M. Worthington, *Inorg. Chem. Acta* **38** (1980) 107
20. M. K. Adwankar, M. P. Chitnis, *Chemotherapy* **28** (1982) 291
21. A. A. Sittie, E. Lemmich, C. E. Olsen, L. Hviid, A. Kharazmi, F. K. Nkrumah, S. B. Christensen, *Planta Med.* **65** (1999) 259
22. G. Bath, M. Ndonzao, K. Hostettmann, *Int. J. Pharmacog.* **33** (1995) 107
23. C. Younos, A. Rolland, M. Fleurentin, M.-C. Lanher, R. Misslin, F. Mortier, *Planta Med.* **56** (1990) 430
24. K. Koumaglo, M. Gbeassor, O. Nikabu, C. D. Souza, W. Werner, *Planta Med.* **58** (1992) 533
25. Y. B. Tripathi, M. Sharma, M. Manickam, *Indian J. Biochem. Biophys.* **34** (1997) 302
26. P. Chang, C. Chen, *Chin. Pharm. J. (Taipei)* **47** (1995) 347
27. N. H. Ismail, A. M. Ali, N. Aimi, M. Kitajima, H. Takayama, N. H. Lajja, *Phytochemistry* **45** (1997) 1723
28. H. Weingart, J. P. Chupp, W. A. White, *J. Org. Chem.* **32** (1967) 213
29. J. H. Billman, K. M. Tai, *J. Org. Chem.* **23** (1958) 535
30. A. K. Chakraborti, S. Bhagat, S. Rudrawar, *Tetrahedron Lett.* **45** (2004) 7641



31. S. M. Devidas, S. H. Quadri, S. A. Kamble, F. M. Syed, D. Y. Vyavhare, *J. Chem. Pharm. Res.* **3** (2011) 489
32. A. Mobinikhaledi, N. Fourughifar, M. Kalhor, *Turk. J. Chem.* **34** (2010) 367
33. R. S. Vass, J. Dudas, R. S. Varma, *Tetrahedron Lett.* **40** (1999) 4951
34. B. P. Branchaud, *J. Org. Chem.* **48** (1983) 3531.

