

*Maejo Int. J. Sci. Technol.* 2010, 4(03), 483-495

## *Maejo International Journal of Science and Technology*

ISSN 1905-7873

Available online at [www.mijst.mju.ac.th](http://www.mijst.mju.ac.th)

*Full Paper*

### **Urban ecological characteristics and vascular wall flora on the Anatolian side of Istanbul, Turkey**

**Volkan Altay<sup>1\*</sup>, İbrahim İ. Özyiğit<sup>2</sup> and Celal Yarci<sup>2</sup>**

<sup>1</sup> Mustafa Kemal University, Science and Arts Faculty, Department of Biology, 31040, Antakya, Hatay, Turkey

<sup>2</sup> Marmara University, Science and Arts Faculty, Department of Biology, 34722, Göztepe, Istanbul, Turkey

\*Corresponding author, E-mail: [valtay@mku.edu.tr](mailto:valtay@mku.edu.tr) ; Tel: +90 326 245 58 66; Fax: + 90 326 245 58 67

*Received: 5 May 2010 / Accepted: 16 November 2010 / Published: 27 November 2010*

---

**Abstract:** The aim of this study is to specify the urban ecologic characteristics of Istanbul and to show their reflection to the vascular wall flora of the Anatolian side, which is a distinctive wall habitat. Plants samples of the urban habitat were collected from the top and vertical surfaces of walls during 2005-2007. A total of 101 taxa (81 species, 13 subspecies and 7 varieties) belonging to 74 genera and 33 families were recorded. It was determined that 80 species were Dicotyledones while 1 was Monocotyledone. The families with the largest number of taxa were Asteraceae (18 species, 22.22%), Poaceae (8 species, 9.87%), Lamiaceae and Brassicaceae (5 species, 6.17%), and Polygonaceae and Scrophulariaceae (4 species, 4.93%). The most common plant species on walls were *Parietaria judaica* L. (Urticaceae), *Stellaria media* (L.) Vill. subsp. *media* (Caryophyllaceae), and *Mercurialis annua* L. (Euphorbiaceae). The percentage of phytogeographical elements among the recorded taxa varied as follows: Euro-Siberian (6 taxa, 7.41%), Mediterranean (11 taxa, 13.58%), E. Mediterranean (2 taxa, 2.47%), Irano-Turanian (1 taxon, 1.23%) and unknown (61 taxa, 75.31%). It was found that 6 taxa (7.41%) were cosmopolitan, 12 (14.82%) were widespread while 1 (1.23%) was endemic. The results were compared with some other European wall floras and some similarities and dissimilarities were noted.

**Keywords:** vascular wall flora, Istanbul, urban habitat, urban ecology

---

#### **INTRODUCTION**

Istanbul is one of the most populous cities of Eurasia and is the world's 4<sup>th</sup> largest city proper and 20<sup>th</sup> largest urban area as well as Turkey's cultural and financial centre. The city is located in the

NW part of Turkey ( $41^{\circ} 01.2' N$ ,  $28^{\circ} 58.2' E$ ) and it extends both on European (Thrace) and Asian (Anatolia) sides of the Bosphorus. It is the only metropolis in the world which is situated on two continents [1-5]. Its neighbours are the Black Sea in the north, Marmara Sea in the south, Kocaeli City in the east, and Tekirdağ City in the west (Figure 1). Istanbul has approximately  $5100 \text{ km}^2$  of land area and 12,573,836 population. In its long history, Istanbul served as a capital city of the Roman Empire (AD 330-395), Byzantine Empire (AD 395-1204 and AD1261-1453), Latin Empire (AD 1204-1261), and Ottoman Empire (AD 1453-1922) [3, 6]. In addition to its rich history, high population and productive economy, Istanbul also has a wide variety of ecological features [3].



**Figure 1.** Districts studied in Istanbul during 2005-2007 (Beykoz, Üsküdar, Kadıköy, Kartal, Ümraniye, Maltepe, Tuzla, Pendik, Adalar and Sultanbeyli)

The main topographical feature of Istanbul is a low plateau at about 100-200 m elevation where there are many hills and a few streams in the city. The geological structure of Istanbul is diverse with Silurian, Devonian, Carboniferous, and Tertiary formations. There are different kinds of rocks including granitic plutons, quartzes, grovacs, clayed schists and radiolarites [2, 7].

In Istanbul, many different types of soils are present while brown forest soil covers the most area. Non-calcareous brown soil is the second and is kind of soil is suitable for plant growth as a result of organic matter. The rendzinas, which cover especially the European side of the city, are also important. In addition, alluvial soil is also present [2, 7].

Istanbul is in a kind of transition zone between less rainy Mediterranean and oceanic climates. In the summer, less precipitation and high temperature are characteristic and the annual mean temperature is  $14.5^{\circ}\text{C}$  for the last two decades. Between May and September the temperature is

generally above 30°C and between November and April it is rarely below 0°C. In the vegetation period, the daily mean temperature is approximately 8°C, which lasts for about 280 days (between 15 March and 20 December) [2-5, 8].

The total precipitation for Istanbul averages 640 mm per year and 40% of the total precipitation falls in winter. December and January are the wettest months. Precipitation in summer is more than the typical Mediterranean situation and is more related to the oceanic climate. The minimal amount of rain falls (about 8%) in July and August. Precipitation is less in spring (about 20-21%) while it increases in autumn (about 28-29%), and snow rarely falls in Istanbul. The rain regime is Winter-Autumn-Spring-Summer (W.A.Sp.Su) and the rain type is “Central Mediterranean Rain Type” [2, 9].

The relative humidity is between 73-77% in the city and these values decrease to 65-68% in the summer despite the effect of the seas. The lower relative humidity, especially in the dry period, forms xerophytic vegetation. The dominant wind in the city is from the NE [8-9].

The wide variety of ecological features in Istanbul have resulted in more Pteridophyta and Angiospermae (2500 species) than those in England (250,000 km<sup>2</sup> land area, 1850 plant species) and Netherlands (50,000 km<sup>2</sup> land area, 1600 species) [10]. Istanbul’s rich flora has been studied by many botanists due to its varied climatic and geographic conditions. Many floristic studies in urban and rural areas have been done in Turkey [7, 11-13]. Aksoy [14-15] carried out some studies on the green areas in Istanbul while others have done urban ecological studies, i.e. Şahin [16] in Eminönü and Fatih, Altay et al. [3] in Kartal, Osma et al. [4] in Kadıköy, Mutlu [17] in Üsküdar, and Eskin [18] in Pendik districts.

Buildings and all types of walls are urban features represent a specific environment [19-20]. The colonisation of plants on walls is favoured by the wall ages, the presence of lime mortar, exposure to rain, and such aspects as south and verticality. Most true wall species are only found on vertical walls and as the angle of inclination decreases an ever-widening range of common species colonise [21]. The wall habitat is different from natural habitat and rocks, depending on many different properties related to wall structures. Buildings contain binding materials, which structurally and chemically differ from the original building materials. Usually most of them are cleaned repeatedly; thus, they basically are temporary habitat. Walls are generally isolated, small, and their microclimate is more affected by changes of climatic factors such as temperature, precipitation and irradiation than that of rocks. Wall surfaces resemble each other and have a uniform slope (generally right-angled) and microtopographical characteristic with few microhabitats. Walls are located both in urban and rural areas, so wall flora is influenced by the nearby ruderal and semi-natural vegetation [22].

Wall flora which shows differences from urban and rural flora has attracted many botanists in different countries and cities. Comparison of wall vegetation in southern, western and central Europe was done by Segal [23] and Brandes [24-25]. Carmona et al.[26] in Spain, Hruska [27] in Italy, Gehu [28] in France, Woodell [19] in England, Oberdorfer [29] in Germany, Weretelnik [30] in Poland, Mucina [31] in Austria, Valachovic and Maglocky [32] in Slovakia, Duchoslav [22] in the Czech Republic, and Pavlova and Tonkov [20] in Bulgaria have studied wall flora.

Wall flora studies in Turkey are limited and include those by Gemici et al. [33] in İzmir, Aksoy and Çelik [34] in Kayseri, Yeşilot [35] in Istanbul European side, and Yarcı and Özçelik [36] in Edirne. These studies were mostly on walls of historical buildings.

Our study was carried out in all districts of the Anatolian side of Istanbul, excluding Şile, during 2005-2007 vegetation periods. This study was made on walls of historical buildings as well as those of gardens, houses, hospitals, schools, campuses, train stations and ateliers, and especially on drain walls.

## MATERIALS AND METHODS

The wall vascular flora on the Anatolian side of Istanbul was studied. Bryophyta and ornamental plants were not collected. Plant specimens were identified by using the “Flora of Turkey and the East Aegean Islands Volumes I-XI and Supp.” [37] and were deposited in MÜFE Herbarium (Marmara University, Sciences and Arts Faculty Herbarium). The flora is listed alphabetically in the Appendix according to family, genera and species. Life forms [phanerophytes (Ph); chaemaphytes (Ch); hemicryptophytes (H); therophytes (Th); geophytes (G)] and phytogeographical origins [Euro-Siberian (Euro.-Sib.), Irano-Turanian (Ir.-Tur.), Mediterranean (Medit.), East Mediterranean (E. Medit.)] are included and were determined according to the Raunkier system [38-39].

## RESULTS AND DISCUSSION

The total number of vascular plants growing spontaneously on walls included 101 taxa with 33 families and 74 genera (81 species, 13 subspecies and 7 varieties). Eighty species were Dicotyledones with one monocotyledon species (Table 1).

**Table 1.** Summary of collecting

Taxon	Monocotyledones		Dicotyledones		Total Number
	Number	%	Number	%	
Family	1	3.04	32	96.96	33
Genus	1	1.36	73	98.64	74
Species	1	1.24	80	98.76	81

There was only one monocotyledon species, *Asparagus acutifolius* L. (Liliaceae), which consisted of 1.24% of all species, while Dicotyledones were the prevalent group with 98.76% of all species. The following families were represented by the largest number of species: Asteraceae (18 species, 22.22%), Poaceae (8 species, 9.87%), Lamiaceae (5 species, 6.17%), Brassicaceae (5 species, 6.17%), Polygonaceae (4 species, 4.93%) and Scrophulariaceae (4 species, 4.93%) (Table 2). Although Asteraceae was the most common family, it had the lowest species/genus ratio (1.0). The same ratio was also for Brassicaceae, although it was a low-taxa family in this study. This ratio was higher in Polygonaceae (2.0), Scrophulariaceae (1.33), Lamiaceae (1.25) and Poaceae (1.14).


Previous studies noted that Asteraceae species have a species richness compared with other families [3-4, 20, 22, 24, 40-44]. Pavlova and Tonkov [20] also observed that Asteraceae was high in number of species in Central Europe, demonstrating the remarkable success of this family in terms of dispersal and establishment. Davis [37] also mentioned that Asteraceae represents highest number of

species in the flora of Turkey. Other researchers obtained similar results with other families [20, 35-36].

**Table 2.** Genera and species numbers of the richest families

Family	Genera		Species		Species/Genus ratio
	Number	%	Number	%	
Asteraceae	18	24.32	18	22.22	1
Poaceae	7	9.45	8	9.87	1.14
Lamiaceae	4	5.40	5	6.17	1.25
Brassicaceae	5	6.75	5	6.17	1
Polygonaceae	2	2.70	4	4.93	2
Scrophulariaceae	3	4.05	4	4.93	1.33


In this study, life forms were determined according to the Raunkier system [38] and the largest groups were therophytes (45.72%) and hemicryptophytes (37.00%) (Figure 2). The patterns of these two life forms found on walls were similar to walls in Europe [18, 20-22]. In our study, percentages of other life forms were as follows: chamaephytes (2.46%), geophytes (4.94%) and phanerophytes (9.88%). Therophytes and hemicryptophytes are widespread in areas with a Mediterranean climate [3-4].



**Figure 2.** Pie diagram of the prevalence of life forms (H. = hemicryptophytes; Ph. = phanerophytes; G. = geophytes; Ch. = chamaephytes; Th. = therophytes)

The most common phytogeographical elements were Mediterranean (13.58%) and Euro-Siberian elements (7.41%) (Figure 3). This is because Istanbul has a mostly Mediterranean climate.

The northern side of Istanbul is partly affected by the oceanic climate [14]. Our study showed that six taxa (7.41%) were cosmopolitan and 12 taxa (14.82%) were widespread. Only one endemic taxon was found in our study. The most common plant species on walls were: *Parietaria judaica* L. (Urticaceae), *Stellaria media* (L.) Vill. subsp. *media* (Caryophyllaceae) and *Mercurialis annua* L. (Euphorbiaceae). We did not find any pteridophytes although in Central Europe and in the Mediterranean area they are present.



**Figure 3.** Pie diagram of the phytogeographical origins of the wall flora (Medit. El. = Mediterranean elements, Euro.-Sib. El. = Euro-Siberian elements, E. Medit. El. = East Mediterranean elements, Ir.-Tur. El. = Irano-Turanian elements)

We also found some archaeophyte and neophyte plants. An archaeophyte is not native to the geographical region, but was introduced in "ancient" times rather than during modern times. A neophyte has been recently introduced to the area. Archaeophytes are considered to be those species first introduced prior to 1500 AD while neophytes came after 1500 AD [45-46]. Archaeophyte and neophyte plants are listed in Table 3.

Plants growing on walls reach these habitats by wind (anemochorous), animals (zoochorous), mostly by birds and by stolon fragments (autochorous) [22, 36], and grow there randomly. Deep-rooted plants can be destructive. Although their roots are weak at the beginning of growth, they become stronger in time and cause widening of cracks. Most of these plants absorb little water from the substrate, but absorb it from the air [36]. Although wall plants are often aesthetically appealing, the local municipalities occasionally clean up the walls to prevent damage by the plants. It would be more preferable if the clean-up was more selective by allowing for plant type and degree of damage.

**Table 3.** Archaeophyte and neophyte plants

Archaeophytes (before 1500 AD)	Neophytes (after 1500 AD)
<i>Lamium purpureum</i> L. (Lamiaceae)	<i>Conyza canadensis</i> (L.) Cronquist (Asteraceae)
<i>Lamium amplexicaule</i> L.	<i>Veronica persica</i> Poiret (Scrophulariaceae)
<i>Euphorbia helioscopia</i> L. (Euphorbiaceae)	<i>Cymbalaria muralis</i> Gaertner
<i>Euphorbia peplus</i> L.	<i>Ailanthus altissima</i> (Miller) Swingle (Simaroubaceae)
<i>Sinapis arvensis</i> L. (Brassicaceae)	<i>Oxalis corniculata</i> L. (Oxalidaceae)
<i>Capsella bursapastoris</i> (L.) Medik.	<i>Diploaxis tenuifolia</i> (L.) DC. (Brassicaceae)
<i>Fumaria officinalis</i> L. (Papaveraceae)	
<i>Stellaria media</i> (L.) Vill. (Caryophyllaceae)	
<i>Cerastium glomeratum</i> Thuill	
<i>Cichorium intybus</i> L. (Asteraceae)	
<i>Solanum nigrum</i> L. (Solanaceae)	
<i>Chenopodium album</i> L. (Chenopodiaceae)	
<i>Setaria viridis</i> (L.) P. Beauv. (Poaceae)	

### ACKNOWLEDGEMENTS

We wish to thank Miss Busecan Aksoydan for rearranging the Tables and Figures and Mr. Ahmet Yılmaz for rechecking the language in this document.

### REFERENCES

1. The official website of Istanbul metropolis municipality, <http://www.ibb.gov.tr>, access date: March 2010.
2. Ü. Yaşar İ. İ. Özyiğit and M. Serin “Judas tree (*Cercis siliquastrum* L. subsp. *siliquastrum*) as a possible biomonitor for Cr, Fe and Ni in Istanbul (Turkey)” *Rom. Biotech. Lett.*, 2010, 15, 4979-4989.
3. V. Altay, İ. İ. Özyiğit and C. Yarcı “Urban flora and ecological characteristics of the Kartal District (Istanbul): A contribution to urban ecology in Turkey” *Sci. Res. Essays*, 2010, 5, 183-200.
4. E. Osma, İ. İ. Özyiğit, V. Altay and M. Serin “Urban vascular flora and ecological characteristics of Kadıköy district, Istanbul, Turkey” *Maejo Int. J. Sci. Technol.*, 2010, 4, 64-87.
5. S. Kaya and P. J. Curran, “Monitoring urban growth on the European side of the Istanbul Metropolitan area: A case study”, *Int. J. Appl. Earth Obs. Geoinf.*, 2006, 8, 18-25.
6. M. O. Bayrak, “İstanbul Tarihi (History of Istanbul)”, İnkılap Kitabevi, Istanbul, 1996.
7. F. Yaltrık, A. Efe and A. Uzun, “Tarih boyunca İstanbul’un park, bahçe ve koruları egzotik ağaç ve calıları (The inventory of exotic tree and bush species of Istanbul’s parks, gardens and groves)”, No. 4, İstanbul Metropolis Municipality, İsfalt Yayınları, Istanbul, 1997.

8. “The Meteorological Report for 2007 Year of Istanbul”, Prime Ministership, General Directorship for Meteorology, the Directorship for Statistics and Publication, Ankara, **2007**.
9. Y. Akman, “İklim ve Biyoiklim (Climate and Bioclimate)”, Palme Yayın-Dağıtım, Ankara, **1990**.
10. N. Özhatay and M. Keskin, “Ömerli Havzasının İstanbul Doğal Bitkileri (Natural Plants of Ömerli Reservoir, Istanbul)”, Doğal Hayatı Koruma Vakfı Yayınlarından, İstanbul, **2007**.
11. A. Baytop, “Aydos ve Kayışdağ havalisinin çiçekli bitkileri (The flowering plants of Aydos and Kayışdağ environs)”, *Türk Biol. Derg.*, **1962**, 12, 75-111.
12. A. Baytop, “İstanbul Yıldız Parkı’nın florası (The flora of Yıldız Park, Istanbul)”, *İst. Üniv. Ecz. Fak. Mec.*, **1985**, 21, 86-97.
13. N. Özhatay, A. Byfield and S. Atay, “Türkiye’nin Önemli Bitki Alanları (The Important Plant Areas of Turkey)”, Doğal Hayatı Koruma Vakfı Yayınlarından, İstanbul, **2003**, p. 87.
14. Y. Aksoy, “İstanbul’da yeşil alan kullanımı üzerine bir araştırma (A study on the use of green areas in Istanbul)”, *MSc Thesis*, **1994**, Istanbul Technical University, Turkey.
15. Y. Aksoy, “İstanbul kenti yeşil alan durumu irdelenmesi (The evaluation of green areas position in Istanbul)”, *PhD Thesis*, **2001**, Istanbul Technical University, Turkey.
16. N. Şahin, “Eminönü ve Fatih İlçeleri’nin (İstanbul) kentsel ekolojik özellikleri (Urban ecological characteristics of Eminönü and Fatih Towns - Istanbul)”, *MSc Thesis*, **2002**, Marmara University, Turkey.
17. P. Mutlu, “Üsküdar İlçesi’nin (İstanbul) kentsel ekolojik özellikleri (Urban ecological characteristics of Üsküdar Town - Istanbul)”, *MSc Thesis*, **2004**, Marmara University, Turkey.
18. B. Eskin, “Pendik İlçesi (İstanbul) kentsel ekolojisi (Urban ecology of Pendik Town - Istanbul)”, *MSc Thesis*, **2005**, Marmara University, Turkey.
19. S. Woodell, in “Nature in Cities: The Natural Environment in the Design and Development of Urban Green Space” (Ed. I. C. Laurie), John Wiley, Chichester, **1979**, pp. 135-157.
20. D. Pavlova and S. Tonkov, “The wall flora of the Nepet Tepe Architectural Reserve in the city of Plovdiv (Bulgaria)”, *Acta Bot. Croat.*, **2005**, 64, 357-368.
21. O. L. Gilbert, “The Ecology of Urban Habitats”, Chapman and Hall, Cambridge, **1991**, p. 251.
22. M. Duchoslav, “Flora and vegetation of stony walls in East Bohemica (Czech Republic)”, *Preslia, Praha*, **2002**, 74, 1-25.
23. S. Segal, “Ecological Notes on Wall Vegetation”, Junk, The Hague, **1969**.
24. D. Brandes, “Flora und Vegetation von Stadtmauern”, *Tuexenia*, **1992**, 12, 315-319.
25. D. Brandes, “*Asplenietea*- Gesellschaften an sekundären Standorten in Mitteleuropa”, *Ber. d. Reinh.-Tüxen-Ges.*, **1992**, 4, 73-93.
26. E. C. Carmona, M. M. Luque and F. V. Tendero, “The plant communities of the *Asplenietea trichomanis* in the SW Iberian Peninsula”, *Folia Geobot. Phytotax.*, **1997**, 32, 361-376.
27. K. Hruska, “Syntaxonomical study of Italian wall vegetation”, *Vegetatio*, **1987**, 73, 13-20.
28. J. M. Gehu, “Les groupements végétaux du bassin de la Sambre Française”, *Vegetatio*, **1961**, 10, 161-208.
29. E. Oberdorfer, in “Süddeutsche Pflanzengesellschaften” (Ed. E. Oberdorfer), Gustav Fischer Verlag, Jena, **1992**, pp. 23-38.



30. E. Weretelnik, "Flora i zbiorowiska roslin murow niektorych miast i zamkow na dolnnyin Slasku", *Acta Univ. Wratislav. - Pr. Bot. Wroclaw*, **1982**, 25, 63-110.
31. L. Mucina, "Die Pflanzengesellschaften Österreichs" (Ed. G. Grabherr and L. Mucina), Gustav Fischer Verlag, Jena, **1993**, pp. 241-275.
32. M. Valachovic and S. Maglocky, in "Rastlinne spolocenstva Sloveska. I, Pionierska Vegetacia", (Ed. M. Valachovic), Veda, Bratislava, **1995**, pp. 85-106.
33. Y. Gemici, Ö. Seçmen and G. Görk, in "Urban Ecology" (Ed. M. Öztürk), Ege University Press, Izmir, **1995**, pp. 408-413.
34. A. Aksoy and A. Çelik, "Studies on the ecology of plants growing on the historical monuments of Kayseri, Türkiye", Proceedings of the V<sup>th</sup> International Symposium, **2000**, Tashkent, Uzbekistan.
35. A. Yeşilot, "İstanbul'da tarihi yapılar üzerinde yetişerek bunları tahrip eden bitkiler (The plants that grow up on historical buildings and destroy them)", *MSc Thesis*, **2000**, Marmara University, Istanbul.
36. C. Yarci and H. Özçelik, "Wall flora of Edirne (Thrace Region)", *Ot Sistematik Botanik Dergisi*, **2002**, 9, 57-66.
37. P. H. Davis (Ed.), "Flora of Turkey and the East Aegean Islands", Vol. 1-11, University Press, Edinburgh, **1965-2000**.
38. J. Braun-Blanquet, "Pflanzensoziologie Grundzüge der Vegetationkunde", Springer-Verlag, Vienna, **1964**.
39. C. Raunkier, "The Life Forms of Plants and Statistical Plant Geography", Oxford University Press, Oxford, **1934**. [Reprinted in "History of Ecology Series" (Ed. F. N. Egerton), Ayer Co., Manchester, 1978.]
40. S. L. Mosyakin and O. G. Yavorska, "The nonnative flora of the Kiev (Kyiv) urban area, Ukraine: A checklist and brief analysis", *Urban Habitats*, **2002**, 1, 45-65. (Electronic version: <http://www.urbanhabitats.org>, access date: September 2009)
41. A. Gonzalvo, F. J. Martinez and M. Boscaiu, "The floristic and vegetation study of an area of interest in Valencia (Spain) proposed as plant micro-reserve", *Notulae Bot. Horti Agrobot. Cluj Napoca*, **2008**, 36, 7-12.
42. C. Yarci, M. Serin and V. Altay, "The segetal vegetation of Kocaeli Province (Turkey)", *Ekoloji*, **2007**, 16, 23-33.
43. J. F. Maxwell, "Vascular flora of the Emerald Pool area, Krabi province, southern Thailand", *Maejo Int. J. Sci. Technol.*, **2009**, 3, 1-25.
44. N. Krigas, E. Lagiou, E. Hanlidou and S. Kokkini, "The vascular flora of the Byzantine walls of Thessaloniki (North Greece)", *Willdenowia*, **1999**, 29, 77-94.
45. H. Schepker and I. Kowarik, "Bekämpfung von Neophyten in Niedersachsen: Ursachen. Umfang, Erfolg", *Neobiota*, **2002**, 1, 343-354.
46. I. Kowarik, "Human agency in biological invasions: Secondary releases foster naturalisation and population expansion of alien plant species", *Biol. Invasions*, **2003**, 5, 293-312.

**APPENDIX (Flora List)**

(Abbreviations: Ph = phanerophytes; Ch = chaemaphytes; H = hemicryptophytes; Th = therophytes; G = geophytes; Euro.-Sib. = Euro-Siberian; Ir.-Tur. = Irano-Turanian; Medit. = Mediterranean; E. Medit. = East Mediterranean; cos. = cosmopolitan; wid. = widespread; end. = endemic)

## ANACARDIACEAE

*Pistacia terebinthus* L. subsp. *terebinthus* Ph, Medit.

## APIACEAE

*Conium maculatum* L. H

## ARALIACEAE

*Hedera helix* L. Ph

## ASTERACEAE

*Anthemis cretica* L. H

*Calendula arvensis* L. Th

*Carlina corymbosa* L. H, Medit.

*Centaurea solstitialis* L. subsp. *solstitialis* H, wid.

*Cichorium intybus* L. Ch, wid.

*Cirsium creticum* (Lam.) d'Urv. subsp. *creticum* H, E. Medit.

*Conyza canadensis* L. Th

*Erigeron acer* L. H

*Helminthotheca echioides* (L.) Holup Th.

*Hypochoeris radicata* L. H, Euro.- Sib.

*Inula viscosa* (L.) Ailton H, Medit.

*Lactuca saligna* L. Th

*Pallenis spinosa* (L.) Cass. Th, Medit.

*Picris strigosa* Bieb. H, Ir.- Tur.

*Senecio vulgaris* L. Th

*Sonchus asper* (L.) Hill subsp. *glaucescens* (Jordan) Ball H, wid.

*Taraxacum officinale* Wiggers Ch

*Tussilago farfara* L. G, Euro.- Sib.

## BRASSICACEAE

*Capsella bursa-pastoris* (L.) Medik. Th, cos.

*Diplotaxis tenuifolia* (L.) DC. H

*Lepidium graminifolium* L. H

*Neslia apiculata* Fisch. Th, wid.

*Sinapis arvensis* L. Th, wid.

## CAMPANULACEAE

*Campanula lyrata* Lam. subsp. *lyrata* H, wid., end.

## CARYOPHYLLACEAE

*Cerastium glomeratum* Thuill Th, cos.*Stellaria media* (L.) Vill. subsp. *media* Th*Telephium imperati* L. subsp. *orientale* (Boiss.) Nyman H

## CHENOPODIACEAE

*Chenopodium album* L. subsp. *album* var. *album* Th

## CONVOLVULACEAE

*Convolvulus arvensis* L. H, cos.

## CUCURBITACEAE

*Eballium elaterium* (L.) A. Rich. H, Medit.

## DIPSACACEAE

*Scabiosa columbaria* L. subsp. *columbaria* var. *columbaria* H

## EUPHORBIACEAE

*Euphorbia helioscopia* L. Th*E. peplus* L. var. *peplus* Th*Mercurialis annua* L. Th

## FABACEAE

*Vicia hybrida* L. Th

## GERANIACEAE

*Geranium purpureum* Vill. Th

## LAMIACEAE

*Ajuga chamaepitys* (L.) Schreber H*Calamintha nepeta* (L.) Savi H*Lamium amplexicaule* L. Th, Wid., Euro.- Sib.*Lamium purpureum* L. var. *purpureum* Th, Euro.- Sib.*Micromeria graeca* L. subsp. *graeca* H, Medit.

## LILIACEAE

*Asparagus acutifolius* L. G, Medit.

## MALVACEAE

*Malva sylvestris* L. H

## MORACEAE

*Ficus carica* L. subsp. *carica* Ph, wid.

## OLEACEAE

*Ligustrum vulgare* L. Ph, Euro.- Sib.*Phillyrea latifolia* L. Ph, Medit.

## OXALIDACEAE

*Oxalis corniculata* L. Th, cos.

## PAPAVERACEAE

*Fumaria officinalis* L. Th*Glaucium flavum* Crantz H*Papaver dubium* L. Th

## PLANTAGINACEAE

*Plantago coronopus* L. Th, Euro.- Sib.*P. lagopus* L. Th, Medit.

## POACEAE

*Agrostis capillaris* L. var. *capillaris* H*Bromus sterilis* L. Th*Cynodon dactylon* (L.) Pers var. *dactylon* G*Dactylis glomerata* L. subsp. *hispanica* (Roth) Nyman H*Hordeum murinum* L. Th*Poa annua* L. Th*Setaria viridis* (L.) P. Beauv. Th, wid.*Setaria verticillata* (L.) P. Beauv. Th

## POLYGONACEAE

*Polygonum aviculare* L. Th, cos.*Polygonum lapathifolium* L. Th.*Rumex conglomeratus* Murray H*Rumex pulcher* L. H

## RESEDACEAE

*Reseda lutea* L. var. *lutea* H

## ROSACEAE

*Rubus canescens* DC. var. *canescens* Ph, wid.*Sarcopoterium spinosum* (L.) Spach H, E. Medit.

## RUBIACEAE

*Galium aparine* L. Th

## SCROPHULARIACEAE

*Antirrhinum majus* L. subsp. *majus* H*Cymbalaria muralis* Gaerth. Th, Medit.*Veronica cymbalaria* Bodara Th, Medit.*Veronica persica* Poiret Th

## SIMAROUBACEAE

*Ailanthus altissima* (Miller) Swingle Ph

## SOLANACEAE

*Hyocyamus niger* L. Th, wid.*Solanum nigrum* L. subsp. *nigrum* Th, cos.

## URTICACEAE

*Parietaria judaica* L. H, wid.

## VALERIANACEAE

*Centranthus ruber* (L.) DC. G

## VITACEAE

*Vitis vinifera* L. Ph