

PRENYLATED DIHYDROSTILBENES FROM *Macaranga rubiginosa* *by* Mulyadi Tanjung

Submission date: 08-May-2018 06:23PM (UTC+0800)


Submission ID: 960731955

File name: Chem_Nat_Compd-publish.pdf (284.76K)

Word count: 3972


Character count: 19300

Biomedical Sciences - Pharmacology & Toxicology | Chemistry of Natural Compounds (Editorial Board)

 www.springer.com

Pharmacology & Toxicology [Home](#) > [Biomedical Sciences](#) > [Pharmacology & Toxicology](#)

SUBDISCIPLINES | JOURNALS | BOOKS | SERIES | TEXTBOOKS | REFERENCE WORKS



Chemistry of Natural Compounds

Editor-in-Chief: Sh. S. Sagdullaev
ISSN: 0009-3130 (print version)
ISSN: 1573-8388 (electronic version)
Journal no. 10600

Read
Online

125,21 € Personal Rate e-only

Get Subscription

- ▶ Online subscription, valid from January through December of current calendar year
- ▶ Immediate access to this year's issues via SpringerLink
- ▶ 1 Volume(-s) with 6 issue(-s) per annual subscription
- ▶ Automatic annual renewal
- ▶ More information: >> [FAQs](#) // >> [Policy](#)

READ THIS JOURNAL ON SPRINGERLINK

All Volumes & Issues

FOR AUTHORS AND EDITORS

2016 Impact Factor 0.460

- Aims and Scope
- Instructions for Authors
- Resources for Journal Authors
- Author Academy: Training for Authors
- Copyright Information

SERVICES FOR THE JOURNAL

- Contacts
- Download Product Flyer
- Shipping Dates
- Order Back Issues
- Article Reprints
- Bulk Orders

ALERTS FOR THIS JOURNAL

Get the table of contents of every new issue published in **Chemistry of Natural Compounds**.

Your E-Mail Address

Please send me information on new Springer publications in **Pharmacy**.

ABOUT THIS JOURNAL | EDITORIAL BOARD

Editor-in-Chief

Sh. Sh. Sagdullaev – S. Yu. Yunusov Institute of the Chemistry of Plant Substances, Academy of Sciences of the Republic of Uzbekistan, Tashkent, Uzbekistan

Secretary

V. I. Vinogradova – S. Yu. Yunusov Institute of the Chemistry of Plant Substances, Academy of Sciences of the Republic of Uzbekistan, Tashkent, Uzbekistan

Editorial Board

N. D. Abdullaev – S. Yu. Yunusov Institute of the Chemistry of Plant Substances, Academy of Sciences of the Republic of Uzbekistan, Tashkent, Uzbekistan

Haji Akber Aisa – Xinjiang Technical Institute of Physics and Chemistry, Chinese Academy of Sciences, Urumqi, Xinjiang, P. R. China

S. M. Adekenov – JSC International Research and Production Holding "Phytochemistry", Karaganda, Republic of Kazakhstan

Sh. S. Azimova – S. Yu. Yunusov Institute of the Chemistry of Plant Substances, Academy of Sciences of the Republic of Uzbekistan, Tashkent, Uzbekistan

T. F. Aripov – A. S. Sadykov Institute of the Bioorganic Chemistry, Academy of Sciences of the Republic of Uzbekistan, Tashkent, Uzbekistan

S. F. Aripova – S. Yu. Yunusov Institute of the Chemistry of Plant Substances, Academy of Sciences of the Republic of Uzbekistan, Tashkent, Uzbekistan

K. Husnu Can Baser – Near East University, Department of Pharmacognosy, Faculty of Pharmacy, Lefkosa (Nicosia), N. Cyprus

S. O. Bachurin – Institute of Physiologically Active Compounds, Russian Academy of Sciences, Moscow, Russia

A. I. Miroshnikov – M. M. Shemyakin and Yu. A. Ovchinnikov Institute of Bioorganic Chemistry, Russian Academy of Sciences, Moscow, Russia

Elmuradov B. Zh. – S. Yu. Yunusov Institute of the Chemistry of Plant Substances, Academy of Sciences of the Republic of Uzbekistan, Tashkent, Uzbekistan

M. S. Yunusov – Ufa Institute of Chemistry, Russian Academy of Sciences, Ufa, Russia

Editorial Advisory Board

S. D. Gusakova – S. Yu. Yunusov Institute of the Chemistry of Plant Substances, Academy of Sciences of the Republic of Uzbekistan, Tashkent, Uzbekistan


B. T. Ibragimov – A. S. Sadykov Institute of the Bioorganic Chemistry, Academy of Sciences of the Republic of Uzbekistan, Tashkent, Uzbekistan

V. G. Kartsev – InterBioScreen Ltd., Moscow, Russia

E. P. Kemertelidze – I. Kutateladze Institute of Pharmacochimistry, Tbilisi State Medical University, Tbilisi, Georgia

A. V. Kutchin – Institute of Chemistry, Komi Scientific Center, Ural Branch, Russian Academy of Sciences, Syktyvkar, Russia

RELATED BOOKS - SERIES - JOURNALS



Journal

Organic and Medicinal Chemistry Letters

Editor-in-Chief: Bimal K. Banik

1/10

A. M. Muzafarov – A. N. Nesmeyanov Institute of Organoelement Compounds, Russian Academy of Sciences, Moscow, Russia

V. G. Nenajdenko – M. V. Lomonosov Moscow State University, Moscow, Russia

R. K. Rakhmanberdieva – S. Yu. Yunusov Institute of the Chemistry of Plant Substances, Academy of Sciences of the Republic of Uzbekistan, Tashkent, Uzbekistan

S. Sh. Rashidova – Institute of Polymers Chemistry and Physics, Academy of Sciences of the Republic of Uzbekistan, Tashkent, Uzbekistan

N. F. Salakhutdinov – Novosibirsk Institute of Organic Chemistry Siberian Branch, Russian Academy of Sciences, Novosibirsk, Russia

V. A. Stonik – G. B. Elyakov Pacific Institute of Bioorganic Chemistry, Far-East Branch, Russian Academy of Sciences, Vladivostok, Russia

A. S. Turaev – A. S. Sadykov Institute of the Bioorganic Chemistry, Academy of Sciences of the Republic of Uzbekistan, Tashkent, Uzbekistan

V. P. Khilya – Taras Shevchenko Kiev National University, Kiev, Ukraine

M. I. Choudhary – International Center for Chemical and Biological Sciences, University of Karachi, Karachi, Pakistan

Xiaojiang Hao – Kunming Institute of Botany, Yunnan, P. R. China

For other enquiries, please contact the Publishing Editor



Chemistry of Natural Compounds

All Volumes & Issues

ISSN: 0009-3130 (Print) 1573-8388 (Online)

In this issue (56 articles)

1.

OriginalPaper

A New Biphenanthrene Glucoside with Cytotoxic Activity from *Cremastra appendiculata*

Xin-Qiao Liu, Wen-ke Yuan, Qiao-Yu Yuan, Xiao-Ping Li... Pages 211-214

2.

OriginalPaper

Prenylated Dihydrostilbenes from *Macaranga rubiginosa*

Mulyadi Tanjung, Euis H. Hakim, Yana M. Syah Pages 215-218

3.

OriginalPaper

Structural Characteristics and Antitumor Activity of Fucoidans from the Brown Alga *Sargassum muticum*

R. V. Usoltseva, Peipei Zhao, M. I. Kusaikin, Aironq Jia... Pages 219-223

4.

OriginalPaper

Secondary Metabolites Produced by the Deep-Sea-Derived Fungus *Engyodontium album*

Weiyl Wang, Shanshan Li, Zhuo Chen, Zengpeng Li... Pages 224-226

5. 

OriginalPaper

New Fatty Acid From a Gorgonian-Derived Xylaria sp. Fungus*Da-Wei Sun, Fei Cao, Min Liu, Fei-Fei Guan...* Pages 227-2306. 

OriginalPaper

[1 + 1]-Condensation of 12-Oxo-Derivatives of Ricinoleic Acid Esters with Hydrazine Hydrate on the Route to Macrocycles*G. Yu. Ishmuratov, M. P. Yakovleva, V. A. Vydrina...* Pages 231-2337. 

OriginalPaper

A New Lactone from the Twigs of Cinnamomum cassia*Zhenlin Li, Zhichen Cai, Shihui Qian, Minghua Chen* Pages 234-2368. 

OriginalPaper

Identification and Antifungal Activity of Metabolites from the Mangrove Fungus Phoma sp. L28*Song Huang, Jiaxin Xu, Fenqi Li, Danli Zhou, Li Xu...* Pages 237-2409. 

OriginalPaper

Hemisynthesis and Spectroscopic Characterization of Three New Chalcone Derivatives from Dorstenia barteri*Bathelemy Ngameni, Ghislain Wabo Fotso, Pantaleon Ambassa...* Pages 241-247

10. 

OriginalPaper

Synthesis and Antioxidant Properties of 5,6,7,8-TetrahydroxyflavoneLinlin Jing, Hui ping Ma, Pengcheng Fan, Zhengping Jia Pages 248-25311. 

OriginalPaper

Three New Phenolic Compounds from Eucommia ulmoidesJia Chen, Xiao-Qing Xu, Xing-Dong Kang, Wen-Ting Zhang... Pages 254-25612. 

OriginalPaper

A New Isoflavan from Abrus precatoriusZhihui Xiao, Shuhong Tao, Yaxian Yang, Yu Zhang Pages 257-25913. 

OriginalPaper

Influenza Antiviral Activity of Br-Containing [2R,4R(S),4aR,7R,8aR]-4,7-Dimethyl-2-(Thiophen-2-YL)Octahydro-2H-Chromen-4-Ols Prepared from (-)-IsopulegolE. V. Nazimova, A. A. Shtro, V. B. Anikin... Pages 260-26414. 

OriginalPaper

New Iridoid from the Stems of Viburnum erosumSeo-Ji In, Kyeong-Hwa Seo, Hyoung-Geun Kim, Na-Young Song... Pages 265-26815. 

OriginalPaper

Phlotuberosides I and II, New Iridoid Glycosides from Phlomis tuberosa

D. N. Olennikov, N. K. Chirikova Pages 269-272

16. 

OriginalPaper

A New Sesquiterpenoid from Acanthopanax senticosus

Zhi-Feng Li, Qi Wang, Gang Chen, Shi-Lin Yang... Pages 273-275

17. 

OriginalPaper

Synthesis of Rupestonic Acid Derivatives with Antiviral Activity

Jiangyu Zhao, Chao Niu, Gen Li, Haji Akber Aisa Pages 276-283

18. 

OriginalPaper

Two New Sesquiterpene Lactones from Artemisia halophila

S. M. Adekenov, Zh. R. Shaimerdenova, Yu. V. Gatilov... Pages 284-289

19. 

OriginalPaper

New Thomimarine E from Marine Isolate of the Fungus Penicillium thomii

Sh. Sh. Afiyatullo, E. V. Leshchenko, M. P. Sobolevskaya... Pages 290-294

20. 

OriginalPaper

Two New ent-Labdane Diterpenes from the Roots of Euphorbia yinshanica

Ben-Yin Zhang, Heng-Xia Yin, De-Jun Zhang Pages 295-298

We use cookies to improve your experience with our site. [More information](#)


PRENYLATED DIHYDROSTILBENES FROM *Macaranga rubiginosa*

Mulyadi Tanjung,¹ Euis H. Hakim,² and Yana M. Syah^{2*}

Phytochemical isolation of the methanol extract of *Macaranga rubiginosa* leaves afforded five prenylated dihydrostilbenes. Two of them were known dihydrostilbenes laevifolins A (1) and B (2), while the other three were new compounds, trivially named macarubiginosins A–C (3–5). The structures of the new compounds were elucidated based on their UV, 1D and 2D NMR, and HR-ESI-MS spectral data. Compounds 1–5 were tested for their cytotoxicity against P-388 cells, showing that compound 1 was the most active with IC_{50} 4.3 μ M.

Keywords: macarubiginosins A–C, laevifolins A and B, prenylated dihydrostilbenes, *Macaranga rubiginosa*, Euphorbiaceae, cytotoxicity, P-388 cells.

Macaranga is a large genus of Euphorbiaceae (ca. 260 species) distributed in the tropical region of Africa, India, South East Asia, and Pacific islands [1]. The species of this genus are important plants found in the secondary forests of Kalimantan and Sumatra Islands of Indonesia [2, 3]. Our interest in the phytochemistry of *Macaranga* species is motivated by the fact that they contain flavonoid and stilbene derivatives bearing various terpenyl groups, including prenyl, geranyl, farnesyl, and labdanyl groups [4]. Recently, we reported our phytochemical investigation of some *Macaranga* plants growing in Kalimantan Island [5–9]. As further work on Indonesian *Macaranga*, in this paper we report the presence of prenylated derivatives of dihydrostilbenes, trivially named macarubiginosins A–C (3–5), from the leaf extract of *Macaranga rubiginosa* Ridl. collected from Sumatra Island. Along with these compounds, two known prenylated dihydrostilbens, laevifolins A and B (1, 2), were also isolated. As a preliminary biological test, this paper also briefly describes the cytotoxic properties of compounds 1–5 against murine leukemia P-388 cells.



1) Chemistry Department, Airlangga University, Jalan Darmawangsa Dalam, 60222, Surabaya, Indonesia; 2) Natural Products Chemistry Research Group, Organic Chemistry Division, Bandung Institute of Technology, Jalan Ganesha 10, 40132, Bandung, Indonesia, e-mail: yana@chem.itb.ac.id. Published in *Khimiya Prirodnykh Soedinenii*, No. 2, March–April, 2017, pp. 184–187. Original article submitted September 22, 2015.

TABLE 1. NMR Data of Compounds 3–5 (δ , ppm, J/Hz)

C atom	δ_H			δ_C		
	3 ^a	4 ^b	5 ^b	3	4	5
1	–	–	–	140.8	139.3	139.5
2	–	–	–	119.7	117.7	112.0
3	–	–	–	155.1	153.7	153.2
4	6.21 (s)	6.22 (s)	6.21 (s)	102.3	102.7	102.9
5	–	–	–	152.6	153.1	153.8
6	–	–	–	110.9	111.9	117.7
7	2.87 (dd, J = 16.0, 5.7) 2.51 (dd, J = 16.0, 8.3)	2.55 (t, J = 6.8)	2.60 (t, J = 6.8)	29.9	20.1	20.3
8	3.70 (dd, J = 8.3, 5.7)	1.73 (t, J = 6.8)	1.76 (t, J = 6.8)	70.7	33.0	35.8
9	–	–	–	76.6	73.2	73.4
10	1.15 (s)	1.27 (s)	1.29 (s)	19.9	26.6	26.8
11	1.29 (s)	1.27 (s)	1.29 (s)	25.4	26.6	26.8
12	3.33 (d, J = 6.8)	3.31 (d, J = 6.7)	3.34 (d, J = 6.7)	25.5	25.8	25.5
13	5.09 (br.t, J = 6.8)	5.10 (br.t, J = 6.7)	5.14 (br.t, J = 6.7)	125.8	123.3	123.3
14	–	–	–	130.2	133.6	133.5
15	1.73 (br.s)	1.74 (br.s)	1.78 (br.s)	18.1	17.9	18.1
16	1.62 (br.s)	1.70 (br.s)	1.70 (br.s)	25.8	25.7	25.9
α	2.76 (m)	2.79 (m)	2.80 (m)	31.5	31.1	31.8
β	2.67 (m)	2.70 (m)	2.62 (m)	33.3	33.2	33.4
1'	–	–	–	133.0	132.2	136.8
2'	–	–	6.63 (d, J = 2.4)	127.3	125.8	113.3
3'	–	–	–	144.1	142.4	144.4
4'	–	–	–	143.2	142.2	140.2
5'	6.65 (d, J = 8.6)	6.72 (d, J = 8.0)	–	113.3	112.8	121.3
6'	6.36 (d, J = 8.6)	6.66 (d, J = 8.0)	6.48 (d, J = 2.4)	120.2	121.3	122.0
7'	3.39 (d, J = 6.8)	3.34 (d, J = 6.7)	3.33 (d, J = 6.7)	26.1	25.3	30.2
8'	5.09 (br.t, J = 6.8)	5.10 (br.t, J = 6.7)	5.14 (br.t, J = 6.7)	124.6	121.1	122.1
9'	–	–	–	131.2	134.8	134.7
10'	1.75 (br.s)	1.78 (br.s)	1.82 (br.s)	18.2	18.0	18.2
11'	1.64 (br.s)	1.71 (br.s)	1.80 (br.s)	25.8	25.6	26.0

^aIn acetone- d_6 ; ^bin CDCl₃.




Fig. 1. Selected important HMBC correlations of compounds 3–5.

Compounds 1 and 2 were determined to have the structures of laevofilins A and B, respectively, based on comparison of their NMR spectral data with those reported in the literature [10].

Compound 3 (macarubiginosin A) was isolated as a pale yellowish semisolid, and its UV absorption characteristics were very close to those 1 and 2 (see Experimental). Based on high resolution electrospray ionization mass spectrometry (HR-ESI-MS) data, 3 has the molecular formula C₂₉H₃₈O₅ (found [M – H][–] at *m/z* 465.2643, calcd 465.2641), suggesting that

it could be a monooxygenated derivative of either **1** or **2**. The ^{13}C NMR spectrum of **3** (Table 1), however, showed 29 carbon signals, 16 of which were sp^2 -carbon atoms, indicating that the symmetry of ring A in **1** or **2** is lost in **3**. The ^1H NMR spectrum (Table 1) disclosed two mutually coupled methylene signals (δ 2.76 and 2.67) for C- α and C- β , and signals characteristics for only two prenyl groups (δ 3.39 and 3.33, each 2H; 5.09, 2H; 1.75, 1.73, 1.64, and 1.62, each 3H). Accordingly, the third prenyl group in **3** should be oxygenated to form a 3-methyl-2,3-epoxybutyl group (δ 3.70, 2.87, and 2.51, each 1H and dd; 1.29 and 1.15, each 3H and s; δ_{C} 19.9, 25.4, 29.9, 70.7, and 76.6). The aromatic region of the ^1H NMR spectrum showed a singlet at δ 6.21 and a pair of *ortho*-coupled doublets at δ 6.65 and 6.36. From these NMR data, structure **3** was assigned to macarubiginosin A (**1**) as an epoxy derivative of laevofilin A. HSQC and HMBC spectra of **3** were consistent with the structure of macarubiginosin A, and the important HMBC correlations are as shown in Fig. 1. The stereochemistry of the epoxy group in **3** was not determined.

Compounds **4** (macarubiginosin B) and **5** (macarubiginosin C) have the same molecular formula, namely $\text{C}_{29}\text{H}_{38}\text{O}_4$, as determined by their HR-ESI-MS data (see Experimental), suggesting that these compounds are isomers of either laevofilins A (**4**) or B (**5**). Looking at their aromatic signals in their ^1H NMR spectra (Table 1), **4** contained a pair of *ortho*-coupled doublets (δ 6.72 and 6.66), while **5** had a pair of *meta*-coupled doublets (δ 6.63 and 6.48), in addition to singlet signals at δ 6.22 and 6.21, respectively. Therefore, **4** and **5** must have the same arrangements in the aromatic parts as those of **1** and **2**. Further spectral analysis disclosed that the ^1H and ^{13}C NMR data of **4** and **5** were very similar to those of **3**, except that the signals of the 3-methyl-2,3-epoxybutyl group in **3** were replaced by signals characteristic of a dihydrodimethylpyran ring in **4** and **5** (see Table 1). Because the ^1H and ^{13}C NMR spectra of **4** and **5** indicated no symmetry in the molecules, the dimethylpyran ring should be in ring A of **4** and **5**. Therefore, structures **4** and **5** can be assigned to macarubiginosins B and C, respectively. The HMBC correlations found in both compounds, as shown in Fig. 1, confirmed the structural assignments for these two compounds. In addition, the NMR chemical shifts and coupling constants of the dihydrodimethylpyran ring in **4** and **5** were in agreement with those reported in the literature [11, 12].

Preliminary cytotoxic evaluation of compounds **1–5** against murine leukemia P-388 was carried out by the method previously described [13]. These compounds exhibited IC_{50} values of 4.3 ± 0.6 , 12.3 ± 0.6 , 10.8 ± 3.0 , 52.9 ± 3.4 , and 45.8 ± 2.6 μM , respectively. Modification of one of the prenyl groups in ring A of **1** or **2** to an epoxy (compound **3**) or a dimethylpyran (compounds **4** or **5**) derivative reduces its cytotoxicity.

EXPERIMENTAL

General Experimental Procedures. UV spectra were measured with a Varian 100 spectrophotometer. ^1H and ^{13}C NMR spectra were recorded with a JEOL ECA500 spectrometer operating at 500 (^1H) and 125 (^{13}C) MHz, using residual and deuterated solvent peaks of CDCl_3 (δ_{H} 7.26 and δ_{C} 77.0, respectively) or acetone- d_6 (δ_{H} 2.04 and δ_{C} 29.8, respectively) as reference standards. Mass spectra were obtained with an ESI-TOF Waters LCT Premier XE mass spectrometer (negative mode). Vacuum liquid chromatography (VLC) and centrifugal planar chromatography (CPC) were carried out using Merck Si gel 60 GF₂₅₄ and, for TLC analysis, precoated Si gel plates (Merck Kieselgel 60 GF₂₅₄, 0.25 mm) were used. Visualization of TLC chromatogram was done under 254 nm UV light and by spraying with ceric sulfate solution and heating. Solvents used for extraction and separation were technical grades that were distilled before used.

Plant Material. The leaves of *M. rubiginosa* were collected in April 2008 from Cinta Damai Village, Banyuasin District, South Sumatra, Indonesia. The specimen was identified at the Herbarium Bogoriense, Center of Biological Research and Development, National Institute of Science, Bogor, Indonesia.

Extraction and Isolation. Extraction and isolation of the compounds from *M. rubiginosa* were done according to the method previously described [6] with some modifications. The dried and powdered leaves of *M. rubiginosa* (3.0 kg) were macerated in MeOH at room temperature (3×10 L) to give a semisolid extract (300 g) after solvent evaporation under reduced pressure. The extract was redissolved in MeOH–water (9:1) and partitioned into *n*-hexane (228 g) and EtOAc (50 g) fractions. A part of the EtOAc fraction (20 g) was fractionated using VLC (Si gel 160 g) eluted with *n*-hexane–EtOAc (4:1, 7:3, and 1:1) and EtOAc, giving four major fractions A–D. TLC analysis of fraction A (165 mg), detected under UV light 254 nm, showed no phenolic spots, and therefore analysis was not continued. Fraction C (1.3 g) was refractionated using CPC eluted with CHCl_3 and CHCl_3 –EtOAc (9:1) into two subfractions C1 and C2. Fraction B and subfraction C1 were combined into subfraction BC1 (450 mg), and this subfraction was repeatedly purified using PCC eluted with *n*-hexane– CHCl_3 (3:7, 1:4), and CHCl_3 to give macarubiginosins B (**4**, 15 mg) and C (**5**, 3.5 mg). Subfraction C2 (650 mg) was also purified using the

same method [eluents of *n*-hexane-CHCl₃ (3:7, 1:4), CHCl₃, CHCl₃-EtOAc (9:1)] to afford laevifolins A (**1**, 250 mg) and B (**2**, 60 mg). Using the same method, purification of fraction D (250 mg) with eluents of *n*-hexane-EtOAc (4:1 and 7:3) gave macarubiginosin A (**3**, 25 mg).

Laevofilin A (1). Pale yellowish semisolid. UV (MeOH, λ_{\max} , nm) (log ϵ): 209 (4.98), 228 sh (4.52), 284 (4.02), 316 (3.16); (MeOH + NaOH): 208 (4.98), 227 sh (4.71), 290 (4.10), 340 (3.98). ¹H and ¹³C NMR data were in agreement with those reported in [10].

Laevofilin B (2). Pale yellowish semisolid. UV (MeOH, λ_{\max} , nm) (log ϵ): 214 (4.98), 229 sh (4.82), 284 (4.32), 318 (3.35); (MeOH + NaOH): 215 (4.99), 229 sh (4.86), 286 (4.32), 332 (3.98). ¹H and ¹³C NMR data were in agreement with those reported in [12].

Macarubiginosin A (3). Pale yellowish semisolid. UV (MeOH, λ_{\max} , nm) (log ϵ): 215 (4.99), 232 sh (4.91), 284 (4.52), 315 (3.97); (MeOH + NaOH): 217 (4.99), 253 sh (4.69), 292 (4.55), 330 (3.98). For ¹H (500 MHz, acetone-d₆) and ¹³C (125 MHz, acetone-d₆) NMR data, see Table 1. HR-ESI-MS *m/z* 465.2643 [M - H]⁻ (calcd for C₂₉H₃₇O₅, 465.2641).

Macarubiginosin B (4). Pale yellowish semisolid. UV (MeOH, λ_{\max} , nm) (log ϵ): 215 (4.99), 231 sh (4.87), 280 (4.64), 317 (4.14); (MeOH + NaOH): 214 (4.99), 232 sh (4.90), 282 (4.63), 348 (4.11). For ¹H (500 MHz, CDCl₃) and ¹³C (125 MHz, CDCl₃) NMR data, see Table 1. HR-ESI-MS *m/z* 449.2672 [M - H]⁻ (calcd for C₂₉H₃₇O₄, 449.2692).

Macarubiginosin C (5). Pale yellowish semisolid. UV (MeOH, λ_{\max} , nm) (log ϵ): 215 (4.95), 232 sh (4.80), 283 (4.30), 318 (3.37); (MeOH + NaOH): 215 (4.96), 230 sh (4.86), 291 (4.16), 332 (3.96). For ¹H (500 MHz, CDCl₃) and ¹³C (125 MHz, CDCl₃) NMR data, see Table 1. HR-ESI-MS *m/z* 449.2688 [M - H]⁻ (calcd for C₂₉H₃₇O₄, 449.2692).

Cytotoxic Evaluation. Cytotoxic properties of the isolated compounds **1-5** against murine leukemia P-388 cells was evaluated according to the method of MTT assay as previously described [13].

ACKNOWLEDGMENT

Part of this work is supported by the KK Research Grant (Contract No. 1763/I1.B04.1/KU/2015), Bandung Institute of Technology. The authors also thank the staff of Herbarium Bogoriense, Center of Biological Research and Development, National Institute of Science, Bogor, Indonesia, for identification of the plant specimen.

REFERENCES

1. K. K. M. Kulju, S. E. C. Sierra, S. G. A. Draisma, R. Samuel, and P. C. van Welzen, *Am. J. Bot.*, **94**, 1726 (2007).
2. J. W. F. Slik, P. J. A. Kebler, and P. C. van Welzen, *Ecol. Indic.*, **2**, 311 (2003).
3. D. Priatna, K. Kartawinata, and R. Abdulhadi, *Reinwardtia*, **12**, 237 (2004).
4. J. J. Magadula, *J. Med. Plant Res.*, **8**, 489 (2014).
5. Y. M. Syah, E. H. Hakim, S. A. Achmad, M. Hanafi, and E. L. Ghisalberti, *Nat. Prod. Commun.*, **4**, 63 (2009).
6. M. Tanjung, E. H. Hakim, D. Mujahidin, and M. Hanafi, *J. Asian Nat. Prod. Res.*, **11**, 929 (2009).
7. W. Agustina, L. D. Juliawaty, E. H. Hakim, and Y. M. Syah, *ITB J. Sci.*, **44A**, 13 (2012).
8. Y. M. Syah and E. L. Ghisalberti, *Nat. Prod. J.*, **2**, 45 (2012).
9. M. S. Fareza, Y. M. Syah, D. Mujahidin, L. D. Juliawaty, and I. Kurniasih, *Z. Naturforsch.*, **69c**, 375 (2014).
10. N. Ahmat, I. M. Said, J. Latip, L. B. Din, Y. M. Syah, and E. H. Hakim, *Nat. Prod. Commun.*, **2**, 1137 (2007).
11. T. Ersam, S. A. Achmad, E. L. Ghisalberti, E. H. Hakim, and Y. M. Syah, *J. Chem. Res. (S)*, 186 (2002).
12. V. Rukachaisirikul, M. Kamkaew, D. Sukavisit, S. Phongpaichit, P. Sawangchote, and W. C. Taylor, *J. Nat. Prod.*, **66**, 1531 (2003).
13. S. Sahidin, E. H. Hakim, L. D. Juliawaty, Y. M. Syah, L. B. Din, E. L. Ghisalberti, J. Latip, I. M. Said, and S. A. Achmad, *Z. Naturforsch.*, **60c**, 723 (2005).

PRENYLATED DIHYDROSTILBENES FROM *Macaranga rubiginosa*

ORIGINALITY REPORT

18%

SIMILARITY INDEX

11%

INTERNET SOURCES

17%

PUBLICATIONS

2%

STUDENT PAPERS

PRIMARY SOURCES

- 1** Haryoto Saroyobudiono. "Oligostilbenoids from *Shorea gibbosa* and their cytotoxic properties against P-388 cells", *Journal of Natural Medicines*, 04/2008

Publication

2%
- 2** d-nb.info

Internet Source

2%
- 3** jocpr.com

Internet Source

2%
- 4** www.mdpi.com

Internet Source

2%
- 5** Aulia Ilmiawati, Euis H. Hakim, Yana M. Syah. "Prenylated 9,10-dihydrophenanthrenes from *Macaranga javanica*", *Zeitschrift für Naturforschung B*, 2015

Publication

2%
- 6** Andrea Mühlbauer. "Five Novel Taccalonolides from the Roots of the Vietnamese Plant *Tacca paxiana*", *Helvetica Chimica Acta*, 06/2003

1%

-
- 7 www.derpharmachemica.com 1 %
Internet Source
-
- 8 Muhtadi. "Cytotoxic resveratrol oligomers from the tree bark of *Dipterocarpus hasseltii*", *Fitoterapia*, 200612 1 %
Publication
-
- 9 mdpi.com 1 %
Internet Source
-
- 10 Phi-Hung Nguyen, Thi-Ngoc-Anh Nguyen, Trong-Tuan Dao, Hu-Won Kang, Derek-Tantoh Ndinteh, Joseph-Tanyi Mbafor, Won-Keun Oh. "AMP-Activated Protein Kinase (AMPK) Activation by Benzofurans and Coumestans Isolated from ", *Journal of Natural Products*, 2010 1 %
Publication
-
- 11 Vatcharin Rukachaisirikul, Mayuree Kamkaew, Dusadee Sukavisit, Souwalak Phongpaichit, Prakart Sawangchote, Walter C. Taylor. "Antibacterial Xanthonenes from the Leaves of ", *Journal of Natural Products*, 2003 1 %
Publication
-
- 12 znaturforsch.com 1 %
Internet Source
-
- 13 V. U. Ahmad. "A new acylated flavone glycoside

from the fruits of *Stocksia brauhica*", *Journal of Asian Natural Products Research*, 4/2007

1%

Publication

14

Yana M. Syah, Emilio L. Ghisalberti. "Flavanone derivatives from *Macaranga tanarius*", *Biochemical Systematics and Ecology*, 2015

<1%

Publication

15

Mulyadi Tanjung, Tjitjik Srie Tjahjandarie, Mulya Hadi Sentosa. "Antioxidant and cytotoxic agent from the rhizomes of *Kaempferia pandurata*", *Asian Pacific Journal of Tropical Disease*, 2013

<1%

Publication

16

Yana M. Syah. "Cytotoxic prenylated flavones from *Artocarpus champeden*", *Journal of Natural Medicines*, 10/2006

<1%

Publication

17

Saranyoo Klaiklay, Yaowapa Sukpondma, Vatcharin Rukachaisirikul, Souwalak Phongpaichit. "Friedolanostanes and xanthonenes from the twigs of *Garcinia hombroniana*", *Phytochemistry*, 2013

<1%

Publication

18

Iqbal Musthapa. "An oxepinoflavone from *Artocarpus elasticus* with cytotoxic activity against P-388 cells", *Archives of Pharmacal Research*, 02/2009

<1%

Publication

19

Zhang, Lai-Bin, Chun Lei, Li-Xin Gao, Jing-Ya Li, Jia Li, and Ai-Jun Hou. "Isoprenylated Flavonoids with PTP1B Inhibition from *Macaranga denticulata*", *Natural Products and Bioprospecting*, 2016.

Publication

<1%

20

Yana M Syah, Sjamsul A Achmad, Emilio L Ghisalberti, Euis H Hakim, Lukman Makmur, Didin Mujahidin. "Artoindonesianins Q–T, four isoprenylated flavones from *Artocarpus champeden* Spreng. (Moraceae)", *Phytochemistry*, 2002

Publication

<1%

21

Tiphaine Péresse, Gwenaëlle Jézéquel, Pierre-Marie Allard, Van-Cuong Pham et al. " Cytotoxic Prenylated Stilbenes Isolated from ", *Journal of Natural Products*, 2017

Publication

<1%

22

J. Trujillo-Ferrara. "Solvent-Free Synthesis of Arylamides and Arylimides, Analogues of Acetylcholine", *Synthetic Communications*, 8/1/2005

Publication

<1%

23

Yong-Hong Wang. "Prenylated Stilbenes and Their Novel Biogenetic Derivatives from *Artocarpus chama*", *European Journal of Organic Chemistry*, 08/2006

<1%

Publication

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

PRENYLATED DIHYDROSTILBENES FROM *Macaranga rubiginosa*

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10
