

This is the peer-reviewed version of the article:

Kukić, J.; Popović, V.; Petrović, S.; Mucaji, P.; Ćirić, A.; Stojković, D.; Soković, M.  
Antioxidant and Antimicrobial Activity of *Cynara Cardunculus* Extracts. *Food Chem.* **2008**,  
*107* (2), 861–868. <https://doi.org/10.1016/j.foodchem.2007.09.005>



This work is licensed under the  
<https://creativecommons.org/licenses/by-nc-nd/4.0/>

1           Antioxidant and Antimicrobial Activity of  
2                           *Cynara cardunculus* Extracts

3       *Jelena Kukić<sup>1</sup>, Višnja Popović<sup>1</sup>, Silvana Petrović<sup>1,\*</sup>, Pavel Mucaji<sup>2</sup>, Ana Ćirić<sup>3</sup>, Dejan*  
4   *Stojković<sup>3</sup>, Marina Soković<sup>3</sup>*

5  
6       <sup>1</sup> Institute of Pharmacognosy, Faculty of Pharmacy, Vojvode Stepe 450, 11221 Belgrade,  
7   Serbia

8       <sup>2</sup> Department of Pharmacognosy and Botany, Pharmaceutical Faculty, Comenius  
9   University, Odbojarov 10, 83232, Bratislava, Slovak Republic

10       <sup>3</sup> Mycological Laboratory, Department for Plant Physiology, Institute for Biological  
11       Research "Siniša Stanković", Bulevar Despota Stefana 142, 11000 Belgrade, Serbia

---

\* Corresponding author. Tel.: +381-11-39-70-379/722; Fax: +381-11-39-72-840.

*E-mail address:* silvana@pharmacy.bg.ac.yu (S. Petrović)

12 **Abstract**

13 **The whole,** fresh involucre bracts of cardoon, *Cynara cardunculus* L. (Compositae),  
14 were extracted with EtOH and aqueous suspension of obtained EtOH extract was  
15 partitioned successively with CHCl<sub>3</sub>, EtOAc and *n*-BuOH, leaving residual water extract.  
16 All obtained extracts were evaluated on their antioxidant and antimicrobial properties.  
17 The antioxidant potential was evaluated using following *in vitro* methods: FRAP (Ferric  
18 Reducing Antioxidant Power) assay, and scavenging of 2,2-diphenyl-1-picrylhydrazyl  
19 (DPPH) radical. Antimicrobial activity was estimated using microdilution technique  
20 **against food-borne, mycotoxin producers and human pathogenic bacteria and**  
21 **micromycetes.** Following bacteria were tested: *Salmonella typhimurium*, *Escherichia*  
22 *coli*, *Bacillus subtilis*, *Staphylococcus epidermidis*, *Staphylococcus aureus*, as well as  
23 micromycetes: *Aspergillus niger*, *Aspergillus ochraceus*, *Aspergillus flavus*, *Penicillium*  
24 *ochrochloron*, *Penicillium funiculosum*, *Trichoderma viride*, *Fusarium tricinctum* and  
25 *Alternaria alternata*. Results showed that all extracts possess concentration dependent  
26 antioxidant activity. In biological assays, *C. cardunculus* extracts showed antimicrobial  
27 activity comparable with standard antibiotics.

28

29 *Keywords:* *Cynara cardunculus*; Involucre bracts; Antioxidant activity; FRAP; DPPH;  
30 Antimicrobial activity

31 **1. Introduction**

32 Cardoon or wild artichoke (*Cynara cardunculus* L., Compositae) is a perennial plant,  
33 which shares a recent common ancestor with the modern cultivated “globe” artichoke, *C.*  
34 *scolymus* L. Both plants have their origin in edible *Cynara* cultivars used by early  
35 farmers in the Mediterranean region (Kelly & Pepper, 1996). Traditional applications of  
36 *C. cardunculus* consider the usage of the blanched leaves, fleshy leaf petioles and the  
37 receptacle in soups, stews and salads (do Amaral Franco, 1976; Grieve, 1971;  
38 Fernandez, Curt, & Aguado, 2006). There are reports of usage of its petioles and roots if  
39 properly prepared (Kelly & Pepper, 1996). Flowers of *C. cardunculus* are rich in  
40 proteases, namely cardosins A and B, due which aqueous extracts of its flowers have  
41 been used for centuries in the Iberian Peninsula for manufacturing of ovine and/or  
42 caprine milk cheeses (Silva & Malcata, 2005; Fernandez et al., 2006). Cardoon is  
43 traditionally used as a diuretic, choloretic, cardiotonic and an antihemorrhoidal (Koubaa,  
44 Damak, McKillop, & Simmonds, 1999). Cardoon leaves are used for their cholagogue,  
45 choloretic and choliokinetic actions, for treatment of dyspepsia and as antidiabetics (Paris  
46 & Moyses, 1971; Koubaa et al., 1999).

47 Previous chemical investigations have shown the presence of saponins, sesquiterpene  
48 lactones, flavones, sterols, coumarins and lignans in leaves and seeds of *C. cardunculus*  
49 (Valentao, Fernandez, Carvalho, Andrade, Seabra, & Bastos, 2002; Ševčíkova, Glatz, &  
50 Slanina, 2002; Pinelli, Agostini, Comino, Lanteri, Portis, & Romani, 2007; Koubaa &  
51 Damak, 2003). In involucre bracts of the investigated species were identified sterols,  
52 triterpenoid saponins, coumarines, flavonoids and caffeic acid derivatives (Mučaji,  
53 Grančai, Nagy, Višňovská, & Ubik, 2000).

54 The antioxidant activity of lyophilized aqueous extract of cardoon leaves and against  
55 superoxide radical is reported (Valentao et al., 2002). Mono- and dicaffeoylquinic acids  
56 which are present in cardoon extracts showed anti-HIV integrase activity (Slanina,  
57 Taborska, Bochorakowa, Humpa, Robinson, & Schram, 2001). Triterpenoid saponins,  
58 isolated from involucre bracts of *C. cardunculus*, reduce the chemically induced  
59 mutagenesis *in vitro* (Križkova, Mučaji, Nagy, & Krajčovič, 2004) and possess  
60 anticomplement activity (Mučaji, Bukovsky, Grančai, & Nagy, 2003). Recent study  
61 showed that *C. cardunculus* leaf extract prevents the age-associated loss of vasomotor  
62 function (Rossoni, Grande, Galli, & Visioli, 2005).

63 The objectives of this study were to investigate antioxidant and antimicrobial activity  
64 of various extracts from *C. cardunculus* involucre bracts, as well as activity of some  
65 compounds previously isolated from therein.

66

## 67 **2. Material and methods**

### 68 *2.1. Chemicals*

69 Dimethylsulfoxide (DMSO), 2,2-diphenyl-1-picrylhydrazyl (DPPH), Folin-Ciocalteu  
70 reagent, 2,4,6-tris(2-pyridyl)-s-triazine (TPTZ) were obtained from Sigma Chemical Co.  
71 (St. Louis, U.S.A.); L-ascorbic acid from Lachema (Neratovice, Czech Republic);  
72 Müller-Hinton agar (MH), Malt agar (MA) from Institute of Immunology and Virology,  
73 Torlak (Belgrade, Serbia); streptomycin (Streptomycin-sulfat, ampoules 1 g) and  
74 miconazole from Galenika, a.d. (Belgrade, Serbia). **Standard** compounds **1-9**, namely:  
75 apigenin (**1**), cynarasaponins A+H (**2**), luteolin 7-glucoside (**3**), apigenin 7-rutinoside (**4**),  
76 luteolin (**5**), chlorogenic acid (**6**),  $\beta$ -sitosterol (**7**), cynarasaponins B+K (**8**) and apigenin

77 7-glucoside (9), were isolated previously from *C. cardunculus* involucre bracts at the  
78 Department of Pharmacognosy and Botany, Pharmaceutical Faculty, Comenius  
79 University, in Bratislava.

80

## 81 2.2. Plant material

82 The whole involucre bracts of *C. cardunculus* were collected from plants grown at  
83 Medicinal Plants Garden in Bratislava. A voucher specimen was deposited at the  
84 Pharmaceutical Faculty, Comenius University, Bratislava.

85

## 86 2.3. Extraction

87 The whole, fresh involucre bracts were cut in pieces and repeatedly extracted with  
88 EtOH (96%, v/v) at room temperature. Aqueous suspension of the concentrated EtOH  
89 extract was partitioned successively with CHCl<sub>3</sub>, EtOAc and *n*-BuOH, leaving residual  
90 water extract. All obtained extracts, including residual water extract, were evaporated till  
91 dryness and used for all investigations.

92

## 93 2.4. Determination of total phenolics content

94 Total phenolics content was determined using Folin-Ciocalteu (FC) reagent as  
95 previously described (Velioglu, Mazza, Gao, & Oomah, 1998). 100 µl of the extract  
96 dissolved in methanol was mixed with 750 µl of FC reagent (previously diluted 10-fold  
97 with distilled water) and allowed to stand at 22 °C for 5 min; 750 µl of Na<sub>2</sub>CO<sub>3</sub> (60 g/l)  
98 solution was added to the mixture. After 90 min the absorbance was measured at 725  
99 nm. Results were expressed as gallic acid equivalents (mg of gallic acid/mg dry weigh  
100 extract).

101

102 *2.5. Antioxidant activity*

103

104 *2.5.1. Thin-layer chromatography*

105 Each extract and previously isolated compounds (**1-9**) were dissolved in appropriate  
106 solvent, applied on silica gel plates (Merck, Darmstadt, Germany), and developed using  
107 different solvent systems: EtOAc/HCOOH/glacial AcOH/water (100:11:11:26, v/v/v/v),  
108 toluene/EtOAc/HCOOH (5:4:1, v/v/v), and toluene/EtOAc (7:3, v/v) system.  
109 Components were detected by spraying with NP/PEG reagent (flavonoids, phenolic  
110 acids) and with vanillin-sulphuric acid (VS) reagent (saponins and sterols) (Wagner &  
111 Bladt, 1996). DPPH test performed directly on TLC plates (0.2% DPPH in MeOH (w/v)  
112 used as spray reagent) revealed contributions to the antioxidant activity of different  
113 compounds separately (Cuendet, Hostettmann, & Potterat, 1997).

114

115 *2.5.2. FRAP assay*

116 Total antioxidant activity (TAA) was investigated using Ferric Reducing Antioxidant  
117 Power (FRAP) assay, which is based upon reduction of Fe<sup>3+</sup>-TPTZ complex in acidic  
118 conditions. Increase in absorbance of blue colored ferrous form (Fe<sup>2+</sup>-TPTZ complex) is  
119 measured at 593 nm. FRAP reagent was freshly prepared by mixing 25 ml acetate buffer  
120 (300 mM, pH 3.6), 2.5 ml TPTZ solution (10 mM TPTZ in 40 mM HCl) and 2.5 ml  
121 FeCl<sub>3</sub> (20 mM) water solution. 100 µl of each extract dissolved in appropriate solvent  
122 was added in 4.5 ml of FRAP reagent, stirred and incubated for 30 min absorbance was  
123 measured at 593 nm, using FRAP working solution as blank. Calibration curve of ferrous  
124 sulfate (100-1000 µM) was used, and results were expressed in µmol Fe<sup>2+</sup>/mg dry weight

125 extract. The relative activity of the samples was compared to L-ascorbic acid (Pellegrini  
126 et al., 2003).

127

### 128 2.5.3. DPPH radical assay

129 Extracts were dissolved in appropriate solvents, mixed with 1 ml of 0.5 mM 2,2-  
130 diphenyl-1-picrylhydrazyl radical (DPPH) in MeOH, and final volume adjusted up to 5  
131 ml. Mixtures were vigorously shaken and left 30 min in dark. Absorbance was measured at  
132 517 nm using MeOH as blank. 1 ml of 0.5 mM DPPH diluted in 4 ml of MeOH was used  
133 as control. Neutralisation of DPPH radical was calculated using the equation:  
134  $S(\%)=100 \times (A_0 - A_s) / A_0$ , where  $A_0$  is the absorbance of the control (containing all reagents  
135 except the test compound), and  $A_s$  is the absorbance of the tested sample. The  $SC_{50}$   
136 value represented the concentration of the extract that caused 50% of neutralisation  
137 (Cuendet et al., 1997). Results were compared with the activity of L-ascorbic acid.

138

## 139 2.6. Bioassays

### 140 2.6.1. Test on antibacterial activity

141 In order to obtain quantitative data for extracts and previously isolated compounds (**1-**  
142 **9**), the modified microdilution technique was used (Hanel & Raether, 1988; Daouk,  
143 Dagher, & Sattout, 1995). The following bacteria were tested: *Salmonella typhimurium*  
144 (ATCC 13311), *Escherichia coli* (ATCC 35210), *Bacillus subtilis* (ATCC 10907),  
145 *Staphylococcus epidermidis* (ATCC 12228) and *Staphylococcus aureus* (ATCC 29213).  
146 The organisms tested were obtained from Department for Plant Physiology, Institute for  
147 Biological Research “Siniša Stanković”, Belgrade, Serbia.

148 The bacterial suspension was adjusted with sterile saline to a concentration of  
149 approximately  $1.0 \times 10^7$  cell/ml. The inocula were stored at +4 °C for further use.



150 Dilutions of the inocula were cultured on solid Müller-Hinton (MH) agar (Institute of  
151 Immunology and Virology, Torlak, Belgrade, Serbia) to verify the absence of  
152 contamination and to check the validity of the inoculum.

153 Minimum inhibitory concentrations (MICs) determination was performed by a serial  
154 dilution technique using 96-well microtitre plates. The bacterial inocula applied contained  
155 approximately  $1.0 \times 10^5$  cells in a final volume of 100  $\mu$ l/well. The extracts and  
156 compounds tested were dissolved in DMSO (0.1-1.0 mg/ml) and added in broth medium  
157 with bacterial inocula. The microplates were incubated for 24 h at 37 °C. The lowest  
158 concentrations without visible growth (at the binocular microscope) were defined as  
159 concentrations which completely inhibited bacterial growth (MICs). The minimum  
160 bactericidal concentrations (MBCs) were determined by serial subcultivation of a 2  $\mu$ l  
161 into microtitre plates containing 100  $\mu$ l of broth per well and further incubation for 24 h  
162 at 37 °C. The lowest concentration with no visible growth was defined as the MBC,  
163 indicating  $\geq 99.5\%$  killing of the original inoculum. DMSO was used as a negative  
164 control, while streptomycin was used as a positive control (0.5-2.0  $\mu$ g/ml). Dilutions of  
165 the inocula were also cultured on solid MH to verify the absence of contamination and to  
166 check their validity.

167

#### 168 2.6.2. Test on antifungal activity

169 Antifungal activity of the extracts and previously isolated compounds (**1-9**) was  
170 investigated using modified microdilution technique (Hanel & Raether, 1988; Daouk et  
171 al., 1995). For the bioassays eight fungi were tested: *Aspergillus niger* (ATCC 6275),  
172 *Aspergillus ochraceus* (ATCC 12066), *Aspergillus flavus* (ATCC 9643), *Penicillium*  
173 *ochrochloron* (ATCC 9112), *Penicillium funiculosum* (ATCC 36839), *Trichoderma*

174 *viride* (IAM 5061), *Fusarium tricinctum* (CBS 514478) and *Alternaria alternata* (DSM  
175 2006). The organisms tested were obtained from the Mycological Laboratory,  
176 Department of Plant Physiology, Institute for Biological research "Siniša Stanković",  
177 Belgrade, Serbia. The micromycetes were maintained on malt agar (MA) the cultures  
178 were stored at +4 °C and subcultured once a month (Booth, 1971).

179 The fungal spores were washed from the surface of agar plates with sterile 0.85%  
180 saline containing 0.1% Tween 80 (v/v). The spore suspension was adjusted with sterile  
181 saline to a concentration of approximately  $1.0 \times 10^5$  in a final volume of 100 µl/well. The  
182 inocula were stored at +4 °C for further use. Dilutions of the inocula were cultured on  
183 solid MA to verify the absence of contamination and to check the validity of the  
184 inoculum.

185 Minimum inhibitory concentrations (MICs) determination was performed by a serial  
186 dilution technique using 96-well microtitre plates. The compounds and extracts  
187 investigated were dissolved in DMSO (0.1 – 1.0 mg/ml) and added in broth malt medium  
188 with fungal inoculum. The microplates were incubated for 72 h at 28 °C. The lowest  
189 concentrations without visible growth (at the binocular microscope) were defined as  
190 MIC. The minimum fungicidal concentrations (MFCs) were determined by serial  
191 subcultivation of a 2 µl into microtitre plates containing 100 µl of malt broth per well and  
192 further incubation for 72 h at 28 °C. The lowest concentration with no visible growth  
193 was defined as the MFC, indicating  $\geq 99.5\%$  killing of the original inoculum. DMSO was  
194 used as a negative control, while miconazole was used as a positive control (0.1 – 5.0  
195 µg/ml).

196

197 **2.7. Statistical analysis**

198 The results of the experiments were analyzed by two factorial analysis of variance  
199 (ANOVA). The Package program Statistica (release 4.5, Copyright StatSoft, Inc. 1993)  
200 was used for statistical evaluation. Antioxidant activity and determination of total  
201 phenolics content were run in triplicates. Experiments on antimicrobial activity were  
202 replicated twice on same occasions. All analyses were run in triplicate for each replicate  
203 ( $n = 2 \times 3$ ).

204

### 205 3. Results

206 Total phenolics content was 0.203, 0.062, 0.050, 0.046 and 0.026 mg of gallic acid  
207 equivalent/mg dry weigh for EtOAc, *n*-BuOH, EtOH, water and CHCl<sub>3</sub> extracts of *C.*  
208 *cardunculus* whole involucral bracts, respectively (Table 1).

209 Total antioxidant activity (TAA) of the investigated extracts was 0.38, 0.36, 0.35, 0.34  
210 and 0.12  $\mu\text{mol Fe}^{2+}$ /mg dry weigh for EtOAc, *n*-BuOH, EtOH, water and CHCl<sub>3</sub>  
211 extracts, respectively. L-Ascorbic acid used as standard had TAA at 7.41  $\mu\text{mol Fe}^{2+}$ /mg  
212 (Table 1). Scavenging of DPPH radical was concentration-dependent. EtOAc extract  
213 expressed the strongest activity ( $\text{SC}_{50}$ =21.50  $\mu\text{g/ml}$ ), while *n*-BuOH, EtOH and water  
214 extracts showed moderate activity ( $\text{SC}_{50}$ =127.10, 157.00 and 173.15  $\mu\text{g/ml}$ ,  
215 respectively). CHCl<sub>3</sub> extract did not reach 50% of DPPH neutralisation at the highest  
216 concentration applied (Table 1).

217 TLC-DPPH test showed that phenolic compounds were the main antioxidant  
218 components in the investigated extracts. The most prominent anti-DPPH zones were  
219 revealed only few seconds after spraying with DPPH reagent, in chromatograms of  
220 EtOAc, *n*-BuOH and EtOH extracts. According to applied standards, main “scavengers”  
221 were apigenin (1), luteolin (5), apigenin 7-glucoside (9), and luteolin 7-glucoside (3)  
222 previously isolated from EtOAc extract (Grančai, Nagy, Suchý, & Ubik, 1993), as well

223 as apigenin 7-rutinoside (4) and chlorogenic acid (6) from *n*-BuOH extract (Grančai,  
224 Mučaji, Nagy, & Ubik, 1996; Mučaji et al., 2000). Cynarasaponins (2, 8) previously  
225 isolated from *n*-BuOH extract (Mučaji, Grančai, Nagy, Buděšinský, & Ubik, 1999;  
226 Mučaji, Grančai, Nagy, Buděšinský, & Ubik, 2001), and  $\beta$ -sitosterol (7) from CHCl<sub>3</sub>  
227 extract (Grančai, Nagy, Suchý, & Ubik, 1992), did not express any scavenging activity.

228 The results of testing of antibacterial activity of *C. cardunculus* extracts showed that  
229 EtOAc extract was the most effective (with MICs of 1.0-1.5 mg/ml and MBCs 1.5-2.0  
230 mg/ml), followed by EtOH, CHCl<sub>3</sub>, water and *n*-BuOH extracts. *S. typhimurium* was  
231 found to be the most resistant species with MICs of 1.5-2.0 mg/ml and MBCs of 2.0-2.5  
232 mg/ml. *E. coli* was the most sensitive with MICs of 1.0-1.5 mg/ml and MBCs of 1.5-2.0  
233 mg/ml. Commercial drug streptomycin showed higher antibacterial potency than extracts  
234 tested (Table 2). Considering antifungal potential of investigated *C. cardunculus*  
235 extracts, EtOAc extract was also the most effective one with values of MICs and MFCs  
236 in equal range of 1.0-1.5 mg/ml (Table 3). Miconazole showed stronger antifungal  
237 activity than extracts tested. As for the standard compounds, the uppermost antibacterial,  
238 as well as the highest antifungal activity was observed by luteolin (5) with MICs and  
239 MBCs ranged from 0.05-0.10 mg/ml, and MICs and MFCs ranged from 0.03-0.10 mg/ml  
240 (Tables 4 and 5).

241

#### 242 4. Discussion

243 Many studies report the polyphenolic composition of cultivated and wild artichokes.  
244 The major class of polyphenols in *C. scolymus* are caffeic acid derivatives (Mulinacci et  
245 al., 2004) which, in heads, mainly occur as esters with quinic acid; leaves and heads of  
246 globe artichoke have been also found to be rich in mono- and dicaffeoylquinic  
247 compounds and flavonoids (Alamanni & Cossu, 2003; Wang, Simon, Fabiola Aviles, He,

248 Zheng, & Tadmor, 2003; Schutz, Kammerer, Carle & Schieber, 2004; Fratianni, Tucci,  
249 De Palma, Pepe, & Nazzaro, 2007; Pinelli et al., 2007). As for cardoon, *C. cardunculus*,  
250 there are reports on phenolic composition of their leaves: caffeoylquinic acids and  
251 glycosides of luteolin and apigenin were identified using HPLC (Valentao et al., 2002;  
252 Pinelli et al., 2007). In the involucre bracts of this plant various compounds were also  
253 identified:  $\beta$ -sitosterol, sitosteryl-3 $\beta$ -glucoside, sitosteryl-3 $\beta$ -acetate, taraxasterole and  
254 taraxasteryl-3 $\beta$ -acetate (Grančai et al., 1992), apigenin, apigenin 7-glucoside, luteolin  
255 and luteolin 7-glucoside (Grančai et al., 1993), apigenin 7-rutinoside, luteolin 7-  
256 rutinoside (Grančai et al., 1996), and apigenin 7-methylglucuronide (Mučaji, et al.,  
257 2000), scopolin and scopoletin (Grančai, Nagy, Mučaji, Suchý, & Ubik, 1994a), cynarin  
258 (Grančai, Nagy, Suchý, & Novomeský, 1994b) and chlorogenic acid (Mučaji et al.,  
259 2000), cynarasaponins A and H, and their methyl derivatives (Mučaji et al., 1999), and  
260 cynarasaponins B and K (Mučaji et al., 2001).

261 As previously showed, apigenin, luteolin and their glycosides are powerful antioxidants  
262 (Kwon, Kim, Kim, Kim, & Kim, 2002; Müller, Vasconcelos, Coelho, & Biavatti, 2005).  
263 The antioxidant effectiveness of apigenin was determined in models such as the *in vitro*  
264 lipoprotein oxidation model (Vinson, Dabbagh, Serry, & Jang, 1995). The antioxidant  
265 properties of luteolin 7-glucoside and of the respective aglycon, luteolin, have already  
266 been observed against low-density lipoprotein oxidation (Brown & Rice-Evans, 1998),  
267 DPPH free radical scavenging activity and ABTS<sup>•+</sup> radical cation scavenging effect  
268 (Wang et al., 1998).

269 Chlorogenic acid is one of the most abundant phenolic acids in various plant extracts  
270 and also the most active antioxidant constituent. It has been shown that the antioxidant  
271 activities of 3-*O*-caffeoylquinic acid and 4-*O*-caffeoylquinic acid are almost the same as  
272 chlorogenic acid when assayed for scavenging activity on superoxide anion radicals and

273 inhibitory effect against oxidation of methyl linoleate (Takeoka & Dao, 2003). 3,4-Di-*O*-  
274 caffeoylquinic acid, 3,5-di-*O*-caffeoylquinic acid, along with chlorogenic acid inhibited  
275 lipid peroxidation and exhibit neuroprotective activities (Nakajima, Shimazawa, Mishima,  
276 & Hara, 2007).

277  $\beta$ -Sitosterol generally showed low antioxidant activity, comparing to different  
278 phenolics such as flavonoids, caffeic and chlorogenic acid, but it exhibited a higher lipid  
279 peroxidation inhibition rate (Yokota et al., 2006). Antioxidant activity of  $\beta$ -sitosterol  
280 determined by the oxidative stability instrument (OSI) was considerable (Weng & Wang,  
281 2000), and even much stronger than that of  $\alpha$ -tocopherol (Jiang & Wang, 2006). It was  
282 suggested that  $\beta$ -sitosterol, which inhibits active oxygen produced by neutrophils, exerts  
283 its antioxidative action through a preventive action, such as stabilization of the cell  
284 membrane. Caffeic acid derivatives and polyphenols that capture hydroxyl and  
285 superoxyde anion radicals act as radical scavengers, while  $\beta$ -sitosterol exerts a preventive  
286 action by inhibiting the excess production of active oxygen by various cells (Yokota et  
287 al., 2006).

288 Results of our experiments are consistent with previous data reported (Alamanni &  
289 Cossu, 2003). As main antioxidant compounds in investigated *C. cardunculus* extracts  
290 we identified flavones: apigenin and luteolin, and their glycosides, as well as chlorogenic  
291 acid. The highest antioxidant activity of the EtOAc extract could be explained, among  
292 other, by presence of apigenin and luteolin in significantly larger amount than in other  
293 extracts.

294 Our experiments presented substantial antimicrobial activity of *C. cardunculus*  
295 involucre bracts extracts with MICs, MBCs and MFCs of 1.00-2.50 mg/ml. EtOAc  
296 extract was again the most effective.

297 Mossi and Echeverrigaray (1999) found that CH<sub>2</sub>Cl<sub>2</sub> *C. scolyumus* leaf extract, in  
298 concentrations of 5 mg/ml, completely inhibited the growth with a bactericidal effect on  
299 *Staphylococcus aureus*, *Bacillus cereus* and *B. subtilis*. Zhu, Zhang, and Lo (2004)  
300 investigated antimicrobial activity of different extracts of *C. scolyumus* leaf and showed  
301 that the *n*-BuOH fraction was the most active one, followed by CHCl<sub>3</sub> and EtOAc  
302 fractions. Similar investigations were done with successive CHCl<sub>3</sub>, EtOH, and EtOAc  
303 partitions of extracts of *C. scolyumus* leaf, head, and stem. The MIC values for fungi were  
304 at or below 2.5 mg/ml and for bacteria were at or above 2.5 mg/ml (Zhu, Zhang, Lo, &  
305 Lu, 2005).

306 The results of our experiment showed that all standard compounds, previously  
307 isolated from involucre bracts of *C. cardunculus*, possess antimicrobial activity against  
308 all tested strains of bacteria and fungi (MICs, MBCs and MFCs in a range of 0.03-0.10  
309 mg/ml). Among them, luteolin showed the best activity.

310 Similar results were also previously observed with compounds isolated from *C.*  
311 *scolymus* leaves. Among them, chlorogenic acid, cynarin, luteolin 7-rutinoside, and  
312 cynaroside exhibited a relatively higher activity than other compounds and were more  
313 effective against fungi than against bacteria, with MICs ranged from 0.05-0.20 mg/ml  
314 (Zhu et al., 2004). Antimicrobial activity of apigenin, apigenin 7-glucoside, luteolin and  
315 other flavones has been also previously reported (Aljančić et al., 1999; Tshikalange,  
316 Meyer, & Hussein, 2005).

317 Herein obtained results on antioxidant and antimicrobial activity of different extracts of  
318 *C. cardunculus* involucre bracts supported the traditional medicinal use of this plant and  
319 provided grounds for its further establishing as a functional food.

320     **ABBREVIATIONS USED:** ATCC, American Type of Culture Collection; *n*-BuOH,  
321     *n*-butanol; CBS, Centraalbureau voor Schimmelcultures, Baarn, The Netherlands;  
322     CHCl<sub>3</sub>, chloroform; DMSO, dimethylsulfoxide; DPPH, 2,2-diphenyl-1-picrylhydrazyl;  
323     DSM, Deutsche Sammlung von Mikroorganismen; EDTA, ethylenediaminetetraacetic  
324     acid; EtOAc, ethyl acetate; EtOH, ethanol; FC reagent, Folin-Ciocalteu reagent; FRAP  
325     assay, Ferric reducing antioxidant power assay; IAM, Institute of Applied Microbiology,  
326     University of Tokyo, Japan; MA, malt agar; MBC, minimum bactericidal concentration;  
327     MeOH, methanol; MFC, minimum fungicidal concentration; MH, Müller-Hinton; MIC,  
328     minimum inhibitory concentration; NP/PEG reagent, natural products-polyethylene  
329     glycol reagent; TAA, total antioxidant activity; TPTZ, 2,4,6-tris(2-pyridyl)-*s*-triazine.

330

331     **ACKNOWLEDGMENT.** This research was supported by the Ministry of Science  
332     and Environmental Protection of Republic of Serbia (Grants № 143012 and 143041).



333        **LITERATURE CITED**

- 334        Alamanni, M. C., & Cossu, M. (2003). Antioxidant activity of the extracts of the edible  
335        part of artichoke (*Cynara scolimus* L.) var. spinoso sardo. *Italian Journal of Food*  
336        *Science*, *15*, 187-195.
- 337        Aljančić, I., Vajs, V., Menković, N., Karadžić, I., Juranić, N., Milosavljević, S., &  
338        Macura, S. (1999). Flavones and sesquiterpene lactones from *Achillea atrata* subsp.  
339        *multifida*: antimicrobial activity. *Journal of Natural Products*, *62*, 909-911.
- 340        Booth, C. (1971). Fungal Culture Media. In J. R. Norris, & D. W. Ribbons, *Methods in*  
341        *Microbiology*, (pp. 49-94). London & New York: Academic Press.
- 342        Brown, J. E., & Rice-Evans, C. A. (1998). Luteolin-rich artichoke extract protects low-  
343        density lipoprotein from oxidation *in vitro*. *Free Radical Research*, *29*, 247-255.
- 344        Cuendet, M., Hostettmann, K., & Potterat, O. (1997). Iridoid glucosides with free  
345        radical scavenging properties from *Fagraea blumei*. *Helvetica Chimica Acta*, *80*,  
346        1144–1152.
- 347        Daouk, R. K., Dagher, S. M., & Sattout, J. E. (1995). Antifungal activity of the Essential  
348        oil of *Origanum syriacum* L. *Journal of Food Protection*, *58*, 1147-1149.
- 349        do Amaral Franco, J. (1976). *Cynara* L. In T. G. Tutin, V. H. Heywood, N. A. Burges,  
350        D. M. Moore, D. H. Valentine, S. M. Walters, & D. A. Webb, *Flora Europaea*, vol.  
351        4 (pp 248-249). Cambridge: Cambridge University Press.
- 352        Fernandez, J., Curt, M. D., & Aguado, P. L. (2006). Industrial applications of *Cynara*  
353        *cardunculus* L. for energy and other uses. *Industrial Crops and Products*, *24*, 222-  
354        229.

- 355 Fratianni, F., Tucci, M., De Palma, M., Pepe, R., & Nazzaro, F. (2007). Polyphenolic  
356 composition in different parts of some cultivars of globe artichoke (*Cynara*  
357 *cardunculus* L. var. *scolymus* (L.) Fiori). *Food Chemistry*, *104*, 1282-1286.
- 358 Grančai, D., Nagy, M., Suchý, V., & Ubik, K. (1992). Constituents of *Cynara*  
359 *cardunculus* L. I. Sterols and pentacyclic triterpens. *Farmaceutický Obzor*, *61*, 577-  
360 580.
- 361 Grančai, D., Nagy, M., Suchý, V., & Ubik, K. (1993). Constituents of *Cynara*  
362 *cardunculus* L. II. Flavonoids. *Farmaceutický Obzor*, *62*, 31-33.
- 363 Grančai, D., Nagy, M., Mučaji, P., Suchý, V., & Ubik, K. (1994a). Constituents of  
364 *Cynara cardunculus* L. III. Coumarins. *Farmaceutický obzor*, *63*, 447-449.
- 365 Grančai, D., Nagy, M., Suchý, V., & Novomeský, P. (1994b). Cynarin from the fresh  
366 flower buds of *Cynara cardunculus*. *Fitoterapia*, *65*, 282.
- 367 Grančai, D., Mučaji, P., Nagy, M., & Ubik, K. (1996). Constituents of *Cynara*  
368 *cardunculus* L. IV. Flavonoid glycosides. *Farmaceutický obzor*, *65*, 255-256.
- 369 Grieve M. (1971). *A Modern Herbal*. New York: Dover Publications, Inc.
- 370 Hanel, H., & Raether, W. (1998). A more sophisticated method of determining the  
371 fungicidal effect of water-insoluble preparations with a cell harvester, using  
372 miconazole as an example. *Mycoses*, *31*, 148-154.
- 373 Jiang, A., & Wang, C. (2006). Antioxidant properties of natural components from *Salvia*  
374 *plebeia* on oxidative stability of ascidian oil. *Process Biochemistry*, *41*, 1111-1116.

- 375 Kelly, M., & Pepper, A. (1996). Controlling *Cynara cardunculus* (Artichoke Thistle,  
376 Cardoon, etc.). In J. E. Lovich, J. Randall, & M. D. Kelly, *Proceedings of the*  
377 *California Exotic Pest Plant Council Symposium*, vol. 2 (pp. 97-101). San Diego:  
378 California Exotic Pest Plant Council.
- 379 Koubaa, I., Damak, M., McKillop, A., & Simmonds, M. (1999). Constituents of *Cynara*  
380 *cardunculus*. *Fitoterapia*, 70, 212-213.
- 381 Koubaa, I., & Damak, M. (2003). A new dilignan from *Cynara cardunculus*.  
382 *Fitoterapia*, 74, 18-22.
- 383 Križkova, L., Mučaji, P., Nagy, M., & Krajčovič, J. (2004). Triterpenoid cynarasaponins  
384 from *Cynara cardunculus* L. reduce chemicaly induced mutagenesis *in vitro*.  
385 *Phytomedicine*, 11, 673-678.
- 386 Kwon, Y. S., Kim, E. Y., Kim, W. J., Kim, W. K., & Kim, C. M. (2002). Antioxidant  
387 constituents from *Setaria viridis*. *Archives of Pharmacal Research* 25, 300-305.
- 388 Mossi, A. J., & Echeverrigaray, S. (1999). Identification and characterization of  
389 antimicrobial components in leaf extracts of globe artichoke (*Cynara scolymus* L.).  
390 *Acta Horticulturae*, 501, 111-114.
- 391 Mučaji, P., Grančai, D., Nagy, M., Buděšínský, M., & Ubik, K. (1999). Triterpenoid  
392 saponins from *Cynara cardunculus* L. *Pharmazie*, 54, 714-716.
- 393 Mučaji, P., Grančai, D., Nagy, M., Višňovská, Z., & Ubik, K. (2000). Apigenin-7-  
394 methylglucuronide from *Cynara cardunculus* L. *Česká a slovenská farmacie*, 49,  
395 75-77.

- 396 Mučaji, P., Grančai, D., Nagy, M., Buděšínský, M., & Ubik, K. (2001). Monodesmosidic  
397 saponins from *Cynara cardunculus* L. *Česká a slovenská farmacie*, 50, 277-279.
- 398 Mučaji, P., Bukovsky, M., Grančai, D., & Nagy, M. (2003). Anticomplement activity of  
399 saponins from *Cynara cardunculus* L. *Česká a slovenská farmacie*, 52, 306-309.
- 400 Mulinacci, N., Prucher, D., Peruzzi, M., Romani, A., Pinelli, P., Giaccherini, C., &  
401 Vincieri, F. F. (2004). Commercial and laboratory extracts from artichoke leaves:  
402 estimation of caffeoyl esters and flavonoidic compounds content. *Journal of*  
403 *Pharmaceutical and Biomedical Analysis*, 34, 349-357.
- 404 Müller, S. D., Vasconcelos, S. B., Coelho, M., & Biavatti, M. W. (2005). LC and UV  
405 determination of flavonoids from *Passifora alata* medicinal extracts and leaves.  
406 *Journal of Pharmaceutical and Biomedical Analysis*, 37, 399-403.
- 407 Nakajima, Y., Shimazawa, M., Mishima, S., & Hara, H. (2007). Water extract of  
408 propolis and its main constituents, caffeoylquinic acid derivatives, exert  
409 neuroprotective effects via antioxidant actions. *Life Sciences*, 80, 370-377.
- 410 Paris, R., & Moyse, H. (1971). *Précis de matière médicale, tome III*. Paris: Masson et  
411 C<sup>ie</sup>.
- 412 Pellegrini, N., Serafini, M., Colombi, B., Del Rio, D., Salvatore, S., Bianchi, M., &  
413 Brighenti, F. (2003). Total antioxidant capacity of plant foods, beverages and oils  
414 consumed in Italy assessed by three different *in vitro* assays. *Journal of Nutrition*,  
415 133, 2812-2818.

416 Pinelli, P., Agostini, F., Comino, C., Lanteri, S., Portis, E., & Romani, A. (2007).  
417 Simultaneous quantification of caffeoyl esters and flavonoids in wild and cultivated  
418 cardoon leaves, *Food Chemistry*, doi: 10.1016/j.foodchem.2007.05.014.

419 Rossoni, G., Grande, S., Galli, C., & Visioli, F. (2005). Wild artichoke prevents age  
420 associated loss of vasomotor function. *Journal of Agricultural and Food Chemistry*,  
421 53, 10291-10296.

422 Schutz, K., Kammerer, D., Carle, R., & Schieber, A. (2004). Identification and  
423 quantification of caffeoylquinic acids and flavonoids from artichoke (*Cynara*  
424 *scolymus* L.) heads, juice, and pomace by HPLC-DAD-ESI/MSn. *Journal of*  
425 *Agricultural and Food Chemistry*, 52, 4090-4096.

426 Silva, S. V., & Malcata, F. X. (2005). Studies pertaining to coagulant and proteolytic  
427 activities of plant proteases from *Cynara cardunculus*. *Food Chemistry*, 89, 19-26.

428 Slanina, J., Taborska, E., Bochorakowa, H., Humpa, O., Robinson, W. E., & Schram, K.  
429 H. (2001). New and facile method of preparation of the anti-HIV agent 1,3-  
430 dicaffeoylquinic acid. *Tetrahedron Letters*, 42, 3383-3385.

431 Sroka, Z., & Cisowski, W. (2003). Hydrogen peroxide scavenging, antioxidant and anti-  
432 radical activity of some phenolic acids. *Food and Chemical Toxicology*, 41, 753-  
433 758.

434 Ševčíková, P., Glatz, Z., & Slanina, J. (2002). Analysis of artichoke extracts (*Cynara*  
435 *cardunculus* L.) by means of micellar electrokinetics capillary chromatography.  
436 *Electrophoresis*, 23, 249-252.

- 437 Takeoka, G. R., & Dao, L. T. (2003). Antioxidant constituents of almond [*Prunus dulcis*  
438 (Mill.) D.A. Webb] hulls. *Journal of Agricultural and Food Chemistry*, 51, 496-  
439 501.
- 440 Tshikalange, T. E., Meyer, J. J. M., & Hussein, A. A. (2005). Antimicrobial activity,  
441 toxicity and the isolation of a bioactive compound from plants used to treat sexually  
442 transmitted diseases. *Journal of Ethnopharmacology* 96, 515–519.
- 443 Valentao, P., Fernandes, E., Carvalho, F., Andrade, P. B., Seabra, R. M., & Bastos, M.  
444 L. (2002). Antioxidative properties of cardoon (*Cynara cardunculus* L.) infusion  
445 against superoxide radical, hydroxyl radical and hypochlorous acid. *Journal of*  
446 *Agricultural and Food Chemistry*, 50, 4989-4993.
- 447 Velioglu, Y. S., Mazza, G., Gao, L., & Oomah, B. D. (1998). Antioxidant activity and  
448 total phenolics in selected fruits, vegetables, and grain products. *Journal of*  
449 *Agricultural and Food Chemistry*, 46, 4113-4117.
- 450 Vinson, J. A., Dabbagh, Y. A., Serry, M. M., & Jang, J. (1995). Plant flavonoids,  
451 especially tea flavonols, are powerful antioxidants using an in vitro oxidation model  
452 for heart disease. *Journal of Agricultural and Food Chemistry*, 43, 2800-2802.
- 453 Wagner, H., & Bladt, S. (1996). *Plant Drug Analysis. A Thin Layer Chromatography*  
454 *Atlas*, 2nd edition. Berlin-Heidelberg: Springer-Verlag.
- 455 Wang, M., Li, J., Rangarajan, M., Shao, Y., La Voie, E. J., Huang, T.-C., & Ho, C.-T.  
456 (1998). Antioxidative phenolic compounds from sage (*Salvia officinalis*). *Journal of*  
457 *Agricultural and Food Chemistry*, 46, 4869-4873.

- 458 Wang, M., Simon, J. E., Fabiola Aviles, I., He, K., Zheng, Q.-Y., & Tadmor, Y. (2003).  
459 Analysis of antioxidative phenolic compounds in artichoke (*Cynara scolymus* L.).  
460 *Journal of Agricultural and Food Chemistry*, *51*, 601-608.
- 461 Weng, X. C., & Wang, W. (2000). Antioxidant activity of compounds isolated from  
462 *Salvia plebeia*. *Food Chemistry*, *71*, 489-493.
- 463 Zhu, X., Zhang, H., & Lo, R. (2004). Phenolic compounds from the leaf extract of  
464 artichoke (*Cynara scolymus* L.) and their antimicrobial activities. *Journal of*  
465 *Agricultural and Food Chemistry*, *52*, 7272-7278.
- 466 Zhu, X., Zhang, H., Lo, R., & Lu, Y. (2005). Antimicrobial activities of *Cynara*  
467 *scolymus* L. leaf, head, and stem extracts. *Journal of Food Science*, *70*, M149-  
468 M152.

469 **Table 1.** Antioxidant activity and total phenolics content of *Cynara cardunculus* extracts

Extract	FRAP value <sup>a</sup>	DPPH scavenging <sup>b</sup>	Total phenolics content <sup>c</sup>
EtOAc	0.38 ± 0.01	21.50 ± 1.87	0.203 ± 0.018
BuOH	0.36 ± 0.01	127.10 ± 0.88	0.062 ± 0.019
EtOH	0.35 ± 0.01	157.00 ± 0.16	0.050 ± 0.010
H <sub>2</sub> O	0.34 ± 0.01	173.15 ± 0.65	0.046 ± 0.007
CHCl <sub>3</sub>	0.12 ± 0.02	-	0.026 ± 0.002
L-ascorbic acid	7.41 ± 0.05	4.09 ± 0.08	-

470 <sup>a</sup> in μmol Fe<sup>2+</sup>/mg dry weigh extract, <sup>b</sup> SC<sub>50</sub>, μg/ml, <sup>c</sup> mg of gallic acid equivalent/mg  
 471 dry weigh extract



472 **Table 2.** Minimum inhibitory and bactericidal concentrations (MICs and MBCs) of  
 473 *Cynara cardunculus* extracts (mg/ml)

Bacteria		E x t r a c t s					Streptomycin
		BuOH	EtOH	EtOAc	CHCl <sub>3</sub>	H <sub>2</sub> O	
<i>S. typhimurium</i>	MIC	2.0±0.2	1.5±0.0	1.5±0.0	2.0±0.2	2.0±0.1	0.0010±0.0002
	MBC	2.5±0.3	2.0±0.2	2.0±0.2	2.5±0.3	2.0±0.1	0.0010±0.0002
<i>E. coli</i>	MIC	1.0±0.0	1.0±0.0	1.0±0.0	1.5±0.0	1.0±0.0	0.0005±0.0001
	MBC	1.5±0.2	1.5±0.0	1.5±0.0	2.0±0.2	1.5±0.1	0.0010±0.0002
<i>S. epidermidis</i>	MIC	1.5±0.0	1.5±0.0	1.5±0.1	1.5±0.0	1.5±0.0	0.0010±0.0000
	MBC	2.0±0.2	2.0±0.2	2.0±0.2	2.0±0.2	2.0±0.2	0.0010±0.0000
<i>S. aureus</i>	MIC	1.5±0.0	1.5±0.0	1.5±0.0	1.5±0.0	1.5±0.0	0.0010±0.0002
	MBC	2.0±0.2	2.0±0.2	2.0±0.2	2.0±0.2	2.0±0.2	0.0010±0.0003
<i>B. subtilis</i>	MIC	2.0±0.2	2.0±0.2	1.0±0.0	1.5±0.0	1.0±0.0	0.0005±0.0000
	MBC	2.5±0.0	2.0±0.0	1.0±0.0	2.0±0.2	1.0±0.0	0.0005±0.0002

474

475

476 **Table 3.** Minimum inhibitory and fungicidal concentrations (MICs and MFCs) of *Cynara*  
 477 *cardunculus* extracts (mg/ml)

Fungal species		E x t r a c t s					Miconazole
		BuOH	EtOH	EtOAc	CHCl <sub>3</sub>	H <sub>2</sub> O	
<i>A. flavus</i>	MIC	1.5±0.2	1.5±0.0	1.5±0.0	1.5±0.2	1.5±0.2	0.0005±0.0000
	MFC	1.5±0.2	1.5±0.0	1.5±0.2	1.5±0.0	2.0±0.0	0.0020±0.0002
<i>A. niger</i>	MIC	1.5±0.2	1.5±0.2	1.5±0.2	1.5±0.2	1.5±0.2	0.0015±0.0003
	MFC	1.5±0.2	1.5±0.0	1.5±0.0	1.5±0.0	2.0±0.2	0.0040±0.0002
<i>A. ochraceus</i>	MIC	1.0±0.0	1.0±0.0	1.0±0.0	1.0±0.0	1.0±0.0	0.0015±0.0002
	MFC	1.5±0.2	1.5±0.2	1.5±0.2	1.5±0.0	1.0±0.0	0.0040±0.0004
<i>P. funiculosum</i>	MIC	1.5±0.1	1.5±0.2	1.0±0.0	1.0±0.0	1.0±0.0	0.0020±0.0000
	MFC	1.5±0.2	1.5±0.2	1.5±0.2	1.5±0.2	1.5±0.2	0.0050±0.0000
<i>P. ochrachloron</i>	MIC	1.0±0.0	1.0±0.0	1.0±0.0	1.0±0.0	1.0±0.0	0.0020±0.0002
	MFC	1.5±0.2	1.5±0.1	1.5±0.2	1.5±0.2	1.5±0.2	0.0050±0.0000
<i>T. viride</i>	MIC	1.5±0.2	1.5±0.2	1.0±0.0	1.0±0.0	1.0±0.0	0.0020±0.0000
	MFC	1.5±0.0	1.5±0.2	1.0±0.0	1.5±0.2	1.0±0.0	0.0020±0.0000
<i>F. tricinctum</i>	MIC	1.5±0.0	1.5±0.2	1.0±0.0	1.5±0.2	1.0±0.2	0.0002±0.0000
	MFC	1.5±0.0	1.5±0.2	1.5±0.0	1.5±0.2	1.5±0.2	0.0010±0.0002
<i>A. alternata</i>	MIC	1.5±0.2	1.5±0.2	1.0±0.0	1.5±0.2	1.0±0.0	0.0002±0.0000
	MFC	1.5±0.0	1.5±0.1	1.0±0.0	1.5±0.0	1.0±0.0	0.0010±0.0002

478 **Table 4.** Minimum inhibitory and bactericidal concentrations (MICs and MBCs) of the compounds tested\* (mg/ml)

Bacteria		1	2	3	4	5	6	7	8	9	Streptomycin
<i>S. typhimurium</i>	MIC	0.10±0.02	0.10±0.00	0.10±0.00	0.10±0.02	0.05±0.00	0.10±0.02	0.10±0.00	0.10±0.00	0.10±0.02	0.0010±0.0002
	MBC	0.10±0.02	0.10±0.00	0.10±0.00	0.10±0.02	0.05±0.00	0.10±0.02	0.10±0.00	0.10±0.00	0.10±0.02	0.0010±0.0002
<i>E. coli</i>	MIC	0.10±0.00	0.10±0.02	0.10±0.02	0.10±0.00	0.05±0.02	0.10±0.02	0.10±0.02	0.10±0.02	0.10±0.00	0.0005±0.0001
	MBC	0.10±0.00	0.10±0.02	0.10±0.00	0.10±0.00	0.10±0.00	0.10±0.02	0.10±0.02	0.10±0.02	0.10±0.00	0.0010±0.0002
<i>S. epidermidis</i>	MIC	0.15±0.02	0.15±0.00	0.15±0.02	0.15±0.02	0.10±0.01	0.15±0.00	0.15±0.01	0.15±0.01	0.15±0.02	0.0010±0.0000
	MBC	0.20±0.00	0.20±0.02	0.20±0.02	0.20±0.01	0.10±0.02	0.20±0.02	0.20±0.02	0.20±0.02	0.20±0.02	0.0010±0.0000
<i>S. aureus</i>	MIC	0.15±0.00	0.15±0.02	0.15±0.02	0.15±0.01	0.05±0.00	0.15±0.02	0.15±0.02	0.15±0.02	0.15±0.01	0.0010±0.0002
	MBC	0.20±0.02	0.20±0.00	0.20±0.02	0.20±0.01	0.05±0.00	0.20±0.02	0.20±0.00	0.20±0.02	0.20±0.02	0.0010±0.0003
<i>B. subtilis</i>	MIC	0.15±0.02	0.15±0.00	0.15±0.00	0.15±0.02	0.05±0.00	0.10±0.00	0.10±0.02	0.10±0.00	0.10±0.00	0.0005±0.0000
	MBC	0.20±0.02	0.20±0.02	0.20±0.02	0.20±0.00	0.10±0.02	0.15±0.02	0.15±0.02	0.15±0.00	0.15±0.02	0.0005±0.0002

479 \* apigenin (1), cynarasaponins A+H (2), luteolin 7-glucoside (3), apigenin 7-rutinoside (4), luteolin (5), chlorogenic acid (6), β-  
 480 sitosterol (7), cynarasaponins B+K (8) and apigenin 7-glucoside (9)

481

482 **Table 5.** Minimum inhibitory and fungicidal concentrations (MICs and MFCs) of the compounds tested\* (mg/ml)

Fungal species		1	2	3	4	5	6	7	8	9	Miconazole
<i>A. flavus</i>	MIC	0.10±0.02	0.10±0.02	0.10±0.01	0.10±0.02	0.10±0.02	0.10±0.02	0.10±0.02	0.10±0.02	0.10±0.02	0.0005±0.0000
	MFC	0.10±0.02	0.10±0.01	0.10±0.01	0.10±0.02	0.10±0.02	0.10±0.02	0.10±0.02	0.10±0.02	0.10±0.02	0.0020±0.0002
<i>A. niger</i>	MIC	0.10±0.01	0.10±0.02	0.10±0.02	0.10±0.02	0.10±0.02	0.10±0.02	0.10±0.02	0.10±0.02	0.10±0.02	0.0015±0.0003
	MFC	0.10±0.01	0.10±0.02	0.10±0.02	0.10±0.02	0.10±0.02	0.10±0.02	0.10±0.02	0.10±0.02	0.10±0.02	0.0040±0.0002
<i>A. ochraceus</i>	MIC	0.10±0.00	0.10±0.02	0.10±0.00	0.10±0.01	0.05±0.00	0.10±0.02	0.10±0.02	0.10±0.02	0.10±0.00	0.0015±0.0002
	MFC	0.10±0.00	0.10±0.02	0.10±0.00	0.10±0.02	0.10±0.00	0.10±0.01	0.10±0.02	0.10±0.02	0.10±0.02	0.0040±0.0004
<i>P. funiculosum</i>	MIC	0.05±0.02	0.10±0.02	0.10±0.02	0.05±0.00	0.03±0.00	0.05±0.00	0.05±0.02	0.05±0.02	0.05±0.01	0.0020±0.0000
	MFC	0.10±0.02	0.10±0.02	0.10±0.02	0.10±0.02	0.05±0.00	0.10±0.00	0.05±0.02	0.05±0.02	0.05±0.02	0.0050±0.0000
<i>P. ochrachloron</i>	MIC	0.05±0.00	0.10±0.02	0.10±0.02	0.05±0.00	0.05±0.00	0.05±0.00	0.05±0.02	0.05±0.02	0.05±0.01	0.0020±0.0002
	MFC	0.10±0.01	0.10±0.02	0.10±0.02	0.10±0.00	0.05±0.00	0.10±0.01	0.05±0.02	0.05±0.02	0.05±0.01	0.0050±0.0000
<i>T. viride</i>	MIC	0.05±0.00	0.05±0.00	0.05±0.00	0.05±0.00	0.03±0.00	0.05±0.00	0.05±0.00	0.05±0.01	0.05±0.02	0.0020±0.0000
	MFC	0.05±0.00	0.10±0.02	0.10±0.01	0.05±0.00	0.05±0.00	0.10±0.01	0.10±0.00	0.10±0.02	0.10±0.02	0.0020±0.0000
<i>F. tricinctum</i>	MIC	0.05±0.02	0.05±0.00	0.05±0.00	0.05±0.02	0.05±0.00	0.05±0.02	0.05±0.02	0.05±0.00	0.05±0.00	0.0002±0.0000
	MFC	0.10±0.02	0.10±0.02	0.10±0.00	0.10±0.02	0.10±0.01	0.10±0.02	0.10±0.02	0.10±0.00	0.10±0.02	0.0010±0.0002
<i>A. alternata</i>	MIC	0.10±0.02	0.10±0.02	0.10±0.02	0.05±0.02	0.05±0.02	0.10±0.02	0.10±0.01	0.10±0.02	0.10±0.02	0.0002±0.0000
	MFC	0.10±0.02	0.10±0.02	0.10±0.02	0.10±0.01	0.10±0.02	0.10±0.02	0.10±0.02	0.10±0.02	0.10±0.02	0.0010±0.0002

483 \* apigenin (1), cynarasaponins A+H (2), luteolin 7-glucoside (3), apigenin 7-rutinoside (4), luteolin (5), chlorogenic acid (6), β-  
484 sitosterol (7), cynarasaponins B+K (8) and apigenin 7-glucoside (9)