

TITLE: Tracking patterns of body mass index and triceps skinfold thickness from childhood to young adulthood: a 12-year prospective cohort study in Slovenia

Authors:

Gregor Starc¹

Janko Strel¹

¹ Faculty of Sport, University of Ljubljana, Slovenia

Correspondence:

Gregor Starc

Faculty of Sport, University of Ljubljana

Gortanova 22

SI-1000 Ljubljana

Slovenia

E-mail: gregor.starc@guest.arnes.si

Mobile phone: ++386 (0)40 265 765

Fax: ++386 (0)1 520 77 40

Sources of support: The SLOFIT monitoring system is financed by the Slovenian Ministry of Education and Sport and managed by the Faculty of Sport, University of Ljubljana.

Running head: Tracking patterns of body mass index and triceps skinfold thickness from childhood to young adulthood

1 **ABSTRACT**

2 **OBJECTIVE:** To examine tracking patterns of body mass index (BMI), triceps skinfold
3 thickness (TSF), height and weight from childhood to young adulthood in a rapidly
4 developing society.

5 **DESIGN:** Prospective 12-year cohort study of the Slovenian children, born during 1990-
6 1991.

7 **SUBJECTS:** A subsample of 4 833 Slovenian children from the 1990-1991 national cohort
8 ($n = 21\ 777$) who were included in the SLOFIT monitoring system from 1997 to 2008, with
9 complete data at age 7 and 18 y.

10 **MEASUREMENTS:** Height, weight and TSF were measured at ages 7, 11, 14 and 18 y. The
11 IOTF BMI cutoff points were used to identify overweight and obesity.

12 **RESULTS:** Height, weight, and BMI at age 18 y were well predicted from childhood and
13 grew more predictable with age, while TSF was not. Obese and overweight children had the
14 greatest risk of becoming overweight or obese young adults, since tracking of overweight and
15 obesity showed that 53.9% of overweight and 77.7% of obese 7-y old males remained
16 overweight or obese at age 18 y in comparison with 32.1% of overweight and 62% of obese
17 7-y old females. History of obese 18-y olds showed that 40% of males and 48.6% of females
18 had been obese already at age 7 y.

19 **CONCLUSION:** The study confirmed the tracking of BMI from childhood to young
20 adulthood. Overweight or obese Slovenian children are very likely to become overweight or
21 obese adolescents and young adults, which calls for the need of early prevention and
22 treatment of overweight and obesity.

23

24 **KEY WORDS** Child, young adult, body mass index, triceps skinfold thickness,
25 obesity, tracking

1 INTRODUCTION

2 In the last decades overweight and obesity became acknowledged as one of the fastest
3 growing nontransmissible health problems worldwide.¹ An increase of the prevalence of
4 overweight and obesity is a particular concern since it proves to be an important factor for
5 numerous chronic diseases and other health disorders among adults²⁻⁸ as well as among
6 children.⁹⁻¹⁴ Consequently, obesity does not only worsen the quality of life of affected
7 individuals but also increases public and personal health expenditures.¹⁵⁻¹⁹ Obesity is a
8 burdening factor in all respects, it should become a public health priority in the developed and
9 developing countries and it is from this perspective of prevention that the relation between
10 childhood obesity and young adult obesity has been investigated. Several studies showed a
11 tendency of those who are overweight or obese as children to remain overweight or obese as
12 adolescents and adults.²⁰⁻²⁸ Although the evidence is extensive, not many studies have been
13 based on large and unselected population samples or have used other predictors than height
14 and weight. Furthermore, much of the research has been conducted in high-income and
15 predominantly western countries while very limited data is available for middle- or low-
16 income countries and post-socialist countries (like Slovenia), which quickly and abruptly
17 adopted the capitalist economy and consumerist lifestyles, including eating habits and
18 physical activity patterns. Longitudinal research on these topics from rapidly changing
19 societies might provide new insights, enable comparison and consequent prediction of obesity
20 trends among children and adults in these societies.

21 This study focused on tracking of height, weight, triceps skinfold thickness (TSF) and body
22 mass index (BMI) from childhood to young adulthood among the Slovenian schoolchildren
23 who entered primary school in the school year 1996/97 and turned 18 y of age in the school
24 year 2007/08. We used longitudinal data to investigate relations among all four measures at
25 ages 7, 11, 14 and 18 y to asses: 1) associations between the height of children and young

1 adults, 2) the extent to which obese children become obese young adults, 3) whether obese
2 young adults were obese children, and 4) BMI groupings match of the TSF groupings of
3 normal, obese, and overweight. Apart from the first year of life, two critical periods in
4 childhood for the development of obesity have been identified in the literature;^{27, 29–34} the
5 period of adiposity rebound between age 5 and 7 y and of adolescence. This is why our
6 analysis used the measurements at age 7, 11, and 14 y to analyse the overweight and obesity
7 outcome at 18y.

8

9 **METHODS**

10 **Subjects**

11 The 1988/90 Slovenian cohort includes children born between September 1988 and April
12 1990. From a target population of 23 114 children who were enrolled in the first grade of the
13 Slovenian primary schools in the school year 1996/97, information on weight and height in
14 the last year of the secondary school at average age 18.7 ± 0.3 y was obtained on 4 833
15 (20.9%) children in April 2008. The nationally representative population sample included 21
16 777 children (94.2% of the entire generation) enrolled in the SLOFIT monitoring system of
17 physical and motor development of children and youth.³⁵ Children, who have been enrolled in
18 the SLOFIT system by the positive written consent of their parents, have been measured every
19 school year between 15 and 30 April throughout their 11-year schooling period. The analysed
20 subsample included only children whose information on body weight and body height was
21 obtained both in the first grade of primary school and the last year of the 4-year secondary
22 school. Children with incomplete information in the 1997 or 2008 measurements were
23 excluded from the subsample. The subsample also did not include pupils who were enrolled
24 in the 3-year vocational schools and therefore finished their schooling a year earlier in 2007
25 not reaching age 18 y at that time. Additional drop of the sample at the end of the schooling

1 occurred because many pupils refused to participate in the measurements after they turned 18
2 and did not need the written consent of their parents any more. This study presents the data
3 from the 1997 SLOFIT measurement and the 2001, 2004 and 2008 follow-ups. The age of the
4 participating children was calculated to the closest month at the day of the measurements
5 every year and the average age at the 1997 measurement was 92.8 ± 3.7 m for males and 92.7
6 ± 3.9 m for females. The differences in mean values and standard deviations of height,
7 weight, TSF, and BMI among the population sample and the analysed subsample are
8 presented in **Table 1**. Mean values and standard deviations of the population sample and the
9 subsample were very similar, with differences in mean height below 5mm, differences in
10 mean weight below 0.6kg (accept in 11 y females where the average weight of the population
11 sample was 1.1kg larger), differences in mean TSF below 0.4mm, and differences in mean
12 BMI below 0.4kg/m^2 at all ages.

13

14 **Measurements**

15 Heights, weights and TSF were measured by trained physical education teachers at 7, 11, 14
16 and $18\text{ y} \pm 1\text{ y}$. Subjects were weighted barefoot in their shorts and t-shirts to the nearest 0.5
17 kg by calibrated portable scales, height was measured with a stadiometer to the nearest 0.5 cm
18 and triceps skinfold was measured with Holtain-Tanner calliper to the nearest mm. Data were
19 checked to detect coding errors. BMI was calculated at each age. The IOTF cut off points³⁶
20 were used to define overweight and obese children. Age-specific values are given in **Table 2**.

21

22 **Statistics**

23 Partial correlation coefficients, adjusted for date of measurements, were calculated between
24 height, weight, TSF and BMI at age 7, 11, 14, and 18. To reduce skewness of the weight, TSF
25 and BMI distributions, correlations were also performed by using $\text{Lg}(\text{weight})$, $\text{Lg}(\text{TSF})$,

1 Lg(BMI) and Sqrt for all three variables. Differences between correlations using transformed
2 and untransformed variables proved to be trivial (-0.26 to 0.24), so only correlations of
3 untransformed variables are presented. BMI categories at age 7, 11, 14, and 18 were cross-
4 classified. Percentages are shown primarily to show outcome for the obese children and to
5 show the BMI distribution at age 7 for the different BMI groups at age 18. The IOTF-
6 classified normal, overweight and obese BMI categories³⁶ were cross-classified with the TSF
7 normal, overweight and obese categories³⁷ to establish how the first classification related to
8 the latter. All analyses were performed using SPSS 15.0 for Windows (SPSS Inc., Chicago,
9 IL).

10

11 **RESULTS**

12 Mean (\pm SD) heights, weights, triceps skinfold and BMIs for subjects 7 to 18 y for males and
13 females are shown in **Table 3**.

14

15 **Height**

16 Correlations between height at 7 and 18 y in **Table 4** tended to be strong for males ($r = 0.69$)
17 and females ($r = 0.65$). As expected, correlations were greater over shorter intervals. Males had
18 higher correlations than females at ages 7 and 11 y, but lower correlations at age 14. These
19 sex differences reflect the differences in the pubertal growth spurt, which occurs later for
20 males. At age 14 the growth period of females is nearing the end and the difference between
21 average height of females at ages 14 and 18 y, visible in **Table 3**, was only 2cm, while the
22 difference between the average height of males in the same period was four times bigger at
23 8cm. The concluded growth of females at age 14 y is visible also in Table 4 since the
24 correlation between height at age 14 and 18 y among females was considerably stronger ($r =$
25 0.91) than among males ($r = 0.72$).

1

2 **Weight and BMI**

3 In general, correlations for weight were very similar to those of height ($r = 0.64$ for males and
4 females between ages 7 and 18 y). As in height, stronger associations were observed over the
5 shorter periods in both sexes (**Table 4**). As in height, the correlations in weight among shorter
6 periods were slightly stronger among males than among females. Correlations between BMI
7 at different ages were weaker than for weight and height ($r = 0.58$ for males and $r = 0.59$ for
8 females between ages 7 and 18 y) and the correlations among shorter periods were, again,
9 slightly stronger for males than for females. The correlations indicated ($BMI^2 \cdot 100$) that BMI
10 at 11 or 14 y explained 49–69% of the variability in BMI at age 18, while BMI at 7 y
11 explained only around 34% of the variability in BMI at age 18 for males and females.

12

13 **Triceps skinfold fat**

14 In comparison to all other measures, the TSF showed the weakest correlations (**Table 4**)
15 between ages 7 and 33 y ($r = 0.39$ for males and $r = 0.35$ for females). Stronger correlations
16 were, again, observable among males and over the short term. The strongest correlations in
17 TSF were observable between ages 11 and 16 y and were slightly higher among males ($r =$
18 0.70) than among females ($r = 0.62$). The correlations indicated that only 12–25% of the
19 variability in TSF at age 18 y was explained by TSF at 7, 11, or 14 y. Correlations of TSF
20 with height, weight, and BMI (**Table 5**) suggested that TSF correlates best with BMI and that
21 correlations are largest at age 11 y in males ($r = 0.77$) and females ($r = 0.70$). High
22 correlations were observed also between TSF and weight at all ages in both sexes, while
23 correlations of TSF with height proved to be small ($r < 0.29$). Comparison of groupings
24 according to TSF percentile groups³⁷ and IOTF BMI groups³⁶ in **Table 6** showed, that both
25 groupings overlapped in the grouping of normal BMI group with 50th percentile TSF group,

1 while the matching of the overweight BMI group with 85th percentile TSF group, and of the
2 obese BMI group with 95th TSF percentile group was rather incoherent.

3

4 **BMI outcome for obese and overweight children**

5 The data in **Table 7** shows whether overweight and obese 7-y olds become overweight 18-y
6 olds, giving the percentage of children, who were subsequently in each BMI group according
7 to IOTF cutoff points at age 18 y. The general trend was that the higher the BMI group the
8 higher the probability of being obese at age 18 y. For example, 17.3% of 7-y old males with
9 normal weight were obese or overweight at age 18 y compared with 57.1% who were
10 overweight or 77.7 who were obese at age 7 y. The chances of an obese 7-y old becoming an
11 obese 18-y old increased consistently with age. For the obese males at 7 y the percentage of
12 those, who were obese at 18 y increased from 19.8% at age 7 y to 26.4 at age 11 y and 33.8 at
13 age 14 y. The trends among females were similar with 17% of obese 18-y olds at age 7 y,
14 32.7% at age 11 y, and 37.1% at age 14 y.

15

16 **Previous BMI of obese 18-y olds**

17 The data in **Table 8** presents percentages for BMI groups of those 18 y who had been in
18 different BMI groups in childhood and adolescence in order to assess whether obese 18-y olds
19 were obese as children. At every age the majority (around 90%) of 18-y olds with normal
20 weight would be identified as children with normal weight. In contrast, between 41.3 and
21 57.6% of overweight 18-y old males, and between 53.2 and 63.4% of overweight or obese
22 females would be identified as overweight children. In both sexes there was a very high
23 percentage of obese at 18 y who would have been identified as overweight or obese at earlier
24 ages. The percentage grew with age in both sexes; among males 73.3% of obese 18-y olds
25 were overweight or obese at age 7 y, 93.9% at age 11 y and 100% at age 14 y, while among

1 females 85.7% of obese 18-y olds were obese at age 7 y, 93.5% at age 14 y and 97.1% at age
2 14 y.

3

4 **DISCUSSION**

5 It has already been established that tracking childhood obesity beyond early adult age is a
6 serious limitation of the literature to date^{36, 38} but the lack of studies from the developing
7 countries and the countries that experienced abrupt and big socio-political and economical
8 changes in recent years seems to be equally limiting.³⁹ Although our study has a limitation of
9 reaching only the school-going population, Slovenia is a country which experienced
10 enormous socio-political and economical changes in the last 20 years that strongly influenced
11 the lifestyles, nutritional habits of children and youth, and their physical development.⁴⁰⁻⁴⁷

12 Our study is the most recent cohort study including young adults who turned 18 y in 2008 and
13 were born at the beginning of the socio-political and economical transition of Slovenia in
14 early 1990s.

15 Studies in the United States and Europe that tracked obesity from childhood to adulthood
16 generally found that about one-third of overweight and obese children remained overweight
17 and obese as adults, but the rate varies because of differences in how obesity is defined, the
18 length of follow-up, and children's initial age.^{24, 38, 48-52} In our cohort study, the percentage of
19 overweight and obese 7-y olds who became overweight or obese 18-y olds was considerably
20 higher (over 65% among males and below 50% among females). Our findings, which are
21 more alerting, might be the outcome of the recent cohort since most of the studies used the
22 cohorts from the 1950s and 1960s. This means that young adults, included in our cohort, lived
23 in a very different environment as children than the ones in the older cohorts, especially
24 regarding the abundance of fat- and sugar-rich food, availability of media and information
25 technology, and the growingly sedentary lifestyles.

1 In addition, we found that overweight and obesity tracks slightly more consistently in females
2 than it does in males since obese and overweight females were more likely to remain in the
3 same BMI group over time or move to a lower BMI group (obese to overweight and
4 overweight to normal, respectfully). The reason for this phenomenon remains unclear but it
5 could be that body image and consequent weight management is more important for girls than
6 for boys and they put more effort into maintaining or lowering their weight, while boys who
7 much often moved from the lower to the higher BMI group, show much lower concern in this
8 matter. Another possible reason that might explain this phenomenon is the use of the IOTF
9 standard, which could misclassify some Slovenian children and adolescents because this
10 standard was constructed on the data from other countries than Slovenia.

11 In comparison to some other studies,^{25, 38, 48} BMI in our population is well predicted from
12 childhood. This is suggested by large correlations (above $r = 0.50$) between childhood and
13 young adulthood BMI and a large proportion of obese and overweight 18-y olds identified
14 from childhood. According to the IOTF cutoff points for obesity, 40% of Slovenian 18-y old
15 obese males and 48.6% females would have been identified from their BMI at age 7 y. The
16 percentage identified increased with years and it is notable that at age 14 y a large majority of
17 the obese 18-y olds had already been obese. The data suggests that children who today enter
18 secondary schools obese are almost certain to remain so at the end of their schooling.

19 Although Serdula et al³⁸ estimated that less than a half of adult obesity can be attributed to
20 childhood obesity, and our data supports this finding, it should be noted that taking into
21 account overweight and obese BMI groups combined, over 73% of males and over 85% of
22 18-y old obese females would have been classified as overweight or obese already at age 7 y,
23 and the percentage identified would have increased to over 90% with age in both sexes. Also
24 the correlations of BMI at age 18 y with BMI at earlier ages suggest that between one third (at

1 age 7 y) and two thirds (at age 14 y) of the variability in BMI at age 18 y was explained by
2 BMI in childhood.

3 In comparison with correlations between BMI from childhood to young adulthood,
4 correlations for height were even larger ($r = 0.69$ for males and $r = 0.65$ for females at age 7
5 y). As it has been confirmed in other studies,^{25,53} the observed correlations between female
6 height at age 11 y and height at age 18 y were slightly lower than the ones in height between
7 ages 7 and 18 y. This was expected since correlations around the time of maturation in
8 females may be slightly poorer than for prepubertal measures.⁵³ Earlier maturation of females
9 in comparison to males was observed in correlation between heights at age 14 y and at age 18
10 y with $r = 0.91$ for females and 0.72 for males. Similarly high correlations were found also
11 among childhood and young adulthood weight.

12 Finally, in contrast with the large correlations of height, weight, and BMI between childhood
13 and adulthood, those for TSF were only medium ($r < 0.50$), except the correlation of TSF at
14 age 11 and 14 y, when correlation was large among females ($r = 0.62$) and even larger among
15 males ($r = 0.70$). This finding is conclusive with other findings which suggested that precision
16 of obesity prediction by TSF is very low up to late childhood and becomes somewhat better in
17 adolescence, and that correlations are usually higher for males.²² Our data suggested that TSF
18 was not well predicted from childhood and that although the BMI and TSF groupings did
19 match well in the normal BMI group, they behaved much poorer in the obese, and especially
20 in the overweight BMI group. This phenomenon could be attributed to the known
21 methodological problems.³⁷ It has been reported previously that skinfold-thickness
22 measurements in obese subjects have poor reliability,⁵⁴ and that skin thickness and skinfold
23 compressibility vary by age, by site, and possibly also by sex.⁵⁵⁻⁵⁸ This suggests that using
24 TSF curves for classification of overweight and obese children should be used with all these

1 problems in mind and that use of BMI curves are more practical for tracking of obesity from
2 childhood to adulthood.

3 Child to adult relations for obesity will continue to be of interest to researchers and policy
4 makers, since in many countries policies to prevent childhood and adult obesity are being
5 discussed.⁵⁹ The evidence presented here suggests that prevention of adulthood obesity should
6 focus on identification and treatment of a high risk group in childhood and adolescence. As
7 children and adolescents with normal weight are very likely to remain normal, and according
8 to the evidence which suggests that environmental influences determine overweight and
9 obesity among children in a much larger extent than genetics⁶⁰ it seems especially important
10 to prevent and treat overweight and obesity already at young age. On the other hand, the
11 relative risk for becoming an obese adult is much increased for those who are already obese or
12 overweight as children or adolescents. From an epidemiological viewpoint these are the
13 critical periods when preventive measures such as nutrition and physical activity management
14 could be applied.

15

16 **ACKNOWLEDGEMENTS**

17 This study was funded by the Slovenian Ministry of Education and Sport.

18

1 REFERENCES

- 2
3 1 World Health Organization. *Obesity: preventing and managing the global epidemic*. World
4 Health Organization: Geneva 1998.
- 5 2 Lew EA, Garfinkel L. Variations in mortality by weight among 750,000 men and women. *J*
6 *Chronic Dis* 1979; **32**: 563–576.
- 7 3 Hu G, Tuomilehto J, Silventoinen K, Barengo NC, Peltonen M, Jousilahti P. The effects of
8 physical activity and body mass index on cardiovascular, cancer and all-cause mortality
9 among 47 212 middle-aged Finnish men and women. *Int J Obes* 2005; **29**: 894–902.
- 10 4 El-Serag H. The association between obesity and GERD: a review of the epidemiological
11 evidence. *Dig Dis Sci* 2008; **53**: 2307–2312.
- 12 5 Cho E, Manson JE, Stampfer MJ, Solomon CG, Colditz GA, Speizer FE, Willett WC, Hu
13 FB. A prospective study of obesity and risk of coronary heart disease among diabetic women.
14 *Diabetes Care* 2002; **25**: 1142–1148.
- 15 6 Mokdad AH, Ford ES, Bowman BA, Dietz WH, Vinicor F, Bales VS, Marks JS. Prevalence
16 of obesity, diabetes, and obesity-related health risk factors, 2001. *JAMA* 2003; **289**: 76–79.
- 17 7 Michaud DS, Giovannucci E, Willett WC, Colditz GA, Stampfer MJ, Fuchs CS. Physical
18 activity, obesity, height, and the risk of pancreatic cancer. *JAMA* 2001; **286**: 921–929.
- 19 8 Thomsen SF, Ulrik CS, Kyvik KO, Sørensen TI, Posthuma D, Skadhauge LR, Steffensen I,
20 Backer V. Association between obesity and asthma in a twin cohort. *Allergy* 2007; **62**: 1199–
21 1204.
- 22 9 Zimmet P, Alberti KG, Shaw J. Global and societal implications of the diabetes epidemic.
23 *Nature* 2001; **414**: 782–787.
- 24 10 Arslanian S. Type 2 diabetes in children: clinical aspects and risk factors. *Horm Res Suppl*
25 2002; **57**: 19–28.

- 1 11 Freedman DS, Dietz WH, Srinivasan SR, Berenson GS. The relation of overweight to
2 cardiovascular risk factors among children and adolescents: the Bogalusa Heart Study.
3 *Pediatrics* 1999; **103**: 1175–1182.
- 4 12 Harrell JS, Jessup A, Greene N. Changing our future: obesity and the metabolic syndrome
5 in children and adolescents. *J Cardiovasc Nurs* 2006; **21**: 322–330.
- 6 13 Weiss R, Dziura J, Burgert TS, Tamborlane WV, Taksali SE, Yeckel CW, Allen K, et al.
7 2004. Obesity and the metabolic syndrome in children and adolescents. *N Engl J Med* 2004;
8 **350**: 2362–2374.
- 9 14 Gilliland FD, Berhane K, Islam T, McConnell R, Gauderman WJ, Gilliland SS, Avol E,
10 Peters JM. Obesity and the risk of newly diagnosed asthma in school-age children. *Am J*
11 *Epidemiol* 2003; **158**: 406–415.
- 12 15 Colditz GA. Economic costs of obesity and inactivity. *Med Sci Sports Exerc Suppl* 1999;
13 **31**: 663–667.
- 14 16 Wolf AM, Colditz GA. Current estimates of the economic cost of obesity in the United
15 States. *Obes Res* 1998; **6**: 97–106.
- 16 17 Thompson D, Wolf AM. The medical-care cost burden of obesity. *Obes Rev* 2001; **2**: 189–
17 197.
- 18 18 Wang G, Dietz WH. Economic burden of obesity in youths aged 6 to 17 years: 1979-1999.
19 *Pediatrics* 2002; **109**: E81–81.
- 20 19 Seidell JC. Societal and personal costs of obesity. *Exp Clin Endocrinol Diabetes Suppl*
21 1998; **106**: 7–9.
- 22 20 Whitaker RC, Wright JA, Pepe MS, Seidel KD, Dietz WH. Predicting obesity in young
23 adulthood from childhood and parental obesity. *N Engl J Med* 1997; **337**: 869–873.
- 24 21 Kotani K, Nishida M, Yamashita S, Funahashi T, Fujioka S, Tokunaga K, Ishikawa K,
25 Tarui S, Matsuzawa Y. Two decades of annual medical examinations in Japanese obese

- 1 children: do obese children grow into obese adults?. *Int J Obes Relat Metab Disord* 1997; **21**:
2 912–921.
- 3 22 Clarke WR, Lauer RM. Does childhood obesity track into adulthood?. *Crit Rev Food Sci*
4 *Nutr* 1993; **33**: 423–430.
- 5 23 Magarey AM, Daniels LA, Boulton TJ, Cockington RA. Predicting obesity in early
6 adulthood from childhood and parental obesity. *Int J Obes Relat Metab Disord* 2003; **27**:
7 505–513.
- 8 24 Guo SS, Huang C, Maynard LM, Demerath E, Towne B, Chumlea WC, Siervogel RM.
9 Body mass index during childhood, adolescence and young adulthood in relation to adult
10 overweight and adiposity: the Fels Longitudinal Study. *Int J Obes Relat Metab Disord* 2000;
11 **24**: 1628–1635.
- 12 25 Power C, Lake JK, Cole TJ. Body mass index and height from childhood to adulthood in
13 the 1958 British born cohort. *Am J Clin Nutr* 1997; **66**: 1094–1101.
- 14 26 Kemper HC, Post GB, Twisk JW, and van Mechelen W. Lifestyle and obesity in
15 adolescence and young adulthood: results from the Amsterdam Growth And Health
16 Longitudinal Study (AGAHLS). *Int J Obes Relat Metab Disord Suppl* 1999; **23**: 34–40.
- 17 27 Williams S, Davie G, Lam F. Predicting BMI in young adults from childhood data using
18 two approaches to modelling adiposity rebound. *Int J Obes Relat Metab Disord* 1999; **23**:
19 348–354.
- 20 28 Kvaavik E, Tell GS, Klepp KI. Predictors and tracking of body mass index from
21 adolescence into adulthood: follow-up of 18 to 20 years in the Oslo Youth Study. *Arch*
22 *Pediatr Adolesc Med* 2003; **157**: 1212–1218.
- 23 29 Dietz WH. Periods of risk in childhood for the development of adult obesity—what do we
24 need to learn?. *J Nutr* 1997; **127**: 1884–1886.

- 1 30 Dietz WH. Critical periods in childhood for the development of obesity. *Am J Clin Nutr*
2 1994; **59**: 955–959.
- 3 31 Rolland-Cachera MF, Deheeger M, Bellisle F, Sempe M, Guilloud-Bataille M, Patois E.
4 Adiposity rebound in children: a simple indicator for predicting obesity. *Am J Clin Nutr* 1984;
5 **39**: 129–135.
- 6 32 Charney E, Goodman HC, McBride M, Lyon B, Pratt R. Childhood antecedents of adult
7 obesity. Do chubby infants become obese adults? *N Engl J Med* 1976; **295**: 6–9.
- 8 33 Stettler N, Zemel BS, Kumanyika S, Stallings VA. Infant weight gain and childhood
9 overweight status in a multicenter, cohort study. *Pediatrics* 2002; **109**: 194–199.
- 10 34 Whitaker RC, Pepe MS, Wright JA, Seidel KD, and Dietz WH. Early adiposity rebound
11 and the risk of adult obesity. *Pediatrics* 1998; **101**: E5–E5.
- 12 35 Strel J, Ambrožič F, Kondrič M, Kovač M, Leskošek B, Štihec J, Šturm J. *Sports*
13 *educational chart*. Ljubljana, Slovenia: Ministry of Education and Sport, 1997.
- 14 36 Cole TJ, Bellizzi MC, Flegal KM, Dietz WH. Establishing a standard definition for child
15 overweight and obesity worldwide: international survey. *BMJ* 2000; **320**: 1240–1243.
- 16 37 Must A, Dallal GE, Dietz WH. Reference data for obesity: 85th and 95th percentiles of
17 body mass index (wt/ht²) and triceps skinfold thickness. *Am J Clin Nutr* 1991; **53**: 839–846.
- 18 38 Serdula MK, Ivery D, Coates RJ, Freedman DS, Williamson DF, Byers T. Do obese
19 children become obese adults? A review of the literature. *Prev Med* 1993; **22**: 167–177.
- 20 39 Wang Y, Ge K, Popkin BM. Tracking of body mass index from childhood to adolescence:
21 a 6-y follow-up study in China. *Am J Clin Nutr* 2000; **72**: 1018–1024.
- 22 40 Koch V, Pokorn D. Some differences in nutritional habits between males and females in
23 the Republic of Slovenia. *Food Technol Biotechnol*, 1999; **37**: 181–186.
- 24 41 Koch V, Pokorn D. Comparison of nutritional habits among various adult age groups in
25 Slovenia. *Nutr Res* 1999; **19**: 1153–1164.

- 1 42 Kovač M. When social becomes biological: the effect of different physical education
2 curricula on motor and physical development of high-school girls. *Anthropol Noteb* 2006; **12**:
3 97–112.
- 4 43 Leskošek B, Kovač M, Strel J. A comparison of the physical characteristics and motor
5 abilities of boys and girls attending different high school. *Acta Univ Carol Kinanthropol*
6 2006; **42**: 85–101.
- 7 44 Leskošek B, Strel J, Kovač M. Differences in physical fitness between normal-weight,
8 overweight and obese children and adolescents. *Kinesiol Slov* 2007; **13**: 21–30.
- 9 45 Kovač M, Leskošek B, Strel J. Overweight and obesity trends in Slovenian boys from
10 1991 to 2006. *Acta Univ Palacki Olomuc Gymn* 2008; **38**: 17–26.
- 11 46 Jurak G. Sporting lifestyle vs. 'cigarettes & coffee' lifestyle of Slovenian high school
12 students. *Anthropol Noteb* 2006; **12**: 79–95.
- 13 47 Jurak G, Kovač M, Strel J. How Slovene primary school pupils spend their summer
14 holidays. *Kinesiol Slov* 2002; **8**: 35–43.
- 15 48 Lake JK, Power C, and Cole TJ. Child to adult body mass index in the 1958 British birth
16 cohort: associations with parental obesity. *Arch Dis Child* 1997; **77**: 376–381.
- 17 49 Rolland-Cachera MF, Deheeger M, Guilloud-Bataille M, Avons P, Patois E, Sempé M.
18 Tracking the development of adiposity from one month of age to adulthood. *Ann Hum Biol*
19 1987; **14**: 219–229.
- 20 50 Rolland-Cachera MF, Bellisle F, and Sempe M. The prediction in boys and girls of the
21 weight/height index and various skinfold measurements in adults: a two-decade follow-up
22 study. *Int J Obes* 1989; **13**: 305–311.
- 23 51 Guo SS, Chumlea WC, Roche AF, and Siervogel RM. Age- and maturity-related changes
24 in body composition during adolescence into adulthood: the Fels Longitudinal Study. *Int J*
25 *Obes Relat Metab Disord* 1997; **21**: 1167–1175.

- 1 52 DiPietro L, Mossberg HO, and Stunkard AJ. A 40-year history of overweight children in
2 Stockholm: life-time overweight, morbidity, and mortality. *Int J Obes Relat Metab Disord*
3 1994; **18**: 585–590.
- 4 53 Tanner JM. *Growth at adolescence*. Oxford, United Kingdom: Blackwell Scientific
5 Publications, 1962.
- 6 54 Forbes GB. Lean body mass and fat in obese children. *Pediatrics* 1964; 34: 308–314.
- 7 55 Martin AD, Ross WD, Drinkwater DT, Clarys JP. Prediction of body fat by skinfold
8 caliper: assumptions and cadaver evidence. *Int J Obes Suppl* 1985; **9**: 31–39.
- 9 56 Roche AF. Some aspects of the criterion methods for the measurement of body
10 composition. *Hum Biol* 1987; **59**: 209–220.
- 11 57 Himes JH, Roche AF, Siervogel RM. Compressibility of skinfolds and the measurement of
12 subcutaneous fatness. *Am J Clin Nutr* 1979; **32**: 1734–40.
- 13 58 Clegg EJ, and Kent C. Skinfold compressibility in young adults. *Hum Biol* 1967; **39**: 418–
14 429.
- 15 59 Commission of the European Communities. *White Paper on a Strategy for Europe on*
16 *Nutrition, Overweight and Obesity Related Health Issues*. Commission of the European
17 Communities: Brussels 2007.
- 18 60 Kral TV, Berkowitz RI, Stunkard AJ, Stallings VA, Brown DD, Faith MS. Dietary energy
19 density increases during early childhood irrespective of familial predisposition to obesity:
20 results from a prospective cohort study. *Int J Obes* 2007; **31**: 1061–1067.

21

1 **TABLE 1**

2 Differences in mean values and standard deviations between the population sample and the
 3 analysed subsample

Age	7 y	11 y	14 y	18 y
Males				
Height (m)	0.000 [0.001] ¹	0.002 [0.002]	0.000 [0.001]	0.000 [0.000]
Weight (kg)	0.060 [0.154]	0.591 [0.474]	0.403 [0.155]	-0.058 [0.085]
TSF (mm)	0.002 [0.078]	0.252 [0.274]	0.223 [0.146]	0.134 [0.148]
BMI (kg/m ²)	0.022 [0.048]	0.200 [0.119]	0.147 [0.069]	-0.025 [0.035]
Females				
Height (m)	0.002 [0.000]	0.006 [0.001]	0.001 [0.002]	0.000 [0.000]
Weight (kg)	0.345 [0.311]	1.140 [0.573]	0.770 [0.777]	0.225 [0.208]
TSF (mm)	0.161 [0.125]	0.458 [0.461]	0.370 [0.217]	0.172 [0.143]
BMI (kg/m ²)	0.153 [0.143]	0.356 [0.263]	0.257 [0.273]	0.084 [0.087]

4 ¹ SD differences in brackets

5

1 **TABLE 2**2 Percentage of normal, overweight and obese children at age 7, 11, 14, and 18¹

Age	Males			Females		
	Normal	Overweight	Obese	Normal	Overweight	Obese
7 y	81.1 [1712]	13.2 [279]	5.7 [121]	82.5 [1871]	13.1 [296]	4.4 [100]
11 y	76.8 [1471]	18.7 [358]	4.5 [87]	81.8 [1686]	15.9 [327]	2.4 [49]
14 y	78.7 [1663]	17.6 [372]	3.6 [77]	87.2 [1977]	11.2 [255]	1.5 [35]
18 y	74.5 [1573]	22.7[479]	2.8 [60]	87.4 [1982]	11.0 [250]	1.5 [35]

3 ¹ n in brackets

4

5

1 **TABLE 3**

2 Mean height, weight, triceps skinfold thickness and BMI at ages 7–18 y

	Height <i>m</i>	Weight <i>kg</i>	TSF <i>mm</i>	BMI <i>kg/m²</i>
Males				
7 y (n = 2112)	1.29 ± 0.05	28.01 ± 5.30	9.76 ± 4.22	16.62 ± 2.35
11 y (n = 1917)	1.51 ± 0.07	43.54 ± 9.77	12.63 ± 6.10	18.93 ± 3.30
14 y (n = 2112)	1.72 ± 0.08	62.04 ± 11.70	10.88 ± 5.84	20.86 ± 3.18
18 y (n = 2112)	1.80 ± 0.06	75.64 ± 11.13	10.74 ± 5.49	23.27 ± 3.06
Females				
7 y (n = 2267)	1.28 ± 0.06	27.04 ± 5.03	10.98 ± 4.15	16.34 ± 2.27
11 y (n = 2065)	1.52 ± 0.07	43.33 ± 9.57	12.85 ± 5.61	18.63 ± 3.18
14 y (n = 2267)	1.65 ± 0.06	55.72 ± 8.66	14.17 ± 5.25	20.56 ± 2.81
18 y (n = 2267)	1.67 ± 0.06	60.59 ± 9.08	15.54 ± 5.33	21.81 ± 2.89

3

4

5

1 **TABLE 4**2 Partial correlation matrix of height, weight, TSF and BMI at ages 7–18 y¹

	Males				Females			
	7 y	11 y	14 y	18 y	7 y	11 y	14 y	18 y
Height								
7 y	1.00 [2112]	0.87 [1917]	0.78 [2112]	0.69 [2112]	1.00 [2267]	0.84 [2065]	0.74 [2267]	0.65 [2267]
11 y		1.00 [1917]	0.86 [1917]	0.70 [1917]		1.00 [2065]	0.74 [2065]	0.60 [2065]
14 y			1.00 [2112]	0.72 [2112]			1.00 [2267]	0.91 [2267]
18 y				1.00 [2112]				1.00 [2267]
Weight								
7 y	1.00 [2112]	0.86 [1917]	0.78 [2112]	0.64 [2112]	1.00 [2267]	0.85 [2065]	0.73 [2267]	0.64 [2267]
11 y		1.00 [1917]	0.87 [1917]	0.72 [1917]		1.00 [2065]	0.83 [2065]	0.68 [2065]
14 y			1.00 [2112]	0.78 [2112]			1.00 [2267]	0.82 [2267]
18 y				1.00 [2112]				1.00 [2267]
TSF								
7 y	1.00 [2071]	0.66 [1882]	0.52 [2071]	0.39 [2071]	1.00 [2222]	0.62 [2029]	0.49 [2222]	0.35 [2219]
11 y		1.00 [1917]	0.70 [1917]	0.49 [1917]		1.00 [2065]	0.62 [2065]	0.45 [2064]
14 y			1.00 [2112]	0.50 [2112]			1.00 [2267]	0.49 [2264]
18 y				1.00 [2112]				1.00 [2264]
BMI								
7 y	1.00 [2112]	0.80 [1917]	0.71 [2112]	0.58 [2112]	1.00 [2267]	0.78 [2065]	0.66 [2267]	0.59 [2267]
11 y		1.00 [1917]	0.84 [1917]	0.70 [1917]		1.00 [2065]	0.82 [2065]	0.70 [2065]
14 y			1.00 [2112]	0.78 [2112]			1.00 [2267]	0.79 [2267]
18 y				1.00 [2112]				1.00 [2267]

3 ¹ n in brackets

4

1 **TABLE 5**2 Partial correlation matrix of TSF with height, weight, and BMI at ages 7–18 y¹

TSF at age	Males			Females		
	Height	Weight	BMI	Height	Weight	BMI
7 y	0.26 [2071]	0.66 [2071]	0.70 [2071]	0.26 [2222]	0.61 [2222]	0.64 [2222]
11 y	0.25 [1917]	0.71 [1917]	0.77 [1917]	0.16 [2065]	0.61 [2065]	0.70 [2065]
14 y	0.01 [2112]	0.56 [2112]	0.68 [2112]	0.05 [2267]	0.58 [2267]	0.63 [2267]
18 y	0.05 [2112]	0.56 [2112]	0.60 [2112]	0.04 [2264]	0.56 [2264]	0.62 [2264]

3 ¹ n in brackets

4

5

1 **TABLE 6**

2 Comparison of BMI and TSF grouping

TSF percentile groups ¹	Males			Females		
	50 th	85 th	95 th	50 th	85 th	95 th
BMI group						
Normal	93.2	5.8	1.0	93.8	5.5	0.7
Overweight	39.6	30.8	29.7	54.9	25.2	19.9
Obese	12.7	18.6	68.6	19.4	30.6	50.0

3 ¹ TSF 50th, 85th, and 95th percentiles correspond to normal, overweight, and obese TSF groups

4

5

6

1 **TABLE 7**2 BMI outcome at age 18 for the normal, overweight and obese 7-, 11-, and 14-year olds¹

BMI group	Males at age 18 y				Females at age 18 y			
	Normal	Overweight	Obese	Total (n)	Normal	Overweight	Obese	Total (n)
Age 7 y	[1573]	[479]	[60]		[1982]	[250]	[35]	
Normal	82.7	16.4	0.9	100.0 [1712]	93.5	6.3	0.3	100.0 [1871]
Overweight	47	45.9	7.2	100.0 [279]	65.9	29.7	4.4	100.0 [296]
Obese	22.3	57.9	19.8	100.0 [121]	38	45	17	100.0 [100.0]
Age 11 y	[1441]	[426]	[49]		[1807]	[224]	[31]	
Normal	86.5	13.3	0.2	100.0 [1471]	95	4.9	0.1	100.0 [1686]
Overweight	43	50.6	6.4	100.0 [358]	59.9	36.1	4	100.0 [327]
Obese	16.1	57.5	26.4	100.0 [87]	18.4	49	32.7	100.0 [49]
Age 14 y	[1573]	[479]	[60]		[1982]	[250]	[35]	
Normal	87.8	12.2	0.0	100.0 [1663]	95.1	4.8	0.1	100.0 [1977]
Overweight	28.0	62.9	9.1	100.0 [372]	38.4	53.3	8.2	100.0 [255]
Obese	11.7	54.5	33.8	100.0 [77]	8.6	54.3	37.1	100.0 [35]

3 ¹ n in brackets

4

1 **TABLE 8**

2 Percentages within BMI groups at earlier ages according to BMI status at age 33 y

BMI group ¹	Males at age 18			%	Females at age 18		
	Normal	Overweight	Obese		Normal	Overweight	Obese
Age 7 y	[1573] ²	[479]	[60]		[1982]	[250]	[35]
Normal	90.0	58.7	26.7		88.2	46.8	14.3
Overweight	10.0	41.3	73.3		11.8	53.2	85.7
Obese	1.7	14.6	40.0		1.9	18.0	48.6
Age 11 y	[1441]	[426]	[49]		[1807]	[224]	[31]
Normal	88.3	45.8	6.1		88.7	36.6	6.5
Overweight	11.7	54.2	93.9		11.3	63.4	93.5
Obese	1.0	11.7	46.9		0.5	10.7	51.6
Age 14 y	[1573]	[479]	[60]		[1982]	[250]	[35]
Normal	92.8	42.4	0.0		94.9	38.0	2.9
Overweight	7.2	57.6	100.0		5.1	62.0	97.1
Obese	0.6	8.8	43.3		0.2	7.6	37.1

3 ¹ BMI groups are not mutually exclusive, ie, Overweight includes also Obesity4 ² n in brackets

5

6