




# Computationally efficient modeling and data assimilation of near-surface variability

## Observed Near-Surface Variability

Process studies (e.g., TOGA COARE, Arabian Sea Experiment, SPURS1) clearly depict a diurnal cycle in surface temperature. Data from WHOI <http://uop.whoi.edu/projects/projects.html>



**Remarks:**

- Notice largest diurnal amplitude ( $\geq 2C$ ) in the first few meters (about 2m; red and blue lines).
- Some days show a small ( $\leq 0.25C$ ) diurnal warming even at 10- 15m depth, typically around 20m there is none.

**Issues:**

- How to model such variability? Parameterize or *sufficient* vertical resolution in the model?
- Much of this variability is driven by solar radiation, hence *frequent* shortwave radiation flux is needed.
- Momentum stress due to surface winds and/or waves dissipate this variability, hence high frequency/resolution data is needed.
- Satellite measurements and turbulent air-sea fluxes are sensitive to near-surface temperature.

## MORE INFO

Akella and Suarez, 2018  
NASA GMAO Tech Memo.

E-mail: [santha.akella@nasa.gov](mailto:santha.akella@nasa.gov)

Web: [gmao.gsfc.nasa.gov](http://gmao.gsfc.nasa.gov)



## Atmosphere-Ocean Interface Layer (AOIL) of the NASA GMAO GCM

The NASA GMAO GCM is used for:


- Near-real time Weather Analysis and Prediction [https://gmao.gsfc.nasa.gov/weather\\_prediction/](https://gmao.gsfc.nasa.gov/weather_prediction/)
- Seasonal-Decadal Analysis and Prediction [https://gmao.gsfc.nasa.gov/cgi-bin/products/climateforecasts/geos5/S2S\\_2/index.cgi](https://gmao.gsfc.nasa.gov/cgi-bin/products/climateforecasts/geos5/S2S_2/index.cgi)
- Reanalysis (atmospheric) <https://gmao.gsfc.nasa.gov/reanalysis/MERRA-2/>

Since 01/2017 the Weather Analysis and Prediction system resolves this surface variability (via modeling and assimilation of AVHRR infrared brightness temperatures). See Akella et al, 2017

<http://onlinelibrary.wiley.com/doi/10.1002/qj.2988/full> for description and Gentemann and Akella, 2018 <https://doi.org/10.1002/2017JCO13186> for validation. The diurnal variability model was formulated and implemented for atmospheric GCM, with prescribed *foundation* SST that does not have diurnal variability.

The present AOIL is a reformulation:


- Of the exchange of variables and fluxes across the air-sea interface, designed to work *seamlessly* within atmospheric and coupled GCMs.
- It models the near-surface variability; amplitude of the diurnal warming is improved from Akella et al., 2017 model.



We solve for  $\sigma_T = T_w - T_f$

$$\frac{\partial \sigma_T}{\partial t} = \frac{Q_\sigma}{d \rho_w c_w} - \frac{1}{\tau_\sigma} \sigma_T,$$

$$Q_\sigma = (SW_{top} - SW_d + Q^l) - \frac{\epsilon}{1 - \epsilon} (SW_d - SW_D) \quad \tau_\sigma = \frac{d(1 - \epsilon)\phi(\zeta)}{\kappa u_* (1 + \mu)}$$




$\sigma_T$  versus Local Mean Time (LMT) for an ideal day with constant friction velocity ( $u_*$ ) and  $Q^l = -150 W/m^2$ .




Solid (dashed) lines are with AOIL (Akella et al., 2017) model. The AOIL better models the gradual decay of diurnal warming.

## Validation

Offline model runs using the COARE bulk fluxes and measured temperatures from WHOI <http://uop.whoi.edu/projects/projects.html>



Online model runs using within GMAO atmospheric data assimilation system. Diurnal cycle at select stations



## SUMMARY

The AOIL has been implemented in the NASA GMAO models: both uncoupled (AGCM) and coupled GCM.

Improvements to model near surface salinity are underway.

National Aeronautics and Space Administration

