

Quark confinement as nonlinear phenomenon in QCD vacuum

V. Kuvshinov

JIPNR-Sosny, NAS of Belarus, Belarus

In the framework of stochastic QCD vacuum model we show that colour particle loss its colour propagating through QCD vacuum medium. At large distances colour disappear completely.

Testing of quark-hadron duality via the Adler D-function

O.P. Solovtsova, D.S. Babanin

Gomel State Technical University, Gomel, Belarus

A key problem in making precise perturbative QCD predictions, which connects with the requirement of renormalization group invariance, is the uncertainty in fixing the renormalization scale. We study the scale dependence of fresh four-loop QCD corrections to some processes both within the standard perturbative and singularity-free analytic perturbation theory. We demonstrate that within the analytic approach the results are practically renormalization scale independent and lead to rather different Q^2 evolution than the standard perturbative correction possesses.

Berwald-Lagrange scalar curvature in the structure process of LB-monolayer

N. Krylova¹, A. Oana², V. Balan³, G. Grushevskaya¹, M. Neagu²

¹*Faculty of Physics, Belarussian State University, 4 Nezavisimosti Ave., 220030 Minsk, Belarus, E-mail: grushevskaja@bsu.by, nina-kr@tut.by*

²*University Transilvania of Brasov, 50 Iuliu Maniu Str.,*

500091 Brasov, Romania, E-mail: alexandruo@gmail.com

³*University Politehnica of Bucharest, Splaiul Independentei 313, 060042 Bucharest, Romania, E-mail: vladimir.balan@upb.ro*

In this talk we examine a geometric 2D-dynamics of Langmuir-Blodgett monolayer (LB-monolayer). To do it the vertical component of Berwald curvature tensor is calculated, and the Berwald-Lagrange flag curvature K is determined. A simulation is performed, and possible physical interpretations are also given. It was established that sign change of K from positive to negative occurs at the compressing. The dependence of $(-K)$ on an area of monolayer behaves similar to an isotherm of compressing in the neighbourhood of phase transition. Convolution of the vertical component of Berwald curvature tensor -- Douglas tensor B^i_{jkl} with metric tensor was calculated. An anomalous behavior of the convolution is similar to a behavior of susceptibility at phase transition.