

Type II Aerobic Methane Oxidizing Bacteria (AMOB) Drive Methane Oxidation in Pulsed Wetlands as Indicated by ¹³C-Phospholipid Fatty Acid Composition



Taniya Roy Chowdhury^a and Richard P. Dick^a

^aSchool of Environment & Natural Resources, The Ohio State University, Columbus, OH, U.S.A.

INTRODUCTION

> The magnitude of climate change predicted for the United States over the next 100 years will cause significant impacts on temperature and precipitation patterns

> Wetlands are a major source (~25 %) of methane globally¹

> Aerobic Methane Oxidizing Bacteria (AMOB) have the unique ability to utilize CH₄ as the sole source of carbon and energy by the Methane Mono Oxygenase (MMO) mediated reaction

> A variety of environmental determinants have been implicated in the dominance of AMOB type. Generally, type I AMOB outcompete type II AMOB in low CH₄ (~2ppmv), high oxygen environments, while type II AMOB are favored in high CH₄ (≥40ppmv) environments²

> AMOB are active at the oxic sediment - water interface ("pulsing fringe", Fig. 2)

> AMOB consume ~30 Tg-CH₄-yr⁻¹, and potentially can offset CH₄ losses to the atmosphere¹

> Very little is known about the effects of pulsing hydrology and season on the AMOB ecology

> Therefore, the Objectives of this Study were:


❑ Experiment 1: Determine the effects of Season and Landscape position on Potential Methane Oxidation (PMO) in the two wetlands

❑ Experiment 2: Determine the effects of Season and Landscape position on the Aerobic Methane Oxidizing Bacteria (AMOB) as reflected by their biomarker Phospholipid Fatty Acid (PLFA) Compositions

EXPERIMENTAL SITE

THE OLENTANGY RIVER "Kidney" WETLANDS, Ohio, U.S.A. Soil Sampling Locations (Fig. 3) and Factors:

Factor	Number	Name
Landscape	2	ORW1, ORW2
Hydrology	3	Permanently Flooded (PF) Intermittently Flooded (IF) Upland (UP)
Flow zones	2	Inflow (I), Outflow (O)
Soil Depth	2	0-8 cm 816 cm
Season	4	Fall, Winter, Spring, Summer



RESULTS: ¹³C-Phospholipid Fatty Acids






Fig. 6. Compound-specific carbon isotope values of PLFAs incorporating ¹³C following incubation of Winter soil samples under Yppmv ¹³CH₄ :

A: 0-8cm, PF Site, 2ppmv B: 0-8cm, IF Site, 2ppmv
C: 0-8cm, PF Site, 60ppmv D: 0-8cm, IF Site, 60ppmv

WETLAND HYDROLOGY EXPERIMENT

Effect of Pulsing on Methane Emission in the Olentangy River Wetlands⁴ (Altor and Mitsch, 2006)



REFERENCES


- Whalen, S.C. (2005). Environ. Eng. Sc. 22(1):73-92
- Hanson, R.S., and Hanson, T.E. (1996). Methanotrophic bacteria. Microbiol. Rev. 60:439-471
- Jahnke, L. L., R. E. Summons, J. M. Hope, and Des Marais, D.J. (1999). Geochim. Cosmochim. Acta 63:79-93
- Altor, A. E. and Mitsch, W.J.. (2006). Ecological Engineering 28: 224-234.
- Crossman, Z.M., Abraham, F., Evershed, R.P. (2004). Environ.Sc. & Tech. 38: 359-1367
- Maxfield, P. J., E. R. C. Hornibrook and Evershed, R.P. (2006). Appl. Environ. Microbiol. 72: 3901-3907

Figs. 4 & 5: Seasonal average PMO across landscape positions:

- > PMO in Winter significantly higher than in Summer (p < 0.01)
- > PMO statistically higher at 0-8 cm depth of soil in the PF and IF sites (p < 0.05)

METHODS

• Soil samples injected with known concentration of 99.99% pure CH₄ gas




• PMO reported in nmol-CH₄-gdwt.⁻¹Soil.h⁻¹

Method 1: Potential Methane Oxidation (PMO), Crossman et al. (2004)

Soils Incubated under two different concentrations of ¹³CH₄ : 2ppmv and 60ppmv


- i. "Type I AMOB": [CH₄] < 1.8 ppm
- ii. "Type II AMOB": [CH₄] > 40 ppm




Fingerprint AMOB PLFAs:
Type I AMOB: 16:1ω5c, 16:1ω7c
Type II AMOB: 18:1ω7c, 18:1ω9c, 18:1ω8c

Method 2: ¹³CH₄ Incubation Pulse-Chase Experimental Setup: Stable Isotope Probing of PLFA using GC-c-IRMS, Maxfield et al., 2006

RESULTS: Potential Methane Oxidation



SUMMARY: AMOB PLFA Conc. & PMO



CONCLUSIONS

- > The concentration profiles of AMOB signature PLFAs, as detected by Stable Isotope Probing (SIP) completely corroborates with the Potential Methane Oxidation (PMO) values at all study sites
- > The highest PMO values in the Permanently Flooded site can be entirely attributed to the AMOB, as reflected by the relative abundance of the signature PLFAs
- > Type I AMOB are dominant under the oxygen rich conditions in contrast to the Type II that are abundant under submerged conditions
- > The three hydrologically distinct landscape positions each present characteristic microbial community structure, as evident in this study

ACKNOWLEDGEMENT

- > Dr. W.J. Mitsch, Director, Wilma H. Schiermeier Olentangy River Wetland Research Park, OSU, Columbus, OH, U.S.A.
- > Dr. Andrea Grotoli, Biogeochemistry Lab, School of Earth Sciences, OSU
- > DR. Kary Green-Church, Director, Campus Chemical Instrument Center, Mass Spectrometry & Proteomics Facility, OSU