



PERGAMON

www.elsevier.nl/locate/poly

Polyhedron 19 (2000) 1663–1672



POLYHEDRON

Chemistry of ruthenium with some phenolic ligands: synthesis, structure and redox properties

Falguni Basuli^a, Anjan Kumar Das^a, Golam Mostafa^b, Shie-Ming Peng^c,
Samaresh Bhattacharya^{a,*}

^a Department of Chemistry, Inorganic Chemistry Section, Jadavpur University, Calcutta 700032, India

^b Department of Physics, Krishnath College, Berhampur, West Bengal, India

^c Department of Chemistry, National Taiwan University, Taipei, Taiwan, ROC

Received 11 January 2000; accepted 11 April 2000

Abstract

Reaction of three phenolate ligands, viz. salicylaldehyde (HL¹), 2-hydroxyacetophenone (HL²) and 2-hydroxynaphthylaldehyde (HL³), (abbreviated in general as HL, where H stands for the phenolic proton) with [Ru(PPh₃)₃Cl₂] in 1:1 mole ratio gives complexes of the type [Ru(PPh₃)₂(L)Cl₂]. The structure of the [Ru(PPh₃)₂(L²)Cl₂] complex has been solved by X-ray crystallography. The coordination sphere around ruthenium is O₂P₂Cl₂ with a *cis-trans-cis* geometry, respectively. The [Ru(PPh₃)₂(L)Cl₂] complexes are one-electron paramagnetic (low-spin d⁵, *S* = 1/2) and show rhombic ESR spectra in 1:1 dichloromethane–toluene solution at 77 K. In dichloromethane solution the [Ru(PPh₃)₂(L)Cl₂] complexes show several intense LMCT transitions in the visible region. Reaction between the phenolic ligands and [Ru(PPh₃)₃Cl₂] in 2:1 mole ratio in the presence of a base affords the [Ru(PPh₃)₂(L)₂] complexes in two isomeric forms. ¹H NMR spectra of one isomer shows that it does not have any C₂ symmetry and has the *cis-cis-cis* disposition of the three sets of donor atoms. ¹H NMR spectra of the other isomer shows that it has C₂ symmetry. The structure of the isomer of the [Ru(PPh₃)₂(L)₂] complex has been solved by X-ray crystallography. The coordination sphere around ruthenium is O₄P₂ with a *cis-trans-cis* disposition of the carbonylic oxygens, phenolate oxygens and phosphorus atoms, respectively. The [Ru(PPh₃)₂(L)₂] complexes are diamagnetic (low-spin d⁶, *S* = 0) and show intense MLCT transitions in the visible region. Cyclic voltammetry on the [Ru(PPh₃)₂(L)Cl₂] complexes shows a ruthenium(III)–ruthenium(II) reduction near –0.3 V versus SCE and a ruthenium(III)–ruthenium(IV) oxidation in the range 1.08–1.24 V versus SCE. Cyclic voltammetry on both isomers of the [Ru(PPh₃)₂(L)₂] complexes shows a ruthenium(II)–ruthenium(III) oxidation within 0.09–0.41 V versus SCE, followed by a ruthenium(III)–ruthenium(IV) oxidation within 1.31–1.52 V versus SCE. © 2000 Elsevier Science Ltd. All rights reserved.

Keywords: Ruthenium; Phenolic ligands; Synthesis; Structure; Redox properties



1. Introduction

The chemistry of ruthenium has currently been receiving a lot of attention [1–10] primarily because of the fascinating electron-transfer and energy-transfer properties displayed by the complexes of this metal. Ruthenium offers a wide range of oxidation states and the reactivities of the ruthenium complexes depend on the stability and interconvertibility of these oxidation

states, which in turn depend on the nature of ligands bound to the metal. Complexation of ruthenium by ligands of different types has thus been of particular interest. In the present study, which has originated from our interest in the chemistry of ruthenium in different coordination environments [11–18], we have chosen phenolic ligands of type **1** as the principal ligand, which are abbreviated in general as HL where H stands for the dissociable phenolic proton. The deprotonated ligands are known to coordinate metal ions as bidentate O,O-donor forming six-membered chelate rings (**2**) [19,20]. Three different phenolic ligands have been used in this study, which are shown in **1** along with their specific abbreviations.

* Corresponding author. Fax: +91-33-473-4266.

E-mail address: samaresh_b@hotmail.com (S. Bhattacharya).



While phenolate oxygen is a recognized hard donor, and coordination by it is known to stabilize the higher oxidation states of ruthenium [21–23], coordination of metals by carbonylic oxygen has also recently been of considerable interest in bioinorganic chemistry [24–28]. It may be mentioned here that ruthenium chemistry of these ligands has not been explored much [19]. Stability of different oxidation states of ruthenium is expected to depend on the number of phenolate and carbonylic oxygen in the coordination sphere and in the case of mixed-ligand complexes, on the nature of coligands also. Herein we have restricted our studies to some mono- and bis-phenolate complexes of ruthenium, which have been synthesized by reacting the phenolic ligands with $[\text{Ru}(\text{PPh}_3)_3\text{Cl}_2]$ under different experimental conditions. The chemistry of complexes of the type $[\text{Ru}(\text{PPh}_3)_2(\text{L})\text{Cl}_2]$ and $[\text{Ru}(\text{PPh}_3)_2(\text{L})_2]$ has been described in this paper with special reference to synthesis, stereoisomerism and electron-transfer properties.

2. Experimental

2.1. Materials

Commercial ruthenium trichloride was purchased from Arora Matthey, Calcutta, India, and was converted into $\text{RuCl}_3 \cdot 3\text{H}_2\text{O}$ by repeated evaporation with concentrated hydrochloric acid. Triphenylphosphine (PPh_3), triethylamine (NEt_3) and salicylaldehyde (HL^1) were obtained from SD, India. 2-Hydroxyacetophenone (HL^2) and 2-hydroxynaphthaldehyde (HL^3) were purchased, respectively, from Spectrochem, India and Aldrich. All other chemicals and solvents were reagent grade commercial materials and were used as received. $[\text{Ru}(\text{PPh}_3)_3\text{Cl}_2]$ was prepared following a reported procedure [29]. Purification of acetonitrile and preparation of tetraethylammonium perchlorate (TEAP) for electrochemical work were performed as before [30,31].

2.2. Preparation of complexes

2.2.1. $[\text{Ru}(\text{PPh}_3)_2(\text{L}^1)\text{Cl}_2]$

HL^1 (15 mg, 0.12 mmol) was added to a suspension of $[\text{Ru}(\text{PPh}_3)_3\text{Cl}_2]$ (100 mg, 0.10 mmol) in ethanol (40 cm^3) and the mixture was stirred for 5 h to produce a green solution. On partial evaporation of the solvent, $[\text{Ru}(\text{PPh}_3)_2(\text{L}^1)\text{Cl}_2]$ separated out as a green crystalline

solid, which was collected by filtration, washed with ethanol and dried in air. The yield was 55 mg (64%).

2.2.2. $[\text{Ru}(\text{PPh}_3)_2(\text{L}^2)\text{Cl}_2]$

This was synthesized by following the same procedure above using HL^2 instead of HL^1 . The yield was 60 mg (69%).

2.2.3. $[\text{Ru}(\text{PPh}_3)_2(\text{L}^3)\text{Cl}_2]$

Dichloromethane (40 cm^3) was added to a solid mixture of $[\text{Ru}(\text{PPh}_3)_3\text{Cl}_2]$ (100 mg, 0.10 mmol) and HL^3 (20 mg, 0.12 mmol) and the solution was stirred for 3 h to afford a green solution. On evaporation of the solvent, a green solid was obtained, which was washed with ethanol and dried in air. Purification of this product was achieved by chromatography through a silica gel column. Using toluene as the eluent a green band resulted, which was collected. Evaporation of the eluate gave $[\text{Ru}(\text{PPh}_3)_2(\text{L}^3)\text{Cl}_2]$ as a green microcrystalline solid. The yield was 55 mg (61%).

2.2.4. *ctc* and *ccc* isomers of $[\text{Ru}(\text{PPh}_3)_2(\text{L}^1)_2]$

2.2.4.1. Method A. $[\text{Ru}(\text{PPh}_3)_3\text{Cl}_2]$ (100 mg, 0.10 mmol) was added to a hot solution of HL^1 (30 mg, 0.24 mmol) in ethanol (40 cm^3), followed by NEt_3 (30 mg, 0.30 mmol). The mixture was refluxed for 2 h to produce a red solution. On partial evaporation of solvent, a microcrystalline red solid precipitated, which was collected by filtration, washed thoroughly with water and dried in vacuo over P_4O_{10} . Purification was achieved by chromatography through silica gel column. Using toluene and 1:4 acetonitrile–toluene as the eluents, two different red bands resulted, which were collected separately and evaporation of the eluates, respectively, gave the *ctc* and *ccc* isomers of $[\text{Ru}(\text{PPh}_3)_2(\text{L}^1)_2]$. The yield was 30 mg (33%) for the *ctc* isomer and 32 mg (35%) for the *ccc* isomer.

2.2.4.2. Method B. $[\text{Ru}(\text{PPh}_3)_2(\text{L}^1)\text{Cl}_2]$ (50 mg, 0.06 mmol) was dissolved in a minimum volume of dichloromethane and to it a solution of HL^1 (10 mg, 0.08 mmol) in ethanol (30 cm^3) was added, followed by NEt_3 (10 mg, 0.10 mmol). The solution was refluxed for 4 h. Upon evaporation of the solution, a red crystalline solid was obtained, which was washed with water and dried in vacuo over P_4O_{10} . The solid was then purified as in Method A. The *ctc* and *ccc* isomers of $[\text{Ru}(\text{PPh}_3)_2(\text{L}^1)_2]$ were obtained in 36 and 32% yields, respectively.

2.2.5. *ctc* and *ccc* isomers of $[Ru(PPh_3)_2(L^2)_2]$

These isomers have been prepared by following the above procedures. In method A, HL² was used instead of HL¹. The *ctc* and *ccc* isomers of $[Ru(PPh_3)_2(L^2)_2]$ were obtained in 30 and 34% yields. In method B $[Ru(PPh_3)_2(L^2)Cl_2]$ and HL² were used instead of $[Ru(PPh_3)_2(L^1)Cl_2]$ and HL¹, respectively. Yields of the *ctc* and *ccc* isomers were 35 and 33%, respectively.

2.2.6. *ctc* and *ccc* isomers of $[Ru(PPh_3)_2(L^3)_2]$

These isomers have been prepared by following the above procedures (Section 2.2.4). In method A, HL³ was used instead of HL¹. The *ctc* and *ccc* isomers of $[Ru(PPh_3)_2(L^3)_2]$ were obtained in 32 and 36% yields. In method B, $[Ru(PPh_3)_2(L^3)Cl_2]$ and HL¹ were used instead of $[Ru(PPh_3)_2(L^1)Cl_2]$ and HL¹, respectively. Yields of the *ctc* and *ccc* isomers were 32 and 34%, respectively.

2.3. Physical measurements

Microanalyses (C, H, N) were performed using a Perkin–Elmer 240C elemental analyzer. IR spectra were obtained on a Perkin–Elmer 783 spectrometer with samples prepared as KBr pellets. Electronic spectra were recorded on Shimadzu UV 240 spectrophotometer. Magnetic susceptibilities were measured using a PAR 155 Vibrating sample magnetometer fitted with a Walker scientific L75FBAL magnet. ¹H NMR spectra

were recorded on a Bruker AC-200 spectrometer using TMS as the internal standard. ESR spectra were recorded on a Varian Model 109C E-line X-band spectrometer fitted with a quartz Dewar for measurements at 77 K (liquid dinitrogen). All spectra were calibrated with the help of DPPH ($g = 2.0037$). Electrochemical measurements were made using a PAR model 273 potentiostat. A platinum disc working electrode, a platinum wire auxiliary electrode and an aqueous saturated calomel reference electrode (SCE) were used in a three electrode configuration. Dinitrogen gas was purified by successively bubbling it through alkaline dithionite and concentrated sulfuric acid. All electrochemical experiments were performed under a dinitrogen atmosphere. All electrochemical data were collected at 298 K and are uncorrected for junction potentials. An RE 0089 X-Y recorder was used to trace the voltammograms.

2.4. Crystallography

Single crystals of $[Ru(PPh_3)_2(L^2)Cl_2]$ were grown by slow diffusion of hexane into a dichloromethane solution of the complex. Selected crystal data and data collection parameters are given in Table 1. Data were collected on a Siemens Smart CCD diffractometer using graphite monochromated Mo K α radiation ($\lambda = 0.71073$ Å) by ω scans within the angular range $1.57 < \theta < 25.00^\circ$. X-ray data reduction, structure solution and refinement were done using SHELXTL-PLUS package. The structure was solved by direct methods.

Single crystals of *ctc*- $[Ru(PPh_3)_2(L^1)_2]$ were grown by slow diffusion of benzene into an acetonitrile solution of the complex. Selected crystal data and data collection parameters are given in Table 1. The unit cell dimensions were determined by a least-squares fit of 25 centered reflections ($10.80 \leq \theta \leq 20.94^\circ$). Data were collected on an Enraf–Nonius CAD-4 diffractometer using graphite monochromated Mo K α radiation ($\lambda = 0.71073$ Å) by $\theta - 2\theta$ scans within the angular range $3.0 < 2\theta < 45.0^\circ$. Three standard reflections, used to check the crystal stability towards X-ray exposure, showed no significant intensity variation over the course of data collection. X-ray data reduction, and structure solution and refinement were carried out using the SHELXS-97 package. The structure was solved by the direct methods.

Table 1
Crystallographic data

	$[Ru(PPh_3)_2(L^2)Cl_2]$	<i>ctc</i> - $[Ru(PPh_3)_2(L^1)_2]$
Formula	C ₄₄ H ₃₇ Cl ₂ O ₂ P ₂ Ru	C ₅₀ H ₄₀ O ₄ P ₂ Ru
Formula weight	831.65	867.87
Space group	orthorhombic, <i>Pnma</i>	triclinic, <i>P</i> $\bar{1}$
<i>a</i> (Å)	10.5844(10)	10.7368(19)
<i>b</i> (Å)	23.817(2)	11.637(4)
<i>c</i> (Å)	15.407(2)	16.732(7)
α (°)	90	96.18(3)
β (°)	90	91.78(3)
γ (°)	90	99.44(3)
<i>V</i> (Å ³)	3884.0(8)	2047.7(12)
<i>Z</i>	4	2
Crystal size (mm)	0.50 × 0.30 × 0.15	0.40 × 0.30 × 0.20
<i>T</i> (K)	293(2)	298
μ (cm ⁻¹)	6.60	4.951
<i>R</i>	$R_1 = 0.0522^a$	$R_f = 0.046^d$
	$wR_2 = 0.1390^b$	$R_w = 0.047^c$
GOF	1.082 ^e	2.01 ^f

$$^a R_1 = \frac{\sum |F_o| - |F_c|}{\sum |F_o|}$$

$$^b wR_2 = \frac{[\sum [w(F_o^2 - F_c^2)^2]]^{1/2}}{[\sum w(F_o^2)^2]^{1/2}}$$

^c GOF = $[\sum [w(F_o^2 - F_c^2)^2]] / (M - N)^{1/2}$, where *M* is the number of reflections and *N* is the number of parameters refined.

$$^d R_f = \frac{\sum |F_o| - |F_c|}{\sum |F_o|}$$

$$^e R_w = \frac{[\sum w(|F_o| - |F_c|)^2]}{[\sum w(F_o)^2]^{1/2}}$$

^f GOF = $[\sum w(|F_o| - |F_c|)^2] / (M - N)^{1/2}$, where *M* is the number of reflections and *N* is the number of parameters refined.

3. Results and discussion

3.1. Preparation and characterization

3.1.1. $[Ru(PPh_3)_2(L)Cl_2]$ complexes

Reaction of $[Ru(PPh_3)_3Cl_2]$ with an equimolar quantity of each phenolic ligand (HL) proceeds smoothly in dichloromethane solution at ambient temperature to

Table 2
Microanalytical, electronic spectral and cyclic voltammetric data

Compound	Microanalytical data ^a		Electronic spectral data ^b λ_{\max} (nm) (ϵ , M ⁻¹ cm ⁻¹)	Cyclic voltammetric data ^c $E_{1/2}$ (V) (ΔE_p , mV)
	%C	%H		
[Ru(PPh ₃) ₂ (L ¹)Cl ₂]	63.5 (63.2)	4.4 (4.3)	656 (423), 492 ^d (966), 396 (4700), 304 (19300), 272 ^d (22 000) 240 (25 600)	-0.28 (120), 1.16 (120)
Ru(PPh ₃) ₂ (L ²)Cl ₂]	62.6 (63.5)	4.9 (4.5)	624 (907), 388 (5200), 300 ^d (15 100), 268 ^d (23 100), 236 (38800)	-0.37 (120), 1.08 (120)
[Ru(PPh ₃) ₂ (L ³)Cl ₂]	65.3 (65.1)	4.5 (4.3)	588 (855), 400 ^d (2000), 312 ^d (3300), 260 ^d (10700), 232 (19300)	-0.30 (120), 1.24 (120)
ctc-[Ru(PPh ₃) ₂ (L ¹) ₂]	69.7 (69.2)	4.7 (4.6)	504 (2300), 404 ^d (7000), 364 (10 400), 232 (78 300)	0.23 (80), 1.35 (100)
ccc-[Ru(PPh ₃) ₂ (L ¹) ₂]	69.5 (69.2)	4.5 (4.6)	500 (1300), 408 ^d (3700), 356 (5000), 232 (38600)	0.28 (80), 1.41 (100)
ctc-[Ru(PPh ₃) ₂ (L ²) ₂]	70.1 (69.7)	5.0 (4.9)	492 (1800), 388 ^d (5000), 340 (7800), 260 (22 900), 236 (23600)	0.09 (80), 1.31 (110)
ccc-[Ru(PPh ₃) ₂ (L ²) ₂]	69.9 (69.7)	4.9 (4.9)	500 ^d (2300), 400 (6300), 336 (8200), 232 (64 600)	0.21 (80), 1.52 (120)
ctc-[Ru(PPh ₃) ₂ (L ³) ₂]	72.1 (72.0)	4.7 (4.6)	490 (3000), 390 (9600), 340 ^d (12 200), 320 ^d (17 700), 280 (32 700), 232 (77 600)	0.35 (80), 1.46 (100)
ccc-[Ru(PPh ₃) ₂ (L ³) ₂]	72.2 (72.0)	4.6 (4.6)	490 (3000), 398 (7400), 340 (11 200), 320 ^d (17 200), 276 (30 000), 232 (61 300)	0.41 (80), 1.52 (110)


^a Calculated values are in parenthesis.

^b Dichloromethane solution.

^c Solvent, acetonitrile; supporting electrolyte, TEAP; reference electrode, SCE; $E_{1/2} = 0.5(E_{pa} + E_{pc})$, where E_{pa} and E_{pc} are anodic and cathodic peak potentials, respectively; $\Delta E_p = E_{pa} - E_{pc}$; scan rate, 50 mV s⁻¹.

^d Shoulder.

afford complexes of the type [Ru(PPh₃)₂(L)Cl₂]. It is interesting to note here that ruthenium has undergone a one-electron oxidation during the course of this synthetic reaction. In view of the ruthenium(III)–ruthenium(II) reduction potentials in these complexes (vide infra), oxygen in air appears to have served as the oxidant. Compositions of the complexes have been confirmed by their microanalytical data (Table 2). As all the three phenolate ligands used in the present study are asymmetric bidentate in nature, the [Ru(PPh₃)₂(L)Cl₂] complexes may exist in three geometric isomeric forms (3–5).



To distinguish between the possible three isomers, molecular structure of a representative complex, viz. [Ru(PPh₃)₂(L²)Cl₂], has been determined by X-ray crystallography. The structure is shown in Fig. 1 and

selected bond distances and angles are presented in Table 3. The O₂P₂Cl₂ coordination sphere around ruthenium is distorted octahedral in nature, which is




Fig. 1. View of the [Ru(PPh₃)₂(L²)Cl₂] molecule.

Table 3
Selected bond distances (Å) and bond angles (°) for
[Ru(PPh₃)₂(L²)Cl₂]

Ru–Cl(1)	2.324(2)	O(2)–C(7)	1.298(12)
Ru–Cl(2)	2.304(2)	O(1)–C(1)	1.267(10)
Ru–P(1)	2.415(2)	C(1)–C(2)	1.405(14)
Ru–O(1)	1.995(5)	C(2)–C(3)	1.45(3)
Ru–O(2)	2.025(6)	C(3)–C(4)	1.39(5)
		C(4)–C(5)	1.22(4)
		C(5)–C(6)	1.44(2)
		C(6)–C(7)	1.42(2)
		C(1)–C(6)	1.38(2)
		C(7)–C(8)	1.494(13)
O(2)–Ru–Cl(2)	176.9(2)		
P(1)–Ru–P(1A)	179.41(8)		
O(1)–Ru–Cl(1)	173.2(2)		
O(1)–Ru–O(2)	86.4(2)	O(1)–Ru–Cl(2)	90.5(2)
O(2)–Ru–Cl(1)	86.9(2)	C(2)–Ru–Cl(1)	96.27(11)
O(1)–Ru–P(1)	89.73(4)	O(2)–Ru–P(1)	90.10(5)
Cl(2)–Ru–P(1)	89.88(5)	Cl(1)–Ru–P(1)	90.28(4)




Fig. 2. Electronic spectra of (a) [Ru(PPh₃)₂(L³)Cl₂] and (b) *cis*-[Ru(PPh₃)₂(L¹)₂] (—) and *ccc*-[Ru(PPh₃)₂(L¹)₂] (-----) in dichloromethane solution.

reflected in the bond parameters. The two bulky PPh₃ ligands are in *trans* positions, as is usually observed in complexes of ruthenium(III) having the Ru(PPh₃)₂ moiety [32–34], while the two chloride ligands have occupied *cis* positions. The observed bond distances are all quite normal as observed in structurally characterized complexes of ruthenium containing similar ligands

[19,32–34]. This structure determination thus shows that [Ru(PPh₃)₂(L²)Cl₂] has structure **3**. As all the three [Ru(PPh₃)₂(L)Cl₂] complexes display similar properties (*vide infra*), the other two [Ru(PPh₃)₂(L)Cl₂] complexes are assumed to have a similar structure as [Ru(PPh₃)₂(L²)Cl₂].

Infrared spectra of the [Ru(PPh₃)₂(L)Cl₂] complexes show many sharp and strong vibrations in the 1600–300 cm⁻¹ region, of which the vibrations near 520, 700 and 745 cm⁻¹ are also observed in [Ru(PPh₃)₂Cl₂] and hence these are attributable to the Ru(PPh₃)₂ fragment of the [Ru(PPh₃)₂(L)Cl₂] complexes. One new band, observed near 1580 cm⁻¹ in all these complexes, is assigned to the ν(C=O) vibration of the coordinated phenolate ligand. The ν(Ru–Cl) stretching vibration appears as a strong band in all the complexes near 330 cm⁻¹. The infrared spectral data thus correspond to the composition of the complexes. The [Ru(PPh₃)₂(L)Cl₂] complexes are soluble in common polar organic solvents, like dichloromethane, chloroform, acetonitrile, etc., producing green solutions. Electronic spectra of these complexes have been recorded in dichloromethane solution. A selected spectrum is shown in Fig. 2 and spectral data are listed in Table 2. Each complex shows few intense absorptions in the visible region together with some very intense absorptions in the ultraviolet region. The absorptions in the ultraviolet region may be assigned to transitions occurring within the ligand orbitals. To have an insight into the nature of transitions appearing in the visible region, qualitative EHMO calculations have been performed [35,36] on a model of the [Ru(PPh₃)₂(L)Cl₂] complexes, which was computer generated from [Ru(PPh₃)₂(L¹)Cl₂] by replacing the phenyl groups of the PPh₃ ligands with hydrogen. A partial MO diagram is shown in Fig. 3. The highest occupied (singly occupied) molecular orbital (MO-1) and the next two filled orbitals (MO-2 and MO-3) of this model are predominantly (≥ 80%) ruthenium t₂ in character. There are two filled molecular orbitals (MO-4 and MO-5) below these metal t₂ orbitals, which are localized almost entirely (≥ 96%) on the phenolate ligand. The intense absorptions observed in the visible region may therefore be assigned to the allowed ligand-to-metal charge-transfer transitions occurring from the filled orbitals of the phenolate ligand (MO-4 and MO-5) to the half-filled ruthenium t₂ orbital (MO-1).

Magnetic susceptibility show that all the three [Ru(PPh₃)₂(L)Cl₂] complexes are one-electron paramagnetic, which is in accordance with the +3 oxidation state of ruthenium (low-spin d⁵, S = 1/2) in these complexes. Electron spin resonance spectra of the [Ru(PPh₃)₂(L)Cl₂] complexes, recorded in 1:1 dichloromethane–toluene solution at 77 K, show rhombic spectra with three distinct signals (g₁, g₂ and g₃ in the order of decreasing magnitude). A representative spectrum is shown in Fig. 4 and the spectral data are




Fig. 3. Qualitative molecular orbital diagram of $[\text{Ru}(\text{PPh}_3)_2(\text{L})\text{Cl}_2]$.

presented in Table 4. The observed rhombicity of the ESR spectra is understandable in terms of the gross molecular symmetry of these complexes containing the three non-equivalent P–Ru–P, O(phenolic)–Ru–Cl and O(carbonylic)–Ru–Cl axes. The rhombic distortion can be thought of a combination of axial distortion (A , which splits t_2 into a and e) and rhombic distortion (V , which splits e). The splitting pattern is illustrated in Fig. 4. Spin-orbit coupling causes further changes in the energy gaps. Thus two electronic transitions (transition energies ΔE_1 and ΔE_2 ; $\Delta E_1 < \Delta E_2$) are possible within these three levels. All these energy parameters have been computed (Table 4) using the observed g values, the g tensor theory of low-spin d^5 complexes and a reported method [37–39]. The axial distortion is observed to be much stronger than the rhombic one. The ΔE_1 transition falls in the infrared region ($3200\text{--}3700\text{ cm}^{-1}$) and could not be detected. The ΔE_2 transition, which is expected to occur near 6000 cm^{-1} ($\sim 1667\text{ nm}$), could not be verified either because of the non-transparency of the solvent in this region. However, the ESR data analysis shows that the $[\text{Ru}(\text{PPh}_3)_2(\text{L})\text{Cl}_2]$ complexes are significantly distorted from ideal octahedral geometry, which was also observed in the structural analysis of $[\text{Ru}(\text{PPh}_3)_2(\text{L}_2)\text{Cl}_2]$.

3.1.2. $[\text{Ru}(\text{PPh}_3)_2(\text{L})_2]$ complexes

These complexes have been prepared in two different ways (Scheme 1). Direct reaction of the phenolic lig-

ands (HL) with $[\text{Ru}(\text{PPh}_3)_3\text{Cl}_2]$ in 2:1 mole ratio proceeds smoothly in refluxing ethanol in the presence of a base to afford the bis complexes of type $[\text{Ru}(\text{PPh}_3)_2(\text{L})_2]$. These complexes can also be prepared from $[\text{Ru}(\text{PPh}_3)_2(\text{L})\text{Cl}_2]$ by reacting them with one equivalent of the respective phenolic ligand (HL) in the presence of a base. Thin layer chromatographic experiments indicated the presence of two isomers (isomer-I and isomer-II) in all $[\text{Ru}(\text{PPh}_3)_2(\text{L})_2]$ complexes, which have




Fig. 4. ESR spectrum of $[\text{Ru}(\text{PPh}_3)_2(\text{L})\text{Cl}_2]$ in 1:1 dichloromethane-toluene solution at 77 K.

Table 4
Magnetic moment and ESR g values ^a and derived parameters ^b of the $[\text{Ru}(\text{PPh}_3)_2(\text{L})\text{Cl}_2]$ complexes


Compound	μ_{eff} (BM)	g_1	g_2	g_3	Δ/λ	V/λ	$\Delta E_1/\lambda$	$\Delta E_2/\lambda$
$[\text{Ru}(\text{PPh}_3)_2(\text{L}^1)\text{Cl}_2]$	1.85	2.497	2.212	1.791	4.313	-2.450	3.236	5.801
$[\text{Ru}(\text{PPh}_3)_2(\text{L}^2)\text{Cl}_2]$	1.93	2.428	2.233	1.828	4.574	-1.976	3.691	5.830
$[\text{Ru}(\text{PPh}_3)_2(\text{L}^3)\text{Cl}_2]$	1.97	2.407	2.258	1.795	4.065	-1.264	3.515	5.029

^a In 1:1 dichloromethane–toluene solution at 77 K.

^b Spin-orbit coupling constant (λ) for complexed ruthenium(III) is $\sim 1000 \text{ cm}^{-1}$.

been separated by column chromatography. Microanalytical data of these complexes are in good agreement with their compositions (Table 2). Both isomers of these complexes are diamagnetic, which corresponds to the bivalent state of ruthenium (low-spin d^6 , $S = 0$) in these complexes.

As all three phenolate ligands used in the present study are asymmetric, the $[\text{Ru}(\text{PPh}_3)_2(\text{L})_2]$ complexes may exist in five geometrical isomeric forms (6–10). Isomers 6–9 have a C_2 axis, while isomer 10 does not have any C_2 symmetry. ^1H NMR spectra have been recorded on both isomers of all three $[\text{Ru}(\text{PPh}_3)_2(\text{L})_2]$ complexes in CDCl_3 solution. Isomer-I of all three complexes clearly shows the presence of a C_2 axis, while isomer-II indicates the absence of C_2 symmetry. For example, isomer-I of $[\text{Ru}(\text{PPh}_3)_2(\text{L}^1)_2]$ shows only one aldehydic proton signal at 8.68 ppm, while isomer-II of $[\text{Ru}(\text{PPh}_3)_2(\text{L}^1)_2]$ shows two aldehydic proton signals (1H each) at 8.36 and 8.53 ppm. This shows that isomer-II has structure 10, where the three sets of donor atoms are in *cis* positions and henceforth this isomer will be labeled as the *cis-cis-cis* or *ccc* isomer.



It has not been possible to assign specific stereochemistry of isomer-I on the basis of ^1H NMR spectral results alone. However, as complexes of ruthenium with the $\text{Ru}(\text{PPh}_3)_2$ are known to prefer *cis*-disposition of PPh_3 ligands [33,34,40], these $[\text{Ru}(\text{PPh}_3)_2(\text{L})_2]$ complexes may be assumed to have either structure 8 or 9. To sort out this problem of stereochemistry of isomer-I,

the molecular structure of isomer-I of $[\text{Ru}(\text{PPh}_3)_2(\text{L}^1)_2]$ has been determined by X-ray crystallography. The structure is shown in Fig. 5 and selected bond distances and angles are presented in Table 5. Ruthenium has a distorted octahedral O_4P_2 coordination sphere with the two PPh_3 ligands in *cis* positions, the two phenolate oxygens in *trans* positions and the two carbonylic oxygens in *cis* positions. Therefore isomer-I of $[\text{Ru}(\text{PPh}_3)_2(\text{L}^1)_2]$ has structure 9 (henceforth referred to as the *ctc* isomer to indicate the *cis-trans-cis* dispositions of the carbonylic oxygens, phenolate oxygens and phosphines, respectively). The observed bond parameters are all quite normal. In view of the similarity in synthetic procedure and properties, isomer-I of the other two $[\text{Ru}(\text{PPh}_3)_2(\text{L})_2]$ complexes are assumed to have similar *ctc* structure.




Fig. 5. View of the *ctc*- $[\text{Ru}(\text{PPh}_3)_2(\text{L}^1)_2]$ molecule.

Table 5
Selected bond distances (Å) and bond angles (°) for $[\text{Ru}(\text{PPh}_3)_2(\text{L}^1)_2]$

Ru–P(1)	2.2929(18)	O(1)–C(1)	1.295(6)
Ru–P(2)	2.3031(17)	O(2)–C(7)	1.239(7)
Ru–O(1)	2.057(3)	O(3)–C(8)	1.313(6)
Ru–O(2)	2.090(4)	O(4)–C(14)	1.262(7)
Ru–O(3)	2.052(3)		
Ru–O(4)	2.128(4)		
P(1)–Ru–O(2)	169.48(11)		
P(2)–Ru–O(4)	169.69(11)		
O(1)–Ru–O(3)	173.61(14)		
P(1)–Ru–P(2)	99.22(6)	P(2)–Ru–O(3)	87.90(11)
P(1)–Ru–O(1)	90.17(11)	O(1)–Ru–O(2)	90.75(14)
P(1)–Ru–O(3)	93.05(11)	O(1)–Ru–O(4)	83.48(14)
P(1)–Ru–O(4)	91.07(11)	O(2)–Ru–O(3)	85.08(14)
P(2)–Ru–O(1)	97.04(11)	O(2)–Ru–O(4)	78.62(14)
P(2)–Ru–O(2)	91.08(11)	O(3)–Ru–O(4)	90.94(14)

Infrared spectra of *ctc* and *ccc* isomers of the $[\text{Ru}(\text{PPh}_3)_2(\text{L})_2]$ complexes are very similar. Each shows characteristic vibrations near 500, 700, 750 cm^{-1} , indicating the presence of the $\text{Ru}(\text{PPh}_3)_2$ moiety. The $\nu(\text{C}=\text{O})$ vibration is observed as a strong band near 1580 cm^{-1} in all these complexes. The $[\text{Ru}(\text{PPh}_3)_2(\text{L})_2]$ complexes are soluble in common organic solvents like dichloromethane, chloroform, acetone, acetonitrile, etc., forming intense red solutions. Electronic spectra of these complexes, recorded in dichloromethane solution, show several intense absorptions in the visible region and few absorptions of very high intensity in the ultraviolet region (Fig. 2, Table 2). The latter absorptions are assigned to transitions within the ligand orbitals. The former absorptions in the visible region are probably due to allowed metal-to-ligand charge-transfer transitions. Multiple charge-transfer transitions are common in such mixed ligand complexes, primarily

because of the lower symmetry splitting of the metal orbitals and presence of different accepting orbitals. For proper assignment of the absorptions in the visible region, qualitative EHMO calculations have been performed as before on models of the *ctc* and *ccc* isomers of $[\text{Ru}(\text{PPh}_3)_2(\text{L}^1)_2]$, where phenyl groups of the PPh_3 ligands have been replaced by hydrogen. The results of these calculations are qualitatively very similar for both the isomers. A partial MO diagram for one isomer is shown in Fig. 6. The highest occupied molecular orbital (HOMO) and the next two filled orbitals (HOMO-1 and HOMO-2) are basically (> 75%) ruthenium t_2 orbitals. There are two relatively close vacant molecular orbitals above these filled orbitals, the lowest unoccupied molecular orbital (LUMO) and the next unoccupied orbital (LUMO + 1), which are primarily (> 95%) π -orbitals of the phenolate ligands. The intense absorptions observed in the visible region may therefore be assigned to the charge-transfer transitions occurring from the filled ruthenium t_2 orbitals to the vacant π -orbitals of the phenolate ligands.

3.2. Electron-transfer properties

Electron-transfer properties of the $[\text{Ru}(\text{PPh}_3)_2(\text{L})\text{Cl}_2]$ and $[\text{Ru}(\text{PPh}_3)_2(\text{L})_2]$ have been studied in acetonitrile solution by cyclic voltammetry. Representative voltammograms are shown in Fig. 7 and voltammetric data are presented in Table 2.

Each $[\text{Ru}(\text{PPh}_3)_2(\text{L})\text{Cl}_2]$ complex shows a reductive response on the negative side of SCE and an oxidative response on the positive side. Both responses are quasi-reversible in nature. The reduction, observed near -0.3 V (all potentials are referenced to SCE), is tentatively assigned to ruthenium(III)-ruthenium(II) reduction and the oxidation, which occurs within 1.08–




Fig. 6. Qualitative molecular orbital diagram of $[\text{Ru}(\text{PPh}_3)_2(\text{L})_2]$.




Fig. 7. Cyclic voltammograms of (a) $[\text{Ru}(\text{PPh}_3)_2(\text{L}^3)\text{Cl}_2]$ and (b) $\text{ctc-}[\text{Ru}(\text{PPh}_3)_2(\text{L}^1)_2]$ (—) and $\text{ccc-}[\text{Ru}(\text{PPh}_3)_2(\text{L}^1)_2]$ (-----) in acetonitrile solution (0.1 M TEAP) at a scan rate of 50 mV s^{-1} .

1.24 V, is tentatively assigned to ruthenium(III)–ruthenium(IV) oxidation. Both potentials have shifted to a little more negative values in $[\text{Ru}(\text{PPh}_3)_2(\text{L}^2)\text{Cl}_2]$ relative to $[\text{Ru}(\text{PPh}_3)_2(\text{L}^1)\text{Cl}_2]$, which may be attributed to the electron donating character of the methyl group in L^2 .

Both isomers of $[\text{Ru}(\text{PPh}_3)_2(\text{L})_2]$ complexes show two oxidative responses on the positive side of SCE. The first oxidation, which is observed within 0.09–0.41 V, is reversible in nature and is assigned to ruthenium(II)–ruthenium(III) oxidation. The second quasi-reversible oxidation appears within 1.31–1.52 V and is tentatively assigned to the ruthenium(III)–ruthenium(IV) oxidation. Potentials of both oxidations have been observed to shift to lower values in the ctc isomer than the ccc isomer, which indicates that the lower oxidation states are more comfortable in the ccc isomer than in the ctc isomer. This is in accordance with the higher probability of $d\pi\text{-}p\pi$ backbonding in the ccc isomer [41–43]. The oxidation potentials are observed to be a little lower in $[\text{Ru}(\text{PPh}_3)_2(\text{L}^2)_2]$ than in $[\text{Ru}(\text{PPh}_3)_2(\text{L}^1)_2]$, as before. The cyclic voltammetric studies indicate that the +3 state of ruthenium is stable in the $\text{O}_2\text{P}_2\text{Cl}_2$ coordination sphere in the $[\text{Ru}(\text{PPh}_3)_2(\text{L})\text{Cl}_2]$ complexes and so is the +2 state of ruthenium in the O_4P_2 coordination sphere in the $[\text{Ru}(\text{PPh}_3)_2(\text{L})_2]$ complexes.

4. Supplementary data

Supplementary data are available from the Cambridge Crystallographic Data Center, 12 Union Road, Cambridge, CB2 1EZ, UK, on request, quoting the deposition numbers CCDC 138541 and 138542.

Acknowledgements

Financial assistance received from the Council of Scientific and Industrial Research, New Delhi [Grant No. 01(1408)/96/EMR-II] is gratefully acknowledged. Thanks are also due to the Third World Academy of Sciences for financial support for the purchase of an electrochemical cell system. F.B. thanks the University Grants Commission, New Delhi, for her fellowship.

References

- [1] S.-M. Lee, W.-T. Wong, *Coord. Chem. Rev.* 164 (1997) 415.
- [2] I. Ortmanes, C. Moucheron, A.K.-D. Mesmaeker, *Coord. Chem. Rev.* 168 (1998) 233.
- [3] A. Islam, N. Ikeda, K. Nozak, Y. Okamoto, B. Gholamkhash, A. Yoshimura, T. Ohno, *Coord. Chem. Rev.* 171 (1998) 355.
- [4] J.W.-S. Hui, Y.-T. Wong, *Coord. Chem. Rev.* 172 (1998) 389.
- [5] P.J. Dyson, B.F.G. Johnson, C.M. Martin, *Coord. Chem. Rev.* 175 (1998) 59.
- [6] D.J. Stufkens, A. Vlcek Jr., *Coord. Chem. Rev.* 177 (1998) 127.
- [7] L.D. Cola, P. Belsler, *Coord. Chem. Rev.* 177 (1998) 301.
- [8] S. Sabo-Etienne, B. Chaudret, *Coord. Chem. Rev.* 178–180 (1998) 381.
- [9] D. Touchard, P.H. Dixneuf, *Coord. Chem. Rev.* 178–180 (1998) 409.
- [10] G. Jia, C.-P. Lau, *Coord. Chem. Rev.* 190–192 (1999) 83.
- [11] N.C. Pramanik, S. Bhattacharya, *J. Chem. Res. (S)* (1997) 98.
- [12] F. Basuli, S.M. Peng, S. Bhattacharya, *Inorg. Chem.* 36 (1997) 5645.
- [13] N.C. Pramanik, K. Pramanik, P. Ghosh, S. Bhattacharya, *Polyhedron* 17 (1998) 1525.
- [14] F. Basuli, M. Ruf, C.G. Pierpont, S. Bhattacharya, *Inorg. Chem.* 37 (1998) 6113.
- [15] N.C. Pramanik, S. Bhattacharya, *Transition Met. Chem.* 24 (1999) 95.
- [16] K. Sui, S.M. Peng, S. Bhattacharya, *Polyhedron* 18 (1999) 631.
- [17] A.K. Das, S.M. Peng, S. Bhattacharya, *J. Chem. Soc., Dalton Trans.* (2000) 181.
- [18] F. Basuli, S.M. Peng, S. Bhattacharya, *Inorg. Chem.* 39 (2000) 1120.
- [19] N. Bag, G.K. Lahiri, S. Bhattacharya, L.R. Falvello, A. Chakravorty, *Inorg. Chem.* 27 (1988) 4396.
- [20] F. Basuli, S.M. Peng, S. Bhattacharya, *Polyhedron* 17 (1998) 2191.
- [21] G.K. Lahiri, S. Bhattacharya, B.K. Ghosh, A. Chakravorty, *Inorg. Chem.* 26 (1987) 4324.
- [22] S. Bhattacharya, S.R. Boone, G.A. Fox, C.G. Pierpont, *Inorg. Chem.* 29 (1990) 1088.
- [23] S. Bhattacharya, *Polyhedron* 12 (1993) 235.
- [24] B. Muller, H. Vahrenkamp, *Chem. Eur. J.* (1998) 117.
- [25] B. Muller, H. Vahrenkamp, *Chem. Eur. J.* (1998) 129.
- [26] B. Muller, H. Vahrenkamp, *Chem. Eur. J.* (1998) 137.
- [27] B. Muller, A. Schneider, M. Tesmer, H. Vahrenkamp, *Inorg. Chem.* 38 (1999) 1900.
- [28] T.R. Ward, A. Lutz, S.P. Parel, J. Ensling, P. Gutlich, P. Buglyo, C. Orvig, *Inorg. Chem.* 38 (1999) 5007.
- [29] T.A. Stephenson, G. Wilkinson, *J. Inorg. Nucl. Chem.* 28 (1966) 945.
- [30] D.T. Sawyer, J.L. Roberts Jr., *Experimental Electrochemistry for Chemists*, Wiley, New York, 1974, pp. 167–215.
- [31] M. Walter, L. Ramaley, *Anal. Chem.* 45 (1973) 165.
- [32] S. Chattopadhyay, N. Bag, G.K. Lahiri, A. Chakravorty, *J. Chem. Soc., Dalton Trans.* (1990) 3389.

- [33] S. Bhattacharya, C.G. Pierpont, *Inorg. Chem.* 30 (1991) 1511.
- [34] M. Menon, A. Pramanik N. Bag, A. Chakravorty, *J. Chem. Soc., Dalton Trans.* (1995) 1417.
- [35] C. Mealli, D.M. Proserpio, CACAO Version 4.0, Firenze, Italy, July 1994.
- [36] C. Mealli, D.M. Proserpio, *J. Chem. Educ.* 67 (1990) 399.
- [37] B. Bleany, M.C.M. O'Brien, *Proc. Phys. Soc. Lond., Sect B* 69 (1956) 1216.
- [38] J.S. Griffith, *The Theory of Transition Metal Ions*, Cambridge University Press, London, 1961, p. 364.
- [39] S. Bhattacharya, A. Chakravorty, *Proc. Indian Acad. Sci.* 95 (1985) 159.
- [40] A. Pramanik, N. Bag, G.K. Lahiri, A. Chakravorty, *J. Chem. Soc., Dalton Trans.* (1990) 3823.
- [41] R.A. Krause, K. Krause, *Inorg. Chem.* 19 (1980) 2600.
- [42] S. Goswami, A.R. Chakravarty, A. Chakravorty, *Inorg. Chem.* 20 (1981) 2246.
- [43] R.A. Krause, K. Krause, *Inorg. Chem.* 21 (1982) 1714.