

Contents lists available at [ScienceDirect](http://www.sciencedirect.com)

Biochemical and Biophysical Research Communications

journal homepage: www.elsevier.com/locate/ybbrc

Membrane cholesterol depletion enhances ligand binding function of human serotonin_{1A} receptors in neuronal cells

Rajesh Prasad¹, Yamuna Devi Paila¹, Amitabha Chattopadhyay^{*}

Centre for Cellular and Molecular Biology, Council of Scientific and Industrial Research, Uppal Road, Hyderabad 500 007, India

ARTICLE INFO

Article history:

Received 12 September 2009

Available online 23 September 2009

Keywords:

Membrane cholesterol
Ligand binding function
Neuronal cells
Serotonin_{1A} receptor
Fluorescence anisotropy

ABSTRACT

Membrane lipid composition of cells in the nervous system is unique and displays remarkable diversity. Cholesterol metabolism and homeostasis in the central nervous system and their role in neuronal function represent important determinants in neuropathogenesis. The serotonin_{1A} receptor is an important member of the G-protein coupled receptor superfamily, and is involved in a variety of cognitive, behavioral, and developmental functions. We report here, for the first time, that the ligand binding function of human serotonin_{1A} receptors exhibits an increase in membranes isolated from cholesterol-depleted neuronal cells. Our results gain pharmacological significance in view of the recently described structural evidence of specific cholesterol binding site(s) in GPCRs, and could be useful in designing better therapeutic strategies for neurodegenerative diseases associated with GPCRs.

© 2009 Elsevier Inc. All rights reserved.

Introduction

The G-protein coupled receptor (GPCR) superfamily is the largest and most diverse protein family in mammals, involved in signal transduction across membranes [1]. GPCRs are increasingly recognized as major targets for the development of novel drug candidates in all clinical areas [2,3]. The serotonin_{1A} (5-HT_{1A}) receptor is an important neurotransmitter receptor belonging to the GPCR superfamily. It is the most extensively studied of the serotonin receptors for a number of reasons [4,5]. Serotonergic signaling plays a key role in the generation and modulation of various cognitive, behavioral and developmental functions [6]. This is supported by the fact that the agonists and antagonists of the serotonin_{1A} receptor represent major classes of molecules with potential therapeutic effects in anxiety- and stress-related disorders. Interestingly, mutant (knockout) mice lacking the serotonin_{1A} receptor exhibit enhanced anxiety-related behavior, and represent an important animal model for genetic vulnerability to complex traits such as anxiety disorders and aggression in higher animals [7,8]. In the context of increasing pharmacological relevance of the serotonin_{1A} receptor, a transmembrane protein, its interaction with sur-

rounding lipids assumes significance in modulating the function of the receptor in healthy and diseased states [5].

Cholesterol is an essential and representative lipid in higher eukaryotic cellular membranes and is crucial in membrane organization, dynamics, function, and sorting [9–11]. It is often found distributed non-randomly in domains in biological and model membranes. Many of these domains (sometimes termed as ‘lipid rafts’) are believed to be important for the maintenance of membrane structure and function. Nonetheless, characterizing the spatiotemporal resolution of these domains has proven to be challenging [11–14]. The idea of such specialized membrane domains assumes relevance in the cellular context since physiologically important functions such as membrane sorting and trafficking [15] signal transduction processes [16], and the entry of pathogens [17–19] have been attributed to these domains. We have earlier shown that membrane cholesterol is essential for the function of the serotonin_{1A} receptor (reviewed in [5] and [20]).

Mammalian cells in culture heterologously expressing membrane receptors represent convenient systems to address problems in receptor biology due to higher expression levels of receptors [21]. An important consideration in such expression systems is selecting a cell type which is derived from the tissue of natural occurrence of the receptor. This is particularly true for receptors of neural origin since the membrane lipid composition of cells in the nervous system is unique and displays remarkable diversity [22–24]. Lipids found in neuronal membranes are often necessary for maintaining the structure and function of neuronal receptors. Keeping this in mind, we earlier reported the pharmacological and functional characterization of the human serotonin_{1A} receptor stably expressed in HN2 cells [25], which are a hybrid cell line between hippocampal

Abbreviations: 5-HT_{1A} receptor, 5-hydroxytryptamine-1A receptor; 8-OH-DPAT, 8-hydroxy-2-(di-*N*-propylamino)tetralin; BCA, bicinechonic acid; DMPC, 1,2-dimyristoyl-*sn*-glycero-3-phosphocholine; DPH, 1,6-diphenyl-1,3,5-hexatriene; GPCR, G-protein coupled receptor; M β CD, methyl- β -cyclodextrin; PMSF, phenylmethylsulfonyl fluoride.

^{*} Corresponding author. Fax: +91 40 2716 0311.

E-mail address: amit@cmb.res.in (A. Chattopadhyay).

¹ These authors contributed equally to the work.

cells and mouse neuroblastoma. Our results showed that the human serotonin_{1A} receptor expressed in HN2 cells displays characteristic features found in the native receptor isolated from bovine hippocampus and represents a realistic model system for the receptor. In this paper, we monitored the function of the human serotonin_{1A} receptor stably expressed in HN2 cells upon depletion of membrane cholesterol from live cells. Cholesterol depletion from cell membranes was achieved using methyl- β -cyclodextrin (M β CD). The corresponding changes in membrane dynamics were monitored by fluorescence anisotropy of the membrane probe DPH. Importantly, our results show that the ligand binding function of serotonin_{1A} receptors is enhanced in membranes isolated from cholesterol-depleted neuronal cells. This constitutes the first report describing functional changes in serotonin_{1A} receptors upon membrane cholesterol depletion in neuronal cells.

Materials and methods

Materials. Cholesterol, M β CD, DMPC, DPH, EDTA, MgCl₂, MnCl₂, Na₂HPO₄, PMSF, serotonin, polyethylenimine, and Tris were obtained from Sigma Chemical Co. (St. Louis, MO). DMEM (Dulbecco's modified Eagle's medium), fetal calf serum, and geneticin (G 418) were from Invitrogen Life Technologies (Carlsbad, CA). Amplex Red cholesterol assay kit was from Molecular Probes (Eugene, OR). BCA reagent for protein estimation was from Pierce (Rockford, IL). [³H]8-OH-DPAT (sp. activity 106 Ci/mmol) was purchased from DuPont New England Nuclear (Boston, MA). GF/B glass microfiber filters were from Whatman International (Kent, UK). Water was purified through a Millipore (Bedford, MA) Milli-Q system and used throughout. All other chemicals and solvents used were of the highest available purity.

Cells and cell culture. The intronless human genomic clone G-21 [26] which encodes the human serotonin_{1A} receptor was used to generate stable transfectants in HN2 cells which are a hybrid cell line between hippocampal cells and mouse neuroblastoma [27]. These cells stably expressing the human serotonin_{1A} receptor are referred to as HN2-5-HT_{1A}R cells [25]. HN2-5-HT_{1A}R cells were maintained as described earlier [25,28].

Cholesterol depletion of neuronal cells in culture. Cells at density of 2×10^6 in 150 cm² flasks were grown for 3 days in DMEM supplemented with 10% serum, followed by incubation in serum-free DMEM for 3 h at 37 °C. Cholesterol depletion was carried out by treating cells with increasing concentrations of M β CD in serum-free DMEM for 30 min at 37 °C, followed by wash with 10 mM sodium phosphate, 150 mM sodium chloride, pH 7.4 buffer.

Cell membrane preparation. Cell membranes were prepared as described earlier [25]. Total protein concentration in isolated membranes was determined using the BCA assay [29].

Radioligand binding assay. Receptor binding assays in membranes isolated from control and cholesterol-depleted HN2-5-HT_{1A}R cells were carried out as described earlier [25,30] with ~200 μ g total protein. The concentration of [³H]8-OH-DPAT used was 0.29 nM.

Analysis of cholesterol and phospholipid contents. Cholesterol content in cell membranes was estimated using the Amplex Red cholesterol assay kit [31]. Total phospholipid content of membranes was determined subsequent to digestion with perchloric acid [32] using Na₂HPO₄ as standard. DMPC was used as an internal standard to assess lipid digestion. Samples without perchloric acid digestion produced negligible readings.

Fluorescence anisotropy measurements. Steady state fluorescence anisotropy measurements were carried out in a Hitachi F-4010 spectrofluorometer using the fluorescent probe DPH incorporated in membranes isolated from control and cholesterol-depleted HN2-5-HT_{1A}R cells, as mentioned earlier [33]. Fluorescence

anisotropy measurements were performed using a Hitachi polarization accessory. Anisotropy values were calculated from the equation [34]:

$$r = \frac{I_{VV} - GI_{VH}}{I_{VV} + 2GI_{VH}}$$

where I_{VV} and I_{VH} are the measured fluorescence intensities (after appropriate background subtraction) with the excitation polarizer vertically oriented and the emission polarizer vertically and horizontally oriented, respectively. G is the grating correction factor and is equal to I_{HV}/I_{HH} . Experiments were done with multiple sets of samples and average values of fluorescence anisotropy are shown in Fig. 4.

Statistical analysis: Significance levels were estimated using Student's two-tailed unpaired t -test using Graphpad Prism software version 4.0 (San Diego, CA).

Results and discussion

We monitored specific binding of the serotonin_{1A} receptor agonist [³H]8-OH-DPAT to membranes isolated from HN2-5-HT_{1A}R cells. Fig. 1 shows that the binding of the [³H]8-OH-DPAT is linear over a broad range (30–200 μ g) of total protein. These results suggest that under the conditions of the assay, there is no depletion of the radiolabel during the course of the assay. These conditions are therefore appropriate for analyzing binding parameters of the receptor using [³H]8-OH-DPAT [35].

M β CD is a water-soluble cyclic oligosaccharide, and has earlier been shown to extract cholesterol from membranes in a selective and efficient manner by including it in a central nonpolar cavity [36]. Fig. 2 shows the cholesterol content in membranes isolated from control and cholesterol-depleted HN2-5-HT_{1A}R cells. Treatment of cells with increasing concentrations of M β CD results in a progressive depletion of membrane cholesterol. When membranes were treated with 10 mM M β CD, the cholesterol content was reduced to ~35% of that of control (without treatment). The concentration range of M β CD was carefully chosen to minimize any possible change in membrane phospholipid content. The phospholipid content remains invariant under these conditions (see inset of Fig. 2).

Fig. 3 shows the increase in specific [³H]8-OH-DPAT binding in membranes isolated from HN2-5-HT_{1A}R cells upon treatment with increasing concentrations of M β CD. For example, specific agonist binding is enhanced by ~47% of the control (in the absence of




Fig. 1. Fold change in specific binding of the agonist [³H]8-OH-DPAT to serotonin_{1A} receptors from HN2-5-HT_{1A}R cell membranes with increasing amounts of total membrane protein. Values have been normalized with respect to specific binding obtained with 110 μ g total protein in the assay. Concentration of [³H]8-OH-DPAT used in the assay was 0.29 nM. Data represent means \pm SE from three independent measurements. See Materials and methods for other details.




Fig. 2. Lipid contents of membranes isolated from HN2-5-HT_{1A}R cells upon cholesterol depletion. The figure depicts cholesterol contents in membranes isolated from control cells and cells treated with increasing concentrations of MβCD. The inset shows the corresponding phospholipid content in membranes isolated from cholesterol-depleted HN2-5-HT_{1A}R cells. Values are expressed as percentages of the respective lipid content in control (without MβCD treatment) cell membranes. Data shown are means ± SE from at least four independent measurements. See Materials and methods for other details.




Fig. 3. Effect of cholesterol depletion on specific binding of the agonist to human serotonin_{1A} receptors in membranes isolated from control and MβCD-treated HN2-5-HT_{1A}R cells. The figure shows specific [³H]-OH-DPAT binding to receptors in membranes isolated from cells treated with increasing concentrations of MβCD. Values are expressed as percentages of specific binding obtained in control cells without MβCD treatment. Data represent means ± SE from at least six independent measurements performed in duplicate. See Materials and methods for other details.

MβCD treatment), when 10 mM MβCD was used. This shows that removal of cholesterol from intact HN2-5-HT_{1A}R cells leads to enhancement of specific agonist binding to serotonin_{1A} receptors. To the best of our knowledge, these results constitute the first report on change in ligand binding function of serotonin_{1A} receptors upon cholesterol modulation in neuronal cells.

The observed increase in specific ligand binding function upon reduction in membrane cholesterol content could be due to an alteration in overall membrane organization. In order to monitor the change in overall membrane order, we monitored the steady state fluorescence anisotropy of the membrane probe, diphenyl-hexatriene (DPH). DPH is a rod-like molecule and partitions into the bilayer, irrespective of the phase (ordered/disordered) of the membrane [37]. Fluorescence anisotropy is correlated to the rotational diffusion of membrane embedded probes such as DPH [34], which is sensitive to the packing of lipid acyl chains. This is due to the fact that fluorescence anisotropy depends on the degree to which the probe is able to reorient after excitation, and probe




Fig. 4. Steady state fluorescence anisotropy of the membrane probe DPH in membranes isolated from cholesterol-depleted HN2-5-HT_{1A}R cells. Fluorescence anisotropy measurements were carried out with membranes containing 50 nmol phospholipid at a probe to phospholipid ratio of 1:100 (mol/mol) at room temperature (~23 °C). The excitation wavelength was 358 nm and emission was monitored at 430 nm. Data represent means ± SE from at least four independent measurements. See Materials and methods for other details.

reorientation is a function of local lipid packing. Fig. 4 shows the steady state fluorescence anisotropy of DPH in membranes isolated from control and cholesterol-depleted HN2-5-HT_{1A}R cells. Interestingly, the fluorescence anisotropy does not exhibit any appreciable change in membranes isolated from cholesterol-depleted cells. This implies that the overall membrane order is not significantly ($p > 0.05$) altered upon cholesterol depletion.

A useful approach for performing pharmacological studies on GPCRs is to use a functional receptor system that converts receptor–ligand interaction into a cellular signal which allows to monitor the relationship between concentration and response [38]. In the past few years, there have been an increasing number of genetically engineered recombinant receptor systems for monitoring drug–receptor interactions. This has led to a corresponding increase in the testing of new drugs in recombinant receptor systems. However, differences in host membrane lipid composition often complicate interpretation of drug testing results in such systems, and can lead to receptors with characteristics different from native receptors. For example, it has earlier been reported that although the rat cortical serotonin_{1A} receptor exists only in the high affinity state in its native environment, it displays both high and low affinity when expressed in HEK293 cells [39]. It is therefore judicious to work with cellular systems in which the membrane lipid composition closely mimics the native lipid environment. Our choice of HN2 cells for exploring the interaction between membrane cholesterol and serotonin_{1A} receptors is based on the observation that cell lines of neural origin represent realistic models for understanding signal transduction in neuronal cells.

Cholesterol metabolism and homeostasis in the central nervous system and their role in neuronal function is only beginning to be addressed [40]. Previous work from our laboratory has shown that depletion of cholesterol from isolated hippocampal membranes using MβCD leads to the reduction in ligand binding of the serotonin_{1A} receptor [41]. In this context, our present observation of increase in agonist binding of serotonin_{1A} receptors in membranes isolated from cholesterol-depleted HN2-5-HT_{1A}R cells appears interesting. Our present results imply that specific ligand binding to serotonin_{1A} receptors upon cholesterol depletion is influenced by whether cholesterol depletion is carried out in isolated membranes or from intact cells. These results assume additional relevance since they represent the first report on change in ligand binding function of serotonin_{1A} receptors upon cholesterol depletion in neuronal cells. This is important in the context of the recent

observation that cholesterol depletion from neuronal and non-neuronal cells differentially modulates the function of the G-protein coupled δ -opioid receptor [42]. Overall, our results are relevant in understanding the interaction of membrane cholesterol with GPCRs, particularly in view of the recently described structural evidence of specific cholesterol binding site(s) in GPCRs [43].

Acknowledgments

This work was supported by the Council of Scientific and Industrial Research, Government of India. R.P. and Y.D.P. thank the Council of Scientific and Industrial Research for the award of Research Fellowships. We thank Dr. Thomas Pucadyil for preliminary experiments. A.C. is an Adjunct Professor at the Special Centre for Molecular Medicine of Jawaharlal Nehru University (New Delhi, India), and Honorary Professor at the Jawaharlal Nehru Centre for Advanced Scientific Research (Bangalore, India). A.C. gratefully acknowledges support from J.C. Bose Fellowship (Department of Science and Technology, Govt. of India). We thank members of our laboratory for critically reading the manuscript.

References

- [1] D.M. Rosenbaum, S.G.F. Rasmussen, B.K. Kobilka, The structure and function of G-protein-coupled receptors, *Nature* 459 (2009) 356–363.
- [2] P.A. Insel, C.M. Tang, I. Hahntow, M.C. Michel, Impact of GPCRs in clinical medicine: monogenic diseases, genetic variants and drug targets, *Biochim. Biophys. Acta* 1768 (2007) 994–1005.
- [3] R. Heilker, M. Wolff, C.S. Tautermann, M. Bieler, G-protein-coupled receptor-focused drug discovery using a target class platform approach, *Drug Discov. Today* 14 (2009) 231–240.
- [4] T.J. Pucadyil, S. Kalipatnapu, A. Chattopadhyay, The serotonin_{1A} receptor: a representative member of the serotonin receptor family, *Cell. Mol. Neurobiol.* 25 (2005) 553–580.
- [5] S. Kalipatnapu, A. Chattopadhyay, Membrane organization and function of the serotonin_{1A} receptor, *Cell. Mol. Neurobiol.* 27 (2007) 1097–1116.
- [6] C.P. Müller, R.J. Carey, J.P. Huston, M.A. De Souza Silva, Serotonin and psychostimulant addiction: focus on 5-HT_{1A}-receptors, *Prog. Neurobiol.* 81 (2007) 133–178.
- [7] J.A. Gingrich, R. Hen, Dissecting the role of the serotonin system in neuropsychiatric disorders using knockout mice, *Psychopharmacology* 155 (2001) 1–10.
- [8] M. Toth, 5-HT_{1A} receptor knockout mouse as a genetic model of anxiety, *Eur. J. Pharmacol.* 463 (2003) 177–184.
- [9] K. Simons, E. Ikonen, How cells handle cholesterol, *Science* 290 (2000) 1721–1726.
- [10] O.G. Mouritsen, M.J. Zuckermann, What's so special about cholesterol? *Lipids* 39 (2004) 1101–1113.
- [11] S. Mukherjee, F.R. Maxfield, Membrane domains, *Annu. Rev. Cell Dev. Biol.* 20 (2004) 839–866.
- [12] K. Jacobson, O.G. Mouritsen, R.G.W. Anderson, Lipid rafts: at a crossroad between cell biology and physics, *Nat. Cell Biol.* 9 (2007) 7–14.
- [13] S. Munro, Lipid rafts: elusive or illusive? *Cell* 115 (2003) 377–388.
- [14] A.S. Shaw, Lipid rafts: now you see them, now you don't, *Nat. Immunol.* 7 (2006) 1139–1142.
- [15] K. Simons, G. Van Meer, Lipid sorting in epithelial cells, *Biochemistry* 27 (1988) 6197–6202.
- [16] K. Simons, D. Toomre, Lipid rafts and signal transduction, *Nat. Rev. Mol. Cell Biol.* 1 (2000) 31–39.
- [17] P. Goluszko, B. Nowicki, Membrane cholesterol: a crucial molecule affecting interactions of microbial pathogens with mammalian cells, *Infect. Immun.* 73 (2005) 7791–7796.
- [18] J. Riethmüller, A. Riehle, H. Grassmé, E. Gulbins, Membrane rafts in host-pathogen interactions, *Biochim. Biophys. Acta* 1758 (2006) 2139–2147.
- [19] T.J. Pucadyil, A. Chattopadhyay, Cholesterol: a potential therapeutic target in Leishmania infection? *Trends Parasitol.* 23 (2007) 49–53.
- [20] T.J. Pucadyil, A. Chattopadhyay, Role of cholesterol in the function and organization of G-protein coupled receptors, *Prog. Lipid Res.* 45 (2006) 295–333.
- [21] C.G. Tate, R. Grishammer, Heterologous expression of G-protein-coupled receptors, *Trends Biotechnol.* 14 (1996) 426–430.
- [22] P.S. Sastry, Lipids of nervous tissue: composition and metabolism, *Prog. Lipid Res.* 24 (1985) 69–176.
- [23] A. Chattopadhyay, Y.D. Paila, Lipid-protein interactions, regulation and dysfunction of brain cholesterol, *Biochem. Biophys. Res. Commun.* 354 (2007) 627–633.
- [24] M. Mulder, Sterols in the central nervous system, *Curr. Opin. Clin. Nutr. Metab. Care* 12 (2009) 152–158.
- [25] Y.D. Paila, A. Chattopadhyay, The human serotonin_{1A} receptor expressed in neuronal cells: toward a native environment for neuronal receptors, *Cell. Mol. Neurobiol.* 26 (2006) 925–942.
- [26] A. Fargin, J.R. Raymond, M.J. Lohse, B.K. Kobilka, R.J. Lefkowitz, The genomic clone G-21 which resembles a β -adrenergic receptor sequence encodes the 5-HT_{1A} receptor, *Nature* 335 (1988) 358–360.
- [27] H.J. Lee, D.N. Hammond, T.H. Large, J.D. Roback, J.A. Sim, D.A. Brown, U.H. Otten, B.H. Wainer, Neuronal properties and trophic activities of immortalized hippocampal cells from embryonic and young adult mice, *J. Neurosci.* 10 (1990) 1779–1787.
- [28] Md. Jafurulla, A. Chattopadhyay, Solubilization of human serotonin_{1A} receptors expressed in neuronal cells, *Chem. Phys. Lipids* 150 (2007) 244–249.
- [29] P.K. Smith, R.I. Krohn, G.T. Hermanson, A.K. Mallia, F.H. Gartner, M.D. Provenzano, E.K. Fujimoto, N.M. Goeke, B.J. Olson, D.C. Klenk, Measurement of protein using bicinchoninic acid, *Anal. Biochem.* 150 (1985) 76–85.
- [30] S. Kalipatnapu, T.J. Pucadyil, K.G. Harikumar, A. Chattopadhyay, Ligand binding characteristics of the human serotonin_{1A} receptor heterologously expressed in CHO cells, *Biosci. Rep.* 24 (2004) 101–115.
- [31] D.M. Amundson, M. Zhou, Fluorometric method for the enzymatic determination of cholesterol, *J. Biochem. Biophys. Methods* 38 (1999) 43–52.
- [32] C.W.F. McClare, An accurate and convenient organic phosphorus assay, *Anal. Biochem.* 39 (1971) 527–530.
- [33] Y.D. Paila, M.R.V.S. Murty, M. Vairamani, A. Chattopadhyay, Signaling by the human serotonin_{1A} receptor is impaired in cellular model of Smith-Lemli-Optiz syndrome, *Biochim. Biophys. Acta* 1778 (2008) 1508–1516.
- [34] J.R. Lakowicz, Principles of Fluorescence Spectroscopy, third ed., Springer, New York, 2006.
- [35] E.C. Hulme, Receptor binding studies: a brief outline, in: E.C. Hulme (Ed.), Receptor-Effector Coupling: A Practical Approach, IRL Press, New York, 1990, pp. 203–215.
- [36] R. Zidovetzki, I. Levitan, Use of cyclodextrins to manipulate plasma membrane cholesterol content: evidence, misconceptions and control strategies, *Biochim. Biophys. Acta* 1768 (2007) 1311–1324.
- [37] E. London, G.W. Feigenson, Fluorescence quenching in model membranes. An analysis of the local phospholipid environments of diphenylhexatriene and gramicidin A', *Biochim. Biophys. Acta* 649 (1981) 89–97.
- [38] T. Kenakin, Differences between natural and recombinant G protein-coupled receptor systems with varying receptor/G protein stoichiometry, *Trends Pharmacol. Sci.* 18 (1997) 456–464.
- [39] J. Watson, L. Collin, M. Ho, G. Riley, C. Scott, J.V. Selkirk, G.W. Price, 5-HT_{1A} receptor agonist-antagonist binding affinity difference as a measure of intrinsic activity in recombinant and native tissue systems, *Br. J. Pharmacol.* 130 (2000) 1108–1114.
- [40] F.W. Pfrieger, Cholesterol homeostasis and function in neurons of the central nervous system, *Cell. Mol. Life Sci.* 60 (2003) 1158–1171.
- [41] T.J. Pucadyil, A. Chattopadhyay, Cholesterol modulates the ligand binding and G-protein coupling to serotonin_{1A} receptors from bovine hippocampus, *Biochim. Biophys. Acta* 1663 (2004) 188–200.
- [42] P. Huang, W. Xu, S.-I. Yoon, C. Chen, P.L.-G. Chong, L.-Y. Liu-Chen, Cholesterol reduction by methyl- β -cyclodextrin attenuates the delta opioid receptor-mediated signaling in neuronal cells but enhances it in non-neuronal cells, *Biochem. Pharmacol.* 73 (2007) 534–549.
- [43] Y.D. Paila, S. Tiwari, A. Chattopadhyay, Are specific nonannular cholesterol binding sites present in G-protein coupled receptors? *Biochim. Biophys. Acta* 1788 (2009) 295–302.