

Abundance and relationship of bacteria with transparent exopolymer particles during the 1996 summer monsoon in the Arabian Sea

N RAMAIAH, V V S S SARMA, MANGESH GAUNS, M DILEEP KUMAR and M MADHUPRATAP

National Institute of Oceanography, Dona Paula, Goa, India.

Bacterial abundance and production, numbers, sizes and concentrations of transparent exopolymer particles (TEP) and total organic carbon (TOC) were measured during the 1996 summer monsoon to understand the relationship between TEP, the most labile particulate organic carbon, and bacteria. While high regional variability in the vertical distribution of TOC was discernible, TEP concentrations were high in surface waters at 18–20°N along 64°E with concentrations well over 25 mg alginic acid equivalents l⁻¹ due to upwelling induced productivity. Their concentrations decreased with depth and were lower between 200 and 500 m. Bacterial concentrations were up to 1.99 × 10⁸ l⁻¹ in the surface waters and decreased by an order of magnitude or more at depths below 500 m. A better relationship has been found between bacterial abundance and concentrations of TEP than between bacteria and TOC, indicating that bacterial metabolism is fueled by availability of TEP in the Arabian Sea. Assuming a carbon assimilation of 33%, bacterial carbon demand (BCD) is estimated to be 1.017 to 4.035 g C m⁻² d⁻¹ in the surface waters. The observed TEP concentrations appear to be sufficient in meeting the surface and subsurface BCD in the northern Arabian Sea.

1. Introduction

The predominant physical factors influencing the northern Arabian Sea biogeochemistry are strong, semi-annual reversals in monsoonal winds. Such reversing winds bring about very large seasonal changes in primary production (Qasim 1982; Banse 1994; Madhupratap *et al* 1996a), with consequences for the biological pump. The latest findings from the JGOFS disprove many previous long-held notions, and suggest high production during the northeast monsoon (Smith *et al* 1998; Burkill *et al* 1993; Prasanna Kumar *et al* 2000). Aggregation and sinking flux of organic matter following phytoplankton blooms are common in marine ecosystems (Takahashi 1986; Bodungen *et al* 1986), including the Arabian Sea where seasonal changes in sedimentation rates are high (Nair *et al* 1989; Haake *et al* 1993). The Arabian Sea also has an extensive denitrification zone at mid depths (Naqvi 1994; Naqvi *et al* 1998). Bacterioplankton often constitutes a bulk of carbon biomass in pelagic ecosystems

(Biddanda and Benner 1997; Wiebinga *et al* 1997), and utilizes a large fraction of the sinking particulate organic matter, converting it to a non-sinking carbon pool (Smith *et al* 1995). Thus, interaction between sinking organics (fecal pellets and /or post-bloom aggregates) and heterotrophic bacteria is important in supporting life below the euphotic zone.

Under the Joint Global Ocean Flux Study (JGOFS) programmes, there have been a number of recent studies on the abundance (and production) of bacteria in the Arabian Sea (Burkill *et al* 1993; Ducklow 1993; Wiebinga 1994; Ramaiah *et al* 1996; Goosen *et al* 1997; Veldhuis *et al* 1997; Wiebinga *et al* 1997; Campbell *et al* 1998). Seasonal variations are clearly seen with higher abundance and production rates during spring and fall inter-monsoons (Ramaiah *et al* 1996; Campbell *et al* 1998; Pomroy and Joint 1999). However, the relationship of bacteria with available organic carbon *in situ* is not well understood. The presence of free, large, discrete polymer particles, known as transparent exopolymer particles (TEP) which are easily

Keywords. Bacteria; TEP; primary production; organic carbon; zooplankton; Arabian Sea.

assimilable by bacteria (Passow and Alldredge 1995, Smith *et al* 1995) have been demonstrated to be of great significance in the biogeochemistry of the marine regimes (Alldredge *et al* 1993; Kumar *et al* 1998). In the light of this recent discovery, we quantified TEP during the summer monsoon of 1996 for the first time from the Arabian Sea with an aim to understand its relationship with bacteria.

2. Material and Methods

This study was carried out during the cruise 115 of ORV *Sagar Kanya* during the summer (southwest) monsoon (August 4th–30th, 1996) as a part of the JGOFS India Programme. Stations were occupied along 64°E from 13°N to 21°N (figure 1). Water samples were collected from the upper 1000 m using CTD rosette and analysed for bacterial abundance and production and for estimating numbers, sizes and concentrations of TEP. Bacterial production was measured only in the top 150 m.

Immediately after collection aliquots (50ml) of seawater were fixed with 0.22 μ m pre-filtered formaldehyde (to a final concentration of 3.7%) for counting total bacterial cells. These samples were stored in the dark at 4°C as per the JGOFS Protocols (UNESCO 1994) until taken for counting within a week of returning from the cruise. Acridine orange direct counts (AODC) of bacteria were carried out following Parsons *et al* (1984) by using 100X oil immersion objective on an Olympus BH2 epifluorescence microscope. Aliquots of two to six ml (with higher volumes from deeper samples) were stained with acridine orange (0.001% final concentration) for 5 minutes, filtered (0.22 μ m black Nuclepore filters) and prepared for microscopy. Bacterial cells in ca. 25 microscopic fields were counted and the mean cell numbers per

field calculated and used for estimating total abundance by using the relationship given by Parsons *et al* (1984). These numbers were then used to calculate bacterial carbon biomass using the conversion of 20 fg C cell⁻¹ (Ducklow 1993).

Incorporation of tritiated (³H)- methyl thymidine (specific activity 17000 mCi/mmol, Bhabha Atomic Research Centre, Mumbai) by the water column bacteria was estimated following the method described in JGOFS Protocols (UNESCO 1994). Samples from surface to 150 m (usually from 1, 20, 40, 60, 80, 100, 120 and 150 m) were subjected for ³H incorporation. An aliquot of 100 μ l working solution (59 nmole) was added to 20 ml triplicate samples from each depth. These samples in 50 ml polycarbonate centrifuge tubes (Tarsons, Calcutta) were incubated in the dark at 22–24°C (similar to the *in situ* temperatures in the mixed layer) for 60 min. The uptake was terminated by adding 400 μ l formaldehyde. Samples were filtered through 0.22 μ m cellulose acetate filters (Millipore India Ltd, Bangalore) and the thymidine incorporated into macromolecular fractions was extracted/precipitated by three repeated, alternate rinses with cold trichloroacetic acid (5 ml) and cold ethanol (3 ml for each rinse). The filters were stored in minivials, refrigerated and brought to the laboratory. After ensuring that they were moisture free, 4 ml liquid scintillation fluid (cocktail-W, Spectrochem, Mumbai) was added and the samples were radio-assayed in a Packard TR 2500 liquid scintillation counter.

The ³H thymidine incorporated (TTI) was calculated by the formula:

$$TTI(\text{pMh}^{-1} \text{l}^{-1}) = (\text{DPM}_s - \text{DPM}_b) / 2200 * V / 1000 * 60 / T * 1 / \text{SA}$$

where, DPM – disintegration per minute; *s* – sample; *b* – blank; *V* – sample volume (20 ml); *T* – incubation time in minutes; SA – specific activity of added ³H thymidine.

A conversion factor of 2.17×10^{18} cells mole⁻¹ thymidine incorporated was used to calculate bacterial production (Ducklow 1993). To estimate the bacterial carbon demand, we assumed a 33% efficiency of carbon assimilation by bacteria (i.e., one unit of bacterial carbon formed for every three units of organic carbon consumed).

Chemical quantification, microscopic counting and sizing of TEP were made from two sets of samples taken from 15 depths in the top 1000 m. The first set of samples was transferred to clean glass bottles and used for chemical quantification within 30 min of collection. From the other set, 50 ml samples were fixed by adding 1.0 ml 37% formaldehyde and refrigerated until analyzed by microscopy.

Following the method of Passow and Alldredge (1995), the samples were filtered through 0.45 μ m polycarbonate or Durapore filters within a couple of hours of collection and stained with alcian blue (0.1%




Figure 1. Sampling locations along 64°E.

wt/vol). The TEP were then dissolved in 80% sulphuric acid and the absorbance of the solution was read at 785 and/or 747 nm (see Kumar *et al* 1998). Concentrations of TEP were expressed as milligram equivalent of alginic acid (AA) l^{-1} . Although for most samples, the TEP concentrations were calculated from single measurements, we did analyze a few samples from discrete depths in triplicates or quadruplicates and the mean coefficient of variation was within 11%.

For microscopic counting and size measurements of TEP, aliquots of 10 ml samples were filtered through 0.22 μm (25 mm diameter) polycarbonate (Nuclepore, USA) or Durapore (Millipore India Ltd, Bangalore) filters. 100 μl (0.1% wt/vol) of alcian blue (8 GX Sigma, USA) and 20 μl of 0.006% glacial acetic acid were added just before the last millilitre sample was vacuum drained. Filters were transferred to microscopic slides with immersion oil and readied for observation as per Parsons *et al* (1984). TEP were counted from at least 30 microscopic fields at a magnification

of 200X. Some samples were observed under oil immersion at 1250X to note the association of bacterial cells and mineral particles with TEP. The average maximum length of TEP from each sample was calculated to understand the distribution of their sizes.

3. Results

During the summer monsoon of 1996, the mixed layer (ML) along 64°E was around 100–50 m with a shoaling towards the north. Nitrate concentrations were high ($\geq 1.0 \mu mole l^{-1}$) in the ML Primary production in the upper 120 m column was in the range of 347–1782 $mg C m^{-2} d^{-1}$. Phytoplankton cell numbers were invariably high in the range of 10^4 – 10^5 cells l^{-1} . Bloom proportions of *Phaeocystis* were found in the north of 17°N, (Madhupratap *et al* 2000).

Bacteria were abundant in the top 50 m ranging from 0.11 to $1.99 \times 10^8 l^{-1}$ (table 1). Their maximum

Table 1. Bacterial abundance and production in surface waters at various locations during the summer monsoon of 1996.

Station	Depth (m)	TDC $\times 10^8 l^{-1}$	BC $g m^{-3}$	TTI $pMI^{-1} h^{-1}$	BCP $gm^{-3} d^{-1}$	BCD $gm^{-3} d^{-1}$
13°N 64°E	0	1.453	0.029	0.491	0.0051	0.0168
	20	1.371	0.027	0.992	0.0103	0.0339
	40	1.996	0.039	1.850	0.0192	0.0633
	60	0.249	0.0049	0.362	0.0037	0.0122
	80	1.292	0.0258	0.880	0.0092	0.0303
	100	0.411	0.0082	0.678	0.0070	0.0231
15°N 64°E	0	0.822	0.0164	1.182	0.0123	0.0406
	20	1.174	0.0235	2.303	0.0239	0.0788
	40	0.787	0.0157	1.477	0.0153	0.0505
	60	0.811	0.0162	0.455	0.0047	0.0155
	80	0.352	0.0070	1.281	0.0133	0.0439
	90	0.423	0.0085	0.198	0.0020	0.0066
	120	0.141	0.0028	0.642	0.0066	0.0218
17°N 64°E	150	0.106	0.0021	NA		
	0	0.235	0.0047	1.968	0.0204	0.0673
	25	0.188	0.0037	1.352	0.0140	0.0462
	40	0.305	0.0061	1.194	0.0124	0.0409
	50	0.059	0.0012	0.679	0.0070	0.0231
	75	0.411	0.0082	0.156	0.0016	0.0053
	100	0.176	0.0035	0.229	0.0023	0.0076
	120	0.296	0.0059	0.180	0.0018	0.0059
19°N 64°E	150	0.341	0.0068	0.460	0.0047	0.0155
	0	0.904	0.0180	0.651	0.0068	0.0223
	25	0.775	0.0155	0.428	0.0045	0.0146
	40	1.197	0.0239	0.675	0.0071	0.0232
	60	1.057	0.0211	0.349	0.0036	0.0119
	75	0.951	0.0190	0.158	0.0017	0.0054
	100	0.341	0.0068	0.021	0.0022	0.0072
21°N 64°E	120	0.294	0.0058	0.183	0.0019	0.0062
	150	0.199	0.0039	NA		
	0	0.129	0.0025	0.705	0.0073	0.0242
	25	0.106	0.0021	0.529	0.0055	0.0181
	50	0.106	0.0021	0.127	0.0013	0.0043
	75	1.291	0.0250	0.144	0.0015	0.0049
	100	1.045	0.0209	0.112	0.0012	0.0038
21°N 64°E	120	1.221	0.0244	0.062	0.0007	0.0021
	150	1.081	0.0216	0.223	0.0023	0.0765

TDC: total direct (acridine orange) counts; BC: bacterial carbon biomass; TTI: thymidine incorporation rate; BCP: bacterial carbon production rate; BCD: bacterial carbon demand and, NA: no data.




Figure 2. Vertical profiles of transparent exopolymer particles (TEP), bacterial counts and δN along $64^\circ E$. Averages of each of these parameters are shown by solid lines.

concentration at the surface was 1.45×10^8 cells l^{-1} and, as can be discerned from figure 2, decreased to ca. 2.0×10^7 l^{-1} around 400 m and below. On an average, their numbers were 0.64×10^7 l^{-1} in waters below the ML (150–500 m) that further decreased to $< 0.11 \times 10^7$ l^{-1} between 500 and 1000 m. Their abundance was higher in the north of $17^\circ N$. A secondary peak, with counts up to 0.91×10^7 l^{-1} , persisted ca. between 200 and 400 m at most sampling stations along $64^\circ E$. Standing stocks of bacterial carbon (BC) ranged from 2.5 to 39 $mg\ m^{-3}$ in the surface and from 0.4 to 0.82 $mg\ m^{-3}$ at 1000 m. Bacterial carbon production (BCP), with thymidine incorporation rates ranging from 0.02 to 2.30 $pM\ l^{-1}\ h^{-1}$ amounted to 2.2–23.98 $mg\ C\ m^{-3}\ d^{-1}$. Assuming a 33% efficiency of carbon assimilation by bacteria, the calculated BCD was in the range of 7.2 to 78.8 $mg\ C\ m^{-3}\ d^{-1}$ in the ML. As can be derived from the data in table 1, the 0–120 m column integrated values of BC, BCP and BCD from different stations ranged respectively from 0.57 to 2.40 $g\ m^{-2}$; 0.34 to 1.35 $g\ C\ m^{-2}\ d^{-1}$ and 1.02 to 4.04 $g\ C\ m^{-2}\ d^{-1}$.

Vertical profiles of TEP at different sampling locations are depicted in figure 2. TEP concentrations decreased with increasing depth from an average of about 60 $mg\ AA\ eq\ l^{-1}$ in the surface to less than 10 $mg\ AA\ eq\ l^{-1}$ at 1000 m. On the whole, TEP varied from < 5 mg to as high as 102 $mg\ AA\ eq\ l^{-1}$ in the water column (Kumar *et al* 1998). Additionally, their concentrations in surface waters (1 – 50 m) were more (≥ 25 $mg\ AA\ eq\ l^{-1}$) in the north, particularly, from 18 to $20^\circ N$ and, very high concentrations were seen around 600 m at these stations. Vertical profiles of

bacterial counts and nitrate deficit (δN)—a measure of nitrate decrease due to its bacterial reduction to molecular nitrogen—are also included in figure 2 for comparison. It is seen that δN is around 0–7.5 μM between 150 and 200 m where the bacterial numbers on an average were high, but the TEP were low.

Latitudinal distribution of mean lengths and counts of TEP along $64^\circ E$ is shown in figures 3 and 4 respectively. Their sizes were bigger in the north of $17^\circ N$. The numbers of discrete TEP were quite low with an overall range of 130 to 4650 ml^{-1} , owing generally to larger sizes (all samples mean length $47.33 \pm 28.19\ \mu m$, $n = 38$).




Figure 3. Distribution of TEP sizes (μm) along $64^\circ E$ during the 1996 summer monsoon.




Figure 4. Distribution of TEP numbers (ml^{-1}) along 64°E during the 1996 summer monsoon.

Regression analyses of bacterial counts versus TEP concentration, size, counts and total organic carbon (figure 5) suggested that bacterial counts are more robustly correlated with TEP concentration ($r = 0.6$; $n = 57$; $P < 0.01$) than with TEP numbers ($r = 0.50$; $n = 60$; $P < 0.01$), with TOC or TEP sizes. Positive linear relationship ($r = 0.49$; $n = 22$; $P < 0.05$) between TEP concentrations and bacterial production (figure 6) suggests that the TEP is in excess and bacterial production in upper 120 m is more or less uniform. Further, maximum rates of bacterial production appear to have been attained when TEP concentrations were equal to or even $10 \text{ mg AA eq l}^{-1}$. This suggests that bacterial abundance and production are supported directly or indirectly by solubilization of TEP.




Figure 5. Scatter diagrams showing relationships of heterotrophic bacterial (H BAC) counts with: TEP size, TEP numbers, TEP concentrations and TOC during the 1996 summer monsoon.




Figure 6. Regression relationship of bacterial production with TEP concentrations in the surface waters (0–150 m) along 64°E during the summer monsoon, 1996.

4. Discussion

The concentration of bacteria during summer monsoon was low compared to those in spring intermonsoon (April – May) period along 64°E (Ramaiah et al 1996). They are in the same range observed during the summer monsoon months (June – September) in other areas of open Arabian Sea (Goosen et al 1997, Wiebinga et al 1997). Bacterial production also agreed with ranges reported by Goosen et al (1997) and Wiebinga et al (1997) during southwest monsoon. This clearly suggests that a large area in the Arabian Sea experiences low abundance and production of bacteria during the summer monsoon. This fact is augmented by the occurrence of lower ratios of




Figure 7. Living component carbon biomass in the sampling locations along 64°E and a transect along the west coast of India within 200m depth contour. Carbon biomass is in mM C m⁻². PP denotes primary production; MZP, microzooplankton; ZP, mesozooplankton; BC, bacterial carbon standing stock; BP, bacterial production; SpIM, spring intermonsoon; SuM, summer monsoon and, Win denotes winter.

bacterial/phytoplankton production (Wiebinga *et al* 1997; Prasanna Kumar *et al* 2000).

The sinking carbon flux from the surface is a result of either biologically mediated aggregation of photosynthetically formed organic matter or abiotically mediated coagulation of small, discrete particles into larger aggregates. The TEP are formed abiotically from dissolved extracellular polysaccharides (Passow *et al* 1994). Diatoms (Williams 1990) and some phytoplankton species (Lancelot and Methot 1985) produce TEP in abundance comparable to phytoplankton carbon (Passow *et al* 1994). For example, *Phaeocystis* spp are known to copiously exude over 50% of their photosynthate (Lancelot and Methot 1985) which might lead to formation of large concentrations of TEP. It is interesting to note that large sized particles (and concentrations) occurred around 600m mostly to the north of 15°N, in the Arabian Sea. This probably implies firstly, a rapid sinking of TEP immediately after formation from bloom proportions of *Phaeocystis* sp. The *Phaeocystis* bloom was in senescent stage (Madhupratap *et al* 2000). Secondly, it is possible that rapid sinking and lower temperatures (facilitating condensation of polysaccharides) below 400 m may have lead to their post-production aggregation leading to larger sized TEP. This is evident from their low numbers but larger sizes (> 50 μm) at 400–600 m. Their smaller sizes (20–30 μm) in the intermediate layers (200–400 m) may be suggestive of the microbial colonization and sheering by ectoenzyme hydrolysis as suggested by Martinez and Azam (1993). Despite a low level of statistical significance, the positive correlation ($r = 0.136$) between bacterial counts and TEP numbers observed during this study may be suggestive of the sheering of TEP in this depth zone in particular. Heterotrophic bacteria are efficient in direct uptake of low-molecular weight dissolved organic matter. Upon attaching, adsorbing or 'packaging' to TEP, bacteria elaborate ectoenzymes, dissolve the easily assimilable TEP and incorporate this into their cells. Recent studies suggest such a role for water column bacteria (Chrost 1990; Azam *et al* 1993; Smith *et al* 1995). The stronger relationship between the water column bacterial abundance (as also their production) and TEP concentrations in the Arabian Sea corroborates with the earlier suggestions.

Microbial denitrification is a biochemically complex, high energy requiring process (Brock *et al* 1991). Ducklow (1993) and Naqvi and Shailaja (1993) found that bacterial carbon demand in the denitrification zone is greater than the sinking carbon flux and suggested a large scale, lateral import of organic matter into mesopelagic zone. The estimated carbon demands for the Arabian Sea intermediate water denitrification range from ca. 155 $\text{mg C m}^{-2} \text{d}^{-1}$ (Naqvi *et al* 1996) to 346–518 $\text{mg C m}^{-2} \text{d}^{-1}$ (derived from Banse 1994) to 519–2274 $\text{mg C m}^{-2} \text{d}^{-1}$ (calculated from Ducklow

1993). During summer monsoon, bacterial production in 0–300 m column is reported to range from 148 to 223 $\text{mg C m}^{-2} \text{d}^{-1}$ in the Somali Basin (Wiebinga *et al* 1997). Extrapolating our assumption of 33% efficiency of carbon assimilation for such production rates, the BCD amounts to 444–669 $\text{mg C m}^{-2} \text{d}^{-1}$ in the 0–300 m column. Despite seasonal differences (lower during intermonsoon: 322–404 $\text{mg C m}^{-2} \text{d}^{-1}$ and higher during winter: 662–1150 $\text{mg C m}^{-2} \text{d}^{-1}$ (data from Sarin *et al.* 1996)), sediment trap and ^{234}Th export fluxes do suggest a noticeable carbon export at 100 m. In addition, the total organic carbon (TOC) concentrations in the region are high varying from 960 to 1440 mg m^{-3} (Menzel 1964); 1600 mg m^{-3} in the surface layers to > 3600 mg m^{-3} (Kumar *et al* 1990) and from 580 to > 960 mg m^{-3} (Hansell and Peltzer 1998). It may be added here that although the labile proportions of this TOC are not known, it is speculated variously that only less than 10% of the TOC may ultimately be refractory (Azam *et al* 1994). From these trends in TOC, particle fluxes, from our TEP data and from the significant relationship between TEP and BP in the Arabian Sea, we infer that a greater portion of BCD is met from the TEP. Thus, TEP are important in fuelling the Arabian Sea suboxic, subsurface denitrification process. Further, from the measurements of Ducklow (1993) and Wiebinga *et al* (1997), it is evident that the bacterial production rates are lower in the deeper layers. This is also clear from the substantially decreasing bacterial counts observed during this study. Hence, the BCD would be lower in the deeper layers. The cumulative fractions of TEP and slow-to-degrade dissolved organic carbon in the sinking flux must be sufficient for metabolism of bacterial communities.

Azam *et al* (1994) suggest that bacteria utilize DOC with highly variable carbon assimilation efficiency of 10–70% and their communities in the denitrification zone may differ in metabolic capacities than those from the surface. Our assumption of 33% efficiency for bacterial carbon formation is similar to those given by Smith *et al* (1995) but quite low when compared with those assuming 50% efficiency (Ducklow 1993). Even at this 'moderately low' efficiency it appears that there is enough TEP to fuel bacterial metabolism during denitrification. Further, the generation times of bacteria in the surface (0–120 m) waters along 64°E range from 5.5 to 19 days during different seasons (Ramaiah, unpublished) suggesting faster carbon turnover than those estimated by Wiebinga *et al* (1997) for the northwestern Arabian Sea bacteria.

Results of biological studies during the 1996 southwest monsoon together with those in other seasons under the Arabian Sea JGOFS (India) Programme are summarized in figure 7. An overview of the living component of carbon biomass and its seasonal variations along 64°E and coastal stations (occupied

by JGOFS India Programme) is also summarized in this figure. It is evident that the bacterial community is dominant during spring intermonsoon (see also Ramaiah *et al* 1996; Campbell *et al* 1998). During April – May, when chlorophyll concentrations and primary production are low (Bhattathiri *et al* 1996), bacteria appear to be very important in sustaining a greater abundance of microzooplankton (Gauns *et al* 1996) as well as of mesozooplankton through the microbial loop (Madhupratap *et al* 1996a). The latter's biomass in the mixed layer is considered to be invariant (Madhupratap *et al* 1996b). Bacterial production through organic carbon assimilation appears to be of greater significance for microzooplankton both during summer monsoon and spring intermonsoon (Gauns *et al* 1996). Feeding on bacteria which are nutritionally 'more complete', would be a more efficient means of "better nutrition" for microzooplankton. In addition, the heterotrophic bacterial community is the key component breaking down organic substances and making available most inorganic nutrients essential for autotrophic production in euphotic zone. The strong relationship between bacterial abundance and concentrations of TEP indicate that TEP in the Arabian Sea play an important role in sustaining these heterotrophic communities which appear to be of greater significance during spring intermonsoon.

In conclusion, TEP concentrations measured during the summer monsoon appear to be sufficient for fuelling bacterial metabolism in the suboxic denitrification layers of the Arabian Sea. However, we do not yet know the actual carbon content of TEP. Recently, Alldredge (1998) hypothesized that up to 14 – 37% of the "miscellaneous aggregates" (we presume that TEP are a major fraction of this class) is carbon. Assuming that 100 mg alginic acid equivalent of TEP will at least have 20 mg carbon, their carbon equivalent between 200 and 1000 m is about 300 mg m⁻³. Thus, results from this study strongly suggest that TEP concentrations may meet bacterial carbon demand for denitrification in the Arabian Sea. Our measurements on TEP are the first ones from the Arabian Sea and future studies would prove helpful in understanding the annual cycles of their concentrations and sizes, as well as their role in carbon cycling in the Arabian Sea.

Acknowledgements

We thank Dr. E Desa, Director NIO for facilities and encouragement. We also thank Drs. S Raghukumar and N B Bhosle at the NIO for helpful suggestions. Financial support by the Department of Ocean Development, New Delhi for JGOFS (India) Programme under which this study was carried out is gratefully acknowledged. This is NIO contribution no. 3584.

References

- Allredge A L, Passow U and Logan B E 1993 The abundance and significance of a class of large, transparent organic particles in the ocean; *Deep-Sea Res.* **40** 1131–1140
- Allredge A L 1998 The carbon, nitrogen and mass content of marine snow as a function of aggregate size; *Deep-Sea Res.* **45** 529–541
- Azam F, Martinez J and Smith D C 1993 Bacteria-organic matter coupling on marine aggregates; In: *Trends in microbial ecology* (ed) R Guerrero and C Pedros-Alio (Barcelona: Spanish Society for Microbiology) pp 410–414
- Azam F, Steward G F, Smith D C and Ducklow H W 1994 Significance of bacterial in carbon fluxes in the Arabian Sea; *Proc. Indian Acad. Sci., (Earth Planet Sci.)* **103** 341–351
- Banse K 1994 On the coupling of hydrography, phytoplankton, zooplankton offshore in the Arabian Sea and organic particles settling to depth. In: *Biogeochemistry of the Arabian Sea*; Lal D (ed), (Bangalore: Indian Academy of Sciences) pp 125–161
- Bhattathiri P.M.A, Aditi Pant, Sawant S, Gauns M Matonkar S G P and Mohanraju R 1996 Phytoplankton production and chlorophyll distribution in the eastern and central Arabian Sea in 1994–95; *Curr. Sci.* **71** 857–862
- Biddanda B and Benner R 1997 Major contribution from mesopelagic plankton to heterotrophic metabolism in the upper ocean; *Deep-Sea Res.* **14** 2069–2085
- Bodungen B van, Smetacek V, Tilzer MM and Zeitzschel B 1986. Primary production and sedimentation during spring in the Antarctic Peninsula region; *Deep-Sea Res.* **33** 177–194
- Brock J C, McClain M C, Luther M E and Hay W W 1991 The phytoplankton bloom in the northwestern Arabian Sea during the southwest monsoon of 1979; *J. Geophys. Res.* **96** 20623–20642
- Burkill P H, Mantoura R F C and Owens N J P 1993 Biogeochemical cycling in the northwestern Indian Ocean: A brief overview. *Deep-Sea Res.*, special issue on *Biogeochemical cycling in the northwestern Indian Ocean* (ed) P H Burkill, R F C Mantoura, N J P Owens, **40** 643–649
- Campbell L, Landry M R, Constantinour J, Nolla, H A, Liu H, Brozn S L, Caron D A 1998 Response of microbial community structure to environmental forcing in the Arabian Sea; *Deep-Sea Res.* **45** 2301–2325
- Chrost R J 1990 Microbial ectoenzymes in aquatic environments; In: *Aquatic microbial ecology: Biochemical and molecular approaches* (ed) J Overbeck and R J Chrost (New York: Brock/Springer) pp. 47–78
- Ducklow H W 1993. Bacterioplankton distributions and production in the northwestern Indian Ocean and Gulf of Oman, September 1986; *Deep-Sea Res.* **40** 753–771
- Gauns M, Mohanraju R and Madhupratap M 1996 Studies on the microzooplankton from the central and eastern Arabian Sea; *Curr. Sci.* **71** 874–877
- Goosen N K, Van-Rijswijk P, Kromkamp J, Peena J 1997 Regulation of annual variation in heterotrophic bacterial production in the Schelde Estuary (SW Netherlands); *Aquat. Micro. Ecol.* **12** 223–232
- Haake B, Ittekkot V, Rixen T, Ramaswamy V, Nair R R, Curry W B 1993 Seasonality and interannual variability of particle fluxes to the deep of the Arabian Sea; *Deep-Sea Res.* **40** 1323–1344
- Hansell D A and Peltzer E T 1998 Spatial and temporal variations of total organic carbon in the Arabian Sea; *Deep-Sea Res.* **45** 2171–2194
- Kumar M D, Rajendran A, Somasundar K, Haake B, Jenisch A, Shuo Z, Ittekkot V and Desai B N 1990 Dynamics of dissolved organic carbon in the northwestern Indian Ocean; *Mar. Chem.* **30** 299–316

- Kumar M D, Sarma V V S S, Ramaiah N and Gauns M 1998 Biogeochemical significance of transparent exopolymer particles in the Indian Ocean; *Geophys. Res. Lett.* **25** 81–84
- Lancelot C and Methot S 1985 Biochemical fractionation of primary production by phytoplankton in Belgian coastal waters during short and longterm incubations with ^{14}C -bicarbonate II. *Phaeocystis pouchetti* colonial population; *Mar. Biol.* **86** 227–232
- Madhupratap M, Prasanna Kumar S, Bhattathiri P M A, Kumar M D, Raghukumar S, Nair K K C, Ramaiah N 1996a Mechanism of the biological response to winter cooling in the northeastern Arabian Sea; *Nature* **384** 452–459
- Madhupratap M, Gopalakrishnan T C, Haridas P, Nair K K C, Aravindakshan P N, Padmavati G and Shinney Paul 1996b Lack of seasonal and geographic variation in mesozooplankton biomass in the Arabian Sea and its structure in the mixed layer; *Curr. Sci.* **71** 863–868
- Madhupratap M, Sawant S and Gauns M 2000 A first report on a bloom of the marine prymnesiophycaen *Phaeocystis globosa* from the Arabian Sea; *Oceanologica Acta* 2383–2390
- Martinez J and Azam F 1993 Periplasmic aminopeptidase and alkaline phosphatase activities in a marine bacterium: Implications for substrate processing in the sea; *Mar. Ecol. Prog. Ser.* **92** 89–97
- Menzel D W 1964 The distribution of dissolved organic carbon in the western Indian Ocean; *Deep-Sea Res.* **11** 757–766
- Nair R R, Ittekkot V, Managanin S J, Ramaswamy V, Haake B, Degens E T, Desai B N and Honjo S 1989 Increased particle flux to the deep ocean related to monsoons; *Nature* **338** 749–751
- Naqvi S W A 1994 Denitrification processes in the Arabian Sea; *Proc. Indian Acad. Sci. (Earth Planet. Sci.)* **103** 279–300
- Naqvi S W A and Shailaja M S 1993 Activity of the respiratory electron transport system and respiration rates within the oxygen minimum layer of the Arabian Sea; *Deep-Sea Res.* **40** 687–695
- Naqvi S W A, Shailaja M S, Kumar M D and Gupta R S 1996 Respiration rates in subsurface waters of the northern Indian Ocean: Evidence for low decomposition rates of organic matter within the water column in the Bay of Bengal; *Deep-Sea Res.* **43** 73–81
- Naqvi S W A, Yoshinari T, Jayakumar D A, Altabet M A, Narvekar P V, Devol A H, Brandes J A, Codispoti L A 1998 Budgetary and biogeochemical implication of N_2O isotope signatures in the Arabian Sea; *Nature* **394** 462–464
- Parsons T R, Maita Y, Lalli C M 1984 A manual of chemical and biological methods for sea water analysis, (Oxford: Pergamon Press)
- Passow U and Alldredge A L 1994 Distribution, size and bacterial colonization of Transparent Exopolymer Particles (TEP) in the ocean; *Mar. Ecol. Prog. Ser.* **113**, 185–198
- Passow U and Alldredge A L 1995 A dye-binding assay for the spectrophotometric measurement of Transparent Exopolymer Particles (TEP); *Limnol. Oceanogr.* **40** 1326–1335
- Pomroy A J and Joint I R 1999 Bacterioplankton activity in the surface waters of the Arabian Sea during and after the 1994 SW monsoon; *Deep-Sea Res.* **46** 767–794
- Prasanna Kumar S, Ramaiah N, Mangesh Gauns, Sarma V V S S, Muraleedharan P M, Raghukumar S., Dileep Kumar M and Madhupratap M Time-Series analyses of the physical, chemical and biological characteristics of the northeastern Arabian Sea during winter; *Deep-Sea Res.* (in press)
- Qasim S Z 1982 Oceanography of the Northern Arabian Sea; *Deep-Sea Res.* **29** 1041–1068
- Ramaiah N, Raghukumar S and Gauns M 1996 Bacterial abundance and production in the central and eastern Arabian Sea; *Curr. Sci.* **71** 878–882
- Sarin M M, Rengarajan R and Ramaswamy V 1996 ^{234}Th scavenging and particle export fluxes from the upper 100 m of the Arabian Sea; *Curr. Sci.* **71** 888–893
- Smith D C, Steward G F, Long R A and Azam F 1995 Bacterial mediation of carbon fluxes during a diatom bloom in a mesocosm; *Deep-Sea Res.* **42** 75–98
- Smith S L, Codispoti L A, Morrison J M and Barber R T 1998 The 1994–1996 Arabian Sea Expedition: an integrated, interdisciplinary investigation of the northeastern Indian Ocean to monsoonal forcing; *Deep-Sea Res.* **45** 1905–1915
- Takahashi K 1986 Seasonal fluxes of pelagic diatoms in the subarctic Pacific, 1982–1983; *Deep-Sea Res.* **33** 1225–1251
- UNESCO 1994 Protocols for the Joint Global Ocean Flux Study (JGOFS) Core Measurements, IOC Manuals and Guides 29, UNESCO, Paris, 170 pp
- Veldhuis M J W, Kraay G W, van Bleijswijk J D L and Baars M A 1997 Abundance of bacterioplankton in relation to seasonal upwelling in the northwest Indian Ocean; *Deep-Sea Res.* **44** 451–476
- Wiebinga C J 1994 Bacterioplankton; In: *Monsoons and Pelagic Systems*. Netherlands Indian Ocean Programme Cruise Reports. Volume 1 (ed) M A Baars (Leiden: National Museum of Natural History) pp. 55–58.
- Wiebinga C J, Veldhuis M J W and De-Baar H J W 1997 Abundance and productivity of bacterioplankton in relation to seasonal upwelling in the northwest Indian Ocean; *Deep-Sea Res.* **14** 451–476
- Williams P M 1990 Studies of the physical and chemical stability of natural sea surface films; Technical Report, Scripps Institute of Oceanography, La Jolla, CA (USA) 3 pp