

Japanese encephalitis: pathogenesis, prophylactics and therapeutics

Kallol Dutta¹, Pundi N. Rangarajan², Sudhanshu Vрати³ and Anirban Basu^{1,*}

¹National Brain Research Centre, Manesar 122 050, India

²Department of Biochemistry, Indian Institute of Science, Bangalore 560 012, India

³National Institute of Immunology and Vaccine and Infectious Disease Research Center, THSTI, New Delhi 110 067, India

Japanese encephalitis (JE) is one of the most dreaded mosquito-borne viral encephalitis known to afflict humans. The Japanese encephalitis virus (JEV) is a neurotropic flavivirus that affects the CNS, causing extensive damage that may lead to fatality in about one third of patients. Half of the survivors suffer from severe neuropsychiatric sequelae. With nearly 3 billion people living under the current JE-endemic region, recurring incidents of epidemic are being reported at regular intervals. With no established antiviral therapies against JE available, vaccination has been the only way of preventing JE. Two types of JE vaccines are currently in vogue although the safety of administering them is questionable, in certain individuals. Thus, there is a need to develop a safe, affordable and potent JE vaccine and this review addresses the current efforts in this direction. This review also focuses on the pathophysiology of JE and efforts towards a possible breakthrough in anti-JEV therapy.

Keywords: Central neurons system, flavivirus, Japanese encephalitis virus, prophylaxis, vaccine,

Origin and spread of JEV

FROM the 1870s, recurring epidemics of encephalitis have been reported from the islands of Japan, especially during the summer season, with major outbreaks occurring every 10 years. This summer encephalitis was termed as type B encephalitis, to differentiate from von Economo's encephalitis lethargica. The name type B was later dropped and in 1935 the Nakayama strain of Japanese encephalitis virus (JEV) was isolated from the brain of a fatal case¹. The virus was later classified as a member of the genus *Flavivirus* (family *Flaviviridae*) named after the prototype Yellow fever virus (Latin; *flavi* = yellow). The genus consists of over 70 other closely related virus species. The origins of JEV are uncertain, but phylogenetic comparisons with other flaviviruses suggest that it evolved from an African ancestral virus, perhaps as recently as a few centuries ago² and evolved into its present form in the Indonesia–Malaysia region³. Since then,

the virus has spread in South Asia, Southeast Asia, East Asia and the Pacific regions⁴. Rapid globalization, population explosion, changes in global climatic condition, industrialization and deforestation, all seem to correlate with the spread of the virus into newer territories. Today, with approximately three billion people living in the JE-endemic region, there are estimated 35,000–50,000 cases and 10,000–15,000 deaths annually, thereby making JE one of the most dreaded vector-borne viral encephalitis in the world⁵ (Figure 1).

In India, JE as a disease was first reported in 1955 when clinical cases were detected in Vellore and Pondicherry in southern India. The virus was however, not recovered from humans until 1958, when three isolations were made from the brain tissue of infected persons. Until early 1970s, the disease was reported only from southern India with periodic focal reports of its occurrence. A major outbreak resulting in 42.6% case-fatality rate was reported from Bankura district of West Bengal in 1973. Subsequently, the disease spread to other states and caused a series of outbreaks in different parts of the country. In 1978, cases were reported from 21 states and union territories, and from then onwards till 2007 there have been 103,389 reported cases of JE in India that has led to 33,729 deaths⁶. Approximately 597,542,000 people in India live under the JE-endemic regions and there are 1500–4000 reported cases every year⁷. These figures are based on total reported cases and it is quite possible that several cases may go unreported and hence the actual magnitude of the threat of JE may be considerably higher both in the Indian as well as the global context.

Enzootic cycle of JEV

JE is a vector-borne viral disease. JEV is transmitted between vertebrate hosts by mosquitoes belonging to the *Culex* sp. (Figure 2). The virus is able to replicate within the salivary glands of the mosquitoes. Mature JE virions remain entrapped in intracellular vacuoles and are later released into the apical cavity of salivary gland cells through the fusion of these vacuoles with the apical plasma membrane. This process is associated with primary re-synthesis of saliva in mosquitoes following blood feeding activity. Another type of shedding involves

*For correspondence. (e-mail: anirban@nbrc.ac.in)




Figure 1. Japanese encephalitis (JE) endemic regions. The current JE-endemic region consisting of entire Southeast Asia, parts of Russia and Australia. The possible direction of spread of the disease is shown as broken arrows. (Figure not to scale; adapted from Ghosh and Basu, *PLoS Neglected Tropical Diseases*, 2009.)




Figure 2. Enzootic transmission cycle of JE virus. The virus is transmitted to vertebrate hosts by mosquitoes belonging to the *Culex sp*. Pigs serve as amplification hosts and forms a critical link in the transmission cycle. Ardeid water birds and bats serve as virus reservoirs. Humans are dead end hosts as the virus cannot be transmitted from an infected person to another.

virus particles, either singly or in mass, being released directly through the apical plasma membrane⁸. Components of the mosquito saliva may also modulate infection by altering the local cytokine milieu. Feeding by mosquitoes of *Culex* sp. or administration of sialokinin-I, a mosquito salivary protein, have been found to downregulate IFN- γ production and upregulate the TH₂ cytokines, IL-4 and IL-10 (ref. 9).

Human infections are mainly spread by *Culex tritaeniorhynchus* which breeds in pools of stagnant water such as rice paddy fields¹⁰. Because the rice paddy is unavoidable, majority of the population in rural Asia has been infected with the virus by early adulthood¹¹. Wading ardeid water birds (e.g. herons and pond egrets) and bats serve as virus reservoirs or maintenance hosts, but the virus regularly spills over into pigs, members of the family of Equidae (e.g. horses and donkeys) and humans. Pigs are considered as the main amplification hosts as viraemia results with a high titre. Due to the close proximity of pigs with human dwellings these animals are considered main component in the transmission cycle with respect to human infection¹². JEV infection in other domestic animals does not result in high viraemia and thus, they are not expected to transmit the virus to humans.

Molecular architecture of JEV

The JE virion consists of a single strand of positive-sense RNA of around 11 kilobase, inside a nucleocapsid and is surrounded by a glycoprotein-containing envelope. The RNA comprises a short 5' untranslated region (UTR), a longer 3'UTR and a single open reading frame between them. It codes for a single polyprotein, which is translationally and post-translationally cleaved by viral and host proteases into three structural proteins (core-C, pre-membrane-PrM and envelope-E), and seven nonstructural (NS) proteins (NS1, NS2A, NS2B, NS3, NS4A, NS4B, and NS5). The C protein (12–14 kDa) is highly basic and combines with the RNA to form the nucleocapsid¹³. The prM is closely associated with the E protein, forming a heterodimer and is thought to act as a 'chaperone' to it, impairing its function until after virion release. Immediately prior to virion release, the prM protein (18–19 kDa) is cleaved to its mature M protein (8–9 kDa) form. This allows the formation of E protein homodimers, which are thus 'activated'. The prM protein of JEV contains a single N-linked glycosylation site, which is highly conserved among the JEV strains. Researches indicated that this highly conserved N glycosylation motif in prM is crucial for multiple stages of JEV biology: prM biogenesis, virus release and pathogenesis¹⁴. The E protein is the largest structural protein (53–55 kDa), with up to two potential glycosylation sites. It is the major target for the humoral immune response and is thought to be important for viral entry into host cells. It is worth mentioning that low pH is extremely important for viral entry into the cell to trigger

viral membrane fusion with host endosomal membrane, thereby releasing the nucleoplasmid in the cytosol.

Pathophysiology and clinical manifestations of JE

The course of JEV infection, starting from its entry till reaching its site of action, the central nervous system (CNS), is not well-defined. Studies with other flaviviruses have led to the belief that upon entry through mosquito bite, the virus infects Langerhan's dendritic cells in the skin¹⁵ and is carried to nearest draining lymph nodes¹⁶, thereby initiating a round of early immune response. Unfortunately for the host, this is not sufficient to counter the virus and it spreads to secondary lymphoid organs before entering the blood circulation through the efferent lymphatic system. During the ensuing transient viraemia, peripheral organs such as kidney, liver and spleen are known to be infected¹⁷, after which the neurotropic flavivirus spreads to the CNS. How JEV manages to evade the host's peripheral immune response is another matter of speculation, as there are no definitive studies regarding this aspect, till date. Once the virus evades the immune system, it crosses the blood–brain barrier (BBB) to enter the CNS. JEV may cross the BBB by passive transport across the endothelium, by active replication in endothelial cells, or by a 'Trojan horse' mechanism in which the virus is carried into the brain by infected inflammatory cells¹⁸. Monocytes and macrophages are thought to be probable carriers of the virus into the CNS as studies have shown that the virus can survive for a prolonged time and effectively replicate within these cells^{19,20}. The structural and functional integrity of the BBB is critically compromised during JE²¹ which may possibly be caused by matrix metalloproteases released in the infected brain²². Due to the impaired functioning of the BBB, peripheral inflammatory cells are recruited to the infected brain at later stages of JE that exacerbates the damage.

JEV causes extensive neuronal damage in the brain, though in many cases, the virus is probably not directly involved in the destruction of brain tissue but may cause damage indirectly by triggering cell-mediated immune response by activating microglia. Microglia are the resident immune cells of the CNS and have a critical role in host defense against invading microorganisms. Microglial activation is viewed as an adaptive response whereby microglia release neuroprotective factors to facilitate the recovery of injured neurons and they also phagocytose dying or damaged neurons, before they lyse and release toxic agents into surrounding areas. JEV infection has been shown to activate microglia both morphologically and functionally, *in vivo*, that leads to an elevation of pro-inflammatory mediators, such as IL-6, TNF- α , RANTES and MCP-1. These proinflammatory mediators and cytotoxins released from activated microglia are instrumental in inducing neuronal death that accompanies JE²³.

Neuronal death by secreted TNF is mediated by the TNF receptor-associated death domain protein (TRADD)²⁴, which there upon regulates a downstream apoptotic cascade in neurons. However, neuronal death also activates microglia and astrocytes and thus, the inflammatory cycle goes on. Nitric oxide (NO) also plays an important role in inflammation during JE infection, although NO itself is a strong antimicrobial agent researchers have shown that it profoundly inhibits viral RNA synthesis, viral protein accumulation and virus release from infected cells²⁵. Thus, NO may play a crucial role in the innate immunity of the host and its ability to restrict the initial stages of JEV infection in the CNS.

Besides neuronal and microglial cells, studies have shown that astrocytes are also infected by JEV²⁶. Astrocytes are known to maintain homeostasis in the CNS and to support the survival and information processing functions of neurons. They respond promptly to CNS infection and help regulate neuroinflammation²⁷. JEV infection results in astrocyte activation, but the infection overwhelms the capacity of even activated astrocytes to maintain metabolic homeostasis, resulting in an over accumulation of toxic byproducts of metabolism that are detrimental to neuronal viability. Though JEV infection triggers metabolic reprogramming by upregulating the expression of several proteins such as IP-10 (ref. 28), ceruloplasmin and glutamine synthetase, involved in the metabolic pathways crucial for maintaining neuronal health, this increase is insufficient to meet the increased demand that accompanies JEV infection²⁹. Astrocytes, being a component of the BBB, may also help in the transmission of JEV from peripheral tissues to the cerebrospinal fluid. JEV also causes the upregulation of certain host genes in the brain. An *in vivo* study using subtraction hybridization technique has identified several genes whose expression is upregulated by JEV in mouse brain^{30,31}. It has been further demonstrated that several of these genes are also induced in the CNS by rabies virus and Sindbis virus. Thus, it appears that different neurotropic viruses activate common host cell pathways during infection of CNS^{30,31}.

JE typically develops in patients after an incubation period of 5–15 days³². In humans, most JEV infections are asymptomatic, with about one in 300 JEV infections resulting in symptoms ranging from non-specific febrile illness to severe meningoencephalitis, characterized by fever, reduced consciousness, seizures and focal neurological signs³³. At later stages, poliomyelitis-like flaccid paralysis and parkinsonian syndrome develops, which manifests the classic description of JE – dull, flat, mask-like face with wide, unblinking eyes, tremor, generalized hypertonia, cogwheel rigidity, and other abnormalities in movement³⁴. Histopathological and imaging studies have shown JEV infection and inflammation in the anterior horn cells of the spinal cord, which provided an anatomical correlate for these presentations³⁵. Radiological studies also support the earlier observations from pathological

investigations implicating the basal ganglia, especially the substantia nigra and thalamus in the parkinsonian syndromes seen in JE, which include tremor, cogwheel rigidity and mask-like faces³⁶. Recently, there has been further molecular evidence of the involvement of nigrostriatal pathway in JE survivors with movement complications. In the study, (99m) Tc-TRODAT-1 and (123)I-IBZM SPECT imaging revealed that presynaptic dopaminergic neurons in JE patients are more susceptible to JEV than post-synaptic striatal neurons³⁷. Electroencephalographic abnormalities observed in JE patients include the presence of alpha, theta and delta coma, and epileptiform conditions³⁸. Symptoms of brainstem infection include changes in the respiratory pattern, flexor and extensor posturing, and abnormalities in the papillary and oculoccephalic reflexes³⁹. In fatal cases of JE, pathological changes are polymorphic and diffuse, involving various parts of the nervous system where the brain shows a severe degree of vascular congestion, microglial proliferation and formation of gliomesenchymal nodules, focal or confluent areas of cystic necrosis, cerebral oedema and transcompartmental shift⁴⁰. Many survivors of JE acquire neuropsychiatric sequelae with cognitive and language impairment, in which case the disease presents itself not only as a killer but also as a cause of an immense social and financial burden, especially for a developing country³³. Recently, it has been reported that neural progenitor cells (NPC) are permissive to JEV infection both *in vivo* and *in vitro*, which leads to their growth retardation. The pathophysiological relevance of these observations was supported by profound decrement in actively proliferating NPCs in the subventricular zone (SVZ) of JEV-infected animals. Infection of the NPCs and suppression of their proliferation might be primarily responsible for dysregulated neurogenesis and development of cognitive deficits in survivors of JE⁴¹.

Prophylactic strategies against JE

Due to lack of definitive therapeutic countermeasures to combat JE, implementation of prophylactic strategies remains the best way to prevent JE. The most important aspect of the prevention of JE spread is vector control. Paddy fields not only provide an ideal breeding ground for mosquitoes but also attract migratory birds, thereby helping the spread of the virus. Responsible use of larvicides, insecticides, and adopting ecofriendly methods such as using neem cakes and growing larvivorous fish are useful in controlling mosquitoes in paddy fields⁴. As pigs are the amplification host in the transmission cycle of the virus, close monitoring of domesticated animals is necessary. Vaccinating pigs is also a logical way to acquire protection against JEV by breaking the mosquito–pig–human transmission cycle⁴².

Human vaccination remains, till date, the most effective measure to prevent JE. Multiple vaccines exist to

control JE, but all have limitations. Unfortunately, unlike smallpox, it is difficult to eradicate JEV by vaccination, because human beings are actually a dead-end host of the virus. The formalin-inactivated vaccine against JEV was produced from infected mouse brain-derived tissue soon after the virus was discovered. This type of vaccine, manufactured by the Research Foundation for Microbial Diseases of Osaka University, Japan, became commercially available in Japan (as the Japanese Biken vaccine JE-VAX) and was produced in Korea by the Green Cross Vaccine company. These were later licensed to be produced in the United States also⁴³. This is the only JE vaccine recommended by the World Health Organization (WHO), but there have been several concerns with its side effects⁴⁴. These vaccines are expensive and require multiple doses to maintain efficacy and immunity. The Biken vaccine is no longer produced. The Korean Green Cross vaccine needs to be administered also in three doses on day 0, 7 and 28 after birth, with boosters given every 12 months following primary course, then every three years, unless at particular risk in which case annual boosters are recommended. This makes its use difficult for people in developing countries. Researchers have been trying to avoid a multidose requirement of the vaccine by using adjuvants such as biodegradable poly (gamma-glutamic acid) nanoparticles and alum⁴⁵. Another mouse brain inactivated JE vaccine was produced in India by the Central Research Institute (CRI) in Kasauli⁴⁶ but the production of the vaccine was suspended from 2007 as the institute had failed to upgrade its main laboratory according to the GMP norms of WHO⁴⁷. An inexpensive, live attenuated vaccine (SA14-14-2)⁴⁸ was licensed by China in 1988, but WHO does not approve it for human use because it is produced in primary hamster kidney cells. Initially, this vaccine was used almost exclusively in China and parts of Korea, but it is now being widely used in the Indian subcontinent as well as Nepal. The vaccine seems to be highly effective, and very few adverse effects have been observed following its use in Nepal⁴⁹. No information is, however, available on the efficacy and side effects of this vaccine in India. Neuroattenuation of the virulent JE SA14-14-2 strain is reported to be based on 57 nucleotide changes and 24 amino acid substitutions, suggesting that reversion to neurovirulence of the vaccine strain is unlikely. Though the vaccine was adjudged to be safe and efficacious by the WHO's Global Advisory Committee on Vaccine Safety, several parameters such as safety in immunocompromised individuals and pregnant women, viral shedding in vaccines and implications of the shedding, efficacy in infants less than one year of age, remains to be ascertained⁵⁰. The recently developed Vero cell-derived inactivated JE vaccine containing the purified, inactivated JEV strain SA14-14-2 with aluminum hydroxide as adjuvant seems to be a promising candidate and has passed the phase III randomized controlled trial⁵¹. Technology for a Vero cell-derived vaccine based

on an Indian strain of JEV⁵² has been developed and the vaccine is under pre-clinical development.

Several efforts have been and are still being made to develop recombinant vaccines for JEV. A highly attenuated recombinant vaccinia virus, NYVAC⁵³ was used for JE vaccine development which proved to be quite successful⁵⁴. Other strains tested include modified vaccinia virus Ankara strain^{55,56}. Later studies focused on safe replication-competent recombinant viruses such as avipoxviruses and ALVAC, which infect cells but do not replicate^{56,57}. In another research on peptide-based vaccine, a JEV E protein was fused to Johnson grass mosaic virus coat protein by using recombinant DNA technology⁵⁸.

Researchers have also constructed the Yellow fever virus to express JEV protein, the most promising recombinant vaccine under development. Widely known as ChimeriVax-JE, this vaccine incorporates the structural proteins (prM and E) of the JEV SA14-14-2 strain into YF17D, the yellow fever vaccine⁵⁹⁻⁶¹.

Attempts have been made to generate DNA vaccines for JE and efficacy of various plasmid DNA constructs encoding the JEV E protein or prM has been evaluated in murine and primate models⁶²⁻⁶⁵. A single intramuscular injection of recombinant plasmid DNA containing cDNA encoding JEV prM and E proteins induced protective immunity and prevented JE in mice and their progeny⁶³. Again, investigators have used both intramuscular injection and a gene gun to deliver the plasmid gene for E protein and prM protein⁶⁴. DNA immunization with colloidal gold, construction of chimeric DNA vaccine vectors are the focus of some of the other recent promising studies^{65,66}. A novel genetically engineered DNA vaccine against flavivirus has been proposed in which the consensus sequence of structural domain III of the E protein of the virus that is reported to carry dominant epitopes that induce neutralizing antibodies, has been created. This has been administered along with a genetically engineered IL15 DNA vaccine molecular adjuvant for co-stimulating the immune response against DIII clones. This has resulted in a 4-5 fold increased immune response⁶⁷. In another strategy, needle-free jet injection of a mixture of JEV DNA and protein vaccines was shown to enhance the immunogenicity of the DNA vaccine in a murine model⁶⁸. Recently, pseudoinfectious JE viruses (named RepliVAX JE) lacking the JEV capsid protein capable of replicating in normal cells but incapable of producing infectious particles have been shown to protect mice and hamsters from lethal JEV challenge indicating that the RepliVAX platform can be used for producing a safe and potent JE vaccine⁶⁹.

Therapeutic measures against JE

Chemotherapy against JE is at present supportive and not targeted at the virus. Interferon therapy has not met with

Table 1. Promising preventive and therapeutic strategies for Japanese encephalitis

Prophylactic strategies – vaccine development	
ChimeriVax-JE	Incorporates the structural proteins (prM and E) of the JEV SA14-14-2 strain into YF17D, the Yellow fever vaccine
RepliVAX JE	Pseudo-infectious JE viruses lacking the JEV capsid protein capable of replicating in normal cells but incapable of producing infectious particles
Vero cell-derived inactivated JE vaccine	Contain purified, inactivated JEV strain SA14-14-2 with aluminum hydroxide as adjuvant; has passed phase III randomized control trials; Vero cell-derived vaccine based on an Indian strain of JEV has also been developed and is currently under pre-clinical development
Recombinant vaccinia and avipoxvirus	JE vaccines using recombinant strains of vaccinia virus and avipoxvirus have been shown to be quite effective
Peptide-based vaccine	JEV E protein was fused to Johnson grass mosaic virus coat protein by using recombinant DNA technology
DNA vaccines	Efficacy of various plasmid DNA constructs encoding the JEV E protein or prM has been evaluated; DNA immunization with colloidal gold, construction of chimeric DNA vaccine vectors has also been found to be effective; the consensus sequence of structural domain III of the E protein of JEV that is reported to carry dominant epitopes that induce neutralizing antibodies, has been created an administered along with a genetically engineered IL15 DNA vaccine molecular adjuvant for co-stimulating the immune response against DIII clones
Therapeutic strategies – drug development	
Rosmarinic acid	Reduces JEV replication in mice brain and also ameliorates the secondary inflammation resulting from microglial activation
Arctigenin	Confer substantial protection in a murine model from GP-78 strain of JEV infection by markedly decreasing virus-induced neuronal apoptosis, microglial activation, active caspase activity and induction of proinflammatory mediators in the brains
Curcumin	Dysregulates ubiquitin-proteasome pathway to inhibit infective viral particle formation in <i>in vitro</i> studies; reduces JEV-induced cellular stress
<i>N</i> -methylisatin- β -thiosemicarbazone derivative	Inhibits JEV replication
Glucosidase inhibitors	Block the trimming step of N-linked glycosylation, thereby eliminate the production of several endoplasmic reticulum-budding viruses, including JEV
Minocycline	Reduces neuronal apoptosis, microglial activation, active caspase activity, proinflammatory mediators and viral titer; possess antioxidative properties that significantly ameliorate the oxidative stress generated as a result of JEV infection; imparts protection to the blood–brain barrier that is severely compromised in JE
siRNA	Single intracranial administration of lentivirus-delivered short hairpin RNA or lipid-complexed siRNA before viral challenge or siRNA treatment after viral challenge has been found to be sufficient to provide protection against lethal encephalitis

great success⁷⁰. Over-expression of interferon stimulating gene 15 has been found to impart protection against several viral infections^{71–74} and may also work against JEV. A recent clinical trial with oral administration of ribavirin, an antiviral drug that interferes with RNA metabolism required for viral replication, was also not encouraging⁷⁵. A naturally occurring compound discovered to have anti-flaviviral activity and the potential to be developed as an anti-flaviviral drug is rosmarinic acid (RA). RA, a phenolic compound found in various Labiatae herbs, possesses several anti-inflammatory and antioxidative properties. An *in-vivo* study conducted with GP78 strain of JEV in murine model has showed that RA reduces viral replication in mice brain and also ameliorates the secondary inflammation resulting from microglial activation⁷⁶. Arctigenin, a naturally occurring phenylpropanoid dibenzylbutyrolactone lignan is known to possess antioxidant, anti-inflammatory, neuroprotective and antiviral properties. It has been reported that treatment with arctigenin confer substantial protection in a murine model from GP-78 strain of JEV infection by markedly decreasing virus-induced neuronal apoptosis, microglial activation,

active caspase activity and induction of proinflammatory mediators in the brains of arctigenin-treated mice⁷⁷. The study also claims that arctigenin may act directly on brain cells, because neuronal cell line was also salvaged from JEV-induced cell death. Curcumin, a naturally occurring phenolic compound extracted from *Curcuma longa* L., has been shown to impart neuroprotection against JEV infection in an *in-vitro* study. Curcumin possibly acts by decreasing cellular reactive oxygen species level, restoration of cellular membrane integrity, decreasing pro-apoptotic signalling molecules, and modulating cellular levels of stress-related proteins. It has also been shown that the production of infective viral particles from previously infected neuroblastoma cells is reduced, which is achieved by the inhibition of ubiquitin-proteasome system⁷⁸.

A remarkable achievement in anti-flaviviral drug research is the discovery of antiviral effect of minocycline, a member of the broad-spectrum antibiotic tetracycline group⁷⁹. A significant piece of research on minocycline as an anti-JEV drug is an *in-vivo* study⁸⁰ that showed minocycline reduces neuronal apoptosis, microglial activation, active caspase activity, proinflammatory mediators and

viral titer markedly on the 9th day post infection. Minocycline was also shown to possess antioxidative properties that significantly ameliorate the oxidative stress generated as a result of JEV infection⁸¹. Minocycline also imparts protection to the BBB that is severely compromised in JE²². Another compound that has shown inhibition of JEV replication completely *in vitro* is an *N*-methylisatin- β -thiosemicarbazone derivative⁸².

Glucosidase inhibitors of the endoplasmic reticulum, such as *N*-nonyl-deoxyjirimycin, which block the trimming step of *N*-linked glycosylation, have been shown to eliminate the production of several endoplasmic reticulum-budding viruses, including dengue type II (DEN-2) and JEV⁸³. In a novel approach, a synthetic oligonucleotide-based DNAzyme targeting the 3'NCR of JEV was shown to significantly inhibit virus replication in cultured cells and mouse brain and protected the JEV-infected mice from death⁶⁶. In yet another innovative study on mice using RNA interference, a single intracranial administration of lentivirus-delivered short hairpin RNA or lipid-complexed small interfering RNA (siRNA) before viral challenge or siRNA treatment after viral challenge was sufficient to provide protection against lethal encephalitis. Interestingly, when a cross-species conserved sequence of cd-loop coding viral envelope protein was targeted, encephalitis following both JEV and WNV challenge was avoided. This result clearly indicates that by careful drug design of the conserved target site, a single siRNA treatment could suppress viral infection across species, thereby augmenting the treatment of acute viral infections with overlapping clinical symptoms⁸⁴.

Future perspectives

With the advent of modern technologies and advancement of knowledge, research in drug and vaccine development against JE has been going on at a steady rate for the past two decades. Though a lot of progress has been made in unraveling the mysteries of the virus that has helped in planning and designing countermeasure strategies, the real success would be when an effective drug or vaccine would be easily available to the masses exposed to the threat of JE, at an affordable cost. Table 1 summarizes the most promising prophylactic and therapeutic approaches against JE. It has been hypothesized that this virus may spread to greater geographic areas apart from its current endemic region. Thus it would be prudent to be prepared for epidemics of greater magnitudes.

- Solomon, T., Dung, N. M., Kneen, R., Gainsborough, M., Vaughn, D. W. and Khanh, V. T., Japanese encephalitis. *J. Neurol. Neurosurg. Psychiatry.*, 2000, **68**, 405–415.
- Gould, E. A., Evolution of the Japanese serological group viruses. In *Current Topics in Microbiology and Immunology: Japanese Encephalitis and West Nile Virus Infections* (eds Mackenzie, J. S., Barrett, A. D. and Deubel, V.), Springer-Verlag, Berlin, 2002, pp. 391–404.
- Solomon, T., Ni, H., Beasley, D. W., Ekkelenkamp, M., Cardoso, M. J. and Barrett, A. D. T., Origin and evolution of Japanese encephalitis virus in southeast Asia. *J. Virol.*, 2003, **77**, 3091–3098.
- Solomon, T., Control of Japanese encephalitis – within our grasp? *N. Engl. J. Med.*, 2006, **355**, 869–871.
- Tsai, T. F., New initiatives for the control of Japanese encephalitis by vaccination: minutes of a WHO/CVI meeting, Bangkok, Thailand, 13–15 October 1998. *Vaccine*, 2000, **18**, 1–25.
- Dhillon, G. P. and Raina, V. K., Epidemiology of Japanese encephalitis in context with Indian scenario. *J. Indian. Med. Assoc.*, 2008, **106**, 660–663.
- Kabilan, L., Rajendran, R., Arunachalam, N., Rameshm, S., Srinivasan, S., Samuel, P. P. and Dash, A. P., Japanese encephalitis in India: an overview. *Indian J. Pediatr.*, 2004, **71**, 609–615.
- Takahashi, M. and Suzuki, K., Japanese encephalitis virus in mosquito salivary glands. *Am. J. Trop. Med. Hyg.*, 1979, **28**, 122–135.
- Zeidner, N. S., Higgs, S., Happ, C. M., Beaty, B. J. and Miller, B. R., Mosquito feeding modulates Th1 and Th2 cytokines in flavivirus susceptible mice: an effect mimicked by injection of sialokinins, but not demonstrated in flavivirus resistant mice. *Parasite Immunol.*, 1999, **21**, 35–44.
- Innis, B. L., Japanese encephalitis. In *Exotic Viral Infections* (ed. Porterfield, J. S.), Chapman and Hall, London, 1995, pp. 147–174.
- Solomon, T., Recent advances in Japanese encephalitis. *J. Neurovirol.*, 2003, **9**, 274–283.
- Ghosh, D. and Basu, A., Japanese encephalitis – a pathological and clinical perspective. *PLoS Negl. Trop. Dis.* (in press).
- Kaur, R. and Vrati, S., Development of a recombinant vaccine against Japanese encephalitis. *J. Neurovirol.*, 2003, **9**, 421–431.
- Kim, J.-M., Yun, S.-I., Song, B.-H., Hahn, Y.-S., Lee, C.-H., Oh, H.-W. and Lee, Y.-M., A single N-linked glycosylation site in the Japanese encephalitis virus prM protein is critical for cell type-specific prM protein biogenesis, virus particle release, and pathogenicity in mice. *J. Virol.*, 2008, **82**, 7846–7862.
- Ho, L.-J., Wang, J.-J., Shaio, M.-F., Kao, C.-L., Chang, D.-M., Han, S.-W. and Lai, J.-H., Infection of human dendritic cells by dengue virus causes cell maturation and cytokine production. *J. Immunol.*, 2001, **166**, 1499–1506.
- Johnston, L. J., Halliday, G. M. and King, N. J., Langerhans cells migrate to local lymph nodes following cutaneous infection with an arbovirus. *J. Invest. Dermatol.*, 2000, **114**, 560–568.
- Solomon, T. and Vaughn, D. W., Pathogenesis and clinical features of Japanese encephalitis and West Nile virus infections. *Curr. Top. Microbiol. Immunol.*, 2002, **267**, 171–194.
- Diamond, M. S., Evasion of innate and adaptive immunity by flaviviruses. *Immunol. Cell Biol.*, 2003, **81**, 196–206.
- Yang, K. D. *et al.*, A model to study neurotropism and persistency of Japanese encephalitis virus infection in human neuroblastoma cells and leukocytes. *J. Gen. Virol.*, 2004, **85**, 635–642.
- Aleyas, A. G. *et al.*, Functional modulation of dendritic cells and macrophages by Japanese encephalitis virus through MyD88 adaptor molecule-dependent and -independent pathways. *J. Immunol.*, 2009, **183**, 2462–2474.
- Mathur, A., Khanna, N. and Chaturvedi, U. C., Breakdown of blood-brain barrier by virus-induced cytokine during Japanese encephalitis virus infection. *Int. J. Exp. Pathol.*, 1992, **73**, 603–611.
- Mishra, M. K., Dutta, K., Saheb, S. K. and Basu, A., Understanding the molecular mechanism of blood-brain barrier damage in an experimental model of Japanese encephalitis: correlation with minocycline administration as a therapeutic agent. *Neurochem. Int.*, 2009 (in press).
- Ghoshal, A. *et al.*, Proinflammatory mediators released by activated microglia induces neuronal death in Japanese encephalitis. *Glia*, 2007, **55**, 483–496.

24. Swarup, V., Ghosh, J., Das, S. and Basu, A., Tumor necrosis factor receptor-associated death domain mediated neuronal death contributes to the glial activation and subsequent neuroinflammation in Japanese encephalitis. *Neurochem. Int.*, 2008, **52**, 1310–1321.
25. Lin, Y. L. *et al.*, Inhibition of Japanese encephalitis virus infection by nitric oxide: antiviral effect of nitric oxide on RNA virus replication. *J. Virol.*, 1997, **71**, 5227–5235.
26. Chen, C. J., Liao, S. L., Kuo, M. D. and Wang, Y. M., Astrocytic alteration induced by Japanese encephalitis virus infection. *Neuroreport*, 2000, **11**, 1933–1937.
27. Kuhlow, C. J., Krady, J. K., Basu, A. and Levison, S. W., Astrocytic ceruloplasmin expression, which is induced by IL-1 β and by traumatic brain injury, increases in the absence of the IL-1 type 1 receptor. *Glia*, 2003, **44**, 76–84.
28. Bhowmick, S., Duseja, R., Das, S., Appaiahgiri, M. B., Vrati, S. and Basu, A., Induction of IP-10 (CXCL10) in astrocytes following Japanese encephalitis. *Neurosci. Lett.*, 2007, **414**, 45–50.
29. Mishra, M. K., Koli, P., Bhowmick, S. and Basu, A., Neuroprotection conferred by astrocytes is insufficient to protect animals from succumbing to Japanese encephalitis. *Neurochem. Int.*, 2007, **50**, 764–73.
30. Saha, S. and Rangarajan, P. N., Common host genes are activated in mouse brain by Japanese encephalitis and rabies viruses. *J. Gen. Virol.*, 2003, **84**, 1729–1735.
31. Saha, S., Sugumar, P., Bhandari, P. and Rangarajan, P. N., Identification of Japanese encephalitis virus inducible genes in mouse brain and characterization of GARG39/IFIT2 as a microtubule-associated protein. *J. Gen. Virol.*, 2006, **87**, 3285–3289.
32. Burke, D. S., Lersomrudee, W., Leake, C. J., Hoke, C. H., Nisalak, A., Chongsawasdi, V. and Laorakpongse, T., Fatal outcome in Japanese encephalitis. *Am. J. Trop. Med. Hyg.*, 1985, **34**, 203–210.
33. Vaughn, D. W. and Hoke, C. H., The epidemiology of Japanese encephalitis: prospects for prevention. *Epidemiol. Rev.*, 1992, **14**, 197–221.
34. Solomon, T. *et al.*, Poliomyelitis-like illness due to Japanese encephalitis virus. *Lancet*, 1998, **351**, 1094–1097.
35. Kumar, S., Misra, U. K., Kalita, J., Sawani, V., Gupta, R. K. and Gujral, R., MRI in Japanese encephalitis. *Am. J. Med. Sci.*, 1997, **39**, 180–184.
36. Murgod, U. A., Muthane, U. B., Ravi, V., Radhesh, S. and Desai, A., Persistent movement disorders following Japanese encephalitis. *Neurology*, 2001, **57**, 2313–2315.
37. Liao, C. H. *et al.*, Involvement of nigrostriatal pathway in Japanese encephalitis with movement disorders: evidence from (99m)Tc-TRODAT-1 and (123)I-IBZM SPECT imaging. *Mol. Imaging Biol.*, 2009 (in press).
38. Kalita, J. and Misra, U. K., EEG in Japanese encephalitis: a clinico-radiological correlation. *Electroencephalogr. Clin. Neurophysiol.*, 1998, **106**, 238–243.
39. Kumar, R., Mathur, A., Kumar, A., Sharma, S., Chakraborty, S. and Chaturvedi, U. C., Clinical features and prognostic indicators of Japanese encephalitis in children in Lucknow (India). *Indian J. Med. Res.*, 1990, **91**, 321–327.
40. Ishii, T., Matsushita, M. and Hamada, S., Characteristic residual neuropathological features of Japanese B encephalitis. *Acta Neuropathol.*, 1977, **38**, 181–186.
41. Das, S. and Basu, A., Japanese encephalitis virus infects neural progenitor cells and decreases their proliferation. *J. Neurochem.*, 2008, **106**, 1624–1636.
42. Sasaki, O., Karoji, Y., Kuroda, A., Karaki, T., Takenokuma, K. and Maeda, O., Protection of pigs against mosquito-borne Japanese encephalitis virus by immunization with a live attenuated vaccine. *Antiviral Res.*, 1982, **2**, 355–360.
43. Solomon, T., New vaccines for Japanese encephalitis. *Lancet Neurol.*, 2008, **7**, 116–118.
44. Shlim, D. R. and Solomon, T., Japanese encephalitis vaccine for travelers: exploring the limits of risk. *Clin. Infect. Dis.*, 2002, **35**, 183–188.
45. Okamoto, S. *et al.*, Single dose of inactivated Japanese encephalitis vaccine with poly(γ -glutamic acid) nanoparticles provides effective protection from Japanese encephalitis virus. *Vaccine*, 2008, **26**, 589–594.
46. Goyal, D. and Tahlan, A. K., Evaluation of effectiveness of mouse brain inactivated Japanese encephalitis vaccine produced in India. *Indian J. Med. Res.*, 1995, **102**, 267–271.
47. <http://www.expressindia.com/latest-news/Kasauli-CRIs-drug-licence-suspended/266285/>
48. Jia, L., Wang, Z. and Yu, Y., Protection of SA14-14-2 live attenuated Japanese encephalitis vaccine against the wild-type JE viruses. *Chin. Med. J. (Engl.)*, 2003, **116**, 941–943.
49. Hennessy, S. *et al.*, Effectiveness of live-attenuated Japanese encephalitis vaccine (SA14-14-2): a case-control study. *Lancet*, 1996, **347**, 1583–1586.
50. http://www.who.int/vaccine_safety/topics/japanese_encephalitis/live_attenuated/June_2005/en/index.html
51. Tauber, E. *et al.*, Safety and immunogenicity of a Vero-cell-derived, inactivated Japanese encephalitis vaccine: a non-inferiority, phase III, randomised controlled trial. *Lancet*, 2007, **370**, 1847–1853.
52. Appaiahgari, M. B. and Vrati, S., Immunogenicity and protective efficacy in mice of a formaldehyde-inactivated Indian strain of Japanese encephalitis virus grown in Vero cells. *Vaccine*, 2004, **22**, 3669–3675.
53. Tartaglia, J. *et al.*, NYVAC: a highly attenuated strain of vaccinia virus. *Virology*, 1992, **188**, 217–232.
54. Konishi, E., Pincus, S., Paoletti, E., Laegreid, W. W., Shope, R. E. and Mason, P. W., A highly attenuated host range-restricted vaccinia virus strain, NYVAC, encoding the prM, E, and NS1 genes of Japanese encephalitis virus prevents JEV viremia in swine. *Virology*, 1992, **190**, 454–458.
55. Nam, J. H., Wyatt, L. S., Chae, S. L., Cho, H. W., Park, Y. K. and Moss, B., Protection against lethal Japanese encephalitis virus infection of mice by immunization with the highly attenuated MVA strain of vaccinia virus expressing JEV prM and E genes. *Vaccine*, 1999, **17**, 261–268.
56. Konishi, E., Pincus, S., Paoletti, E., Shope, R. E. and Wason, P. W., Avipox virus-vectored Japanese encephalitis virus vaccines: use as vaccine candidates in combination with purified subunit immunogens. *Vaccine*, 1994, **12**, 633–638.
57. Kanesa-athan, N. *et al.*, Safety and immunogenicity of attenuated dengue virus vaccines (Aventis Pasteur) in human volunteers. *Vaccine*, 2001, **19**, 3179–3188.
58. Saini, M. and Vrati, S., A Japanese encephalitis virus peptide present on Johnson grass mosaic virus-like particles induces virus-neutralizing antibodies and protects mice against lethal challenge. *J. Virol.*, 2003, **77**, 3487–3494.
59. Monath, T. P. *et al.*, Recombinant, chimeric live, attenuated vaccine (ChimeriVax) incorporating the envelope genes of Japanese encephalitis (SA14-14-2) virus and the capsid and nonstructural genes of yellow fever (17D) virus is safe, immunogenic and protective in non-human primates. *Vaccine*, 1999, **17**, 1869–1882.
60. Monath, T. P. *et al.*, Chimeric yellow fever virus 17D-Japanese encephalitis virus vaccine: dose-response effectiveness and extended safety testing in rhesus monkeys. *J. Virol.*, 2000, **74**, 1742–1751.
61. Monath, T. P. *et al.*, Clinical proof of principle for ChimeriVax: recombinant live, attenuated vaccines against flavivirus infections. *Vaccine*, 2002, **20**, 1004–1018.
62. Bharati, K., Rani, R. and Vrati, S., Evaluation of Japanese encephalitis virus DNA vaccine candidates in rhesus monkeys [Macaca mulatta]. *Vaccine*, 2009, **27**, 10–16.

63. Chang, G. J., Hunt, A. R. and Davis, B., A single intramuscular injection of recombinant plasmid DNA induces protective immunity and prevents Japanese encephalitis in mice. *J. Virol.*, 2000, **74**, 4244–4252.
64. Kaur, R., Sachdeva, G. and Vratil, S., Plasmid DNA immunization against Japanese encephalitis virus: immunogenicity of membrane-anchored and secretory envelope protein. *J. Infect. Dis.*, 2002, **185**, 1–12.
65. Ashok, M. S. and Rangarajan, P. N., Protective efficacy of a plasmid DNA encoding Japanese encephalitis virus envelope protein fused to tissue plasminogen activator signal sequences: studies in a murine intracerebral virus challenge model. *Vaccine*, 2002, **20**, 1563–1570.
66. Appaiahgari, M. B. and Vratil, S., DNase-mediated inhibition of Japanese encephalitis virus replication in mouse brain. *Mol. Ther.*, 2007, **15**, 1593–1599.
67. Ramanathan, M. P. *et al.*, Coimmunization with an optimized IL15 plasmid adjuvant enhances humoral immunity via stimulating B cells induced by genetically engineered DNA vaccines expressing consensus JEV and WNV E DIII. *Vaccine*, 2009, **27**, 4370–4380.
68. Imoto, J. and Konishi, E., Needle-free jet injection of a mixture of Japanese encephalitis DNA and protein vaccines: a strategy to effectively enhance immunogenicity of the DNA vaccine in a murine model. *Viral Immunol.*, 2005, **18**, 205–212.
69. Ishikawa, T., Widman, D. G., Bourne, N., Konishi, E. and Mason, P. W., Construction and evaluation of a chimeric pseudoinfectious virus vaccine to prevent Japanese encephalitis. *Vaccine*, 2008, **26**, 2772–2781.
70. Solomon, T. *et al.*, Interferon alfa-2a in Japanese encephalitis: a randomised double-blind placebo-controlled trial. *Lancet*, 2003, **361**, 821–826.
71. Zhao, C., Denison, C., Huibregtse, J. M., Gygi, S. and Krug, R. M., Human ISG15 conjugation targets both IFN-induced and constitutively expressed proteins functioning in diverse cellular pathways. *Proc. Natl. Acad. Sci. USA*, 2005, **102**, 10200–10205.
72. Yuan, W. and Krug, R. M., Influenza B virus NS1 protein inhibits conjugation of the interferon (IFN)-induced ubiquitin-like ISG15 protein. *EMBO J.*, 2001, **20**, 362–371.
73. Loo, Y. M. *et al.*, Viral and therapeutic control of IFN-beta promoter stimulator 1 during hepatitis C virus infection. *Proc. Natl. Acad. Sci. USA*, 2006, **103**, 6001–6006.
74. Okumura, A., Lu, G., Pitha-Rowe, I. and Pitha, P. M., Innate antiviral response targets HIV-1 release by the induction of ubiquitin-like protein ISG15. *Proc. Natl. Acad. Sci. USA*, 2006, **103**, 1440–1445.
75. Kumar, R., Tripathi, P., Baranwal, M., Singh, S., Tripathi, S. and Banerjee, G., Randomized, controlled trial of oral ribavirin for Japanese encephalitis in children in Uttar Pradesh, India. *Clin. Infect. Dis.*, 2009, **48**, 400–406.
76. Swarup, V., Ghosh, J., Ghosh, S., Saxena, A. and Basu, A., Antiviral and anti-inflammatory effects of rosmarinic acid in an experimental murine model of Japanese encephalitis. *Antimicrob. Agents Chemother.*, 2007, **51**, 3367–3370.
77. Swarup, V., Ghosh, J., Mishra, M. K. and Basu, A., Novel strategy for treatment of Japanese encephalitis using arctigenin, a plant lignan. *J. Antimicrob. Chemother.*, 2008, **61**, 679–688.
78. Dutta, K., Ghosh, D. and Basu, A., Curcumin protects neuronal cells from Japanese encephalitis virus-mediated cell death and also inhibits infective viral particle formation by dysregulation of ubiquitin-proteasome system. *J. Neuroimmune. Pharmacol.*, 2009, **4**, 328–337.
79. Michaelis, M., Kleinschmidt, M. C., Doerr, H. W. and Cinatl Jr, J., Minocycline inhibits West Nile virus replication and apoptosis in human neuronal cells. *J. Antimicrob. Chemother.*, 2007, **60**, 981–986.
80. Mishra, M. K. and Basu, A., Minocycline neuroprotects, reduces microglial activation, inhibits caspase 3 induction, and viral replication following Japanese encephalitis. *J. Neurochem.*, 2008, **105**, 1582–1595.
81. Mishra, M. K., Ghosh, D., Duseja, R. and Basu, A., Antioxidant potential of Minocycline in Japanese encephalitis virus infection in murine neuroblastoma cells: correlation with membrane fluidity and cell death. *Neurochem. Int.*, 2009, **54**, 464–470.
82. Sebastian, L., Desai, A., Shampur, M. N., Perumal, Y., Sriram, D., Vasanthapuram, R., N-methylisatin-beta-thiosemicarbazone derivative (SCH 16) is an inhibitor of Japanese encephalitis virus infection *in vitro* and *in vivo*. *J. Virol.*, 2008, **5**, 64–75.
83. Wu, S. F., Lee, C. J., Liao, C. L., Dwek, R. A., Zitzmann, N. and Lin, Y. L., Antiviral effects of an iminosugar derivative on flavivirus infections. *J. Virol.*, 2002, **76**, 3596–3604.
84. Kumar, P., Lee, S. K., Shankar, P. and Manjunath, N., A single siRNA suppresses fatal encephalitis induced by two different flaviviruses. *PLoS Med.*, 2006, **3**, e96.