

A direct proof of completeness of squeezed odd-number states

S. Chaturvedi*

Department of Physics, University of Queensland

Brisbane, Australia

Abstract

A direct proof of the resolution of the identity in the odd sector of the Fock space in terms of squeezed number states $D(\xi)|2m+1\rangle$; $D(\xi) = \exp((\xi a^{\dagger 2} - \xi^* a^2)/2)$ is given. The proof entails evaluation of an integral involving Jacobi polynomials. This is achieved by the use of Racah identities.

PACS No. : 42.50 Dv 03.65 Fd

*Permanent address: School of Physics, University of Hyderabad, Hyderabad 500 046 **India**

The squeezed number states [1]

$$|\zeta; n \rangle = D(\xi)|n \rangle ; D(\xi) = \exp\left(\frac{1}{2}(\xi a^{\dagger 2} - \xi^* a^2)\right) , \quad (1)$$

regarded as a generalization of the squeezed vacuum [2] have been thoroughly investigated in the quantum optics literature for their non classical properties. From a group theoretic point of view [3] these may be thought of as generalized coherent states associated with the SU(1,1) group and it can be shown that the squeezed number states $|\zeta; 2n + 1 \rangle = D(\xi)|2n + 1 \rangle$ based on the odd Fock states furnish a resolution of the identity as follows

$$X = \frac{1}{2\pi} \int \frac{d^2\zeta}{(1 - |\zeta^2|)^2} |\zeta; 2n + 1 \rangle \langle \zeta; 2n + 1| = I_{odd} , \quad (2)$$

where ζ and ξ are related to each other in the following manner

$$\xi = |\xi|e^{-i\phi} ; \zeta = \tanh|\xi|e^{-i\phi} . \quad (3)$$

The operator I_{odd} in (2) denotes the unit operator in the odd sector of the Fock space

$$I_{odd} = \sum_{k=0}^{\infty} |2k + 1 \rangle \langle 2k + 1| , \quad (4)$$

and the integration in (2) is over the unit disc centered at the origin in the complex ζ -plane. The proof of this statement makes use of the Schur's Lemma and consists in demonstrating that the operator X in (2) commutes $D(\xi)$ for all ξ and hence must be equal to some constant c times the unit operator. (For the squeezed states based on even Fock states the constant c turns out to be infinite and hence one does not have a resolution of the identity similar to that in (2). Another elegant proof of (2) in which only the completeness of $D(\xi)|1 \rangle$ is made use of to derive the general result in (2) for any odd Fock state may be found in [4]. In this letter we show that when one attempts a direct proof of (2) one encounters an integral involving Jacobi polynomials and in order to complete the proof one must show that this integral is equal to one. We show that this is indeed so by calculating the integral in question by making use of the Racah identities [5] in the intermediate steps of the calculation.

The operator X in (2), on using the resolution of the identity in terms of the Fock states and the fact that the matrix elements $\langle 2p|D(\xi)|2n + 1 \rangle$ vanish, may be written as

$$X = \frac{1}{2\pi} \sum_{p,q=0}^{\infty} \int \frac{d^2\zeta}{(1-|\zeta^2|)^2} \langle 2p+1|\zeta; 2n+1 \rangle \langle \zeta; 2n+1|2q+1 \rangle |2p+1 \rangle \langle 2q+1| . \quad (5)$$

The explicit expressions for $\langle 2m+1|\zeta; 2n+1 \rangle$ are as follows. For $m \geq n$

$$\langle 2m+1|\zeta; 2n+1 \rangle = e^{-i(m-n)\phi} \sqrt{\frac{\Gamma(n+1)\Gamma(m+3/2)}{\Gamma(m+1)\Gamma(n+3/2)}} (|\zeta|)^{(m-n)} (1-|\zeta|^2)^{3/4} P_n^{(m-n,1/2)}(1-2|\zeta|^2) , \quad (6)$$

For $m \leq n$

$$\langle 2m+1|\zeta; 2n+1 \rangle = e^{-i(n-m)\phi} \sqrt{\frac{\Gamma(m+1)\Gamma(n+3/2)}{\Gamma(n+1)\Gamma(m+3/2)}} (-|\zeta|)^{(n-m)} (1-|\zeta|^2)^{3/4} P_m^{(n-m,1/2)}(1-2|\zeta|^2) . \quad (7)$$

Here $P_n^{(\alpha,\beta)}(x)$ denote the Jacobi Polynomials [6,7]. The expressions above can also be written in terms of the Gegenbauer polynomials $C_n^\alpha(x)$ or in terms of the associated Legendre polynomials $P_n^\alpha(x)$ using the following relations [6,7]

$$P_n^{(\alpha,1/2)}(x) = \frac{\Gamma(n+3/2)\Gamma(\alpha+1/2)}{\Gamma(1/2)\Gamma(\alpha+n+3/2)} \frac{1}{\sqrt{(1+x)/2}} C_{2n+1}^{(\alpha+1/2)}(\sqrt{(1+x)/2}) , \quad (8)$$

$$P_n^{(\alpha,1/2)}(x) = \frac{1}{2^\alpha} \frac{\Gamma(n+3/2)}{\Gamma(\alpha+n+3/2)} \frac{((1-x)/2)^{-\alpha/2}}{\sqrt{(1+x)/2}} P_{2n+\alpha+1}^\alpha(\sqrt{(1+x)/2}) . \quad (9)$$

Using the expressions in (6) and (7) in (5) and performing the integration over ϕ one obtains

$$X = \sum_{p=0}^{\infty} \frac{1}{2} \int_0^1 \frac{d|\zeta|^2}{(1-|\zeta^2|)^2} |\langle 2p+1|\zeta; 2n+1 \rangle|^2 |2p+1 \rangle \langle 2p+1| , \quad (10)$$

which may be compactly rewritten as

$$X = \sum_{p=0}^n \mathcal{I}_{n,p} |2p+1 \rangle \langle 2p+1| + \sum_{p=n+1}^{\infty} \mathcal{I}_{p,n} |2p+1 \rangle \langle 2p+1| , \quad (11)$$

where

$$\mathcal{I}_{p,n} = \frac{1}{2} \int_0^1 \frac{d|\zeta|^2}{(1-|\zeta^2|)^2} |\langle 2p+1|\zeta; 2n+1 \rangle|^2 ; p \geq n . \quad (12)$$

Thus to prove the resolution of of the identity in (2) we need to show that $\mathcal{I}_{p,n}$ is equal to one for all integer values of p and n with $p \geq n$. To show this we proceed as follows. Substituting the expression for $\langle 2p+1|\zeta; 2n+1 \rangle$ given by (6) into (12) and defining $x = |\zeta|^2$ we find that $\mathcal{I}_{p,n}$ can be written as

$$\mathcal{I}_{p,n} = \frac{1}{2} \frac{\Gamma(n+1)\Gamma(p+3/2)}{\Gamma(p+1)\Gamma(n+3/2)} \int_0^1 \frac{dx}{\sqrt{(1-x)}} x^{(p-n)} \left[P_n^{(p-n, 1/2)}(1-2x) \right]^2. \quad (13)$$

We now make use of the fact that the Jacobi polynomials $P_n^{(p-n, 1/2)}(1-2x)$ can be explicitly written out in two different ways [6,7]

$$P_n^{(p-n, 1/2)}(1-2x) = \sum_{l=0}^n \binom{n}{l} \binom{n+1/2}{n-l} (-1)^{(n-l)} x^{(n-l)} (1-x)^l, \quad (14)$$

$$P_n^{(p-n, 1/2)}(1-2x) = \frac{p!}{n! \Gamma(p+3/2)} \sum_{m=0}^n (-1)^m \binom{n}{m} \frac{\Gamma(p+m+3/2)}{(p-n-m)!} x^m. \quad (15)$$

Substituting the expression (14) for one of the Jacobi polynomials in the integral in (13) and the expression (15) for the other and carrying out the integral over x using

$$\int_0^1 dx x^{p+m-l} (1-x)^{l-1/2} = \frac{\Gamma(l+1/2)\Gamma(p+m-l+1)}{\Gamma(p+m+3/2)}, \quad (16)$$

one obtains

$$\mathcal{I}_{p,n} = \frac{1}{2} n! p! \sum_{l=0}^n \frac{(-1)^l \Gamma(l+1/2)}{l!(p-l)!(n-l)! \Gamma(l+3/2)} \sum_{m=0}^n \frac{(-1)^m (p+m-l)!}{m!(n-m)!(p-n+m)!}. \quad (17)$$

Changing m to $n-m$, (17) may be written as

$$\mathcal{I}_{p,n} = \frac{1}{2} n! p! \sum_{l=0}^n \frac{(-1)^l \Gamma(l+1/2)}{l!(p-l)!(n-l)! \Gamma(l+3/2)} \sum_{m=0}^n \frac{(-1)^m (p+n-l-m)!}{m!(n-m)!(p-m)!}. \quad (18)$$

Using the Racah identity

$$\sum_m \frac{(-1)^m (u-m)!}{m!(x-m)!(z-m)!} = \frac{(-1)^z (u-z)!(u-x)!(x+z-u)! \sin(\pi(x-u))}{x!z! \pi}, \quad (19)$$

with $u = p+n-l$, $x = p$, and $z = n$ to carry out the sum over m in (18) one gets

$$\mathcal{I}_{p,n} = \frac{1}{2} \sum_{l=0}^n \frac{(-1)^{n+l} \sin(\pi(l-n))}{(l+1/2) \pi l} = \frac{1}{2} \sum_{l=0}^n \frac{(-1)^l \sin(\pi l)}{(l+1/2) \pi l} . \quad (20)$$

The factor $\sin(\pi l)/\pi l$ ensures that only the $l = 0$ term survives yielding

$$\mathcal{I}_{p,n} = 1 . \quad (21)$$

This result, written in terms of Gegenbauer polynomials or in terms of associated Legendre functions, reads as follows

$$\int_0^1 \frac{dx}{x^2} (1-x^2)^{p-n} [C_{2n+1}^{(p-n+1/2)}(x)]^2 = \frac{p! \Gamma(p+3/2)}{n! \Gamma(n+3/2)} \left[\frac{\Gamma(1/2)}{\Gamma(p-n+1/2)} \right]^2 , \quad (22)$$

$$\int_0^1 \frac{dx}{x^2} [P_{p+n+1}^{p-n}]^2 = \frac{(2p+1)!}{(2n+1)!} . \quad (23)$$

Here $p \geq n$. The integrals (13), (22) and (23), to the best of our knowledge, are not listed in any of the standard tables of integrals.

Acknowledgements: I wish to thank Dr B. Bambah for bringing the Racah identities to my attention. I am grateful to the University of Queensland for the award of the University of Queensland Travel Grant and to the Physics department for hospitality.

References

1. F.A.M. de Oliveira, M.S.Kim, P.L.Knight and V.Bužek, Phys.Rev.**A41**, 2645 (1990).
2. H.P.Yuen, Phys.Rev.**A13**, 2226 (1976).
3. A.Perelemov, *Generalized coherent states and their applications*, (Springer, Berlin, 1986).
4. G.S.Agarwal and S.Chaturvedi, submitted to J.Phys **A**.
5. G. Racah, Phys.Rev.**61**, 186 (1942) ; ibid **62**,432 (1942).
6. M.A.Abramowitz and I.A.Stegun *Handbook of mathematical functions* , (Dover, N.Y., 1970)
7. I.S.Gradsteyn and I.M.Ryzhik *Tables of integrals, series and products*, (Academic Press, N.Y., 1965).