

Binary cloning vectors for efficient genetic transformation of rice

Surekha Katiyar-Agarwal, Avnish Kapoor and Anil Grover*

Department of Plant Molecular Biology, University of Delhi South Campus, Benito Juarez Road, Dhaula Kuan, New Delhi 110 021, India

The availability of effective vector systems is a prerequisite for genetic manipulation of plants through recombinant DNA technology. We report here construction of a series of binary vectors that have cauliflower mosaic virus 35S promoter-driven genes encoding either resistance to hygromycin or phosphinothricin for selection of the transformants, and high strength constitutive promoters of either *ubiquitin1* or *actin1* genes for efficient expression of the transgenes. The efficacy of the constructs is tested in stably transformed Pusa Basmati 1 rice plants through β -glucuronidase reporter gene activity. Availability of vectors with variable promoters and selectable marker genes provides flexibility in stacking two genes. The vectors constructed in this study are suitable for both particle gun and *Agrobacterium*-based transformation protocols.

RICE is the most important food crop. Of late, rice has emerged as a model crop for the analysis of genome and proteome of cereals^{1,2}. There is a great deal of progress in stable genetic transformation of rice plants. Compared to conventional breeding, genetic engineering is a relatively faster means for varietal improvement that allows trans-

fer of genes from within as well as outside the primary gene pool³. Genetic transformation of rice through use of *Agrobacterium* is a favoured approach as it enables transfer of DNA with defined ends, minimal rearrangement, integration of a small number of copies of the gene and more importantly, the possibility that even large segments of DNA can be efficiently transferred⁴⁻⁸.

Success achieved in genetic improvement of most of the agronomic characteristics in rice (as well as in other cereals and dicot plants) is limited. Despite the massive support given to rice biotechnology programme in recent years, there is not much success in improving resistance against biotic and abiotic stresses that cause significant damage to rice yield^{3,9}. An important reason for this failure is that mostly the stress resistance-related traits are mediated by a number of biochemical reactions/physiological processes, and methods for co-integration of multiple transgenes into the plant genome are not well established.

Earlier studies have shown that *Agrobacterium*-mediated gene pyramiding can possibly be brought about by either sequential transformation or co-transformation with multiple T-DNAs¹⁰⁻¹². The multiple T-DNAs can be delivered to plant cells either from a mixture of strains (mixture method) or from a single strain which contains more than one T-DNA (single-strain method)¹⁰. However, the important parameters that must be noted while designing a co-transformation experiment are that (a) there should be different promoters available to drive different transgenes in the same host, so as to avoid homology-based recombination and gene silencing¹³, and (b) there should be different selectable markers available to select each transgene separately. We have used two different promoters obtained from *ubiquitin1* and *actin1*

*For correspondence. (e-mail: grover_anil@hotmail.com)

1. Esson, V. S., *Curr. Opin. Pulm. Med.*, 1977, **3**, 198–202.
2. Yeung, C. M., *Am. Rev. Resp. Dis.*, 1977, **116**, 1023.
3. Nylander, L. A. and Dement, J. M., *Am. J. Ind. Med.*, 1993, **5**, 619–647.
4. Demers, P. A. *et al.*, *Scand. J. Work, Environ. Health*, 1995, **3**, 179–190.
5. Wolf, J., Schmezer, P., Fengel, D., Schroeder, H. G., Scheithauer, H. and Woeste, P., *Acta Oto-Laryngol., Suppl.*, 1998, **535**, 1–16.
6. Palus, J., Dziuba touska, E. and Rydzyski, K., *Mutat. Res.*, 1999, **444**, 61–74.
7. Vaughan, T. L., Stewart, P. A., Teschke, K., Lynch, C. F., Swanson, G. M., Lyon, J. L. and Berwick, M., *Occup. Environ. Med.*, 2000, **6**, 376–384.
8. Fenech, M., *Prog. Clin. Biol. Res.*, 1990, **340**, 195–206.
9. Lamberti, L., Bigatti Pponzetto, P. and Ardito, G., *Mutat. Res.*, 1983, **120**, 193–199.
10. Holmstrom, M., Granstrand, P., Nylander, L. A. and Rosin, G., *Am. J. Ind. Med.*, 1995, **2**, 207–220.
11. Hebert, F. A., Hessel, P. A., Melenka, L. S., Yoshida, K. and Nakaza, M., *Arch. Environ. Health*, 1994, **4**, 465–470.
12. Norrish, A. E., Beasley, R., Hodgkinson, E. J. and Pearce, N., *N. Z. Med. J.*, 1992, **105**, 185–187.
13. Palus, J., Dziuba touska, E. and Rydzyski, K., *Mutat. Res.*, 1998, **408**, 91–101.
14. Palus, J., Dziuba touska, E. and Rydzyski, K., *Acta Biochim. Pol.*, 1998, **45**, 605–610.
15. Liou, S. H., Cheng, S. Y., Lao, F. M. and Yang, J. L., *Am. J. Ind. Med.*, 1996, **3**, 293–299.
16. Peace, B. E., Livingston, G., Silberstein, E. B. and Loper, J. C., *Mutat. Res.*, 1999, **430**, 109–119.
17. Jiang, Z. *et al.*, *Biomed. Environ. Sci.*, 1997, **4**, 396–401.
18. Maciejewska, A., Wojtczak, J., Bielichowska cybula, G., Domanska, A., Dutkiewicz, J. and Molocznik, A., *Med. PR.*, 1993, **3**, 277–288.

ACKNOWLEDGEMENTS. We thank Prof. C. V. Ramakrishnan and Prof. N. R. Rajagopal for technical advice, suggestions and discussions. We are grateful to M. Revathi and P. Suprabha for assistance. This study was financially supported by the Madras Medical Mission.

Received 31 August 2001; revised accepted 11 January 2002

Binary cloning vectors for efficient genetic transformation of rice

Surekha Katiyar-Agarwal, Avnish Kapoor and Anil Grover*

Department of Plant Molecular Biology, University of Delhi South Campus, Benito Juarez Road, Dhaula Kuan, New Delhi 110 021, India

The availability of effective vector systems is a prerequisite for genetic manipulation of plants through recombinant DNA technology. We report here construction of a series of binary vectors that have cauliflower mosaic virus 35S promoter-driven genes encoding either resistance to hygromycin or phosphinothricin for selection of the transformants, and high strength constitutive promoters of either *ubiquitin1* or *actin1* genes for efficient expression of the transgenes. The efficacy of the constructs is tested in stably transformed Pusa Basmati 1 rice plants through β -glucuronidase reporter gene activity. Availability of vectors with variable promoters and selectable marker genes provides flexibility in stacking two genes. The vectors constructed in this study are suitable for both particle gun and *Agrobacterium*-based transformation protocols.

RICE is the most important food crop. Of late, rice has emerged as a model crop for the analysis of genome and proteome of cereals^{1,2}. There is a great deal of progress in stable genetic transformation of rice plants. Compared to conventional breeding, genetic engineering is a relatively faster means for varietal improvement that allows trans-

fer of genes from within as well as outside the primary gene pool³. Genetic transformation of rice through use of *Agrobacterium* is a favoured approach as it enables transfer of DNA with defined ends, minimal rearrangement, integration of a small number of copies of the gene and more importantly, the possibility that even large segments of DNA can be efficiently transferred⁴⁻⁸.

Success achieved in genetic improvement of most of the agronomic characteristics in rice (as well as in other cereals and dicot plants) is limited. Despite the massive support given to rice biotechnology programme in recent years, there is not much success in improving resistance against biotic and abiotic stresses that cause significant damage to rice yield^{3,9}. An important reason for this failure is that mostly the stress resistance-related traits are mediated by a number of biochemical reactions/physiological processes, and methods for co-integration of multiple transgenes into the plant genome are not well established.

Earlier studies have shown that *Agrobacterium*-mediated gene pyramiding can possibly be brought about by either sequential transformation or co-transformation with multiple T-DNAs¹⁰⁻¹². The multiple T-DNAs can be delivered to plant cells either from a mixture of strains (mixture method) or from a single strain which contains more than one T-DNA (single-strain method)¹⁰. However, the important parameters that must be noted while designing a co-transformation experiment are that (a) there should be different promoters available to drive different transgenes in the same host, so as to avoid homology-based recombination and gene silencing¹³, and (b) there should be different selectable markers available to select each transgene separately. We have used two different promoters obtained from *ubiquitin1* and *actin1*

*For correspondence. (e-mail: grover_anil@hotmail.com)

terminator and CaMV35S promoter-*hyg^r*-*polyA* terminator. The recombinant clones were checked using *Bam*HI, *Eco*RI, *Hind*III, *Kpn*I, *Pst*I, *Sac*I, *Sall*, *Sma*I, *Xba*I, *Xho*I, *Eco*RV and *Sac*II-*Xma*I. These vectors were referred to as pAH-1 and pAH-2 (depending upon the orientation of the *actin* promoter cassette).

5. *Xho*I-*Cla*I-cut 1.6 kb fragment from pActcas containing *act1* promoter (along with *act1* first exon and first intron), MCS and *nos* gene terminator was treated with T₄ DNA polymerase and subsequently ligated to blunted *Eco*RI-*Hind*III-cut pCAMBIA3300, resulting in construction of vectors with *act1* promoter-MCS-*nos* terminator and CaMV35S promoter-*bar*-*polyA* terminator. The recombinant clones were checked using *Bam*HI, *Eco*RI, *Hind*III, *Kpn*I, *Pst*I, *Sac*I, *Sall*, *Sma*I, *Xba*I, *Xho*I and *Eco*RV. These vectors were referred to as pAB-1 and pAB-2 (depending upon the orientation of the *actin* promoter cassette).

6. *Afl*III-*Nco*I-cut 2.1 kb fragment from pCAMBIA1201 was blunted using T₄ DNA polymerase and subsequently ligated with *Sma*I-cut pAH-1, resulting in the construction of a vector with *act1* promoter-*gus*-*nos* terminator and CaMV35S promoter-*hyg^r*-*polyA* terminator. The recombinant clones were checked using *Eco*RV and *Xho*I. This vector is referred to as pAH-1/*gus*.

The restriction maps and other details of the above vectors are provided in Figure 1 (refs 15, 23–25). The pUH, pAH-1 and pAH-2 plasmids have the *hyg^r* gene of pCAMBIA1300 for selection, whereas pUB, pAB-1 and pAB-2 have the *bar* gene from pCAMBIA3300 for

selection. Both *hyg^r* (ref. 26) and *bar²⁷* selectable markers are routinely employed for rice transformation. Both these genes are under the control of CaMV35S promoter which has been shown to work reasonably well with the rice system²⁸. The maize *ubiquitin1* and the rice *actin1* gene promoters have a high strength of expression and are often used for rice transformation^{16,17,29}. Besides the promoter and selectable markers, the other useful feature of these vectors is the availability of several unique restriction sites in the MCS for cloning purpose. The presence of unique *Sma*I site in both the vectors provides flexibility in cloning any gene fragment after blunting through suitable enzymatic reaction. As the foundation vectors used here belong to pCAMBIA series, the cloning vectors designed in this study inherit several useful features. These include kanamycin-based selection in bacteria, high copy number plasmid replication, Kozak sequences for improved expression, minimal extraneous DNA sequences and stability of plasmids under non-selection conditions¹⁵. Importantly, we have the *act1* promoter-MCS-*nos* terminator and CaMV35S promoter-selection marker-*polyA* terminator configurations in same (pAH-1, pAB-1) and different (pAH-2, pAB-2) directions in our vectors. This should enable cloning of the transgene in either way to avoid the possible influence of one promoter on the activity of the other promoter in the vector cassette.

To further check the efficiency of the vectors designed, we subcloned *gus* gene in the MCS of both pUH and pAH-1 to obtain pUH/*gus* and pAH-1/*gus* (Figure 2 a). These constructs were employed to transform mature




Figure 2. a, Diagrammatic representation of binary vectors containing *gus* as a reporter gene used for rice transformation. *gus*, gene for β -glucuronidase with a catalase intron. b, Histochemical staining for GUS expression in (1), control callus; (2), callus transformed with pUH/*gus*; (3), leaf segment of control plant; (4), leaf segment of transgenic plant transformed with pUH/*gus*.

seed-derived calli of Pusa Basmati 1 rice cultivar. Further, subculturing of the transformed calli on selection medium resulted in healthy, vigorously proliferating white daughter calli, indicating that they possess cells with hygromycin resistance gene. These hygromycin-selected calli showed positive reaction for *gus* expression when histochemically stained, while untransformed calli did not show the reaction (Figure 2 b). The calli transformed with pUH/*gus* showed blue colour within a minute, while pAH-1/*gus* showed GUS reaction after nearly 2 h, indicating that the *ubiquitin1* promoter more strongly regulates the expression than the *actin1* promoter. Similar observations have also been reported earlier¹⁷. Moreover when leaf segments from transgenic rice plants (pUH/*gus*) were tested for GUS expression, intense blue colour was noted indicating stable integration of the transgene (Figure 2 b). Subsequent to this work, we have subcloned several agronomically-useful genes in pUH and have noted the expression of the transgenes in stably-transformed rice lines³⁰.

The advantage of having vectors with two different selectable markers and two different promoters in different orientations as achieved in this study provides flexibility in gene-stacking experiments. We will further aim at bringing in additional promoters and selectable markers so that more than two genes can also be pyramided. While the present study is limited to rice, it is to be noted that *ubiquitin1* and *actin1* promoters are shown to work in several other monocot crops³¹. Likewise, there are several reports indicating that the resistance to hygromycin and phosphinothricin is effective in selection of transformed tissues, generally in monocots³²⁻³⁶. We believe that the vectors constructed in this study would have general applicability in genetic transformation of monocots.

1. Goff, S. A., *Curr. Opin. Plant Biol.*, 1999, **2**, 86–89.
2. Dubey, H. and Grover, A., *Curr. Sci.*, 2001, **80**, 262–269.
3. Grover, A. and Minhas, D., *Proc. Indian Natl. Sci. Acad.*, 2000, **B66**, 13–32.
4. Hellens, R., Mullineaux, P. and Klee, H., *Trends Plant Sci.*, 2000, **5**, 446–451.
5. Shibata, D. and Liu, Y-G., *ibid*, 354–357.
6. Hiei, Y., Ohta, S., Komari, T. and Kumashiro, T., *Plant J.*, 1994, **6**, 271–282.
7. Dong, J., Teng, W., Buchholz, W. G. and Hall, T. C., *Mol. Breed.*, 1996, **2**, 267–276.
8. Rashid, H., Yokoi, S., Toriyama, K. and Hinata, K., *Plant Cell Rep.*, 1996, **15**, 727–730.
9. Mohanty, H. K., Mallik, S. and Grover, A., *Curr. Sci.*, 2000, **78**, 132–137.
10. Komari, T., Hiei, Y., Saito, Y., Murai, N. and Kumashiro, T., *Plant J.*, 1996, **10**, 165–174.

11. DeNeve, M., DeBuck, S., Jacobs, A., Van Montagu, M. and Depicker, A., *ibid*, 1997, **11**, 15–29.
12. DeBuck, S., Jacobs, A., Van Montagu, M. and Depicker, A., *Mol. Plant Microb. Intact.*, 1998, **11**, 449–457.
13. Meyer, P., *Gene Silencing in Higher Plants and Related Phenomena in Other Eukaryotes*, Springer-Verlag, Heidelberg, 1995, p. 232.
14. Sambrook, J. and Russell, D. W., *Molecular Cloning, A Laboratory Manual*, Cold Spring Harbor Laboratory Press, New York, 2001.
15. McElroy, D., Chamberlain, D. A., Moon, E. and Wilson, K. J., *Mol. Breed.*, 1995, **1**, 27–37.
16. McElroy, D., Blowers, A. D., Jenes, B. and Wu, R., *Mol. Gen. Genet.*, 1991, **231**, 150–160.
17. Li, Z., Upadhaya, N. M., Meena, S., Gibbs, A. J. and Waterhouse, P. M., *Mol. Breed.*, 1997, **3**, 1–14.
18. Wang, M. B., Upadhaya, N. M., Brettell, R. I. S. and Waterhouse, P. M., *J. Genet. Breed.*, 1997, **51**, 325–334.
19. Mandel, M. and Higa, A., *J. Mol. Biol.*, 1970, **53**, 159–162.
20. An, G., Ebert, P. R., Mitra, A. and Ha, S. B., in *Plant Molecular Biology Manual A3* (eds Gelvin, S. B. and Shilperoot, R. A.), Kluwer Academic Press, Dordrecht, 1988, pp. 1–19.
21. Minhas, D., Ph D thesis, University of Delhi, 1999.
22. Jefferson, R. A., *Plant Mol. Biol. Rep.*, 1987, **5**, 387–405.
23. Depicker, A., Stachel, S., Dhaese, P., Zambryski, P. and Goodman, H. M., *J. Mol. Appl. Genet.*, 1982, **1**, 561–573.
24. McElroy, D., Zhang, W., Cao, J. and Wu, R., *Plant Cell*, 1990, **1**, 163–171.
25. Christensen, A. H., Sharrock, R. A. and Quail, P. H., *Plant Mol. Biol.*, 1992, **18**, 675–689.
26. Aldemita, R. R. and Hodges, T. K., *Planta*, 1996, **199**, 612–617.
27. Rathore, K. S., Chowdhury, V. K. and Hodges, T. K., *Plant Mol. Biol.*, 1993, **21**, 871–884.
28. Mitsuhara, I. *et al.*, *Plant Cell Physiol.*, 1996, **37**, 49–59.
29. Christensen, A. H. and Quail, P. H., *Transgenic Res.*, 1996, **5**, 213–218.
30. Grover, A. *et al.*, in Proceedings of the Third International Rice Genetics Symposium, IRRI, Manila, the Philippines, 2000 (in press).
31. Schledzewski, T. and Mendel, R. R., *Transgenic Res.*, 1994, **3**, 249–255.
32. Horn, M. E., Shillito, R. D., Conger, B. V. and Harms, C. T., *Plant Cell Rep.*, 1988, **7**, 469–472.
33. Fromm, M. E., Taylor, L. P. and Walbot, V., *Nature*, 1986, **319**, 791–793.
34. Vasil, V., Castillo, A. M., Fromm, M. E. and Vasil, I. K., *Bio/Tech.*, 1992, **10**, 667–674.
35. Cassas, A. M. *et al.*, *Proc. Natl. Acad. Sci. USA*, 1993, **90**, 11212–11216.
36. Sukhapinda, K., Kozuch, M. E., Rubin-Wilson, B., Ainley, W. M. and Merlo, D. J., *Plant Cell Rep.*, 1993, **13**, 63–68.

ACKNOWLEDGEMENTS. We thank Dr Richard Jefferson, CAMBIA, Canberra, Australia, Dr Narayanan Upadhayaya, Dr Zhongyi Li, CSIRO Division of Plant Industry, Canberra, Australia and Dr Peter Quail, University of California, Berkeley, USA, for pCAMBIA, pLZUbicas and pActcas plasmids. This research was supported by Department of Biotechnology, Government of India (BT/AB/05/02/91-Vol IV). S.K.-A. thanks the Council of Scientific and Industrial Research for fellowship.

Received 2 August 2001; revised accepted 15 January 2002