

Indian J Med Res 126, November 2007, pp 440-446

Effect of pretreatment with chromium picolinate on haematological parameters during dengue virus infection in mice

Richa Shrivastava*, R. Nagar**, G.A. Ravishankar*, R.K. Upreti⁺ & U.C. Chaturvedi⁺

***Department of Microbiology, K.G. Medical University & ⁺Industrial Toxicology Research Institute Lucknow, India*

Received April 26, 2007

Background & objectives: Dengue virus (DV) has caused severe epidemics of dengue fever (DF) and dengue haemorrhagic fever (DHF) and is endemic all over India. We have earlier reported that exposure of mice to hexavalent chromium [Cr(VI)] compounds increased the severity of dengue virus infection. Trivalent chromium picolinate (CrP) is used worldwide as micronutrient and nutritional supplement. The present study was therefore, carried out to investigate the effects of CrP on various haematological parameters during DV infection of mice.

Methods: The Swiss Albino mice were inoculated with dengue virus (1000 LD₅₀, intracerebrally) and fed with chromium picolinate (CrP) in drinking water (100 and 250 mg/l) for 24 wk. Peripheral blood leucocytes and other haematological parameters, and spleens were studied on days 4 and 8 after virus inoculations and the findings were compared with those given only CrP and the normal control age matched mice.

Results: CrP in drinking water for 24 wk had no significant effects on peripheral blood cells of mice. On the other hand, there was significant decrease in different haematological parameters following inoculation of normal mice with DV. In CrP fed mice the effects of DV infection were abolished on most of the haematological parameters.

Interpretation & conclusions: The findings of present study showed that the adverse effects of DV infection, specially on platelets and leucocytes, were abrogated by pretreatment of mice with CrP. The therapeutic utility of CrP in viral infections including dengue needs to be studied in depth.

Key words Blood cells - chromium picolinate - dengue virus - haematological indices - leucopenia - thrombocytopenia

Dengue virus (DV) produces a benign self-limiting illness, the dengue fever (DF) or a life-threatening serious illness, the dengue haemorrhagic fever (DHF). Dengue virus is endemic in India with frequent

epidemics of DF and DHF¹⁻⁶. The dengue disease is characterized by leucopenia, thrombocytopenia, bone marrow suppression, increased haematocrit value and increased capillary permeability, which may result in

*Present address: Plant Cell Biotechnology Department, Central Food Technological Research Institute (CFTRI), Mysore 570020, India

haemorrhage and shock⁷⁻⁹. Immunopathological mechanisms appear to be responsible for the pathogenesis of DHF. A rapid increase in the levels of cytokines and chemical mediators apparently induced by a unique cytokine, cytotoxic factor (CF), plays a key role in inducing plasma leakage, shock and haemorrhagic manifestations^{10,11}.

Pollution of environment by chromium (Cr) is common in industrialized areas as a consequence of effluent discharge from tanneries and other industries, which include metal plating, manufacturing industries and ferrochrome production. This poses serious problem for environmental quality. In aqueous environments chromium has two oxidation states: hexavalent chromium (Cr-VI) and trivalent chromium (Cr-III). Cr (VI) compounds are generally soluble over a wide pH range and have been shown to exert toxic and carcinogenic effects in humans and experimental animals, also induce DNA damage such as DNA single-strand breaks and DNA-protein crosslinks *in vivo* and in cultured cells. On the other hand, Cr (III) compounds are mainly non-toxic¹²⁻¹⁵.

DV infection is endemic all over India, so is the occupational and non occupational exposure to Cr (VI). The effect of chromium compromises the immune response of the host^{16,17}. It is, therefore, possible that the chromium toxicity may adversely affect the disease process during DV infection. In our earlier study we have reported that Cr (VI) in the form of potassium dichromate drinking led to reduction in lymphocytes, haemoglobin and the haematocrit values while the granulocyte, monocyte and platelet counts were increased. The most significant finding of these experiments was that the DV-induced reduction of platelet counts was cancelled in Cr (VI)-fed mice¹⁸.

Chromium picolinate monohydrate, (CrP) is a synthetic Cr (III) compound that has widespread use as a nutritional supplement. It promotes a variety of beneficial health effects including weight loss, serum cholesterol reduction, treatment of diabetes, and increased muscle mass¹⁹. CrP administered to mice in the diet for 3 months up to the maximal possible dose (approximately 20,000-30,000 times that of the average daily human supplement 200 - 400 µg) to show any adverse effects²⁰⁻²³. No effects of CrP exposure on clinical chemistry or haematology parameters, gross lesions, or microscopic findings have been observed²⁴⁻²⁶. We, therefore, undertook this study to investigate the effects of dengue virus infection in mice fed with non

toxic trivalent chromium in the form of chromium picolinate (CrP).

Material & Methods

Animals: The study was carried out on Swiss Albino mice weighing 25-30 g, aged 6-8 wk obtained from the animal breeding facility of the Industrial Toxicology Research Centre (ITRC), Lucknow. Study protocol was approved by the Animal Ethical Committee of the ITRC, Lucknow. Mice were maintained on pellet diet supplied by local supplier.

Dengue virus: Dengue type 2 virus, strain P23085 (received from National Institute of Virology, Pune) was used in the form of infected infant mouse brain suspension. Mice were given light ether anaesthesia and then inoculated with 1000 LD₅₀ of DV intracerebrally (ic) in volume of 30 µl.

CrP treatment: Mice were divided in three groups of 18 animals each. One group was given *ad libitum* drinking water containing 100 mg/l chromium picolinate, second group was given drinking water containing 250 mg/l chromium picolinate and third group (control) was given plain water to drink. After 24 wk of CrP drinking, 12 animals in each group were inoculated i.c. with dengue virus in doses of 1000 LD₅₀. The mice were sacrificed by cervical dislocation the 4th (6 animals, clinical signs of illness appear) and 8th (6 animals, paralysis of hind limbs set in) day post virus inoculation (pi).

Haematological study: Animals were sacrificed and 1.5 ml blood was collected from the jugular vein in tubes with 0.1 ml freshly prepared 10 per cent K₃ EDTA solution as anticoagulant. The samples were shaken gently. The following estimations were performed immediately with the help of a fully automatic haematology analyzer (Bayer; Technicon H1*E, Germany): total leucocyte count (TLC), differential leucocyte count (DLC), total RBC count, haematocrit (Hct), haemoglobin (Hb), mean corpuscular volume (MCV) and platelet counts.

Preparation of spleen cell culture: The spleen cells were teased out in cold minimal essential medium (MEM, Hi-Media, Mumbai) and a single cell suspension was prepared. The viable nucleated cell count as ascertained by trypan blue dye exclusion test²⁷ was more than 95 per cent.

*Spleen cell proliferation assay*²⁸: The cell suspension of normal mice spleen was adjusted to 2x10⁶ cells/ml

in Eagle's MEM supplemented with 10 per cent foetal calf serum. Cell cultures were set up in 96 wells U-bottom microtitre plates in triplicate. Each well contained 0.2 ml cell suspension with or without 5 µg/ml concanavalin A (Con A). The cultures were incubated for 72 h at 37°C in presence of 5 per cent CO₂. Five hours before the termination of the cultures, 20 µl of MTT3-(4,5-dimethyl thiazole-2-yl)-2,5-diphenyl tetrazolium bromide (5mg/ml PBS, phosphate buffer saline) was added to each well. Cells were harvested and supernatant discarded and 200 µl of dimethyl sulfoxide was added; readings were recorded after 10 min at 570 nm using 630 nm as reference OD. All the chemicals used were procured from Sigma, USA.

Preparation of spleen homogenate: Spleens were washed free of blood, homogenized in a tissue homogenizer giving a 10 per cent solution (w/v) in chilled PBS (pH 7.2). The homogenate was centrifuged in the cold at 8000 g for 10 min and clear supernatant was stored in small aliquots at -20°C. Normal mouse spleen homogenates were prepared similarly and used as control.

Cytotoxicity test: Cytotoxic factor (CF) is present in the homogenates prepared from the DV-infected mouse spleen and has the capacity to kill normal mouse spleen cells in one hour time. The cytotoxic activity of the spleen homogenate isolated from DV infected mice was tested using normal mouse spleen cells as target. The test was carried out in 96 wells plates, using 0.1 ml volume of each preparation. The target single cell suspension of normal mouse spleen was prepared. Then the cell viability was measured by MTT assay. The MTT assay was done in triplicate and the mean value of % of non-viable cell as obtained in repeated experiments are presented²⁴. All procedures were carried out at 4°C in ice bath.

Statistical analysis: Two-way analysis of variance (Two-way ANOVA) was done by Systat Software, Inc. (USA), to compare the mean values of the outcome variables considering chromium exposure and dengue virus infection to animals as independent variables. Prior to this homogeneity of variance between the treated groups were ascertained. The two-group comparison was done by calculating least significant difference at 5 per cent level of significance.

Results

Effect CrP feeding on blood cells during DV infection:

Effects on leucocytes: In normal control mice the total

leucocyte count was $11.5 \pm 4.6 \times 10^3$ /cubic millimeter (Cu mm) with 87 per cent lymphocytes and about 9 per cent granulocytes. By giving CrP (100mg/l and 250mg/l) in drinking water for 24 wk the total as well as differential leucocyte counts were not affected significantly (Table I). In the normal mice DV infection caused a significant reduction in the total leucocyte at day 4 ($P < 0.05$) and 8 ($P < 0.01$) compared to uninfected mice. The predominant cells in the blood remained lymphocytes in all the groups (Table I).

Effects on red blood cells: Feeding CrP had no effect on the red blood cell count, haemoglobin and MCV, where as haematocrit concentration was significantly increased at 250 mg/l dose of CrP ($P < 0.05$) (Table II). Normal mice inoculated with DV showed reduced total RBC counts ($P < 0.01$), haematocrit ($P < 0.05$) and haemoglobin ($P < 0.05$) at day 8 of DV inoculation (Table II). In CrP fed mice DV inoculation caused no significant change in these indices.

Effect on platelet count: The platelet count in the blood of normal control mice was $406 \pm 81 \times 10^6$ /Cu mm, whereas after 100 mg/l CrP drinking it significantly increased to $484 \pm 51 \times 10^6$ /Cu mm ($P < 0.05$) but with 250 mg/l CrP feeding count was slightly decreased to $366 \pm 52 \times 10^6$ /µl compared to normal controls. When normal control mice were inoculated with the dengue virus, a significant reduction in the platelet count was seen on days 4 and 8 being $179 \pm 57 \times 10^6$ ($P < 0.001$) and $312 \pm 51 \times 10^6$ ($P < 0.01$) respectively. In CrP fed mice DV inoculation caused no significant change (Fig. 1).




Fig. 1. Effects of CrP feeding (24 wk) on platelet count during dengue virus infection. CrP, chromium picolinate; DV, dengue virus; 4, 8, days after the virus infection compared to controls. (n=6 in each group). Values are mean \pm SD.

Table I. Leucocytes count in cromium picolinate fed mice after dengue virus (DV) infection

CrP treated groups	TLC ($10^3/\text{Cu mm}$)			Lymphocytes (%)			Granulocytes (%)			Monocytes (%)		
	Control	DV4d	DV8d	Control	DV4d	DV8d	Control	DV4d	DV8d	Control	DV4d	DV8d
Normal	11.5 ± 4.6	8.0 ± 3.2*	5.8 ± 2.3**	87 ± 1.9	82 ± 6.0	80 ± 3.0	9.5 ± 1.8	14 ± 5.6	15 ± 1.4**	3.3 ± 0.3	4.0 ± 0.8	4.5 ± 0.9
100ppm CrP	10.8 ± 1.5	10.5 ± 2.1	9.7 ± 3.0	90 ± 3.5	88 ± 3.5	82 ± 0.5	8.0 ± 2.9	8.0 ± 2.7	12 ± 1.0*	2.3 ± 0.6	3.9 ± 0.8	6.0 ± 0.6
250ppm CrP	11.2 ± 2.3	10.8 ± 0.3	9.3 ± 1.2	86 ± 6.0	85 ± 1.0	87 ± 9.0	10 ± 3.0	11 ± 1.0	12 ± 1.7*	3.2 ± 1.5	4.1 ± 3.2	3.3 ± 1.1

Control, normal mice; DV4d, 4th day after infection DV; DV8d, 8th day after infection with DV; TLC, total leucocyte count
 Values are mean ± SD of 6 animals in each group
 $P^* < 0.05$; $** < 0.001$ compared to normal

Table II. RBC indices in cromium picolinate fed mice after dengue virus (DV) infection

CrP treated groups	RBC ($10^6/\text{Cu mm}$)			Hb (g/l)			HCt (%)			MCV (fl)		
	Control	DV4d	DV8d	Control	DV4d	DV8d	Control	DV4d	DV8d	Control	DV4d	DV8d
Normal	8.8 ± 0.5	6.2 ± 0.3	5.9 ± 0.5**	14 ± 1.9	11 ± 0.8	9.0 ± 1.2*	29 ± 3.8	21 ± 2.0	23 ± 2.3*	36 ± 1.7	37 ± 2.3	34 ± 1.8
100ppm CrP	8.0 ± 0.5	7.3 ± 1.2	7.7 ± 1.0	14 ± 0.9	15 ± 0.9	13 ± 2.5	29 ± 3.3	26 ± 5.6	31 ± 4.5	36 ± 2.2	36 ± 2.7	37 ± 1.2
250ppm CrP	8.7 ± 0.6	7.7 ± 1.7	7.6 ± 0.5	16 ± 0.9	14 ± 3.4	14 ± 0.9	37 ± 1.0*	31 ± 0.7	30 ± 2.5	39 ± 2.7	35 ± 3.5	37 ± 1.7

Control, normal mice; DV4d, 4th day after infection with DV; DV8d, 8th day after infection with DV; RBC, red blood cell; MCV, mean corpuscular volume;
 HCt, haematocrit; Hb, haemoglobin
 $P^* < 0.01$ $** < 0.01$ compared to control
 $^+ P < 0.05$ compared to control of normal unfed group
 Value are mean ± SD of 6 animals in each group

Effect on weight of spleen: The weight of the spleen in the control normal mice was 477 ± 59 mg/100 g body weight. Significant reduction in the weight of the spleen with 100 mg/l CrP was observed [388 ± 55 mg/100 g ($P \leq 0.05$) body weight] whereas with 50 mg/l CrP it was increased significantly ($P \leq 0.05$) being (587 ± 79 mg/100g). The weight of the spleen of normal mice inoculated with DV was 439 ± 30 and 340 ± 25 mg/100 g body weight on day 4 and 8 post-inoculation, respectively (Figs 2 & 3). When CrP treated mice were inoculated with DV, only with 250 mg/l CrP dose a slight protection was observed against DV induced reduction in spleen weight in comparison to DV alone.

Effects on production of cytotoxic factor (CF): The homogenate from DV-infected mice spleen produced 13 to 25 per cent cytotoxicity (Fig. 4). On the other hand, no cytotoxic activity was seen in the spleen




Fig. 2. Effects of CrP on spleen weight (in mg/100g body weight) during dengue virus infection. CrP, chromium picolinate; DV 4, 8 d, dengue virus 4, 8, days after the virus infection compared to controls. Values are mean \pm SD of n=6 mice in each group.




Fig. 3. Cytotoxic activity of the homogenates of spleens obtained from various groups of mice. Values are mean \pm SD of n=6 mice in each group.




Fig. 4. Effects of CrP feeding on mitogenic stimulation of spleen cells during dengue virus infection. Value are mean \pm SD (n=6).

homogenates obtained from CrP fed mice in comparison to normal control mice. A significant decrease ($P < 0.001$) in cytotoxic activity of spleen homogenate of DV infected mice was observed after pretreatment with CrP at both doses.

Effect on mitogenic stimulation: When normal mice spleen cells were treated with mitogen the observed stimulation was 0.171 ± 0.05 . When normal mice were inoculated with DV, maximum proliferation occurred at the fourth day (0.375 ± 0.030) and it was suppressed significantly on day 8 ($P < 0.05$). On the other hand, after 24 wk of drinking of 100 mg/l CrP, spleen cell proliferation in mice increased by two-fold of normal mice spleen cells. When CrP (100 mg/l) fed mice were inoculated with DV, on day 8 of infection there was no DV induced suppression in mitogenic stimulation in comparison to the DV alone mice group.

Discussion

The present study showed abrogation of DV-induced adverse effects on different haematological parameters in mice after feeding CrP for 24 wk. The characteristic leucopenia, thrombocytopenia and increased haematocrit are common findings in clinical dengue disease and together with combination of clinical picture are used as markers for early diagnosis of dengue infection^{7,9,29-31}. Constant haematological abnormalities occurring in severe dengue disease include bone marrow suppression³². La Russa & Innis³³ have reviewed data from experimental dengue infections of volunteers and bone marrow studies from patients with severe dengue virus infection which showed that marrow suppression evolves rapidly through several phases and recovers. The early blast

cells and the differentiated haematopoietic elements are abortively infected, killed and eliminated by phagocytosis by specialized marrow dendritic cells. Cytokine production by virus-infected stromal cells is also altered³³. The combined effect of these changes is transient bone marrow suppression resulting in leucopenia and thrombocytopenia.

Our present findings are supported by those on human subjects that the haematocrit, haemoglobin, RBC count, MCV, MCH, MCHC, red blood cell distribution width, platelet count, and mean platelet volume are within normal clinical ranges and are unchanged by chromium picolinate supplementation for 12 wk²⁸. Recent studies^{35,36} demonstrate that chromium picolinate supplementation significantly improves insulin sensitivity and glucose control in subjects with type 2 diabetes and also in obese subjects with polycystic ovary syndrome.

In this study haemoglobin concentration and the haematocrit value decreased day 8 after DV inoculation in normal mice. Pimpan & Prasert⁷ reported that in patients with DHF haematocrit value is increased initially but later on it decreased. On the other hand, DV-induced fall in RBC count, haemoglobin as well as on haematocrit % was abolished in CrP fed mice. During DV infection of mice a significant reduction in the weight of the spleen occurred, which was associated with a sharp decline in the proportion of T-lymphocytes and macrophages in the spleen²⁷. CrP feeding had no effect on the spleen weight of normal mice which was similar to the findings reported by Rhodes *et al*²⁶. When CrP fed mice were inoculated with DV, a significant protection against reduction of spleen weight was observed on day 8 of the virus infection. Further, the production of DV-induced cytokine, CF was significantly reduced in CrP fed mice.

These findings showed that the mitogenic response of splenic lymphocytes to Con A was increased in CrP fed mice similar to the findings reported in cows³⁷ and in pigs fed supplemental chromium³⁸. Organic chromium supplements are known to increase immune function in animals. The exact mechanism by which chromium picolinate enhances the immune system is not known. However, chromium supplementation has been shown to reduce serum cortisol levels. Glucocorticoids, which include cortisol, are known to suppress the immune system³⁹.

Pollution of environment by Cr (VI) poses serious problem due to its toxicity. On the other hand, Cr (III)

compounds are mainly non-toxic^{16,17}. We had earlier shown that Cr (VI) drinking led to reduction of most of the haematological parameters (except platelet counts) enhancing the severity of dengue illness in mice⁴⁰.

In conclusion, abrogation of DV-induced adverse effects on the haematological parameters by feeding CrP, indicates a need for in depth investigation before it is used as supportive therapy in dengue and other viral infections.

Acknowledgment

Authors thank the Director, National Institute of Virology, Pune for providing dengue type 2 virus strain P23085. The first author (RS) acknowledges the financial support received in form of Research Associate from the Indian Council of Medical Research, New Delhi. The last author (UCC) was the former Professor of Microbiology, K.G. Medical College, Lucknow.

References

1. Agarwal R, Kapoor S, Nagar R, Misra A, Tandon R, Mathur A, *et al*. A clinical study of the patients with dengue haemorrhagic fever during the epidemic of 1996 at Lucknow, India. *Southeast Asian J Trop Med Public Health* 1999; 30 : 735-40.
2. Shah I, Deshpande GC, Tardeja PN. Outbreak of dengue in Mumbai and predictive markers for dengue shock syndrome. *J Trop Pediatr* 2004; 50 : 301-5.
3. Dash PK, Saxena P, Abhyankar A, Bhargava R, Jana AM. Emergence of dengue virus type-3 in northern India. *Southeast Asian J Trop Med Public Health* 2005; 36 : 370-7.
4. Singh NP, Jhamb R, Agarwal SK, Gaiha M, Dewan R, Daga MK, *et al*. The 2003 outbreak of dengue fever in Delhi, India. *Southeast Asian J Trop Med Public Health* 2005; 36 : 1174-8.
5. Dash PK, Parida MM, Saxena P, Abhyankar A, Singh CP, Tewari KN, *et al*. Reemergence of dengue virus type-3 (subtype-III) in India: implications for increased incidence of DHF & DSS. *Virol J* 2006; 6 : 55.
6. Gupta E, Dar L, Kapoor G, Broor S. The changing epidemiology of dengue in Delhi, India. *Virol J* 2006; 5 : 92.
7. Pimpan P, Prasert T. *Clinical laboratory investigations. Monograph on dengue/dengue haemorrhagic fever*, WHO SEARO No.22. New Delhi: WHO SEARO; 1990. p. 62-71.
8. Chaturvedi UC, Shrivastava R. Dengue haemorrhagic fever: A global challenge. *Indian J Med Microbiol* 2004; 22 : 5-6.
9. Chaturvedi UC. The curse of dengue. *Indian J Med Res* 2006; 124 : 467-70.
10. Chaturvedi UC, Raghupathy R, Pacsa AS, Elbishbishi EA, Agarwal R, Nagar R, *et al*. Shift from a Th1-type response to Th2-type in dengue hemorrhagic fever. *Curr Sci* 1999; 76 : 63-9.
11. Chaturvedi UC, Agarwal R, Elbishbishi EA, Mustafa AS. Cytokine cascade in dengue hemorrhagic fever: Implications for pathogenesis. *FEMS Immunol Med Microbiol* 2000; 28 : 183-8.

12. Hamilton JW, Wetterhahn KE. Chromium. In: Seiler HG, Sigel H, editors. *Handbook on toxicity of inorganic compounds*. New York: Marcel Dekker, Inc;1988. p. 239-50.
13. Losi ME, Amrhein C, Frankenberger WTJ. Environmental biochemistry of chromium. *Rev Environ Contam Toxicol* 1994; *136* : 91-131.
14. Kerger BD, Paustenbach DJ, Corbett GE, Finley BL. Absorption and elimination of trivalent and hexavalent chromium in humans following ingestion of a bolus dose in drinking water. *Toxicol Appl Pharmacol* 1996; *141* : 145-58.
15. U.S. Environmental Protection Agency. *Health effects assessment for hexavalent chromium*. Prepared by the Office of Health and Environmental Assessment, Environmental Criteria; 1984. p. 37, EPA/540/1-86-019 Cincinnati, Washington DC, updated 1998.
16. Shrivastava R, Upreti RK, Seth PK, Chaturvedi UC. Effects of chromium on the immune system. *FEMS Immunol Med Microbiol* 2002; *34* : 1-7.
17. Shrivastava R, Upreti RK, Chaturvedi UC. Various cells of immune system and intestine differ in their capacity to reduce hexavalent chromium. *FEMS Immunol Med Microbiol* 2003; *38* : 65-70.
18. Shrivastava R, Srivastava S, Upreti RK, Chaturvedi UC. Effects of dengue virus infection on peripheral blood cells of mice exposed to hexavalent chromium with drinking water. *Indian J Med Res* 2005; *122* : 111-9.
19. Campbell WW, Joseph LJ, Anderson RA, Davey SL, Hinton J, Evans WJ. Effects of resistive training and chromium picolinate on body composition and skeletal muscle size in older women. *Int J Sport Nutr Exerc Metab* 2002; *12* : 125-35.
20. Anderson RA, Bryden NA, Polansky MM, Deuster PA. Exercise effects on chromium excretion of trained and untrained men consuming a constant diet. *J App Physiol* 1988; *64* : 249-52.
21. Anderson RA, Bryden NA, Polansky MM, Gautschi K. Dietary chromium effects on tissue chromium concentrations and chromium absorption in rats. *J Trace Elem Exp Med* 1996; *9* : 11-25.
22. Anderson RA, Bryden NA, Polansky MM. Lack of toxicity of chromium chloride and chromium picolinate in rats. *J Am Coll Nut* 1997; *16* : 273-9.
23. Bagchi D, Stohs SJ, Downs BW, Bagchi M, Preuss HG. Cytotoxicity and oxidative mechanisms of different forms of chromium. *Toxicology* 2002; *180* : 5-22.
24. Pittler MH, Stevinson C, Ernst E. Chromium picolinate for reducing body weight: meta-analysis of randomized trials. *Int J Obes Relat Metab Disord* 2003; *27* : 522-9.
25. Vincent JB. The potential value and toxicity of chromium picolinate as a nutritional supplement, weight loss agent and muscle development agent. *Sports Med* 2003; *33* : 213-30.
26. Rhodes MC, Hebert CD, Herbert RA, Morinello EJ, Roycroft JH, Travlos GS, et al. Absence of toxic effects in F344/N rats and B6C3F1 mice following subchronic administration of chromium picolinate monohydrate. *Food Chem Toxicol* 2005; *43* : 21-9.
27. Chaturvedi UC, Nagar R, Mathur A. Effect of dengue virus infection on Fc-receptor functions of mouse macrophages. *J Gen Virol* 1983; *64* : 2399-407.
28. Mosmann T. Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *J Immunol Methods* 1983; *65* : 55-63.
29. Kularatne SA, Gawarammana IB, Kumarasiri PR. Epidemiology, clinical features, laboratory investigations and early diagnosis of dengue fever in adults: a descriptive study in Sri Lanka. *Southeast Asian J Trop Med Public Health* 2005; *36* : 686-92.
30. Rigau-Pérez JG. Severe dengue: the need for new case definitions. *Lancet Infect Dis* 2006; *6* : 297-302.
31. Chaturvedi UC, Dhawan R, Mukerjee R. Immunosuppression and cytotoxicity of dengue infection in the mouse model. In: Gubler DJ, Kuno G, editors. *Dengue and dengue hemorrhagic fever*, Wallingford, Oxon, UK: CAB International Press; 1997. p. 289-309.
32. Srichaikul T, Nimmannitya S. Haematology in dengue and dengue haemorrhagic fever. *Baillieres Best Pract Res Clin Haematol* 2000; *13* : 261-76.
33. La Russa VF, Innis BL. Mechanisms of dengue virus-induced bone marrow suppression. *Baillieres Clin Haematol* 1995; *8* : 249-70.
34. Campbell WW, Beard JL, Joseph LJ, Davey SL, Evans WJ. Chromium picolinate supplementation and resistive training by older men: effects on iron-status and hematologic indexes. *Am J Clin Nutr* 1997; *66* : 944-9.
35. Martin J, Wang ZQ, Zhang XH, Wachtel D, Volaufova J, Matthews DE, et al. Chromium picolinate supplementation attenuates body weight gain and increases insulin sensitivity in subjects with type 2 diabetes. *Diabetes Care* 2006; *29* : 1826-32.
36. Lydic ML, McNurlan M, Bembo S, Mitchell L, Komaroff E, Gelato M. Chromium picolinate improves insulin sensitivity in obese subjects with polycystic ovary syndrome. *Fertil Steril* 2006; *86* : 243-6.
37. Mowat DN, Chang X. Chromium and immunity of stressed feeder calves. In: *Proceedings of 28th Annual Conference for feed manufacturers*, April 27-28, 1992 College Inn, Guelph., Ontario, Ottawa, Canada Feed Industry Association (Abstract).
38. Heugten E, Spears JW. Immune response and growth of stressed weanling pigs fed diets supplemented with organic and inorganic forms of chromium. *J Anim Sci* 1997; *75* : 409-16.
39. Muck A, Guyre P, Holbrook N. Physiological function of glucorticoids in stress and their relation to pharmacological actions. *Endocr Rev* 1984; *5* : 25-40.
40. Shrivastava R, Upreti RK, Chaturvedi UC. Effects of dengue virus infection on the spleen cells of mice given hexavalent chromium with drinking. *Toxico Mech Meth* 2005; *15* : 1-7.

Reprint requests: Prof. U.C. Chaturvedi, 201-Annapurna Apartments, No. 1, Bishop Rocky Street, Faizabad Road
Lucknow 226007, India
e-mail: uchaturvedi@yahoo.com