

SCIENTIFIC REPORTS



OPEN

Bell's Nonlocality Can be Detected by the Violation of Einstein-Podolsky-Rosen Steering Inequality

Received: 26 September 2016

Accepted: 16 November 2016

Published: 14 December 2016

Jing-Ling Chen^{1,2}, Changliang Ren³, Changbo Chen⁴, Xiang-Jun Ye^{5,6} & Arun Kumar Pati⁷

Recently quantum nonlocality has been classified into three distinct types: quantum entanglement, Einstein-Podolsky-Rosen steering, and Bell's nonlocality. Among which, Bell's nonlocality is the strongest type. Bell's nonlocality for quantum states is usually detected by violation of some Bell's inequalities, such as Clauser-Horne-Shimony-Holt inequality for two qubits. Steering is a manifestation of nonlocality intermediate between entanglement and Bell's nonlocality. This peculiar feature has led to a curious quantum phenomenon, the one-way Einstein-Podolsky-Rosen steering. The one-way steering was an important open question presented in 2007, and positively answered in 2014 by Bowles *et al.*, who presented a simple class of one-way steerable states in a two-qubit system with at least thirteen projective measurements. The inspiring result for the first time theoretically confirms quantum nonlocality can be fundamentally asymmetric. Here, we propose another curious quantum phenomenon: Bell nonlocal states can be constructed from some steerable states. This novel finding not only offers a distinctive way to study Bell's nonlocality without Bell's inequality but with steering inequality, but also may avoid locality loophole in Bell's tests and make Bell's nonlocality easier for demonstration. Furthermore, a nine-setting steering inequality has also been presented for developing more efficient one-way steering and detecting some Bell nonlocal states.

In 1935, the famous Einstein, Podolsky and Rosen (EPR) paper indicated that quantum mechanics is in conflict with the notion of locality and reality¹. If local realism is correct, then quantum mechanics cannot be considered as a complete theory to describe physical reality. Immediately after the publication of the EPR paper, Schrödinger made a response by conjuring two important notions, namely, the quantum *entanglement* and the quantum *steering*. According to Schrödinger, quantum entanglement is “the characteristic trait of quantum mechanics” that distinguishes quantum theory from classical theory². The notion of “steering” is closely related to the statement of “spooky action at a distance”, which Einstein was disturbed all the time. EPR steering reflects such a “spooky action” feature that manipulating one object seemingly affects another instantaneously, even it is far away.

Different to Schrödinger's response, in 1964, Bell proposed an inequality for local hidden variable (LHV) models³. The violation of Bell's inequality by quantum entangled states implies Bell's nonlocality. This is well-known as Bell's theorem, which has established what quantum theory can tell us about the fundamental features of *Nature*, and been widely regarded as “the most profound discovery of science”⁴. Until now, the fundamental theorem has achieved ubiquitous applications in different quantum information tasks, such as quantum key distribution⁵, communication complexity⁶, and random number generation⁷.

¹Theoretical Physics Division, Chern Institute of Mathematics, Nankai University, Tianjin 300071, People's Republic of China. ²Centre for Quantum Technologies, National University of Singapore, 3 Science Drive 2, Singapore 117543, Singapore. ³Center for Nanofabrication and System Integration, Chongqing Institute of Green and Intelligent Technology, Chinese Academy of Sciences, Chongqing 400714, People's Republic of China. ⁴Chongqing Key Laboratory of Automated Reasoning and Cognition, Chongqing Institute of Green and Intelligent Technology, Chinese Academy of Sciences, Chongqing 400714, People's Republic of China. ⁵Key Laboratory of Quantum Information, University of Science and Technology of China, University of Science and Technology of China, Hefei 230026, People's Republic of China. ⁶Synergetic Innovation Center of Quantum Information and Quantum Physics, University of Science and Technology of China, Hefei 230026, People's Republic of China. ⁷Quantum Information and Computation Group, Harish-Chandra Research Institute, Chhatnag Road, Jhansi, Allahabad 211019, India. Correspondence and requests for materials should be addressed to J.-L.C. (email: chenjl@nankai.edu.cn) or C.R. (email: renchangliang@cigit.ac.cn) or A.K.P. (email: akpati@hri.res.in)




Figure 1. Hierarchical structure of quantum nonlocality. Bell's nonlocality is the strongest type of quantum nonlocality. If a state possesses EPR steerability or Bell's nonlocality, then the state must be entangled. EPR steering is a form of nonlocality intermediate between entanglement and Bell nonlocality.

Unlike quantum entanglement and Bell's nonlocality, the research field of quantum steering has been sterile till 2007, when Wiseman, Jones, and Doherty⁸ reformulated the idea and placed it firmly on a rigorous ground. Since then EPR steering has gained a very rapid development in both theories^{9–16} and experiments^{17–26}. Most research topics as well as research approaches in the field of Bell's nonlocality have been transplanted similarly to the field of EPR steering. For instance, steering inequalities have been proposed to reveal the EPR steerability of quantum states, very similar to the violation of Bell's inequalities reveals Bell's nonlocality.

According to ref. 8, entanglement, EPR steering and Bell's nonlocality are called by a joint name as “quantum nonlocality”, which has an interesting hierarchical structure: quantum entanglement is a superset of steering, and Bell's nonlocality is a subset of steering. However, among the three types of quantum nonlocality, only steering can possess a curious feature of “one-way quantumness”. Suppose Alice and Bob share a pair of two-qubit state, it is not hard to imagine that if Alice entangles with Bob, then Bob must also entangle with Alice. Such a symmetric feature holds for both entanglement and Bell nonlocality. However, the situation is dramatically changed when one turns to a novel kind of quantum nonlocality in the middle of entanglement and Bell nonlocality, the EPR steering. It may happen that for some asymmetric bipartite quantum states, Alice can steer Bob but Bob can never steer Alice. This distinguished feature would be useful for some one-way quantum information tasks, such as quantum cryptography. The “one-way EPR steering” or “asymmetric EPR steering” is an important “open question” first proposed by Wiseman *et al.* in ref. 8. Very recently, the question has been answered by Bowles *et al.*¹⁵, who presented a simple class of one-way steerable states in a two-qubit system with at least 13 projective measurements (a linear 14-setting steering inequality was given explicitly in the work). The inspiring result for the first time theoretically confirms quantum nonlocality can be fundamentally asymmetric. Later on, Bowles *et al.* investigated the one-way steering problem by presenting a sufficient criterion (being a nonlinear criterion) for guaranteeing that a two-qubit state is unsteerable²⁷.

In this work, we focus on another curious quantum phenomenon raised by steering: Bell nonlocal states can be constructed from some EPR steerable states. Explicitly we present a theorem, showing that for any two-qubit state τ , if its corresponding state ρ is EPR steerable, then the state τ must be Bell nonlocal. Bell's nonlocality of the quantum state τ can be detected indirectly by the violation of steering inequality for the quantum state ρ . The novel result not only pinpoints a deep connection between EPR steering and Bell's nonlocality, but also sheds a new light to avoid locality loophole in Bell's tests and make Bell's nonlocality easier for demonstration. In addition, we also present a 9-setting linear steering inequality for developing more efficient one-way steering and detecting some Bell nonlocal states. We find that the new steering inequality can actually improve the result of ref. 15 by detecting the one-way steering with fewer measurement settings but with larger quantum violations, which would be helpful for the experimenters.

Results

Bell's Nonlocal states can be constructed from EPR steerable states. It is well-known that quantum nonlocality possesses an interesting hierarchical structure (see Fig. 1). EPR steering is a weaker nonlocality in comparison to Bell's nonlocality. Here we would like to pinpoint a curious quantum phenomenon directly connecting these two different types of nonlocality. We find that Bell's nonlocal states can be constructed from some EPR steerable states, which indicates that Bell's nonlocality can be detected indirectly through EPR steering (see Fig. 2), and offers a distinctive way to study Bell's nonlocality. The result can be expressed as the following theorem.

Theorem 1: For any two-qubit state τ_{AB} shared by Alice and Bob, define another state

$$\rho_{AB} = \mu \tau_{AB} + (1 - \mu) \tau'_{AB}, \quad (1)$$




Figure 2. Illustration of detecting Bell's nonlocality through EPR steering. If a state ρ_{AB} violates a steering inequality, then it implies that ρ_{AB} possesses the EPR steerability. Traditionally, Bell's nonlocality of the two-qubit state τ_{AB} is revealed by violations of Bell's inequality. Based on Theorem 1, Bell's nonlocality of the state τ_{AB} can be detected through EPR steerability of the state ρ_{AB} , and the relation between ρ_{AB} and τ_{AB} is given in Eq. (1).

with $\tau'_{AB} = \tau_A \otimes 1/2$, $\tau_A = \text{tr}_B[\tau_{AB}] = \text{tr}_B[\rho_{AB}]$ being the reduced density matrix at Alice's side, and $\mu = \frac{1}{\sqrt{3}}$. If ρ_{AB} is EPR steerable, then τ_{AB} is Bell nonlocal.

Proof. The implication of the theorem is that, the EPR steerability of the state ρ_{AB} determines Bell's nonlocality of the state τ_{AB} . Namely, the nonexistence of local hidden state (LHS) model for ρ_{AB} implies the nonexistence of LHV model for τ_{AB} . We shall prove the theorem by proving its converse negative proposition: if the state τ_{AB} has a LHV model description, then the state ρ_{AB} has a LHS model description.

Suppose τ_{AB} has a LHV model description, then by definition for any projective measurements A for Alice and B for Bob, one always has the following relation

$$P(a, b|A, B, \tau_{AB}) = \sum_{\xi} P(a|A, \xi)P(b|B, \xi)P_{\xi}. \tag{2}$$

Here $P(a, b|A, B, \tau_{AB})$ is the joint probability, quantum mechanically it is computed as $P(a, b|A, B, \tau_{AB}) = \text{tr}[(\hat{\Pi}_a^{\hat{n}_A} \otimes \hat{\Pi}_b^{\hat{n}_B})\tau_{AB}]$, $\hat{\Pi}_a^{\hat{n}_A}$ is the projective measurement along the \hat{n}_A -direction with measurement outcome a for Alice, $\hat{\Pi}_b^{\hat{n}_B}$ is the projective measurement along the \hat{n}_B -direction with measurement outcome b for Bob (with $a, b=0, 1$), $P(a|A, \xi)$, $P(b|B, \xi)$ and P_{ξ} denote some (positive, normalized) probability distributions.

Let the measurement settings at Bob's side be picked out as x, y, z . In this situation, Bob's projectors are $\hat{\Pi}_b^x, \hat{\Pi}_b^y, \hat{\Pi}_b^z$, respectively. Since the state τ_{AB} has a LHV model description, based on Eq. (2) we explicitly have (with $\hat{n} = x, y, z$)

$$\begin{aligned} P(a, 0|A, \hat{n}, \tau_{AB}) &= \sum_{\xi} P(a|A, \xi)P(0|\hat{n}, \xi)P_{\xi}, \\ P(a, 1|A, \hat{n}, \tau_{AB}) &= \sum_{\xi} P(a|A, \xi)P(1|\hat{n}, \xi)P_{\xi}. \end{aligned} \tag{3}$$

We now turn to study the EPR steerability of ρ_{AB} . After Alice performs the projective measurement on her qubit, the state ρ_{AB} collapses to Bob's conditional states (unnormalized) as

$$\tilde{\rho}_a^{\hat{n}_A} = \text{tr}_A[(\hat{\Pi}_a^{\hat{n}_A} \otimes \mathbb{1})\rho_{AB}], \quad a = 0, 1. \tag{4}$$

To prove that there exists a LHS model for ρ_{AB} is equivalent to proving that, for any measurement $\hat{\Pi}_a^{\hat{n}_A}$ and outcome a , one can always find a hidden state ensemble $\{\varphi_{\xi} \rho_{\xi}\}$ and the conditional probabilities $\varphi(a|\hat{n}, \xi)$, such that the relation

$$\tilde{\rho}_a^{\hat{n}_A} = \sum_{\xi} \wp(a|\hat{n}_A, \xi) \wp_{\xi} \rho_{\xi} \tag{5}$$

is always satisfied. Here ξ 's are the local hidden variables, ρ_{ξ} 's are the hidden states, \wp_{ξ} and $\wp(a|\hat{n}_A, \xi)$ are probabilities satisfying $\sum_{\xi} \wp_{\xi} = 1$ and $\sum_a \wp(a|\hat{n}_A, \xi) = 1$. If there exist some specific measurement settings of Alice, such that Eq. (5) cannot be satisfied, then one must conclude that the state ρ_{AB} is steerable (in the sense of Alice steers Bob's particle).

Suppose there is a LHS model description for ρ_{AB} , then it implies that, for Eq. (5) one can always find the solutions of $\{\wp(a|\hat{n}_A, \xi), \wp_{\xi}, \rho_{\xi}\}$ if Eq. (3) is valid. The solutions are given as follows:

$$\begin{aligned} \wp(a|\hat{n}_A, \xi) &= P(a|A, \xi), \quad \wp_{\xi} = P_{\xi}, \\ \rho_{\xi} &= \frac{\mathbb{1} + \vec{\sigma} \cdot \vec{r}_{\xi}}{2}, \end{aligned} \tag{6}$$

where $\mathbb{1}$ is the 2×2 identity matrix, $\vec{\sigma} = (\sigma_x, \sigma_y, \sigma_z)$ is the vector of the Pauli matrices, and the hidden state ρ_{ξ} has been parameterized in the Bloch-vector form, with

$$\vec{r}_{\xi} = \mu(2P(0|x, \xi) - 1, 2P(0|y, \xi) - 1, 2P(0|z, \xi) - 1), \tag{7}$$

which is the Bloch vector for density matrix of a qubit. It can be checked that $|\vec{r}_{\xi}| \leq 1$, and this ensures ρ_{ξ} being a density matrix.

By substituting Eq. (6) into Eq. (5), we obtain

$$\tilde{\rho}_a^{\hat{n}_A} = \sum_{\xi} P(a|A, \xi) P_{\xi} \frac{\mathbb{1} + \vec{\sigma} \cdot \vec{r}_{\xi}}{2}. \tag{8}$$

To prove the theorem is to verify the relation (8) is always satisfied if Eq. (3) is valid. The verification can be found in **Methods**.

Remark 1.— In Eq. (7), by requiring the condition $|\vec{r}_{\xi}| \leq 1$ be valid for any probabilities $P(0|x, \xi), P(0|y, \xi), P(0|z, \xi) \in [0, 1]$, in general one can have $\mu \in [0, 1/\sqrt{3}]$. Generally, Theorem 1 is valid for any $\mu \in [0, 1/\sqrt{3}]$. In the theorem we have chosen the parameter μ as its maximal value $1/\sqrt{3}$, because the state τ_{AB} is convexed with a separable state τ'_{AB} , the larger value of μ , the easier to detect the EPR steerability.

In the following, we provide two examples for the theorem, showing that Bell's nonlocality of quantum states can be detected indirectly by the violations of some steering inequalities.

Example 1.— For example, let us detect Bell's nonlocality of the maximally entangled state (with $\tau_{AB} = |\Psi\rangle\langle\Psi|$)

$$|\Psi\rangle = \frac{1}{\sqrt{2}}(|00\rangle + |11\rangle) \tag{9}$$

without Bell's inequality. Based on the theorem, it is equivalent to detect the EPR steerability of the following two-qubit state

$$\rho_{AB} = \frac{1}{\sqrt{3}}|\Psi\rangle\langle\Psi| + \left(1 - \frac{1}{\sqrt{3}}\right)\tau_A \otimes \frac{\mathbb{1}}{2}, \tag{10}$$

with $\tau_A = \mathbb{1}/2$. The state (10) is nothing but the Werner state²⁸ with the visibility equals to $1/\sqrt{3}$, its steerability can be tested by using the steering inequality proposed in ref. 17 as

$$\mathcal{S}_N = \frac{1}{N} \sum_{k=1}^N \langle A_k \bar{\sigma}_k^B \rangle \leq C_N \tag{11}$$

with $N=6$. Here \mathcal{S}_N is the steering parameter for N measurement settings, and C_N is the classical bound, with $C_6 = (1 + \sqrt{5})/6 \simeq 0.5393$. The maximal quantum violation of the steering inequality is $\mathcal{S}_6^{\max} = 1/\sqrt{3} \simeq 0.5774$, which beats the classical bound.

Remark 2.— In a two-qubit system, Bell's nonlocality is usually detected by quantum violation of the Clause-Horne-Shimony-Holt inequality²⁹. Bell's nonlocality is the strongest type of nonlocality, due to this reason Bell-test experiments have encountered both the locality loophole and the detection loophole for a very long time³⁰. As a weaker nonlocality, EPR steering naturally escapes from the locality loophole and is correspondingly easier to be demonstrated without the detection loophole^{19,20}, as stated in ref. 17: "because the degree of correlation required for EPR steering is smaller than that for violation of a Bell inequality, it should be correspondingly easier to demonstrate steering of qubits without making the fair-sampling assumption [i.e., closing the detection loophole]". Indeed, the steerability of the Werner state has been experimentally detected in ref. 17 by the steering inequality (11). Our result shows that the EPR steerability of the state ρ_{AB} determines Bell's nonlocality of the state τ_{AB} , thus may shed a new light to realize a loophole-free Bell-test experiment through the violation of steering inequality.

Example 2.— The theorem naturally provides a steering-based criterion for Bell's nonlocality, which is expressed as follows: given an EPR steerable two-qubit state ρ_{AB} , if the matrix

$$\tau_{AB} = \sqrt{3} \rho_{AB} - (\sqrt{3} - 1) \tau'_{AB}, \quad (12)$$

is a two-qubit density matrix, then τ_{AB} is Bell nonlocal.

Let us consider a two-qubit state ρ_{AB} in the following form

$$\rho_{AB} = \frac{1}{4} \left(\mathbb{1} \otimes \mathbb{1} + \beta \sigma_3 \otimes \mathbb{1} + \gamma \mathbb{1} \otimes \sigma_3 - \alpha \sum_{k=1}^3 \sigma_k \otimes \sigma_k \right). \quad (13)$$

By substituting the state ρ_{AB} as in Eq. (13) into Eq. (12), then one obtains

$$\tau_{AB} = \frac{1}{4} \left(\mathbb{1} \otimes \mathbb{1} + \beta' \sigma_3 \otimes \mathbb{1} + \gamma' \mathbb{1} \otimes \sigma_3 - \alpha' \sum_{k=1}^3 \sigma_k \otimes \sigma_k \right), \quad (14)$$

with

$$\beta' = \beta, \quad \gamma' = \sqrt{3} \gamma, \quad \alpha' = \sqrt{3} \alpha. \quad (15)$$

It is worth to mention that the steering inequality (11) is applicable to show Bell's nonlocality of τ_{AB} for some parameters α', β', γ' . Here we would like to show that the similar task can be done by other new steering inequalities. In the following, we present a 9-setting linear steering inequality as

$$\sum_{i=1}^9 \sum_{j=1}^3 s_{ij} \langle ab \rangle_{ij} + \sum_{i=1}^9 s_i^A \langle a \rangle_i + \sum_{j=1}^3 s_j^B \langle b \rangle_j \leq L, \quad (16)$$

here for convenient we have used the same notations as in ref. 15 (where $(\sigma_1, \sigma_2, \sigma_3)$ is equivalent to $(\sigma_x, \sigma_y, \sigma_z)$). The inequality are characterized by matrices $\{\mathbf{S}, \mathbf{S}^A, \mathbf{S}^B\}$ with real coefficients s_{ij}, s_i^A , and s_j^B , and the local bound is $L = 1$ (see Supplementary Materials). The steering inequality (16) may have other particular application for improving the result ref. 15 by developing more efficient one-way steering, which we shall address in the coming section. But now we use it to detect Bell's nonlocality.

For example, let $\alpha' = 0.96, \beta' = -1/5, \gamma' = 1/6$, ones finds that τ_{AB} is a two-qubit state, and the steering inequality (16) is violated by the state ρ_{AB} (with the violation value 1.0064), hence the Bell's nonlocality of state τ_{AB} can be revealed in this way indirectly by the steerability of the state ρ_{AB} .

More efficient one-way EPR steering. Under local unitary transformation (LUT), any two-qubit state can be written in the following form ref. 31

$$\rho_{AB} = \frac{1}{4} \left(\mathbb{1} \otimes \mathbb{1} + \beta \vec{\sigma} \cdot \hat{u} \otimes \mathbb{1} + \gamma \mathbb{1} \otimes \vec{\sigma} \cdot \hat{v} + \sum_{k=1}^3 t_k \sigma_k \otimes \sigma_k \right), \quad (17)$$

with β, γ, t_k being the real coefficients, and \hat{u}, \hat{v} the unit vectors. Obviously, under LUT, the state ρ_{AB} is said to be symmetric if and only if $\beta = \gamma$ and $\hat{u} = \hat{v}$. Let one consider a simple situation with $t_1 = t_2 = t_3 = -\alpha$, and $\hat{u} = \hat{v} = (0, 0, 1)$, then he obtains the two-qubit state ρ_{AB} as in Eq. (13). In such a case, if ρ_{AB} is a one-way steerable state, then one must have $\beta \neq \gamma$.

In ref. 15, the authors have chosen $\beta = \frac{2(1-\alpha)}{5}, \gamma = -\frac{3(1-\alpha)}{5}$ and used the SDP program to numerically prove that the state ρ_{AB} is a one-way steerable state (with at least 13 projective measurements): for $\alpha \leq 1/2$, the state ρ_{AB} is unsteerable from Bob to Alice, while for $\alpha \gtrsim 0.4983$ the state is steerable from Alice to Bob when Alice performs 14 projective measurements. An explicit 14-setting steering inequality has been also proposed to conform the one-way steerability, although for $\alpha = 1/2$, the quantum violation is tiny (only 1.0004). The inspiring result for the first time confirms that the nonlocality can be fundamentally asymmetric. However, the tiny inequality violation as well as the 14 measurement settings give rise to the difficulty in experimental detection. To advance the study of unidirectional quantum steering, here we present a more efficient class of one-way steerable states by choosing

$$\beta = \frac{4\alpha(1-\alpha)}{3}, \quad \gamma = -2\alpha(1-\alpha), \quad (18)$$

with $\alpha \in [0, 1]$. The state $\rho_{AB}(\alpha)$ is entangled for $\alpha > 0.3279$. With the help of the SDP program, we found that in the range $0.4846 \leq \alpha \leq 1/2$, the state $\rho(\alpha)$ is one-way steerable within 10-setting measurements, thus this is more efficient than the previous result in ref. 15 (For the detail derivation of more efficient one-way EPR steering see Supplementary Materials). Furthermore, we can extract an explicit 9-setting steering inequalities (16) based on the SDP program. It can be verified directly that, for the state $\rho_{AB}(1/2)$, the quantum violation of 9-setting inequality (16) is $\frac{119}{116} \simeq 1.0258 > 1$, hence demonstrating steering from Alice to Bob. Compared to the previous result¹⁵, the amount of violation is much larger but achieved with fewer measurements. To our knowledge, we do not know whether the quantum violation by inequality (16) could be observed with current quantum technology. However, we believe that this result would be interesting and helpful for both theoretical and experimental physicists.

Discussion

In this work, we have presented a theorem showing that Bell nonlocal states can be constructed from some EPR steerable states. This result not only offers a novel and distinctive way to study Bell's nonlocality with the violation of steering inequality, but also may avoid locality loophole in Bell's tests and make Bell's nonlocality easier for demonstration. An interesting and inverse problem is whether one can construct some steerable states τ_{AB} from some Bell nonlocal state ρ_{AB} , because Bell's nonlocality has been researched more deeply in theoretical aspect, so that people can conveniently study steering via known criteria of Bell's nonlocality. Furthermore, an explicit 9-setting linear steering inequality has also been presented for detecting some Bell nonlocal states and developing more efficient one-way steering. This result allows one to observe one-way EPR steering with fewer measurement setting but with larger quantum violations. We hope experimental progress in this direction could be made in the near future.

Methods

Verification of equation (8). Let us calculate the left-hand side of Eq. (8). One has

$$\begin{aligned} \tilde{\rho}_a^{\hat{n}_A} &= \text{tr}_A[(\hat{\Pi}_a^{\hat{n}_A} \otimes \mathbf{1})\rho_{AB}] \\ &= \text{tr}_A[(\hat{\Pi}_a^{\hat{n}_A} \otimes \mathbf{1})(\mu\tau_{AB} + (1 - \mu)\tau'_{AB})] \\ &= \mu \text{tr}_A[(\hat{\Pi}_a^{\hat{n}_A} \otimes \mathbf{1})\tau_{AB}] + (1 - \mu)P(a|A, \tau_{AB})\frac{\mathbf{1}}{2}, \end{aligned}$$

where $P(a|A, \tau_{AB}) = \text{tr}[\hat{\Pi}_a^{\hat{n}_A}\tau_{AB}]$ is the marginal probability of Alice when she measures A and gets the outcome a . For convenient, let us denote the 2×2 matrix $\tilde{\rho}_a^{\hat{n}_A}$ as

$$\tilde{\rho}_a^{\hat{n}_A} = \begin{bmatrix} \nu_{11} & \nu_{12} \\ \nu_{21} & \nu_{22} \end{bmatrix},$$

and calculate its each element. We get

$$\begin{aligned} \nu_{11} &= \text{tr} \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} \nu_{11} & \nu_{12} \\ \nu_{21} & \nu_{22} \end{bmatrix} = \text{tr}[\hat{\Pi}_0^z \tilde{\rho}_a^{\hat{n}_A}] \\ &= \mu P(a, 0|A, z, \tau_{AB}) + (1 - \mu)P(a|A, \tau_{AB})\frac{1}{2}, \end{aligned}$$

and similarly,

$$\begin{aligned} \nu_{22} &= \text{tr} \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} \nu_{11} & \nu_{12} \\ \nu_{21} & \nu_{22} \end{bmatrix} = \text{tr}[\hat{\Pi}_1^z \tilde{\rho}_a^{\hat{n}_A}] \\ &= \mu P(a, 1|A, z, \tau_{AB}) + (1 - \mu)P(a|A, \tau_{AB})\frac{1}{2}. \end{aligned}$$

Note that $\nu_{11} + \nu_{22} = \text{tr}[\tilde{\rho}_a^{\hat{n}_A}] = P(a|A, \tau_{AB})$, we then have

$$\nu_{22} = -\mu P(a, 0|A, z, \tau_{AB}) + (1 + \mu)P(a|A, \tau_{AB})\frac{1}{2}.$$

Because

$$\text{tr} \begin{bmatrix} \frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & \frac{1}{2} \end{bmatrix} \tilde{\rho}_a^{\hat{n}_A} = \frac{1}{2}P(a|A, \tau_{AB}) + \text{Re}[\nu_{12}],$$

with $\text{Re}[\nu_{12}]$ is the real part of ν_{12} , thus,

$$\text{Re}[\nu_{12}] = \text{tr}[\hat{\Pi}_0^x \tilde{\rho}_a^{\hat{n}_A}] - \frac{1}{2}P(a|A, \tau_{AB}) = \mu P(a, 1|A, x, \tau_{AB}) - \frac{\mu}{2}P(a|A, \tau_{AB}).$$

Similarly, because

$$\text{tr} \begin{bmatrix} \frac{1}{2} & -\frac{i}{2} \\ \frac{i}{2} & \frac{1}{2} \end{bmatrix} \tilde{\rho}_a^{\hat{n}_A} = \frac{1}{2}P(a|A, \rho) - \text{Im}[\nu_{12}],$$

with $\text{Im}[\nu_{12}]$ is the imaginary part of ν_{12} , thus,

$$\begin{aligned}\text{Im}[\nu_{12}] &= -\text{tr}[\hat{\Pi}_0^y \hat{\rho}_a^{\hat{n}_A}] + \frac{1}{2}P(a|A, \tau_{AB}) \\ &= -\mu P(a, 1|A, y, \tau_{AB}) + \frac{\mu}{2}P(a|A, \tau_{AB}).\end{aligned}$$

By combining the above equations, we finally have

$$\hat{\rho}_a^{\hat{n}_A} = \begin{bmatrix} \nu_{11} & \nu_{12} \\ \nu_{21} & \nu_{22} \end{bmatrix} = \frac{\nu_{11} + \nu_{22}}{2} \mathbb{1} + \text{Re}[\nu_{12}] \sigma_x - \text{Im}[\nu_{12}] \sigma_y + \frac{\nu_{11} - \nu_{22}}{2} \sigma_z. \quad (19)$$

Let us calculate the right-hand side of Eq. (8). It gives

$$\begin{aligned}\sum_{\xi} P(a|A, \xi) P_{\xi} \frac{\mathbb{1} + \vec{\sigma} \cdot \vec{r}_{\xi}}{2} &= \left(\sum_{\xi} P(a|A, \xi) P_{\xi} \right) \frac{\mathbb{1}}{2} \\ &+ \mu \left(\sum_{\xi} P(a|A, \xi) P(0|x, \xi) P_{\xi} \right) \sigma_x \\ &- \frac{\mu}{2} \left(\sum_{\xi} P(a|A, \xi) P_{\xi} \right) \sigma_x \\ &+ \mu \left(\sum_{\xi} P(a|A, \xi) P(0|y, \xi) P_{\xi} \right) \sigma_y \\ &- \frac{\mu}{2} \left(\sum_{\xi} P(a|A, \xi) P_{\xi} \right) \sigma_y \\ &+ \mu \left(\sum_{\xi} P(a|A, \xi) P(0|z, \xi) P_{\xi} \right) \sigma_z \\ &- \frac{\mu}{2} \left(\sum_{\xi} P(a|A, \xi) P_{\xi} \right) \sigma_z.\end{aligned}$$

With the help of Eq. (3) and using $\sum_{\xi} P(a|A, \xi) P_{\xi} = P(a|A, \tau_{AB})$, we finally have

$$\begin{aligned}\sum_{\xi} P(a|A, \xi) P_{\xi} \frac{\mathbb{1} + \vec{\sigma} \cdot \vec{r}_{\xi}}{2} &= P(a|A, \tau_{AB}) \frac{\mathbb{1}}{2} \\ &+ \mu P(a, 0|A, x, \tau_{AB}) \sigma_x \\ &- \frac{\mu}{2} P(a|A, \tau_{AB}) \sigma_x \\ &+ \mu P(a, 0|A, y, \tau_{AB}) \sigma_y \\ &- \frac{\mu}{2} P(a|A, \tau_{AB}) \sigma_y \\ &+ \mu P(a, 0|A, z, \tau_{AB}) \sigma_z \\ &- \frac{\mu}{2} P(a|A, \tau_{AB}) \sigma_z.\end{aligned} \quad (20)$$

By comparing Eqs (19) and (20), it is easy to see that Eq. (8) holds. Thus, if there is a LHV model description for τ_{AB} , then there is a LHS model description for ρ_{AB} . This completes the proof.

References

1. Einstein, A., Podolsky, B. & Rosen, N. Can Quantum-Mechanical Description of Physical Reality Be Considered Complete? *Phys. Rev.* **47**, 777 (1935).
2. Schrödinger, E. Discussion of probability relations between separated systems. *Naturwiss.* **23**, 807 (1935).
3. Bell, J. S. On the Einstein Podolsky Rosen Paradox. *Physics* (Long Island City, NY) **1**, 195 (1964).
4. Stapp, H. Bell's Theorem and World Process. *Nuovo Cimento* **29B**, 270 (1975).
5. Ekert, A. K. Quantum Cryptography Based on Bell's Theorem. *Phys. Rev. Lett.* **67**, 661 (1991).
6. Brukner, Č., Žukowski, M., Pan, J. W. & Zeilinger, A. Bell's Inequalities and Quantum Communication Complexity. *Phys. Rev. Lett.* **92**, 127901 (2004).
7. Pironio, S. *et al.* Random numbers certified by Bell's theorem. *Nature* (London) **464**, 1021 (2010).
8. Wiseman, H. M., Jones, S. J. & Doherty, A. C. Steering, Entanglement, Nonlocality, and the Einstein-Podolsky-Rosen Paradox. *Phys. Rev. Lett.* **98**, 140402 (2007).
9. Oppenheim, J. & Wehner, S. The Uncertainty Principle Determines the Nonlocality of Quantum Mechanics. *Science* **330**, 1072 (2010).
10. Branciard, C., Cavalcanti, E. G., Walborn, S. P., Scarani, V. & Wiseman, H. M. One-sided device-independent quantum key distribution: Security, feasibility, and the connection with steering. *Phys. Rev. A* **85**, 010301(R) (2012).
11. Chen, J. L. *et al.* All-Versus-Nothing Proof of Einstein-Podolsky-Rosen Steering. *Sci. Rep.* **3**, 2143 (2013).
12. He, Q. Y. & Reid, M. D. Genuine Multipartite Einstein-Podolsky-Rosen Steering. *Phys. Rev. Lett.* **111**, 250403 (2013).
13. Jevtic, S., Pusey, M., Jennings, D. & Rudolph, T. Quantum Steering Ellipsoids. *Phys. Rev. Lett.* **113**, 020402 (2014).
14. Skrzypczyk, P., Navascués, M. & Cavalcanti, D. Quantifying Einstein-Podolsky-Rosen Steering. *Phys. Rev. Lett.* **112**, 180404 (2014).
15. Bowles, J., Vertesi, T., Quintino, M. T. & Brunner, N. One-way Einstein-Podolsky-Rosen Steering. *Phys. Rev. Lett.* **112**, 200402 (2014).
16. Piani, M. & Watrous, J. Necessary and Sufficient Quantum Information Characterization of Einstein-Podolsky-Rosen Steering. *Phys. Rev. Lett.* **114**, 060404 (2015).

17. Saunders, D. J., Jones, S. J., Wiseman, H. M. & Pryde, G. J. Experimental EPR-steering using Bell-local states. *Nature Phys.* **6**, 845 (2010).
18. Smith, D. H. *et al.* Conclusive quantum steering with superconducting transition edge sensors. *Nature Comm.* **3**, 625 (2012).
19. Bennet, A. J. *et al.* Arbitrarily loss-tolerant Einstein-Podolsky-Rosen steering allowing a demonstration over 1 km of optical fiber with no detection loophole. *Phys. Rev. X* **2**, 031003 (2012).
20. Wittmann, B. *et al.* Loophole-free quantum steering. *New J. Phys.* **14**, 053030 (2012).
21. Händchen, V. *et al.* Observation of one-way Einstein-Podolsky-Rosen steering. *Nature Photonics* **6**, 596 (2012).
22. Schneeloch, J., Dixon, P. B., Howland, G. A., Broadbent, C. J. & Howell, J. C. Violation of Continuous-Variable Einstein-Podolsky-Rosen Steering with Discrete Measurements. *Phys. Rev. Lett.* **110**, 130407 (2013).
23. Sun, K. *et al.* Experimental Demonstration of the Einstein-Podolsky-Rosen Steering Game Based on the All-Versus-Nothing Proof. *Phys. Rev. Lett.* **113**, 140402 (2014).
24. Li, C. M. *et al.* Genuine High-Order Einstein-Podolsky-Rosen Steering. *Phys. Rev. Lett.* **115**, 010402 (2015).
25. Wollmann, S., Walk, N., Bennet, A. J., Wiseman, H. M. & Pryde, G. J. Observation of Genuine One-Way Einstein-Podolsky-Rosen Steering. *Phys. Rev. Lett.* **116**, 160403 (2016).
26. Sun, K. *et al.* Experimental Quantification of Asymmetric Einstein-Podolsky-Rosen Steering. *Phys. Rev. Lett.* **116**, 160404 (2016).
27. Bowles, J., Hirsch, F., Quintino, M. T. & Brunner, N. Sufficient criterion for guaranteeing that a two-qubit state is unsteerable. *Phys. Rev. A* **93**, 022121 (2014).
28. Werner, R. F. Quantum states with Einstein-Podolsky-Rosen correlations admitting a hidden-variable model. *Phys. Rev. A* **40**, 4277 (1989).
29. Clauser, J. F., Horne, M. A., Shimony, A. & Holt, R. A. Proposed Experiment to Test Local Hidden-Variable Theories. *Phys. Rev. Lett.* **26**, 880 (1969).
30. Hensen, B. *et al.* Loophole-free Bell inequality violation using electron spins separated by 1.3 kilometres. *Nature* **526**, 682 (2015).
31. Luo, S. Quantum discord for two-qubit systems. *Phys. Rev. A* **77**, 042303 (2008).

Acknowledgements

J.L.C. is supported by the National Basic Research Program (973 Program) of China under Grant No. 2012CB921900 and the Natural Science Foundations of China (Grant No. 11475089). C.R. acknowledges supported by Youth Innovation Promotion Association (CAS) No. 2015317, Natural Science Foundations of Chongqing (No. cstc2013jcyjC00001, cstc2015jcyjA00021) and The Project-sponsored by SRF for ROCS-SEM (No. Y51Z030W10). C.C. was partially supported by NSFC (11301524, 11471307) and CSTC (cstc2015jcyjys40001). A.K.P. is supported by the Special Project of University of Ministry of Education of China and the Project of K. P. Chair Professor of Zhejiang University of China.

Author Contributions

J.L.C. initiated the idea. J.L.C., C.R., C.C. and X.J.Y. derived the results. J.L.C. prepared the figure. J.L.C. and A.K.P. wrote the main manuscript text. All authors reviewed the manuscript.

Additional Information

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Chen, J.-L. *et al.* Bell's Nonlocality Can be Detected by the Violation of Einstein-Podolsky-Rosen Steering Inequality. *Sci. Rep.* **6**, 39063; doi: 10.1038/srep39063 (2016).

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2016