

Statistical isotropy of the cosmic microwave background

TARUN SOURADEEP and AMIR HAJIAN

Inter-University Centre for Astronomy and Astrophysics, Post Bag 4, Ganeshkhind,
Pune 411 007, India

Abstract. The breakdown of statistical homogeneity and isotropy of cosmic perturbations is a generic feature of ultra-large scale structure of the cosmos, in particular, of non-trivial cosmic topology. The statistical isotropy (SI) of the cosmic microwave background temperature fluctuations (CMB anisotropy) is sensitive to this breakdown on the largest scales comparable to, and even beyond the cosmic horizon. We propose a set of measures, κ_ℓ ($\ell = 1, 2, 3, \dots$) which for non-zero values indicate and quantify statistical isotropy violations in a CMB map. We numerically compute the predicted κ_ℓ spectra for CMB anisotropy in flat torus universe models. Characteristic signatures of different models in the κ_ℓ spectrum are noted.

Keywords. Cosmic microwave background; cosmology; observations.

In standard cosmology, the cosmic microwave background (CMB) anisotropy is expected to be statistically isotropic, i.e., statistical expectation values of the temperature fluctuations $\Delta T(\hat{q})$ are preserved under rotations of the sky. In particular, the angular correlation function $C(\hat{q}, \hat{q}') \equiv \langle \Delta T(\hat{q}) \Delta T(\hat{q}') \rangle$ is rotationally invariant. In spherical harmonic space, where $\Delta T(\hat{q}) = \sum_{lm} a_{lm} Y_{lm}(\hat{q})$, this translates to a diagonal $\langle a_{lm} a_{l'm'}^* \rangle = C_l \delta_{ll'} \delta_{mm'}$, where C_l is the widely used angular power spectrum.

In the absence of statistical isotropy, $C(\hat{q}, \hat{q}')$ is estimated by a single product $\Delta T(\hat{q}) \Delta T(\hat{q}')$ and hence is poorly determined from a single realization. Although it is not possible to estimate each element of the full correlation function $C(\hat{q}, \hat{q}')$, some measures of statistical anisotropy of the CMB map can be estimated through suitably weighted angular averages of $\Delta T(\hat{q}) \Delta T(\hat{q}')$. The angular averaging procedure should be such that the measure involves averaging over sufficient number of independent ‘measurements’, but should ensure that the averaging does not erase all the signature of statistical anisotropy. Another important desirable property is that the measure be independent of the overall orientation of the sky. Based on these considerations, we propose a set of measures κ_ℓ of statistical anisotropy given by

$$\kappa_\ell = \int d\Omega \int d\Omega' \left[\frac{(2\ell + 1)}{8\pi^2} \int d\mathcal{R} \chi_\ell(\mathcal{R}) C(\mathcal{R}\hat{q}, \mathcal{R}\hat{q}') \right]^2, \quad (1)$$

where $C(\mathcal{R}\hat{q}, \mathcal{R}\hat{q}')$ is the two-point correlation between $\mathcal{R}\hat{q}$ and $\mathcal{R}\hat{q}'$ obtained by rotating \hat{q} and \hat{q}' by an element \mathcal{R} of the rotation group [1]. The measures κ_ℓ involve angular average of the correlation weighed by the characteristic function of the rotation group $\chi_\ell(\mathcal{R}) = \sum_M D_{MM}^\ell(\mathcal{R})$, where D_{MM}^ℓ are the Wigner D -functions [2]. When SI holds $C(\mathcal{R}\hat{q}, \mathcal{R}\hat{q}') = C(\hat{q}, \hat{q}')$ is invariant under rotation, and eq. (1) gives $\kappa_\ell = \kappa_0 \delta_{\ell 0}$ due to the orthonormality of $\chi_\ell(\mathcal{R})$. Hence, non-zero κ_ℓ for $\ell > 0$ measure violation of statistical isotropy.

The measure κ_ℓ has a clear interpretation in harmonic space. The two-point correlation $C(\hat{q}, \hat{q}')$ can be expanded in terms of the orthonormal set of bipolar spherical harmonics whose coefficients $A_{ll'}^{\ell M}$ are related to ‘angular momentum’ sum over the covariances $\langle a_{lm} a_{l'm'}^* \rangle$ as $A_{ll'}^{\ell M} = \sum_{mm'} \langle a_{lm} a_{l'm'}^* \rangle (-1)^{m'} \mathfrak{C}_{lm' - m'}^{\ell \mathfrak{M}}$, where $\mathfrak{C}_{lm' - m'}^{\ell \mathfrak{M}}$ are Clebsch–Gordan coefficients [2]. The estimation of κ_ℓ from a CMB map is discussed in [1].

The detection of statistical isotropy (SI) violations can have exciting and far-reaching implications for cosmology. The realization that the universe with the same local geometry has many different choices of global topology has been a theoretical curiosity as old as modern cosmology. Motivations for cosmic topology and their consequences have been extensively studied [3]. CMB anisotropy measurements have brought cosmic topology from the realm of theoretical possibility to within the grasp of observations [3,4]. A generic consequence of cosmic topology is the breaking of statistical isotropy in characteristic patterns determined by the photon geodesic structure of the manifold. Global isotropy of space is violated in all multi-connected models (except S^3/Z_2). In cosmology, the Dirichlet domain (DD) constructed around the observer represents the universe as ‘seen’ by the observer. The SI breakdown is apparent in the principal axes present in the shape of the DD constructed with the observer located at the base-point [5].

In this paper we compute and study the κ_ℓ spectrum of SI violation arising in flat (Euclidean) simple torus models with a cubic, cuboidal and more generally, parallelepiped (squeezed) fundamental domain. The CMB anisotropy in torus spaces has been well-studied [3]. We can relate the κ_ℓ spectrum to the principal directions normal to pair of faces of the DD, their relative orientation and the relative importance given by the distance to the faces along them. Along the most dominant axes, the distance is minimum, and equals the in-radius, $R_<$, the radii of largest sphere fully enclosed within the DD [4].

The compact spaces with Euclidean geometry (zero curvature) have been completely classified. In three dimensions, there are known to be six possible topologies that lead to orientable spaces [3,6]. The simple flat torus, $\mathcal{M} = T^3$, is obtained by identifying the universal cover $\mathcal{M}^u = \mathcal{E}^3$ under a discrete group of translations along three non-degenerate axes, $\mathbf{s}_1, \mathbf{s}_2, \mathbf{s}_3$: $\mathbf{s}_i \rightarrow \mathbf{s}_i + \mathbf{n}L_i$, where L_i is the identification length of the torus along s_i and \mathbf{n} is a vector with integer components. In the most general form, the fundamental domain (FD) is a parallelepiped defined by three sides L_i and the three angles α_i between the axes (we call it squeezed torus). If \mathbf{s}_i are orthogonal then one gets cuboidal FD, which for equal L_i reduces to the cubic torus. The cuboid and squeezed spaces which can be obtained by a linear coordinate transformation \mathcal{L} on cubic torus can have distinctly different global symmetry [6a].




Figure 1. The κ_ℓ spectra for flat tori models are plotted. The top row panels are for cubic tori spaces. The left panel shows spaces of volume, $V_{\mathcal{M}}$, larger than the volume V_* contained in the sphere of last scattering (SLS) with $V_{\mathcal{M}}/V_* = 3.7, 1.9, 1.1$, respectively. The right panel shows small spaces with $V_{\mathcal{M}}/V_* = 0.24, 0.07, 0.03$, respectively. Note that $\kappa_2 = 0$ for cubic tori. The middle panels consider cuboid tori with 1 : 5 and 1 : 2 ratio of identification lengths. The bottom panels show κ_ℓ for equal-sided squeezed tori with $\alpha = 45^\circ, 60^\circ$ and 75° . In the middle and bottom rows, the right panels show the case when radius of SLS, $R_* = R_<$ the in-radius of the space. Here, the SLS just touches its nearest images which is at the threshold where CMB anisotropy is multiply imaged for larger R_* . The cases in the left panels of lower two rows have $V_{\mathcal{M}}/V_* = 1$ and are at the divide between large and small spaces.

We restrict attention to the case where CMB anisotropy arises entirely at the sphere of last scattering (SLS) of radius R_* . Invoking the method of images, the CMB correlation pattern on the SLS is known to be dictated by the distribution of nearest ‘images’ of the SLS on the universal cover [4]. The correlations are distorted even when the SLS and its images do not intersect ($R_* < R_<$). When SLS intersects its images the CMB sky is multiply imaged in characteristic correlation pattern of pairs of circles [7].

We compute $\mathbf{C}(\hat{\mathbf{q}}, \hat{\mathbf{q}}')$ for CMB anisotropy in torus space using regularized method of images [4]. Figure 1 plots the predicted κ_ℓ spectrum for a number of cubic, cuboidal and squeezed torus spaces. We note the following interesting results:

- (i) $\kappa_\ell = 0$ for odd ℓ for all torus models. This does not hold for compact space of non-zero curvature, e.g., compact hyperbolic spaces.

- (ii) For cubic torus $\kappa_2 = 0$. κ_2 is non-zero for cuboidal and squeezed torus. This is a clear signature of non-cubic torus where the DD differs from the FD and has more than three principal axes.
- (iii) For equal-sided squeezed torus, κ_4 , decreases as α decreases from 90° to 60° as $R_<$ increases. For $\alpha < 60^\circ$ sharply increases with decreasing α as $R_<$ decreases sharply.
- (iv) $\kappa_2 = 0$ increases monotonically as α decreases from 90° .
- (v) The peak of κ_ℓ shifts to larger ℓ for small spaces.

The results can be understood using the leading order terms of the correlation function in a torus where κ_ℓ can be calculated analytically [5].

Preferred directions and statistically anisotropic CMB anisotropy have been discussed in the literature [8]. When CMB anisotropy is multiply imaged, the κ_ℓ spectrum corresponds to a correlation pattern of matched pairs of circles [7]. The generic features of κ_ℓ spectrum are related to the symmetries of correlation pattern. But κ_ℓ are sensitive to SI violation even when CMB is not multiply imaged. Moreover, κ_ℓ have an advantage of being insensitive to the overall orientation of the correlation features. The $A_{\ell}^{\ell M}$ signature, which was not discussed here, contains more details of the SI violation [8]. The estimation of κ_ℓ from a CMB map is described [1]. Before ascribing the detected breakdown of statistical anisotropy to cosmological or astrophysical effects, one must carefully account for and model out other mundane sources of statistical anisotropy in real data, such as, incomplete and non-uniform sky coverage, beam anisotropy, foreground residuals and statistically anisotropic noise. These observational artifacts will be discussed in future publications.

References

- [1] A Hajian and T Souradeep, IUCAA preprint, astro-ph/xxxx
- [2] D A Varshalovich, A N Moskalev and V K Khersonskii, *Quantum theory of angular momentum* (World Scientific, 1988)
- [3] G F R Ellis, *Gen. Relativ. Gravit.* **2**, 7 (1971)
M Lachieze-Rey and J-P Luminet, *Phys. Rep.* **25**, 136, (1995)
J Levin, *Phys. Rep.* **365**, 251 (2002)
G Starkman, *Class. Quantum Gravit.* **15**, 2529 (1998)
- [4] J R Bond, D Pogosyan and T Souradeep, *Class. Quantum Gravit.* **15**, 2671 (1998);
Phys. Rev. **D62**, 043005 (2000); 043006 (2000)
T Souradeep, in *The universe* edited by N Dadhich and A Kembhavi (Kluwer, 2000)
- [5] A Hajian and T Souradeep, preprint (astro-ph/0301590)
- [6] J A Wolf, *Space of constant curvature* (5th ed.) (Publish or Perish Inc., 1994)
E B Vinberg, *Geometry II - Spaces of constant curvature* (Springer-Verlag, 1993)
- [6a] For cubic torus the Dirichlet domain (DD) matches the fundamental domain (FD). However, for torus spaces with cuboid and parallelepiped FD, the corresponding DD is very different, e.g., hexagonal prism [6,5]
- [7] N J Cornish, D N Spergel and G D Starkman, *Class. Quantum Gravit.* **15**, 2657 (1998)
- [8] P G Ferreira and J Magueijo, *Phys. Rev.* **D56**, 4578 (1997)
E Bunn and D Scott, *Mon. Not. R. Astron. Soc.* **313**, 331 (2000)