

Click-generated triazole based ferrocene-carbohydrate bioconjugates: A highly selective multisignalling probe for Cu(II) ions

ARUNABHA THAKUR, SINJINEE SARDAR and SUNDARGOPAL GHOSH*

Department of Chemistry, Indian Institute of Technology Madras, Chennai 600 036, India

e-mail: sghosh@iitm.ac.in

Abstract. Two Cu²⁺-specific colorimetric sensors, based on ferrocene-carbohydrate bioconjugates, **2**, C₄₆H₅₆O₂₀N₆Fe and **3**, C₂₈H₃₃O₁₀N₃Fe were designed and synthesized in good yields. Both the compounds, **2** and **3**, behave as very selective and sensitive chromogenic and electrochemical chemosensor for Cu²⁺ ion in aqueous environment (CH₃CN/H₂O (2:8, v/v)). The analytical detection limit (ADL) for receptor **2** was 7.5 × 10⁻⁷ M. The considerable changes in their absorption spectra of **2** and **3** are accompanied by the appearance of a new low energy (LE) peak at 630 nm (**2**: ε = 1600 M⁻¹ cm⁻¹ and **3**: 822 M⁻¹ cm⁻¹). This is further accompanied by a strong colour change from yellow to dark green that allows the prospective for ‘naked eye’ detection of Cu²⁺ ion.

Keywords. Organometallic bioconjugates; Cu(II) cation sensor; chromogenic and electrochemical chemosensor.

1. Introduction

In recent years, there has been a growing need for constructing chemosensors for fast and economical monitoring of our environmental samples, especially for heavy metal ions.¹ Copper is one of the heavy metals which is an essential element not only for life in mammals but also for plants. It also plays an important role in carbohydrate and lipid metabolism.² It is the most significant metal ion in biological systems³ and also a significant environmental pollutant.⁴ Copper is implicated in inflammatory disorders⁵ and Alzheimer’s disease.⁶ The Cu²⁺ proteins are involved in oxygen binding, electron transfer and the activation of small molecules.⁷ As a result, a strong interest exists in the development of selective Cu²⁺ sensors for biological and environmental applications. A variety of Cu²⁺ probes exhibiting either fluorescence ‘on-off’ or ‘off-on’ signalling modes have been developed.⁸ However, colorimetric Cu²⁺ sensors offering ratiometric response are rare.⁹ Colorimetric probes are currently attracting area, since they can be tailored to allow ‘naked eye’ detection and ratiometric sensing¹⁰ of the analyte.

The most attractive way of achieving sensor design is to functionalize a receptor capable of both selective substrate binding with a metal centre and reporting on the recognition event through a variety of physical responses. Therefore, the design of redox-active receptors in which a change in electrochemical behaviour

can be used to monitor complexations of guest species is significant in molecular recognition.^{11–20} Thus, from a synthetic standpoint, ferrocene is a very convenient building block for redox-active ligand as it can be easily functionalized and incorporated in many structures. These facts, along with its electrochemical and UV-vis spectroscopic properties, demonstrate that ferrocene is a particularly attractive functional antenna in area of sensor for transition metals, p-block anions and organic molecules. For example, water-soluble ferrocenyl sugars are useful for the development of ferrocene-containing drugs. Further this, also been observed that some ferrocenyl sugars possess anti-malarial activity.²¹ Carbohydrate-based chemosensors are chiral entities with hydroxyl groups and oxygen atoms, that form quite suitable cation binding sites. Thus, in the design of chemosensors, the incorporation of sugar molecules is a good strategy for capturing cations.^{22–24} In this article, we report the host-guest complexation properties of two triazole tethered ferrocenyl carbohydrate bioconjugates towards Cu²⁺ ion.

2. Experimental

2.1 General procedures and instrumentation

Perchlorate salts of Li⁺, Na⁺, K⁺, Ag⁺, Ca²⁺, Mg²⁺, Co²⁺, Cu²⁺, Zn²⁺, Cd²⁺, Ni²⁺, Pb²⁺, and Hg²⁺,

*For correspondence

propargyl bromide, butyl-lithium, tetramethylethylenediamine (TMEDA) purchased from Aldrich were used directly without further purification. Ferrocene, sodium ascorbate, sodium azide, acetonitrile purchased were of analytical grade and used without further purification. DMF purchased from Aldrich and freshly distilled prior to use. Chromatography was carried out on 3 cm of silica gel in a 2.5 cm diameter column. Column chromatography was carried out using 100–200 mesh silica gel. All the solvents were dried by conventional methods and distilled under a N₂ atmosphere before use. Glycosyl azide²⁵ and compounds **1a–b** [Fc(CH₂OCH₂C≡CH)_n] (**1a**: n = 2, **1b**: n = 1, where Fc = ferrocene), were synthesized as per literature procedures.²⁶ The cyclic voltammetry (CV) and differential pulse voltammetry (DPV) were performed with a conventional three-electrode configuration consisting of glassy carbon as working electrode, platinum as an auxiliary electrode and Ag/Ag⁺ as a reference electrode. The experiments were carried out with a 10^{−4} M solution of sample in CH₃CN or CH₃CN/H₂O (2/8) containing 0.1 M [(n-C₄H₉)₄NCIO₄] (TBAP) as supporting electrolyte. Deoxygenation of the solutions was achieved by bubbling nitrogen for at least 10 min, and the working electrode was cleaned after each run. The cyclic voltammograms were recorded at a scan rate of 0.1 V s^{−1}. The UV-vis spectra were carried out in CH₃CN or CH₃CN/H₂O (2/8) solutions at c = 1 × 10^{−4} M.

The ¹H and ¹³C NMR spectra were recorded on Bruker 400 MHz FT-NMR spectrometers, using tetramethylsilane as the internal reference. Electrospray ionization mass spectrometry (ESI-MS) measurements were carried out on a Qtof Micro YA263 HRMS instrument. The absorption spectra were recorded with a JASCO V-650 UV-vis spectrophotometer at 298 K. The CV and DPV measurements were performed on a CH potentiostat model 660 B. *Caution*: Metal perchlorate salts are potentially explosive in certain conditions. All precautions should be taken while handling perchlorate salts.

2.2 Synthesis of ferrocene-carbohydrate conjugates 2–3

To a well-stirred solution of **1a** (0.5 g, 1.55 mmol) and glycosyl azide (3.31 g, 3.1 mmol) in 15 mL acetone/H₂O (2:1), an aqueous solution of CuSO₄·5H₂O (0.077 g, 0.31 mmol) was added. To this resultant mixture freshly prepared sodium ascorbate solution (0.122 g, 0.62 mmol) was added and stirred at room temperature for 12 h. 30 mL of ethyl acetate was added into the

reaction mixture and the organic layer was washed several times with water and finally with brine (15 mL) and dried over anhydrous sodium sulphate. The solvent was removed under reduced pressure and the crude product was purified by silica gel column chromatography. Elution with EtOAc:hexane (8:2 v/v) yielded yellow **2** (1.42 g, 86%).

Compound **3** was prepared in good yield following the procedure adopted for **2** from alkyne, **1b** (0.5 g, 1.96 mmol), glycosyl azide (1.23 g, 1.96 mmol), aqueous CuSO₄·5H₂O (0.097 g, 0.392 mmol) and sodium ascorbate (0.149 g, 0.776 mmol). The crude product was purified by silica gel column chromatography and elution with EtOAc: hexane (7:2 v/v) to yield pure yellow, **3** (0.98 g, 80%).


2: ¹H NMR (CDCl₃, 400 MHz): δ = 7.71 (s, 2H, H_{triazole}), 5.82 (d, 2H, H-1), 5.45 (s, 2H, H-5), 5.17 (s, 2H, H-2), 5.01 (s, 2H, H-3), 4.51 (s, 2H, H-4), 4.21 (dd, 4H, H₆, H_{6'}), 4.18 (s, 4H, OCH₂-triazole), 4.08 (t, 4H, H_{Fc}), 4.0 (t, 4H, H_{Fc}), 3.95 (s, 4H, OCH₂), 1.96–1.81 (d, 12H, OAc); ¹³C NMR (CDCl₃, 100 MHz): δ = 170.5 (CO), 169.9 (CO), 169.4 (CO), 168.7 (CO), 146.0 (C_{triazole}), 121.0 (C_{triazole}), 85.6 (C-1), 75.0 (C-2), 72.6 (C-3), 71.7 (C-5), 70.3 (C-4), 68.8 (C_{Fc}), 68.4 (C_{Fc}), 67.6 (C_{Fc}), 62.9 (C-6), 61.5 (OCH₂), 53.6 (OCH₂), 27.7, 20.5, 20.2, 19.1 (CH₃CO); ESI MS, m/z (relative intensity): 1069 (M⁺ + 1).

3: ¹H NMR (CDCl₃, 400 MHz): δ = 7.68 (s, 1H, H_{triazole}), 5.81 (d, 1H, H-1), 5.30 (m, 1H, H-5), 5.23 (m, 1H, H-2), 5.01 (s, 1H, H-3), 4.53 (s, 1H, H-4), 4.26 (s, 2H, H₆, H_{6'}), 4.21 (s, 2H, OCH₂-triazole), 4.06 (m, 4H, H_{Fc}), 4.0 (s, 5H, H_{Fc}), 3.99 (s, 2H, OCH₂), 2.08 (s, 3H, OAc), 1.98 (s, 3H, OAc), 1.94 (s, 3H, OAc), 1.80 (s, 3H, OAc); ¹³C NMR (CDCl₃, 100 MHz): δ = 170.2 (CO), 169.7 (CO), 169.3 (CO), 168.9 (CO), 146.1 (C_{triazole}), 120.9 (C_{triazole}), 85.6 (C-1), 75.0 (C-2), 72.5 (C-3), 71.7 (C-4), 70.2 (C-5), 69.2 (C_{Fc}), 68.6 (C_{Fc}), 68.5 (C_{Fc}), 67.6 (C_{Fc}), 62.9 (C-6), 61.5 (OCH₂), 53.5 (OCH₂), 29.6, 27.7, 20.6, 19.0 (CH₃CO); ESI MS, m/z (relative intensity): 650 (M⁺ + 23).

3. Results and discussion

3.1 Synthesis

Precursors **1a–b** were obtained following literature procedure.²⁶ As shown in scheme 1, they undergo the ‘click reaction’ with glycosyl azide to generate compounds **2** and **3** in 86% and 80% yields, respectively. Compounds **2** and **3** have been characterized by ¹H, ¹³C NMR spectroscopy and ESI-MS spectrometry. Both the compounds **2** and **3** are moderately stable and



Scheme 1. Synthesis of mono and di-ferrocene-carbohydrate bioconjugates, **2** and **3**. **(a)** 2 equiv. glycosyl azide, 0.1 equiv. CuSO₄·5H₂O, 0.25 equiv. Na Ascorbate, Acetone/H₂O (2:1); **(b)** 1 equiv. glycosyl azide, 0.1 equiv. CuSO₄·5H₂O, 0.25 equiv. Na Ascorbate, Acetone/H₂O (2:1).

could be stored for several months. The complexation properties of the receptors **2** and **3** have been investigated by electrochemistry and UV-vis spectroscopic measurements.

3.2 UV-vis absorption studies

The UV-vis binding interaction studies of receptors **2** and **3** in CH₃CN (1 × 10⁻⁴ M) against cation of environmental relevance, such as of Li⁺, Na⁺, K⁺, Ag⁺, Ca²⁺, Mg²⁺, Zn²⁺, Ni²⁺, Co²⁺, Cd²⁺, Hg²⁺ and Pb²⁺ as perchlorate salts, show selective response to Cu²⁺. The change in the UV-vis absorbance spectra of receptors **2**

and **3** in CH₃CN due to the step-wise addition of Cu²⁺ ion are shown in the figures 1 and 2, respectively. As shown in figures 1–2, a new and weak low-energy (LE) absorption band appeared at λ = 630 nm for both **2** (ε = 1600 M⁻¹ cm⁻¹) and **3** (ε = 822 M⁻¹ cm⁻¹) was developed. These facts are responsible for the change of colour from yellow to dark green. In addition, one well-defined isosbestic point at 408 nm and 415 nm was observed for **2** and **3**, respectively. The Cu²⁺ induced UV-vis response of **2** and **3** was almost unaffected in a background of environmentally relevant metallic cations.

The UV-vis spectral change suggests that the ferrocene moiety is oxidized upon complexation with




Figure 1. (a) Changes in the absorption spectra of **2** (10⁻⁴ M) in CH₃CN upon addition of increasing amounts of Cu²⁺ up to 1 equivalent. (b) Expanded form of part a. (c) Job's plot for **2** and Cu²⁺, indicating the formation of 1:1 binding model. The total [2] + [Cu²⁺] = 1 × 10⁻⁴ M.




Figure 2. (a) Changes in the absorption spectra of **3** (10^{-4} M) in CH_3CN upon addition of increasing amounts of Cu^{2+} up to 1 equivalent. (b) Expanded form of part a. (c) Job's plot for **3** and Cu^{2+} , indicating the formation of 1:1 binding model. The total $[\mathbf{3}] + [\text{Cu}^{2+}] = 1 \times 10^{-4}$ M.

Cu^{2+} ion and the change of colour to deep green is characteristic of the ferrocenium ion formation.²⁷ On the basis of absorption intensity changes at 630 nm as a function of the amount of Cu^{2+} (inset figure 1b), it could be estimated that the stoichiometry of both **2** and **3** with Cu^{2+} is 1:1. This is further supported by the Job's plots (figures 1c for **2** and 2c for **3**) and ESI-MS experiments, where a peak at $m/z = 1131$ corresponds to $[\mathbf{2} + \text{Cu}^{2+} - \text{H}^+]$ and a small peak at $m/z = 1230$ for $[\mathbf{2} + \text{Cu}.\text{ClO}_4]$ was observed. Similarly, for **3** a peak at $m/z = 789$ was observed which corresponds to $[\mathbf{3} + \text{Cu}.\text{ClO}_4]$ (supporting information, figures S3 and S4).

3.3 Electrochemical studies

Chemical sensors bearing ferrocene nuclei as part of the sensing unit have been broadly studied. Earlier, the complexation of ferrocene with a variety of binding ligands have been studied by cyclic voltammetry that shows a positive shift of the Fe(II)/Fe(III) redox couple as a result of metal–ligand complexation.²⁸ The metal-recognition properties of receptors **2** and **3** were evaluated by cyclic (CV) and differential pulse voltammetry (DPV) analysis. The reversibility and relative oxidation potential of the redox process were




Figure 3. Evolution of the CV and DPV of **2** (a and b) and **3** (c and d) (10^{-4} M) in CH_3CN upon addition of increasing amounts of Cu^{2+} metal cation up to 1 equivalent using $[(n\text{-Bu})_4]\text{ClO}_4$ as supporting electrolyte. Arrow indicates the movement of the wave during the experiments.




Figure 4. Evolution of LSV of **2** (a) and **3** (b) (10^{-4} M) in CH_3CN upon addition of with Cu^{2+} ion using $[(n\text{-Bu})_4\text{N}]\text{ClO}_4$ as supporting electrolyte and scanned at 0.1 V s^{-1} .

determined by CV and DPV in CH_3CN solutions containing $0.1 \text{ M } [(n\text{-Bu})_4\text{N}]\text{ClO}_4$ as supporting electrolyte. Both the compounds **2** and **3** display a reversible one-electron oxidation process at $E_{1/2} = 0.433$ and 0.415 V , respectively due to the ferrocene/ferrocenium redox couple. No perturbation of the CV and DPV voltammograms of **2** and **3** were observed in the presence of several metal cations such as Li^+ , Na^+ , K^+ , Ca^{2+} , Mg^{2+} , Zn^{2+} , Ag^+ , Ni^{2+} , Co^{2+} , Cd^{2+} , Hg^{2+} and Pb^{2+} as their appropriate salts, even with large excess. However, as shown in figure 3a–d, the original peak gradually decreased upon step-wise addition of Cu^{2+} ion towards more cathodic current which indicate that free receptor is getting oxidized upon interaction with Cu^{2+} ion.

In addition, linear sweep voltammetry (LSV) studies carried out upon addition of Cu^{2+} to the CH_3CN solution of receptor **2**. As shown in the figure 4, a significant shift of the voltammetric wave towards more cathodic current was observed, indicating that this metal cation promotes the oxidation of the free receptor with its concomitant reduction to Cu^+ . This is in agreement with the CV and DPV (figure 3a–d). Remarkably, the redox response towards Cu^{2+} is also preserved in the presence of an aqueous environment ($\text{CH}_3\text{CN}/\text{H}_2\text{O}$ (2:8, v/v)).




Figure 5. Visual features observed in CH_3CN solution of **2** (10^{-4} M) after addition of 10 equivalent of different metal cation tested.

3.4 Visual detection of Cu^{2+} ion

When an excess of different metal cations (Li^+ , Na^+ , K^+ , Ag^+ , Ca^{2+} , Mg^{2+} , Cr^{2+} , Zn^{2+} , Ni^{2+} , Fe^{2+} , Co^{2+} , Cd^{2+} , Hg^{2+} and Pb^{2+}) as their perchlorate salt were separately added to a solution of **2** and **3** in $\text{CH}_3\text{CN}:\text{H}_2\text{O}$ (10^{-4} M), no significant colour change observed, except for Cu^{2+} . As shown in figure 5, Cu^{2+} shows a drastic colour change from yellow to dark green. The sensing potential of **3** toward Cu^{2+} in solution is very similar to **2**. This indicates that both **2** and **3** are highly selective colorimetric sensors for Cu^{2+} ion.

4. Conclusion

In this study, we have designed and synthesized two ferrocene-carbohydrate based organometallic bioconjugates, **2**, $\text{C}_{46}\text{H}_{56}\text{O}_{20}\text{N}_6\text{Fe}$ and **3**, $\text{C}_{28}\text{H}_{33}\text{O}_{10}\text{N}_3\text{Fe}$ in good yields. They behave as selective and sensitive electrochemical as well as chromogenic receptors for the determination Cu^{2+} ion. These receptors showed high selectivity towards Cu^{2+} ion not only through spectrochemical and electrochemical probe but also amenable to the facile colorimetric sensing of Cu^{2+} ion, thus allowing the potential for ‘naked-eye’ detection over some other cations.

Supplementary information

The ^1H , ^{13}C and ESI-MS data of **2** and **3**; electrochemical data for **2** and **3** upon titration with different metal ions; UV-Vis spectra upon titration with different metal ions; ESI-MS spectrum of $[\mathbf{2}.\text{Cu}^{2+}]$, $[\mathbf{3}.\text{Cu}^{2+}]$ are given as Supplementary information. Figures S1–S6 as supporting material are available on www.ias.as.in/chemsci.

Acknowledgements

Generous support of the Indo–French Centre for the Promotion of Advanced Research (IFCPAR-CEFIPRA), No. 4405–1, New Delhi is gratefully acknowledged. AT is grateful to the Council of Scientific and Industrial Research (CSIR), India for research fellowships.

References

1. Yari A and Afshari N 2006 *Sens. Actuators B* **199** 531
2. (a) Scheninberg H J and Morell G A 1973 **1** *Inorganic biochemistry* (New York: Elsevier); (b) Dieter H H, Schimmelpfenning W and Mayer E 1999 *Eur. J. Med. Res.* **4** 233; (c) Miller T, Sluis V B, Miller W and Pearson P 1999 *Eur. J. Med. Res.* **4** 293
3. (a) Linder M C and Hazegh-Azam M 1996 *Am. J. Clin. Nutr.* **63** 797S; (b) Uauy R, Olivares M and Gonzalez M 1998 *Am. J. Clin. Nutr.* **67** 952S; (c) Frausto da Silva J J R and Williams R J P 1993 *The biological chemistry of elements: The inorganic chemistry of life* (Oxford: Clarendon Press)
4. (a) Sarkar B 1981 *Met. Ions Biol. Syst.* **12** 233; (b) Yardim M F, Budinova T, Ekinici E, Petrov N, Razvigorova M and Minkova V 2003 *Chemosphere* **52** 835
5. Alan J L 1984 *Inflamm. Res.* **15** 513
6. Barnham K J, Masters C L and Bush A I 2004 *Nat. Rev. Drug Discovery* **3** 205
7. Frieden E, Osaki S and Kobayashi H 1965 *J. Gen. Physiol.* **49** 213
8. (a) Ghosh P, Bharadwaj P K, Mandal S and Ghosh S 1996 *J. Am. Chem. Soc.* **118** 1553; (b) Kramer R 1998 *Angew. Chem. Int. Ed.* **37** 772; (c) Rurack K, Kollmannsberger M, Resch-Genger U and Daub J 2000 *J. Am. Chem. Soc.* **122** 968; (d) Kaur S and Kumar S 2002 *Chem. Commun.* 2840; (e) Roy B C, Chandra B, Hromas D and Mallik S 2003 *Org. Lett.* **5** 11; (f) Xiao Y and Qian X 2003 *Tetrahedron Lett.* **44** 2087; (g) Zheng Y, Orbulescu J, Ji X, Andreopoulos F M, Pham S M and Leblanc R M 2003 *J. Am. Chem. Soc.* **125** 2680; (h) Wu Q and Anslyn E V 2004 *J. Am. Chem. Soc.* **126** 14682
9. (a) Gunnlaugsson T, Leonard J P and Murray N S 2004 *Org. Lett.* **6** 1557; (b) Royzen M, Dai Z and Canary J W 2005 *J. Am. Chem. Soc.* **127** 1612; (c) Kaur N and Kumar S 2006 *Tetrahedron Lett.* **47** 4109; (d) Martinez R, Zapata F, Caballero A, Espinosa A, Tarraga A and Molina P 2006 *Org. Lett.* **8** 3235; (e) Mu H, Gong R, Ma Q, Sun Y and Fu E 2007 *Tetrahedron Lett.* **48** 5525; (f) Weng Y-Q, Yue F, Zhong Y-R and Ye B-H 2007 *Inorg. Chem.* **46** 7749
10. Kubo Y M, Ikeda Y M, Takeuchi M, Shinkai S, Yamaguchi S and Tamao K 2003 *Angew. Chem. Int. Ed.* **42** 2036; (b) Raker J and Glass T E 2002 *J. Org. Chem.* **67** 6113; (c) Fu H, Loo B H, Xiao D, Xie R, Ji X, Yao J, Zhang B and Zhang L 2002 *Angew. Chem. Int. Ed.* **41** 962; (d) Mello J V and Finney N S 2001 *Angew. Chem. Int. Ed.* **40** 1536; (e) Takakusa H, Kikuchi K, Urano Y, Sakamoto S, Yamaguchi K and Nagano T 2002 *J. Am. Chem. Soc.* **124** 1653
11. Gokel G W 1992 *Chem. Soc. Rev.* **21** 39
12. Beer P D 1992 *Adv. Inorg. Chem.* **39** 79
13. Jorgensen T, Hansen T K and Becher J 1994 *Chem. Soc. Rev.* **23** 41
14. Fabrizzi L and Poggi A 1995 *Chem. Soc. Rev.* 197
15. Kaifer A E and Mendoza S, 1996 **1** 701 In: *Comprehensive supramolecular chemistry*: G W Gokel (ed) (Oxford: Pergamon)
16. Boulas P L and Echegoyen M L 1998 *Angew. Chem. Int. Ed. Engl.* **37** 216
17. Beer P D and Gale P A 1998 *Adv. Phys. Org. Chem.* **31** 1
18. Beer P D 1998 *Acc. Chem. Res.* **31** 71
19. Beer P D, Gale P A and Chen G Z 1999 *Coord. Chem. Rev.* **3** 185
20. Beer P D and Gale P A 2001 *Angew. Chem. Int. Ed. Engl.* **40** 485
21. Itoh T, Shirakami S, Ishida N, Yamashita Y, Yoshida T, Kim H S and Wataya Y 2000 *Bioorg. Med. Chem. Lett.* **10** 1657
22. Yuasa H, Miyagawa N, Izumi T, Nakatani M, Izumi M and Hashimoto H 2004 *Org. Lett.* **6** 1489
23. Singhal N K, Ramanujam B, Mariappanadar V and Rao C P 2006 *Org. Lett.* **8** 3525
24. Xie J, Ménand M, Maisonneuve S and Métivier R 2007 *J. Org. Chem.* **72** 5980
25. Kumar R, Tiwari P, Maulik P R and Misra A K 2006 *Eur. J. Org. Chem.* 74
26. Thakur A, Adarsh N N, Chakraborty A, Devi M and Ghosh S 2010 *J. Organomet. Chem.* **695** 1059
27. Lloveras V, Caballero A, Tárraga A, Velasco M D, Espinosa A, Wurst K, Evans D J, Vidal-Gancedo J, Rovira C, Molina P and Veciana J 2005 *Eur. J. Inorg. Chem.* 2436
28. (a) López J L, Tárraga A, Espinosa A, Velasco M D, Molina P, Lloveras V, Vidal-Gancedo J, Rovira C, Veciana J, Evans D J and Wurst K 2004 *Chem. Eur. J.* **10** 1815; (b) Martínez R, Espinosa A, Tárraga A and Molina P 2005 *Org. Lett.* **7** 5869; (c) Zapata F, Caballero A, Espinosa A, Tárraga A and Molina P 2007 *Org. Lett.* **9** 2385