

## Review paper

# Integrating Nutrition and Immunology: a New Frontier

Fleur Ponton<sup>a\*</sup>, Kenneth Wilson<sup>b</sup>, Andrew J. Holmes<sup>c</sup>, Sheena C. Cotter<sup>d</sup>, David Raubenheimer<sup>e</sup>, and Stephen J. Simpson<sup>a</sup>

<sup>a</sup>School of Biological Sciences and the Charles Perkins Centre, The University of Sydney, NSW 2006, Australia

<sup>b</sup>Lancaster Environment Centre, Lancaster University, Lancaster, LA1 4YQ, UK

<sup>c</sup>School of Molecular Bioscience, The University of Sydney, NSW 2006, Australia

<sup>d</sup>School of Biological Sciences, Queen's University Belfast, BT9 7BL, UK.

<sup>e</sup>Institute of Natural Sciences, Massey University, Albany, New Zealand

\*Corresponding author

Dr Fleur Ponton

School of Biological Sciences

Heydon-Laurence Building, A08

University of Sydney

Sydney, New South Wales 2006

Phone: +61 2 9036 6262

Fax: +61 2 9351 4119

Email: [fleur\\_ponton@hotmail.com](mailto:fleur_ponton@hotmail.com)

Keywords : Geometric Framework; Gut; Immunity; Microbes; Nutrition; Parasites

## **Introduction**

A source of food and somewhere to live are basic requirements for every organism, and achieving these essentials involves interacting with other organisms. By far the majority of these interactions involve microorganisms, and throughout evolutionary history there has been strong selective pressure upon organisms to manage and control these interactions. As a result, key elements of the immune system emerged very early in evolution, including both induced and constitutive defences, allowing an array of complex and effective immune mechanisms (Hamilton et al., 2008; Vilmos and Kurucz, 1998). The function of the immune system is to regulate the full spectrum of interactions with microorganisms; not only the exclusion of organisms that are harmful (henceforth termed parasites) and the clearing of infections, but also limiting the cost of responding to organisms that can be tolerated and allowing (even encouraging) microbes that are beneficial. Collectively this means that immune mechanisms are complex and rely on a range of components that are triggered by different types of signals and may be regulated independently (Beckage, 2008; Forsman et al., 2008).

It has long been recognized that the immune response is modulated not only by host (and parasite) genetics, but also by host nutrition, (Lazzaro and Little, 2009; Schmid-Hempel, 2011), yet there remain important gaps in our knowledge. Gaining a fuller understanding of the interface between nutrition and immunity is particularly important for three reasons. First, immune function is affected by host nutrition, which may greatly affect the outcome of infection (Lazzaro and Little, 2009; Schmid-Hempel, 2011). Host nutrition influences both constitutive and inducible immune function, with consequences for morbidity and mortality (Cunningham-Rundles et al., 2005; Klasing, 2007) (Adams and Hewison, 2008; Amar et al., 2007; Calder, 2006; Cohen et al., 2008; Kelley and Bendich, 1996; Klasing, 2007; Kolb, 1997; Kristan, 2007; Samartin and Chandra, 2000; Sorci and Faivre, 2009). Second,

nutrition-based interactions are one of the major sources of microbial benefits to animals (Bäckhed et al., 2005; Douglas, 2010; Hooper et al., 2002; Kau et al., 2011; Topping and Clifton, 2001). Third, the host's nutrient digesting and absorbing organ, the gut, is home to the highest density of microbial cells – both beneficial and potentially harmful - and is thus the site of greatest intensity of microbe-animal interactions.

Nutrition is also a complex and multi-dimensional trait, and immunity and nutrition interact via multiple direct and indirect pathways, including the involvement of the host's endogenous microbiota (Chambers and Schneider, 2012; Ponton et al., 2011; Simpson and Raubenheimer, 2012). The challenge remains to capture these interactions and complexities to better understand nutritional immunology. In this review, several aspects to this complexity are explored. We first give an overview of the effects of nutritional state on response to microbes in invertebrates. We then propose a framework to measure the simultaneous and interactive effects of multiple food components on immune functions. To further characterize interactions between nutrition and immunity, we also describe how host nutrition can affect the dynamics of pathogen and mutualist populations, notably the gut microbiota. In each section, we detail findings from recent studies that highlight the importance of adopting an integrative and multi-dimensional approach to nutritional immunology.

## **1. Nutrition and the consequences of immune trade-offs**

A common concept in life history theory is that, when resources are limiting, organisms must balance some traits against one another. The idea that disease resistance is costly and is traded off against other traits, such as reproductive effort and longevity, is fundamental to the field of ecological immunology (e.g. (Lochmiller and Deerenberg, 2000; Owens and Wilson, 1999; Schulenburg et al., 2009; Sheldon and Verhulst, 1996; Wilson, 2005) Norris & Evans

2000; Wilson 2005). In order to test this hypothesis, immune-related costs must be experimentally distinguished from other pathological processes associated with infection. This internal competition for resources has been illustrated in workers of the bumblebee, *Bombus terrestris* (Moret and Schmid-Hempel, 2000). To generate distinct immune challenges on different nutritional states, fed or starved worker bees were injected with lipopolysaccharides to simulate bacterial presence or micro-latex beads to simulate macroparasite infection. The survival of challenged and control bees was then followed. Survival time was reduced for challenged workers that were starved, but not when they were well-fed. This implies that simply activating the immune system (no live microbe added) uses resources that would otherwise keep the animal alive, but when sufficient resources are available, hosts can compensate for this cost (Moret and Schmid-Hempel, 2000).

As in the previous example, starvation and energy restriction have typically been used to measure the effects of nutrition on immunity (Kristan, 2007; Murray and Murray, 1979). In insects, experimental studies have demonstrated that food deprivation of the host leads to reduced immune responsiveness (e.g. (Ayres and Schneider, 2009; Siva-Jothy and Thompson, 2002)). For example, short-term starvation resulted in decreased phenoloxidase activity in adult mealworm beetles (Siva-Jothy and Thompson, 2002) and increase mortality in *Rhodnius prolixus* bugs when challenged by bacteria (Feder et al., 1997). Nutrition not only affects host resistance to infection but also host tolerance. Disease tolerance is a defense strategy that reduces the negative impacts of the infection on host fitness without reducing the parasite load. The term employed here is different to immunological tolerance (i.e., the process by which the immune system fails to attack an antigen), but rather captures the idea that the costs of the infection can be reduced through reducing the damage caused by the infection and activation of the immune system on host tissues (Ayres and Schneider, 2012; Medzhitov et al., 2012). For example, Ayres and Schneider, (2009) found that during

*Salmonella* infections, food-restricted *Drosophila* and flies mutant for the gustatory receptor, *gr28b*, had similar levels of bacteria to wild-type individuals but they lived longer. This result suggested that resistance was unchanged but tolerance was increased.

At a genomic level, dietary restriction induces changes in the expression of several immune genes in *Drosophila* (Pletcher et al., 2005; Pletcher et al., 2002). Molecular studies of the interactions between metabolic pathways and innate immunity have provided a new understanding of the interactions between nutrition and immune defense in insects (Castillo et al., 2011; DiAngelo et al., 2009). Mutations of genes in the insulin signaling pathway have considerable effects on immunity. For example, (Libert et al., 2008) investigated the effect of the *chico* mutation on resistance of flies infected with either a gram-negative or a gram-positive bacterium (*Pseudomonas aeruginosa* and *Enterococcus faecalis*, respectively). *Chico* is an adaptor protein, homologous to vertebrate insulin receptor substrates (IRS). Flies homozygous for the *chico* mutation had superior pathogen resistance to that of wild-type controls and heterozygous siblings. (Becker et al., 2010) demonstrated that anti-microbial peptides (AMP) in non-infected flies are activated in response to nuclear FOXO activity. FOXO is a transcription factor playing a pivotal role in adapting metabolism to nutrient conditions and is one of the most evolutionarily ancient downstream effectors of the insulin-signaling pathway (Hay, 2011; Kapahi et al., 2010). Also, *in vivo* studies indicate that the FOXO-dependent regulation of AMPs is evolutionarily conserved (see also (Becker et al., 2010; Garsin et al., 2003; Troemel et al., 2006)). Becker et al. (2010) found that AMP expression was inhibited in *FOXO* null mutants but enhanced when FOXO was over-expressed. FOXO can directly induce the expression of the peptide by binding to the regulatory region of one of the AMP promoters (i.e., *Drosomycin*). FOXO interacts with TOR and AMPK (Hay, 2011). Target of rapamycin (TOR) and AMP-Activated Protein Kinase (AMPK) integrate information on cellular nutritional status by sensing both

qualitative and quantitative changes in nutrients, particularly branch-chain amino acids and glucose (Kapahi et al., 2010; Simpson and Raubenheimer, 2009). Disruption of FOXO activity can have different effects on host survival depending on the nature of the infection (Dionne et al., 2006), which probably reflects the fact that infections by different types of pathogens can trigger different immune pathways (Hoffmann and Reichhart, 2002; Hultmark, 2003; Lemaitre et al., 1997).

Such advances in our understanding at the transcriptional level indicate that the interaction between nutrition and immune function is mediated by nutrient signaling pathways that involve more than the monitoring of energy status, but instead monitor specific nutrients and metabolites. However, starvation and food deprivation protocols do not usually consider the nutritional composition of experimental foods, or include consideration of the animal's multiple nutritional needs. Identifying the nutrients and, critically, the interactions, that modulate immunity remain central challenges for nutritional immunology (Ponton et al., 2011).

## **2. Nutritional immunology: taking a multi-dimensional approach**

Recent experiments have explored the single and interactive effects of nutrients in the diet on immune function, using experimental designs derived from nutritional geometry (Raubenheimer and Simpson, 1993; Simpson and Raubenheimer, 1993, 2012). In an initial study, Lee et al. (2006) (Lee et al., 2006) measured the effects of the dietary ratio of protein to digestible carbohydrate (P:C) on *Spodoptera littoralis* caterpillars infected with a nucleopolyhedrovirus (NPV). Susceptibility to NPV infection decreased as dietary P:C rose. In contrast, the performance of control insects peaked on an intermediate P:C diet (Figure 1). Insects on high-P:C diets had significantly higher levels of constitutive immune function (i.e., antimicrobial activity, encapsulation capacity and total haemocyte count) than those on low-

P:C diets. When insects were allowed to self-compose their diet, the ones that survived the viral challenge had demonstrated an increased consumption of protein compared with uninfected controls and those dying of infection. Povey et al. 2009 (Povey et al., 2009) found similar results for the African armyworm, *Spodoptera exempta*, infected by the bacterium *Bacillus subtilis*. In addition, as dietary P:C increased, antimicrobial activity, phenoloxidase activity and protein levels in the haemolymph also increased. Larvae injected with a sub-lethal dose of bacteria increased their protein intake relative to control in a self-selection test. The results of Lee et al. (2006) and Povey et al. (2009) indicate that dietary protein is a key nutritional component affecting immunity (see also (Alaux et al., 2010; Fellous and Lazzaro, 2010; Peck et al., 1992), and that caterpillars are able to self-medicate for infection by selecting a dietary composition that best supports immune defense (see also (Raubenheimer and Simpson, 2009; Singer et al., 2009).

Recent advances in functional genomics and molecular biology have greatly expanded our understanding of the details of the immune mechanisms that enable insects to defend themselves against parasites (Siva-Jothy et al., 2005; Welchman et al., 2009), and have emphasized the complexity of innate immunity (see also (Hergannan and Rechhart, 1997; Lemaitre and Hoffmann, 2007; Lemaitre et al., 1997). Key questions now are whether these different component mechanisms share similar or different nutritional requirements, and whether they compete for limiting host-derived resources (Cotter et al., 2004; Moret and Schmid-Hempel, 2001).

Povey et al. (2009) found that as the ratio of protein to carbohydrate in the diet increased, caterpillars had elevated their antibacterial activity but reduced their phenoloxidase activity, suggesting a physiological trade-off between these immune traits. However, an alternative explanation, that the traits simply have different nutritional optima, cannot be excluded from these experiments. That immune components do indeed differ in their

nutritional requirements was demonstrated by Cotter et al. (2011) in caterpillars of *S. littoralis* fed one of 20 diets varying in the ratio and amounts of protein and carbohydrate (Cotter et al., 2011). Nutrient-mediated effects on several immune traits were visualized as response surfaces mapped onto nutrient intake arrays for challenged and unchallenged insects. Immune traits were all affected by dietary macronutrient composition (some more so than others), but immune traits peaked at different locations on the nutritional landscape (Figure 2). Hence, no single diet simultaneously optimized all the immune components. It follows that the insect could potentially adjust its dietary choices to achieve a nutrient balance that best meets a particular immune challenge (Cotter et al. 2011).

### **3. Nutritional interactions between hosts, parasites and mutualists**

Hosts are not the only ones facing nutritional challenges. Parasites feed on their host by either hijacking food or feeding on the host's tissues and fluids. The host can, therefore, effectively be considered as a parasite growth medium, with nutrient supply influencing not only within-host pathogen population dynamics, but also the degree of pathogenicity of the infection through the competition for resources with the host. The host could, in turn, adjust its acquisition of nutrients to alleviate resource competition with parasites and to accommodate the extra nutritional demands of fighting the infection (see above) - if the nutritional environment allows (Bedhomme et al., 2004; De Roode et al., 2008; Ebert et al., 2000; J. Ryder et al., 2007; Nesheim et al., 1978; Smith and Holt, 1996). Because the nutritional requirements of hosts and parasites are likely not to be the same, discovering whether hosts can compensate for infection by altered feeding behaviour may require experimental protocols with more than one dietary treatment.

In a protocol using multiple nutritional treatments, Ponton et al. (2011) showed that mealworm beetles, *Tenebrio molitor*, modify their food intake when infected by cysticercoids



of the tapeworm *Hymenolepis diminuta*. More particularly, infected insects increased their carbohydrate consumption during the first few days post-infection, i.e. when the parasites are growing and developing into mature forms. Despite consuming more nutrients, infected individuals deposited less body lipid and were less efficient at converting ingested protein to growth. However, infected insects sustained high levels of reproductive output despite the infection, unless confined to foods that were nutritionally dilute. Furthermore, there was no indication that increased carbohydrate intake promoted host immunity. We might then conclude from these results that beetles modified their feeding behaviour to ameliorate the nutritional demands of the infection.

### **3.2. Microbiota: a key component of nutritional immunology**

Interpreting nutritional interactions between hosts and parasites is made significantly more complex by the normal microbial communities inhabiting the host. The normal microbiota encompasses a spectrum of lifestyles from commensalism (i.e., one partner benefits and the other is apparently unaffected) to mutualism (i.e., both partners experience increased fitness). They all receive their nutrition from the host, but may vary in their contribution of nutrients that are integral to host physiology and ecological adaptations (Brune and Ohkuma, 2011; Douglas, 2010). Obligate or primary symbionts (i.e., intracellular mutualists) are vertically-transmitted between hosts and are essential for host survival in resource-limited environments (see for instance (Douglas, 1998)). Such obligate mutualistic relationships are typically ancient and neither of the partners can survive in the absence of the other. However, most animal-microbial interactions are flexible and facultative, and it is likely that all animals are associated with a complex and ever-changing microbial community that consists predominantly of non-pathogenic, horizontally-acquired bacteria (i.e., facultative or secondary symbionts). The digestive tract of metazoans is particularly rich in such facultative

microbes, where their activity may influence nutrient quality and absorption, as well as immunological challenge. The significance of understanding this is best illustrated in the vertebrates.

The gut microbiota of vertebrates are of particularly high density ( $>10^{11}$  cells/ml) and diversity ( $>1000$  species) (Ley et al., 2008). The metabolic activity of this extensive microbial community is comparable to an organ such as the liver and, via conversion of polysaccharides to short chain fatty acids (SCFAs), it directly contributes up to 70% of a vertebrate herbivore's SCFAs needs. A key point is that in vertebrates this nutritional benefit is an emergent property of the activity of the total microbiota, rather than a benefit derived from one or two primary symbionts. Furthermore, the vertebrate gut microbiota are involved in many other aspects of host health and development (Bäckhed et al., 2005; Kau et al., 2011; Shin et al., 2011; Turnbaugh et al., 2008; Vijay-Kumar et al., 2010). Analyses of commensal host-microbial relationships in the intestine of mammalian models have identified microbial roles in the regulation of genes in many host systems, including development, differentiation, immunity and metabolism (Backhed et al., 2004; Hooper et al., 2001). Thus, in vertebrates, the relationship between immunity and nutrition is remarkably intertwined. The presence of microbes is required for many aspects of the immune system to develop at all. Which microbes are present determines the state of the vertebrate immune system by strongly influencing T-cell maturation patterns. The composition of the microbiota is in turn strongly influenced by long-term diet patterns. The increased incidence of modern lifestyle diseases such as allergies, diabetes, inflammatory bowel diseases and obesity to epidemic levels strongly links changes in nutrition, gut microbiota and immune function.(Backhed et al., 2004; Ley et al., 2006; Noverr and Huffnagle, 2004; Vijay-Kumar et al., 2010; Wen et al., 2008).

Gut microbes also play a role in invertebrate biology (Dillon and Dillon, 2003) and digestive process [Brune, 2011 #1962][Lundgren, 2010 #1980], and recently the composition of microbe gut populations has been described in a variety of insect species, including bees (Jeyaprakash et al., 2003; Mohr and Tebbe, 2006), beetles (Egert et al., 2005; Lehman et al., 2009; Nardi et al., 2006; Zhang and Jackson, 2008), flies (Cox and Gilmore, 2007; Ren et al., 2007; Ryu et al., 2008; Shin et al., 2011; Wong et al., 2011), lepidopterans (Pauchet et al., 2010; Xiang et al., 2006) and termites (Hongoh et al., 2003). In *Drosophila*, the microbiome regulates host metabolic homeostatic and developmental programs by modulating the insulin/insulin-like growth factor (Shin et al., 2011).

Gut microbiota are also key to the infection process itself (Boissière et al., 2012; Borriello, 1990; Broderick et al., 2006; Charroux and Royet, 2012; Cirimotich et al., 2011; Harp et al., 1992; Weiss and Aksoy, 2011; Wilks and Golovkina, 2012). Microbes from the gut can directly interact with parasites through the secretion of inhibitory compounds. Alternatively, the gut microbiota can indirectly affect the development and persistence of parasites by inducing the host's immune system (Buchon et al., 2009; Douglas, 2010; Feldhaar and Gross, 2008; Kau et al., 2011; Lazzaro and Little, 2009; Ryu et al., 2010; Ryu et al., 2008; Wen et al., 2008). For instance, in mosquitoes, commensals can modulate *Plasmodium* infection (Cirimotich et al., 2011; Gonzalez-Ceron et al., 2003; Meister et al., 2009; Pumpuni et al., 1996). Gut bacteria within the mosquito interfere with *Plasmodium* development before invasion of the midgut epithelium, by stimulating the production of basal levels of effector molecules that control the proliferation of the bacterial populations as well as *Plasmodium* populations (Dong et al., 2009). Global transcription profiling of germ-free mosquitoes identified a subset of immune genes that were mostly down-regulated, including several anti-*Plasmodium* factors (Dong et al., 2009). In flies, gut microbiota modulate the immune system, and hence presumably susceptibility to invading parasites, by activating the

Imd pathway transcription factor Relish (Ryu et al., 2008), which triggers the production of AMPs (Feldhaar and Gross, 2008; Ryu et al., 2008).

Commensal bacterial populations can vary greatly in their persistence, abundance and species composition within the host gut, with a major determinant being host diet composition, notably the macronutrient balance (Faith et al., 2011). (Chandler et al., 2011) assessed the importance of host diet and host species in shaping microbiome composition in flies. They showed that whereas taxonomically- and geographically-distant fly populations, collected from various food sources, have very different microbiome compositions, when maintained on the same type of food they developed similar microbiomes. Diet has also been shown to influence the bacterial community in the midgut of larval gypsy moths, *Lymantria dispar* (Broderick 2004) and cotton bollworms, *Helicoverpa armigera* (Xiang et al., 2006). The reasons why host diet has a strong impact on the gut microbial composition are still not well understood (De Filippo et al., 2010; Muegge et al., 2011; Turnbaugh et al., 2009), but presumably reflects a combination of influences on the physical and chemical milieu of the gut (Clissold et al., 2010; Duncan et al., 2008; Faith et al., 2011; Flint et al., 2008; Ley et al., 2008; Sorensen et al., 2010), and effects on immune responses (see above). Also, the diet itself is a vector of commensals, and different diets will provide microbial inoculates of different community compositions. Defining the relationships between diet and the composition and function of the gut microbiome is fundamental to a better understanding the effects of nutrition on immunity and the outcome of host-pathogens interactions.

## **Conclusions**

Unravelling the inter-relationship between host nutrition, host immune function, pathogen population growth and the structure and function of the gut microbiome is essential to predicting the outcome of parasitic infections (Figure 3). Ecological immunology has been

underpinned by the concept of nutrition-dependent condition, with nutrition influencing immunity, resistance and tolerance to pathogens. Geometric nutritional designs offer a powerful yet tractable approach for studying these interactions, allowing quantitative predictions about the consequences of nutrition on immunity, health and disease. Insects and their pathogens show great promise as model systems in the study of the relationships between nutrition, innate immunity and gut microbiota. They are experimentally amenable to large-scale dietary studies (see for instance (Lee et al., 2008)), in certain cases offer substantial molecular genetic resources (Chambers and Schneider, 2012), and have an homologous yet simpler immune system to vertebrates. In particular, insect models do not have the disadvantage of confounding effects due to individual differences in adaptive immune response. Insects possess relatively simple microbial communities, which aids the quantification and manipulation of microbiota. In addition, recent findings concerning *D. melanogaster* intestinal pathology suggest that this model is well suited as a model for the study of intestinal physiology during aging, stress and infection (Apidianakis and Rahme, 2011). With the advent of nutritional genomics (Afacan et al., 2012; Becker et al., 2010; Fellous and Lazzaro, 2010; Grayson, 2010), opportunities exist now to explore the interaction between nutrients and gene expression and their products to determine the mechanism behind disease development. This will provide significant insights into nutritional regulation of the innate immune system, the gut microbiota and pathogenesis.

## References

- Adams, J.S., Hewison, M., 2008. Unexpected actions of vitamin D: new perspectives on the regulation of innate and adaptive immunity. *Nat Clin Pract End Met* 4, 80-90.
- Afacan, N.J., Fjell, C.D., Hancock, R.E.W., 2012. A systems biology approach to nutritional immunology – Focus on innate immunity. *Molecular Aspects of Medicine* 33, 14-25.
- Alaux, C., Ducloz, F., Crauser, D., Le Conte, Y., 2010. Diet effects on honeybee immunocompetence. *Biology Letters* 6, 562-565.
- Amar, S., Zhou, Q., Shaik-Dasthagirisaheb, Y., Leeman, S., 2007. Diet-induced obesity in mice causes changes in immune responses and bone loss manifested by bacterial challenge. *Proceedings of the National Academy of Sciences* 104, 20466-20471.
- Apidianakis, Y., Rahme, L.G., 2011. *Drosophila melanogaster* as a model for human intestinal infection and pathology. *Disease Models & Mechanisms* 4, 21-30.
- Ayres, J.S., Schneider, D.S., 2009. The Role of Anorexia in Resistance and Tolerance to Infections in *Drosophila*. *PLoS Biol* 7, e1000150.
- Ayres, J.S., Schneider, D.S., 2012. Tolerance of infections. *Annu Rev Immunol* 30, 271-294.
- Backhed, F., Ding, H., Wang, T., Hooper, L.V., Koh, G.Y., Nagy, A., Semenkovich, C.F., Gordon, J.I., 2004. The gut microbiota as an environmental factor that regulates fat storage. *PNAS* 101, 15718-15723.
- Bäckhed, F., Ley, R.E., Sonnenburg, J.L., Peterson, D.A., Gordon, J.I., 2005. Host-Bacterial Mutualism in the Human Intestine. *Science* 307, 1915-1920.
- Beckage, N.E., 2008. *Insect Immunology*. Academic Press, San Diego.
- Becker, T., Loch, G., Beyer, M., Zinke, I., Aschenbrenner, A.C., Carrera, P., Inhester, T., Schultze, J.L., Hoch, M., 2010. FOXO-dependent regulation of innate immune homeostasis. *Nature* 463, 369-373.
- Bedhomme, S., Agnew, P., Sidobre, C., Michalakis, Y., 2004. Virulence reaction norms across a food gradient. *Proc Biol Sci* 271, 739-744.
- Boissière, A., Tchioffo, M.T., Bachar, D., Abate, L., Marie, A., Nsango, S.E., Shahbazkia, H.R., Awono-Ambene, P.H., Levashina, E.A., Christen, R., Morlais, I., 2012. Midgut Microbiota of the Malaria Mosquito Vector *Anopheles gambiae* and Interactions with *Plasmodium falciparum* Infection. *PLoS Pathog* 8, e1002742.
- Borriello, S.P., 1990. The influence of the normal flora on *Clostridium difficile* colonisation of the gut. *Ann Med* 22, 61-67.
- Broderick, N.A., Raffa, K.F., Handelsman, J., 2006. Midgut bacteria required for *Bacillus thuringiensis* insecticidal activity. *Proc Natl Acad Sci U S A* 103, 15196-15199.
- Brune, A., Ohkuma, M., 2011. Role of the Termite Gut Microbiota in Symbiotic Digestion

Biology of Termites: a Modern Synthesis, in: Bignell, D.E., Roisin, Y., Lo, N. (Eds.). Springer Netherlands, pp. 439-475.

Buchon, N., Broderick, N.A., Poidevin, M., Pradervand, S., Lemaitre, B., 2009. *Drosophila* Intestinal Response to Bacterial Infection: Activation of Host Defense and Stem Cell Proliferation. *Cell Host & Microbe* 5, 200-211.

Calder, P.C., 2006. n-3 Polyunsaturated fatty acids, inflammation, and inflammatory diseases. *The American Journal of Clinical Nutrition* 83, S1505-S1519S.

Castillo, J., Brown, M.R., Strand, M.R., 2011. Blood Feeding and Insulin-like Peptide 3 Stimulate Proliferation of Hemocytes in the Mosquito *Aedes aegypti*. *PLoS Pathog* 7, e1002274.

Chambers, M.C., Schneider, D.S., 2012. Pioneering immunology: insect style. *Current Opinion in Immunology* 24, 10-14.

Chandler, J.A., Morgan Lang, J., Bhatnagar, S., Eisen, J.A., Kopp, A., 2011. Bacterial Communities of Diverse *Drosophila* Species: Ecological Context of a Host-Microbe Model System. *PLoS Genet* 7, e1002272.

Charroux, B., Royet, J., 2012. Gut-microbiota interactions in non-mammals: What can we learn from *Drosophila*? *Seminars in Immunology* 24, 17-24.

Cirimotich, Chris M., Ramirez, Jose L., Dimopoulos, G., 2011. Native Microbiota Shape Insect Vector Competence for Human Pathogens. *Cell Host & Microbe* 10, 307-310.

Clissold, F.J., Tedder, B.J., Conigrave, A.D., Simpson, S.J., 2010. The gastrointestinal tract as a nutrient-balancing organ. *Proceedings of the Royal Society B: Biological Sciences*, -.

Cohen, Alan A., McGraw, Kevin J., Wiersma, P., Williams, Joseph B., Robinson, W.D., Robinson, Tara R., Brawn, Jeffrey D., Ricklefs, Robert E., 2008. Interspecific Associations between Circulating Antioxidant Levels and Life-History Variation in Birds. *The American Naturalist* 172, 178-193.

Cotter, S.C., Kruuk, L.E., Wilson, K., 2004. Costs of resistance: genetic correlations and potential trade-offs in an insect immune system. *J Evol Biol* 17, 421-429.

Cotter, S.C., Simpson, S.J., Raubenheimer, D., Wilson, K., 2011. Macronutrient balance mediates trade-offs between immune function and life history traits. *Functional Ecology* 25, 186-198.

Cox, C.R., Gilmore, M.S., 2007. Native Microbial Colonization of *Drosophila melanogaster* and Its Use as a Model of *Enterococcus faecalis* Pathogenesis. *Infect. Immun.* 75, 1565-1576.

Cunningham-Rundles, S., McNeeley, D.F., Moon, A., 2005. Mechanisms of nutrient modulation of the immune response. *Journal of Allergy and Clinical Immunology* 115, 1119-1128.

De Filippo, C., Cavalieri, D., Di Paola, M., Ramazzotti, M., Poullet, J.B., Massart, S., Collini, S., Pieraccini, G., Lionetti, P., 2010. Impact of diet in shaping gut microbiota revealed by a comparative study in children from Europe and rural Africa. *Proceedings of the National Academy of Sciences* 107, 14691-14696.

De Roode, J.C., Pedersen, A.B., Hunter, M.D., Altizer, S., 2008. Host plant species affects virulence in monarch butterfly parasites. *Journal of Animal Ecology* 77, 120-126.

DiAngelo, J.R., Bland, M.L., Bambina, S., Cherry, S., Birnbaum, M.J., 2009. The immune response attenuates growth and nutrient storage in *Drosophila* by reducing insulin signaling. *Proceedings of the National Academy of Sciences* 106, 20853-20858.

Dillon, R.J., Dillon, V.M., 2003. THE GUT BACTERIA OF INSECTS: Nonpathogenic Interactions. *Annual Review of Entomology* 49, 71-92.

Dionne, M.S., Pham, L.N., Shirasu-Hiza, M., Schneider, D.S., 2006. Akt and foxo Dysregulation Contribute to Infection-Induced Wasting in *Drosophila*. *Current Biology* 16, 1977-1985.

Dong, Y., Manfredini, F., Dimopoulos, G., 2009. Implication of the Mosquito Midgut Microbiota in the Defense against Malaria Parasites. *PLoS Pathog* 5, e1000423.

Douglas, A.E., 1998. Nutritional Interactions in Insect-Microbial Symbioses: Aphids and Their Symbiotic Bacteria *Buchnera*. *Annual Review of Entomology* 43, 17-37.

Douglas, A.E., 2010. *The symbiotic Habit*. Princeton University Press New Jersey

Duncan, S.H., Loble, G.E., Holtrop, G., Ince, J., Johnstone, A.M., Louis, P., Flint, H.J., 2008. Human colonic microbiota associated with diet, obesity and weight loss. *Int J Obes* 32, 1720-1724.

Ebert, D., Zschokke-Rohringer, C.D., Carius, H.J., 2000. Dose effects and density-dependent regulation of two microparasites of *Daphnia magna*. *Oecologia* 122, 200-209.

Egert, M., Stingl, U., Dyhrberg Bruun, L., Pommerenke, B., Brune, A., Friedrich, M.W., 2005. Structure and Topology of Microbial Communities in the Major Gut Compartments of *Melolontha melolontha* Larvae (Coleoptera: Scarabaeidae). *Applied and Environmental Microbiology* 71, 4556-4566.

Faith, J.J., McNulty, N.P., Rey, F.E., Gordon, J.I., 2011. Predicting a Human Gut Microbiota's Response to Diet in Gnotobiotic Mice. *Science* 333, 101-104.

Feder, D., Mello, C.B., Garcia, E.S., Azambuja, P., 1997. Immune responses in *Rhodnius prolixus*: influence of nutrition and ecdysone. *Journal of Insect Physiology* 43, 513-519.

Feldhaar, H., Gross, R., 2008. Immune reactions of insects on bacterial pathogens and mutualists. *Microbes Infect* 10, 1082-1088.

Fellous, S., Lazzaro, B.P., 2010. Larval food quality affects adult (but not larval) immune gene expression independent of effects on general condition. *Molecular Ecology* 19, 1462-1468.

Flajnik, M.F., Du Pasquier, L., 2004. Evolution of innate and adaptive immunity: can we draw a line? *Trends in Immunology* 25, 640-644.

Flint, H.J., Bayer, E.A., Rincon, M.T., Lamed, R., White, B.A., 2008. Polysaccharide utilization by gut bacteria: potential for new insights from genomic analysis. *Nat Rev Micro* 6, 121-131.

Forsman, A.M., Vogel, L.A., Sakaluk, S.K., Grindstaff, J.L., Thompson, C.F., 2008. Immune-challenged house wren broods differ in the relative strengths of their responses among different axes of the immune system. *Journal of Evolutionary Biology* 21, 873-878.



Garsin, D.A., Villanueva, J.M., Begun, J., Kim, D.H., Sifri, C.D., Calderwood, S.B., Ruvkun, G., Ausubel, F.M., 2003. Long-Lived *C. elegans* *daf-2* Mutants Are Resistant to Bacterial Pathogens. *Science* 300, 1921.

Gonzalez-Ceron, L., Santillan, F., Rodriguez, M.H., Mendez, D., Hernandez-Avila, J.E., 2003. Bacteria in Midguts of Field-Collected *Anopheles albimanus* Block *Plasmodium vivax* Sporogonic Development. *Journal of Medical Entomology* 40, 371-374.

Grayson, M., 2010. Nutrigenomics. *Nature* 468, S1-S1.

Hamilton, R., Siva-Jothy, M., Boots, M., 2008. Two arms are better than one: parasite variation leads to combined inducible and constitutive innate immune responses. *Proceedings of the Royal Society B: Biological Sciences* 275, 937-945.

Harp, J.A., Chen, W., Harmsen, A.G., 1992. Resistance of severe combined immunodeficient mice to infection with *Cryptosporidium parvum*: the importance of intestinal microflora. *Infect Immun* 60, 3509-3512.

Hay, N., 2011. Interplay between FOXO, TOR, and Akt. *Biochimica et Biophysica Acta (BBA) - Molecular Cell Research* 1813, 1965-1970.

Hergannan, J.A., Reichhart, J.-V., 1997. *Drosophila* immunity. *Trends in Cell Biology* 7, 309-316.

Hoffmann, J.A., Reichhart, J.-M., 2002. *Drosophila* innate immunity: an evolutionary perspective. *Nat Immunol* 3, 121-126.

Hongoh, Y., Ohkuma, M., Kudo, T., 2003. Molecular analysis of bacterial microbiota in the gut of the termite *Reticulitermes speratus* (Isoptera; Rhinotermitidae). *FEMS Microbiology Ecology* 44, 231-242.

Hooper, L.V., Midtvedt, T., Gordon, J.I., 2002. HOW HOST-MICROBIAL INTERACTIONS SHAPE THE NUTRIENT ENVIRONMENT OF THE MAMMALIAN INTESTINE. *Annual Review of Nutrition* 22, 283-307.

Hooper, L.V., Wong, M.H., Thelin, A., Hansson, L., Falk, P.G., Gordon, J.I., 2001. Molecular Analysis of Commensal Host-Microbial Relationships in the Intestine. *Science* 291, 881-884.

Hultmark, D., 2003. *Drosophila* immunity: paths and patterns. *Current Opinion in Immunology* 15, 12-19.

J. Ryder, J., Hathway, J., J. Knell, R., 2007. Constraints on parasite fecundity and transmission in an insect-STD system. *Oikos* 116, 578-584.

Jeyaprakash, A., Hoy, M.A., Allsopp, M.H., 2003. Bacterial diversity in worker adults of *Apis mellifera capensis* and *Apis mellifera scutellata* (Insecta: Hymenoptera) assessed using 16S rRNA sequences. *Journal of Invertebrate Pathology* 84, 96-103.

Kapahi, P., Chen, D., Rogers, A.N., Katewa, S.D., Li, P.W., Thomas, E.L., Kockel, L., 2010. With TOR, less is more: a key role for the conserved nutrient-sensing TOR pathway in aging. *Cell Metab* 11, 453-465.

Kau, A.L., Ahern, P.P., Griffin, N.W., Goodman, A.L., Gordon, J.I., 2011. Human nutrition, the gut microbiome and the immune system. *Nature* 474, 327-336.

Kelley, D., Bendich, A., 1996. Essential nutrients and immunologic functions. *The American Journal of Clinical Nutrition* 63, 994S-996S.

Klasing, K.C., 2007. Nutrition and the immune system. *British Poultry Science* 48, 525-537.

Kolb, E., 1997. Vitamins and the immune system. 1-73.

Kristan, D.M., 2007. Chronic calorie restriction increases susceptibility of laboratory mice (*Mus musculus*) to a primary intestinal parasite infection. *Aging Cell* 6, 817-825.

Lazzaro, B.P., Little, T.J., 2009. Immunity in a variable world. *Philosophical Transactions of the Royal Society B-Biological Sciences* 364, 15-26.

Lee, K.P., Cory, J.S., Wilson, K., Raubenheimer, D., Simpson, S.J., 2006. Flexible diet choice offsets protein costs of pathogen resistance in a caterpillar. *Proc Biol Sci* 273, 823-829.

Lee, K.P., Simpson, S.J., Clissold, F.J., Brooks, R., Ballard, J.W., Taylor, P.W., Soran, N., Raubenheimer, D., 2008. Lifespan and reproduction in *Drosophila*: New insights from nutritional geometry. *Proc Natl Acad Sci U S A* 105, 2498-2503.

Lehman, R., Lundgren, J., Petzke, L., 2009. Bacterial Communities Associated with the Digestive Tract of the Predatory Ground Beetle, *Poecilus chalcites*, and Their Modification by Laboratory Rearing and Antibiotic Treatment. *Microb. Ecol.* 57, 349-358.

Lemaitre, B., Hoffmann, J., 2007. The host defense of *Drosophila melanogaster*. *Annu Rev Immunol* 25, 697-743.

Lemaitre, B., Reichhart, J.-M., Hoffmann, J.A., 1997. *Drosophila* host defense: Differential induction of antimicrobial peptide genes after infection by various classes of microorganisms. *Proceedings of the National Academy of Sciences of the United States of America* 94, 14614-14619.

Ley, R.E., Hamady, M., Lozupone, C., Turnbaugh, P.J., Ramey, R.R., Bircher, J.S., Schlegel, M.L., Tucker, T.A., Schrenzel, M.D., Knight, R., Gordon, J.I., 2008. Evolution of mammals and their gut microbes. *Science* 320, 1647-1651.

Ley, R.E., Turnbaugh, P.J., Klein, S., Gordon, J.I., 2006. Microbial ecology: human gut microbes associated with obesity. *Nature* 444, 1022-1023.

Libert, S., Chao, Y., Zwiener, J., Pletcher, S.D., 2008. Realized immune response is enhanced in long-lived *puc* and *chico* mutants but is unaffected by dietary restriction. *Molecular Immunology* 45, 810-817.

Lochmiller, R.L., Deerenberg, C., 2000. Trade-Offs in Evolutionary Immunology: Just What Is the Cost of Immunity? *Oikos* 88, 87-98.

Medzhitov, R., Schneider, D.S., Soares, M.P., 2012. Disease Tolerance as a Defense Strategy. *Science* 335, 936-941.

Meister, S., Agianian, B., Turlure, F., Relógio, A., Morlais, I., Kafatos, F.C., Christophides, G.K., 2009. *Anopheles gambiae* PGRPLC-Mediated Defense against Bacteria Modulates Infections with Malaria Parasites. PLoS Pathog 5, e1000542.

Mohr, K.I., Tebbe, C.C., 2006. Diversity and phylotype consistency of bacteria in the guts of three bee species (Apoidea) at an oilseed rape field. Environmental Microbiology 8, 258-272.

Moret, Y., Schmid-Hempel, P., 2000. Survival for Immunity: The Price of Immune System Activation for Bumblebee Workers. Science 290, 1166-1168.

Moret, Y., Schmid-Hempel, P., 2001. Entomology: Immune defence in bumble-bee offspring. Nature 414, 506-506.

Muegge, B.D., Kuczynski, J., Knights, D., Clemente, J.C., González, A., Fontana, L., Henrissat, B., Knight, R., Gordon, J.I., 2011. Diet Drives Convergence in Gut Microbiome Functions Across Mammalian Phylogeny and Within Humans. Science 332, 970-974.

Murray, M.J., Murray, A.B., 1979. Anorexia of infection as a mechanism of host defense. The American Journal of Clinical Nutrition 32, 593-596.

Nardi, J.B., Bee, C.M., Miller, L.A., Nguyen, N.H., Suh, S.-O., Blackwell, M., 2006. Communities of microbes that inhabit the changing hindgut landscape of a subsocial beetle. Arthropod Structure & Development 35, 57-68.

Nesheim, M.C., Crompton, D.W.T., Arnold, S., Barnard, D., 1978. Host Dietary Starch and Moniliformis (Acanthocephala) in Growing Rats. Proceedings of the Royal Society of London. Series B. Biological Sciences 202, 399-408.

Noverr, M.C., Huffnagle, G.B., 2004. Does the microbiota regulate immune responses outside the gut? Trends in Microbiology 12, 562-568.

Owens, I.P.F., Wilson, K., 1999. Immunocompetence: a neglected life history trait or conspicuous red herring? Trends in Ecology & Evolution 14, 170-172.

Pauchet, Y., Wilkinson, P., Vogel, H., Nelson, D.R., Reynolds, S.E., Heckel, D.G., Ffrench-Constant, R.H., 2010. Pyrosequencing the Manduca sexta larval midgut transcriptome: messages for digestion, detoxification and defence. Insect Molecular Biology 19, 61-75.

Peck, M.D., Babcock, G.F., Alexander, J.W., 1992. The Role of Protein and Calorie Restriction in Outcome From Salmonella Infection in Mice. Journal of Parenteral and Enteral Nutrition 16, 561-565.

Pletcher, S.D., Libert, S., Skorupa, D., 2005. Flies and their Golden Apples: The effect of dietary restriction on Drosophila aging and age-dependent gene expression. Ageing Research Reviews 4, 451-480.

Pletcher, S.D., Macdonald, S.J., Marguerie, R., Certa, U., Stearns, S.C., Goldstein, D.B., Partridge, L., 2002. Genome-Wide Transcript Profiles in Aging and Calorically Restricted Drosophila melanogaster. Current Biology 12, 712-723.

Ponton, F., Wilson, K., Cotter, S.C., Raubenheimer, D., Simpson, S.J., 2011. Nutritional immunology: a multi-dimensional approach. PLoS Pathog 7, e1002223.

Povey, S., Cotter, S.C., Simpson, S.J., Lee, K.P., Wilson, K., 2009. Can the protein costs of bacterial resistance be offset by altered feeding behaviour? *J Anim Ecol* 78, 437-446.

Pumpuni, C.B., Demaio, J., Kent, M., Davis, J.R., Beier, J.C., 1996. Bacterial population dynamics in three anopheline species: The impact on Plasmodium sporogonic development. *American Journal of Tropical Medicine and Hygiene* 54, 214-218.

Raubenheimer, D., Simpson, S.J., 1993. The Geometry of Compensatory Feeding in the Locust. *Animal Behaviour* 45, 953-964.

Raubenheimer, D., Simpson, S.J., 2009. Nutritional PharmEcology: Doses, nutrients, toxins, and medicines. *Integr. Comp. Biol.* 49, 329-337.

Ren, C., Webster, P., Finkel, S.E., Tower, J., 2007. Increased Internal and External Bacterial Load during Drosophila Aging without Life-Span Trade-Off. *Cell Metabolism* 6, 144-152.

Ryu, J.-H., Ha, E.-M., Lee, W.-J., 2010. Innate immunity and gut-microbe mutualism in Drosophila. *Developmental & Comparative Immunology* 34, 369-376.

Ryu, J.-H., Kim, S.-H., Lee, H.-Y., Bai, J.Y., Nam, Y.-D., Bae, J.-W., Lee, D.G., Shin, S.C., Ha, E.-M., Lee, W.-J., 2008. Innate Immune Homeostasis by the Homeobox Gene Caudal and Commensal-Gut Mutualism in Drosophila. *Science* 319, 777-782.

Samartin, S., Chandra, R.K., 2000. Obesity, overnutrition and the immune system. *Nutrition Research* 21, 243-262.

Schmid-Hempel, P., 2011. *Evolutionary Parasitology*. Oxford University Press, Oxford.

Schulenburg, H., Kurtz, J., Moret, Y., Siva-Jothy, M.T., 2009. Introduction. *Ecological immunology*. *Philosophical Transactions of the Royal Society B: Biological Sciences* 364, 3-14.

Sheldon, B.C., Verhulst, S., 1996. Ecological immunology: costly parasite defences and trade-offs in evolutionary ecology. *Trends in Ecology & Evolution* 11, 317-321.

Shin, S.C., Kim, S.-H., You, H., Kim, B., Kim, A.C., Lee, K.-A., Yoon, J.-H., Ryu, J.-H., Lee, W.-J., 2011. Drosophila Microbiome Modulates Host Developmental and Metabolic Homeostasis via Insulin Signaling. *Science* 334, 670-674.

Simpson, S.J., Raubenheimer, D., 1993. A Multilevel Analysis of Feeding-Behavior - the Geometry of Nutritional Decisions. *Philos T Roy Soc B* 342, 381-402.

Simpson, S.J., Raubenheimer, D., 2009. Macronutrient balance and lifespan. *Aging (Albany NY)* 1, 875-880.

Simpson, S.J., Raubenheimer, D., 2012. *The Nature of Nutrition: a unifying framework from animal adaptation to human obesity*. Princeton University Press, Princeton.

Singer, M.S., Mace, K.C., Bernays, E.A., 2009. Self-Medication as Adaptive Plasticity: Increased Ingestion of Plant Toxins by Parasitized Caterpillars. *PLoS One* 4, e4796.

Siva-Jothy, M.T., Moret, Y., Rolff, J., 2005. Insect immunity: An evolutionary ecology perspective. *Advances in Insect Physiology*, Vol 32 32, 1-48.

Siva-Jothy, M.T., Thompson, J.J.W., 2002. Short-term nutrient deprivation affects immune function. *Physiological Entomology* 27, 206-212.

Smith, V.H., Holt, R.D., 1996. Resource competition and within-host disease dynamics. *Trends in Ecology & Evolution* 11, 386-389.

Sorci, G., Faivre, B., 2009. Inflammation and oxidative stress in vertebrate host–parasite systems. *Philosophical Transactions of the Royal Society B: Biological Sciences* 364, 71-83.

Sorensen, A., Mayntz, D., Simpson, S.J., Raubenheimer, D., 2010. Dietary ratio of protein to carbohydrate induces plastic responses in the gastrointestinal tract of mice. *J Comp Physiol B* 180, 259-266.

Topping, D.L., Clifton, P.M., 2001. Short-Chain Fatty Acids and Human Colonic Function: Roles of Resistant Starch and Nonstarch Polysaccharides. *Physiol. Rev.* 81, 1031-1064.

Troemel, E.R., Chu, S.W., Reinke, V., Lee, S.S., Ausubel, F.M., Kim, D.H., 2006. p38 MAPK Regulates Expression of Immune Response Genes and Contributes to Longevity in *C. elegans*. *PLoS Genet* 2, e183.

Turnbaugh, P.J., Bäckhed, F., Fulton, L., Gordon, J.I., 2008. Diet-Induced Obesity Is Linked to Marked but Reversible Alterations in the Mouse Distal Gut Microbiome. *Cell Host & Microbe* 3, 213-223.

Turnbaugh, P.J., Ridaura, V.K., Faith, J.J., Rey, F.E., Knight, R., Gordon, J.I., 2009. The Effect of Diet on the Human Gut Microbiome: A Metagenomic Analysis in Humanized Gnotobiotic Mice. *Science Translational Medicine* 1, 6ra14.

Vijay-Kumar, M., Aitken, J.D., Carvalho, F.A., Cullender, T.C., Mwangi, S., Srinivasan, S., Sitaraman, S.V., Knight, R., Ley, R.E., Gewirtz, A.T., 2010. Metabolic Syndrome and Altered Gut Microbiota in Mice Lacking Toll-Like Receptor 5. *Science*, science.1179721.

Vilmos, P., Kurucz, É., 1998. Insect immunity: evolutionary roots of the mammalian innate immune system. *Immunology Letters* 62, 59-66.

Weiss, B., Aksoy, S., 2011. Microbiome influences on insect host vector competence. *Trends in Parasitology* 27, 514-522.

Welchman, D.P., Aksoy, S., Jiggins, F., Lemaitre, B., 2009. Insect Immunity: From Pattern Recognition to Symbiont-Mediated Host Defense. *Cell Host & Microbe* 6, 107-114.

Wen, L., Ley, R.E., Volchkov, P.Y., Stranges, P.B., Avanesyan, L., Stonebraker, A.C., Hu, C., Wong, F.S., Szot, G.L., Bluestone, J.A., Gordon, J.I., Chervonsky, A.V., 2008. Innate immunity and intestinal microbiota in the development of Type 1 diabetes. *Nature* 455, 1109-1113.

Wilks, J., Golovkina, T., 2012. Influence of Microbiota on Viral Infections. *PLoS Pathog* 8, e1002681.

Wilson, K., 2005. Evolutionary ecology of insect host-parasite interactions: an ecological immunology perspective. CABI Publishing, Wallingford, pp. 289-246.

Wong, C.N.A., Ng, P., Douglas, A.E., 2011. Low-diversity bacterial community in the gut of the fruitfly *Drosophila melanogaster*. *Environmental Microbiology* 13, 1889-1900.

Xiang, H., Wei, G.-F., Jia, S., Huang, J., Miao, X.-X., Zhou, Z., Zhao, L.-P., Huang, Y.-P., 2006. Microbial communities in the larval midgut of laboratory and field populations of cotton bollworm (*Helicoverpa armigera*). *Canadian Journal of Microbiology* 52, 1085-1092.

Zhang, H., Jackson, T.A., 2008. Autochthonous bacterial flora indicated by PCR-DGGE of 16S rRNA gene fragments from the alimentary tract of *Costelytra zealandica* (Coleoptera: Scarabaeidae). *Journal of Applied Microbiology* 105, 1277-1285.

## Figure legends

**Figure 1:** Results from an experiment in which control and nucleopolyhedrovirus infected caterpillars were fed 5 chemically defined diets varying in the ratio and concentration of protein (P) and carbohydrate (C). Performance of the caterpillars was estimated by multiplying the survival by the average biomass gain per day. The arrow indicates performance loss for infected caterpillars when fed on high- and low-P:C (data from Lee et al. 2006).

**Figure 2:** Response surfaces showing the effects of protein (P) and carbohydrate (C) intake on the lysozyme activity, PO activity and cuticular melanism for caterpillars that were fed individually 20 diets varying in the ratio and the quantity of P and C (modified from Cotter et al., 2011).

**Figure 3:** Inter-relationship between host nutrition, host immune function, pathogen population growth, the structure and function of the gut microbiome and host fitness (modified from Ponton et al., 2011).

Figure 1






Figure 2




Figure 3

