

# Immune Modulation by Different Types of $\beta$ 2→1-Fructans Is Toll-Like Receptor Dependent

Leonie Vogt<sup>1\*</sup>, Uttara Ramasamy<sup>2</sup>, Diederick Meyer<sup>3</sup>, Gerdie Pullens<sup>4</sup>, Koen Venema<sup>5</sup>, Marijke M. Faas<sup>1</sup>, Henk A. Schols<sup>2</sup>, Paul de Vos<sup>1</sup>

**1** Department of Pathology and Medical Biology, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands, **2** Laboratory of Food Chemistry, Wageningen University, Wageningen, The Netherlands, **3** Scientific and Regulatory Affairs, Sensus B.V., Roosendaal, The Netherlands, **4** Product Development, Cosun Food Technology Centre, Roosendaal, The Netherlands, **5** Department of Biosciences, TNO, Zeist, The Netherlands

## Abstract

**Introduction:**  $\beta$ 2→1-fructans are dietary fibers. Main objectives of this study were 1) to demonstrate direct signalling of  $\beta$ 2→1-fructans on immune cells, 2) to study whether this is mediated by the pattern recognition receptors Toll-like receptors (TLRs) and nucleotide-binding oligomerisation domain-containing proteins (NODs), and 3) to relate the observed effects to the chain length differences in  $\beta$ 2→1-fructans.

**Methods:** Four different  $\beta$ 2→1-fructan formulations were characterised for their chain length profile. Human peripheral blood mononuclear cells (PBMCs) were stimulated *in vitro* with  $\beta$ 2→1-fructans, and production of IL-1Ra, IL-1 $\beta$ , IL-6, IL-10, IL-12p70, and TNF- $\alpha$  was analysed. Reporter cells for TLRs and NODs were incubated with  $\beta$ 2→1-fructans and analysed for NF- $\kappa$ B/AP-1 activation.

**Results:** Cytokine production in human PBMCs was dose- and chain length-dependent. Strikingly, short chain enriched  $\beta$ 2→1-fructans induced a regulatory cytokine balance compared to long chain enriched  $\beta$ 2→1-fructans as measured by IL-10/IL-12 ratios. Activation of reporter cells showed that signalling was highly dependent on TLRs and their adapter, myeloid differentiation primary response protein 88 (MyD88). In human embryonic kidney reporter cells, TLR2 was prominently activated, while TLR4, 5, 7, 8, and NOD2 were mildly activated.

**Conclusions:**  $\beta$ 2→1-fructans possess direct signalling capacity on human immune cells. By activating primarily TLR2, and to a lesser extent TLR4, 5, 7, 8, and NOD2,  $\beta$ 2→1-fructan stimulation results in NF- $\kappa$ B/AP-1 activation. Chain length of  $\beta$ 2→1-fructans is important for the induced activation pattern and IL-10/IL-12 ratios.

**Citation:** Vogt L, Ramasamy U, Meyer D, Pullens G, Venema K, et al. (2013) Immune Modulation by Different Types of  $\beta$ 2→1-Fructans Is Toll-Like Receptor Dependent. *PLoS ONE* 8(7): e68367. doi:10.1371/journal.pone.0068367

**Editor:** Markus M. Heimesaat, Charité, Campus Benjamin Franklin, Germany

**Received:** May 1, 2013; **Accepted:** May 28, 2013; **Published:** July 5, 2013

**Copyright:** © 2013 Vogt et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Funding:** Within the framework of the Carbohydrate Competence Center, this research has been financially supported by the European Union, the European Regional Development Fund, and The Northern Netherlands Provinces (Samenwerkingsverband Noord-Nederland), KOERS NOORD. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing Interests:** The authors declare the affiliations to commercial companies in the form of employment of Diederick Meyer by Sensus B.V., the employment of Gerdie Pullens by COSUN Food Technology Centre, and the employment of Koen Venema by TNO. There are no patents, products in development or marketed products to declare. This does not alter the authors' adherence to all the PLOS ONE policies on sharing data and materials.

\* E-mail: L.M.Vogt@umcg.nl

## Introduction

High fiber intake is associated with lower mortality in subjects suffering from circulatory, digestive, and non-cardiovascular non-cancerous inflammatory diseases [1-3]. These associations are similar for men and women and are observed in most countries even after careful adjustment for potentially confounding lifestyle and dietary differences [4]. The mechanisms by which fibers contribute to reduced mortality remain to be identified but have been suggested to be of both metabolic and immunological nature [5]. Cereal fibers have been associated with lower concentrations of inflammatory biomarkers [4], but there is an urgent need for studies addressing specific fibers [4] to obtain a better understanding of mechanisms and components underlying protective associations.

One of the many types of dietary fibers which have been reported to elicit health benefits are linear  $\beta$ 2→1-fructans (also named inulin-type fructans, ITFs, or fructooligosaccharides). These fructans are made up of fructose subunits, can vary considerably in chain length, and occur with or without the presence of a terminal glucose moiety [6]. The fructose oligomers are denoted as  $F_n$  or  $GF_n$  ( $n$  = number of fructose subunits,  $GF_n$  = fructose chain terminated with a glucose molecule) [6]. The chain length can also be denoted by the Degree of Polymerisation (DP), representing the number of monosaccharide subunits in the chain [6]. Whether these variations in composition cause different physiological responses is currently unknown. Ingestion of  $\beta$ 2→1-fructans induces specific effects on the immune system, as recently reviewed [7-11]. Changes in immunological parameters have been reported in the gut lumen, Peyer's patches, spleen, and blood [12-14]. These effects could be induced via a prebiotic mechanism by

stimulating beneficial (“probiotic”) bacteria in the intestine, such as bifidobacteria and lactobacilli [10]. This can coincide with modulation of fermentation products such as increased production of short chain fatty acids [15,16], which affect the recruitment of leukocytes to inflammatory sites [17,18], and suppress the production of pro-inflammatory cytokines and chemokines [18–20]. Besides through these more indirect effects on the microbiota and its fermentation products, another pathway of modulation is gaining attention, i.e. through direct signalling on immune cells. This theory is supported by new insights into the structural and functional makeup of the intestine [21,22]. Some dietary fibers such as  $\beta$ -glucans, modulate the human immune system by binding to pattern recognition receptors (PRRs) on cells of the innate immune system, such as Dectin-1 and Toll-like receptors (TLRs) [23–25]. These PRRs recognise so-called pathogen-associated molecular patterns (PAMPs), which are small molecular motifs found on groups of pathogens but also on many microorganisms with beneficial or no effects on host health [26]. Because of their ability to recognise specific carbohydrate moieties and elicit immune responses, we hypothesised that PRRs, and more specifically TLRs and NODs are activated by  $\beta$ 2→1-fructans, and that this is one of the pathways by which  $\beta$ 2→1-fructans can influence the human immune system [27,28].

As little knowledge is available on the specific immunomodulating effects of different types of  $\beta$ 2→1-fructans, we investigated the effect of  $\beta$ 2→1-fructans with different chain lengths on cytokine release by human Peripheral Blood Mononuclear Cells (PBMCs) and we studied whether and which PRRs are involved in  $\beta$ 2→1-fructan recognition. Our studies support the concept that ITFs can signal directly on immune cells via TLRs and that chain length differences in ITFs can differentially affect immunological parameters.

## Materials and Methods

### Ethics Statement

Chemical analyses (HPAEC and HPSEC) were performed at Wageningen University. Other experiments, including blood sampling of human volunteers, were conducted within the University Medical Center Groningen, in the Netherlands. Written informed consent was obtained, and data were analysed and presented anonymously. This research and consent procedure have been approved by the ethical review board of the University Medical Center, Medisch Ethische Toetsingscommissie University Medical Center Groningen, as documented in the approved application “2007/255”. All clinical investigation was conducted according to the principles expressed in the Declaration of Helsinki.

### High Performance Anion Exchange Chromatography

Four different  $\beta$ 2→1-fructan formulations were selected and their specific chain length profiles (range and distribution) were characterised by High Performance Anion Exchange Chromatography (HPAEC). The applied  $\beta$ 2→1-fructans were extracted from chicory root and comprise a mixture of different chain lengths varying from DP 2 to 60, referred to as inulin or native inulin [29]. These chicory root  $\beta$ 2→1-fructan chains are terminated by a glucose molecule [6]. Fructans of varying DP can be acquired by partial enzymatic hydrolysis of this extract resulting in  $G\text{F}_n$  and  $F_n$  chains with  $G\text{F}_n$  ranging from DP 2 to 10 [6]. Another way to obtain  $\beta$ 2→1-fructans is to synthesise them from sucrose, resulting in  $G\text{F}_n$  type fructans with DP 2–4 [30].

The following  $\beta$ 2→1-fructan formulations were analysed: ITF I (Frutalose®OFF) ITF II (Frutafit®CLR), ITF III (Frutafit®IQ),

and ITF IV (Frutafit®TEX!).  $\beta$ 2→1-fructan formulations were provided by Sensus B.V., Roosendaal, The Netherlands. Endotoxin levels (endotoxin units, EU) of all used  $\beta$ 2→1-fructan samples were assessed by Toxikon (Leuven, Belgium) and were below  $0.3 \times 10^{-3} \mu\text{g}^{-1}$ .

For HPAEC, a Dionex ICS 3000 system (Dionex), equipped with a Dionex CarboPac PA-1 column (2 x 250 mm) in combination with a CarboPac PA-1 guard column (2 x 50 mm) was used. Sample concentrations of 0.05–0.1 mg/ml and partial-loop injection of 10  $\mu\text{l}$  were applied. The system was equipped with pulsed amperometric detection. Elution was performed at 0.3 ml/min, and oligomers were separated using a gradient as follows: 0–400mM NaOAc in 100 mM NaOH within 40 min., followed by a 5 min. washing step (1M NaOAc in 100 mM NaOH), and 15 min. equilibration (100 mM NaOH).

### High Pressure Size Exclusion Chromatography

$\beta$ 2→1-fructan formulations were tested for their elution profile using High Performance Size Exclusion Chromatography (HPSEC) on Dionex Ultimate 3000 HPLC (Dionex, Sunnyvale, CA, USA). The analysis was performed on three TSK-Gel columns connected in series (4000–3000–2500 SuperAW; 150 x 6 mm). The columns were preceded by a TSK Super AW-L guard column (35 x 4.6 mm). All columns were from Tosoh Bioscience (Tokyo, Japan). Elution was performed at a flow rate of 0.6 ml/min. using sodium nitrate (0.2 M) as the eluent. A volume of 20  $\mu\text{l}$  of the sample (2.5 mg/ml) in millipore water was injected and eluted at 55°C. Solubles were detected using a refractive index detector, Shodex type RI 101 (Showa Denko, Japan). The software used for acquiring the data was Chromeleon version 7. Molecular weight distribution of polysaccharides was determined using pullulan standards (Polymer Laboratories, Varian Inc., Palo Alto, CA, USA) in the molecular mass range of 0.18 - 790 kDa.

### Isolation Human Peripheral Blood Mononuclear Cells

Human PBMCs were used to study whether  $\beta$ 2→1-fructans can signal on immune cells, especially since PBMCs express many PRRs such as TLRs [31,32] and NODs [33]. Peripheral blood from human volunteers was collected in heparinised tubes (15 IU/ml lithium-heparin, Becton Dickinson B.V., Breda, The Netherlands) and PBMCs were isolated by Ficoll density gradient separation (Lymphoprep, Axis-Shield, Oslo, Norway). Cells were kept in RPMI1640 medium (Gibco, Life Technologies, Bleiswijk, The Netherlands) supplemented with 10% Fetal Bovine Serum (FBS, HyClone, Thermo Scientific, Breda, The Netherlands) and 50  $\mu\text{g}/\text{ml}$  gentamicin (Gibco).

### Cytokine Expression

As a read out for PBMC activation, cytokine production in the medium was analysed after stimulation of PBMCs with a concentration series of  $\beta$ 2→1-fructans. To evaluate whether  $\beta$ 2→1-fructans induce a more anti-inflammatory or pro-inflammatory effect, and whether chain length profile would affect this balance, the IL-10/IL-12 ratio was calculated for each  $\beta$ 2→1-fructan formulation at 1  $\mu\text{g}/\text{ml}$  or 100  $\mu\text{g}/\text{ml}$ . In this context, a higher IL-10/IL-12 ratio is representative of a more anti-inflammatory effect [34,35]. Human PBMCs ( $n = 6$ ; 3 males, 3 females, age 25–42 yr., healthy non-smoker volunteers) were seeded in a 24 wells plate at a density of  $2 \times 10^6$  cells/well with a final volume of 1 ml/well. To characterise possible effects of different chain length profiles on cytokine expression, the  $\beta$ 2→1-fructan formulations ITF I to IV were dissolved in culture medium and added to the PBMCs at final concentrations ranging from 1 to 100  $\mu\text{g}/\text{ml}$ . After 24h of incubation (37°C, 5%  $\text{CO}_2$ ) cytokine

levels in the supernatant were measured using a Bio-Plex™ premixed cytokine assay, human 6-plex group I; cat.#: M5000B6CPS, control 5022016, according to the manufacturer's instructions. (Bio-rad Laboratories, Veenendaal, The Netherlands). This customised kit simultaneously measured human IL-1Ra, IL-1 $\beta$ , IL-6, IL-10, IL-12p70, and TNF- $\alpha$ . Concentration series of cytokine standards were prepared for the appropriate concentration range, and coupled beads were diluted ten times, resuspended, and added to a pre-wetted filter plate. After washing the plate twice, standards, negative controls, and samples (all in duplicate) were transferred into the plate (50  $\mu$ l per well), and the plate was sealed and incubated on a shaker at room temperature (RT) for 30 min. After incubation, the plate was washed three times, detection antibodies were resuspended and diluted ten times and 25  $\mu$ l was added to each well. The plate was incubated on a shaker at RT for 30 min., and after washing three times, 50  $\mu$ l of streptavidin-phycoerythrin was added to each well and the plate was incubated on a shaker at RT for 10 min. After washing the plate three times, 125  $\mu$ l of assay buffer was added per well, the plate was incubated on a shaker for 5 min. and fluorescence was measured using a Luminex 100 System and StarStation software. All procedures as of the incubation with detection antibodies were performed in the dark.

### Cell Culture of Reporter Cell Lines

Selection media, Normocin antibiotic, Quanti-blue reagent, TLR agonists and the following TLR-, and NOD-reporter cell lines were acquired from InvivoGen (InvivoGen, Toulouse, France). Two THP-1 human acute monocytic leukemia reporter cell lines were acquired from InvivoGen, both endogenously expressing human TLRs and with inserted construct for Secreted Embryonic Alkaline Phosphatase (SEAP) coupled to the NF- $\kappa$ B/AP-1 promoter. The first of these THP-1 cell lines carries extra inserts for MD2 and CD14 to boost TLR signalling, and the second THP-1 cell line expresses only a truncated, non-functional form of the TLR adapter MyD88. Nine different human Embryonic Kidney (HEK293)-Blue reporter cell lines were purchased from InvivoGen, each with a different inserted construct for either human TLR2, TLR3, TLR4, TLR5, TLR7, TLR8, TLR9, NOD1, or NOD2, and all nine cell lines carrying an inserted construct for Secreted Embryonic Alkaline Phosphatase (SEAP) coupled to the NF- $\kappa$ B/AP-1 promoter. Both THP-1 cell lines were maintained in RPMI1640 containing 10% heat inactivated FBS, NaHCO<sub>3</sub> (Boom B.V. Meppel, The Netherlands; 1,5 g/l), L-glutamine (2 mM), glucose (4,5 g/l), HEPES (10 mM), sodium pyruvate (1 mM) penicillin/streptomycin (50 U/ml and 50  $\mu$ g/ml), all from Sigma-Aldrich Chemie B.V., Zwijndrecht, The Netherlands, and Normocin (100  $\mu$ g/ml). Both THP-1 cell lines were kept at a concentration of  $5 \times 10^5$  cells/ml. HEK-Blue cells were maintained in DMEM (Life Technologies Europe B.V.) containing 10% heat inactivated FBS, L-glutamine (2 mM), glucose 4,5 g/l, penicillin/streptomycin (50 U/ml and 50  $\mu$ g/ml), and Normocin (100  $\mu$ g/ml). HEK cells were grown to ~80% confluency. After culturing for 3 passages, all reporter cell lines were maintained in selection media according to the manufacturer's protocol.

### Reporter Cell Stimulations and Quanti-Blue™ Analysis

THP-1 cells were centrifuged for 5 min. at 300 *g* to collect the cells and resuspended to the cell density specified by the manufacturer's protocol (Table 1). In a flat bottom 96 wells plate, 100  $\mu$ l of this cell suspension per well was stimulated for 24 h (37°C, 5% CO<sub>2</sub>) with 10  $\mu$ l of stimulus, i.e. concentration series (1  $\mu$ g/ml - 2 mg/ml) of  $\beta$ 2→1-fructan formulations ITF I to IV,

or the relevant positive control as indicated by the manufacturer's protocol (Table 1). Plain culture medium and endotoxin-free water were applied as negative controls. After incubation, 20  $\mu$ l of cell culture medium of stimulated reporter cells was incubated with 180  $\mu$ l of Quanti-blue reagent in a new flat bottom 96 wells plate for 45 min at 37°C and SEAP activity (absorbance), representing activation of NF- $\kappa$ B/AP-1, was measured at 650 nm on a VersaMax microplate reader (Molecular Devices GmbH, Biberach an der Riss, Germany) using SoftMax Pro Data Acquisition & Analysis Software. HEK-Blue reporter cells were rinsed with medium to detach them from the culture flask and cells were resuspended to the cell density specified by the manufacturer's protocol (Table 1). 180  $\mu$ l of cell suspension per well was stimulated for 24 h (37°C, 5% CO<sub>2</sub>) with 20  $\mu$ l of  $\beta$ 2→1-fructan, endotoxin free water, plain culture medium or positive control (Table 1) in a flat bottom 96 wells plate. For all HEK cell lines, after incubation, analysis of SEAP was performed in the same way as described for the THP-1 cells.

### Statistical Analysis

Significance levels were determined by parametric Student's *t*-test for unpaired observations (two-tailed) or by non-parametric Mann-Whitney U-test for unpaired observations (two-tailed). Results are expressed as mean  $\pm$  SEM or mean  $\pm$  SD respectively. A *P*-value <0.05 was considered statistically significant. *P*-values <0.05 are denoted with \*, *P*-values <0.01 are denoted with \*\*, and *P*-values <0.001 are denoted with \*\*\*.

## Results

### Chemical Characterisation of $\beta$ 2→1-fructan Formulations

The oligomer profiles of four  $\beta$ 2→1-fructan formulations were characterised using HPAEC and HPSEC. Figure 1A depicts the oligosaccharide range and relative response per oligomer of the  $\beta$ 2→1-fructan formulations tested with HPAEC. Figure 1B represents the HPSEC analysis of the four different  $\beta$ 2→1-fructan formulations, showing the elution patterns as a measure for the molecular size distributions of these compounds in kDa. The HPSEC analysis corroborated the observed chain length profiles from the HPAEC analysis and in addition this method rendered a visual representation of the degree in which the different chain lengths are present per analysed formulation. The most important differences can be observed between ITF I and II as compared to ITF III and IV. ITF I can be described as fructooligosaccharide (FOS) with mostly chain lengths of DP<10, and ITF II can be described as a FOS-enriched inulin, with a large proportion of chains smaller than DP10, but also containing chains with DP up to 25. ITF III and IV are described as "inulin" due to their broad range of chain lengths present (up to DP60). ITF I and II contain mostly chains of the type GF<sub>3</sub>, GF<sub>4</sub>, and GF<sub>5</sub>, (i.e. starting with a glucose molecule followed by 3, 4, or 5 fructose subunits) and F<sub>3</sub>, F<sub>4</sub>, or F<sub>5</sub> (chains consisting only of 3, 4, or 5 fructose moieties). Between ITF I and II, ITF I contains relatively more F<sub>n</sub> type oligosaccharides and ITF II contains relatively more GF<sub>n</sub> type molecules. ITF III and IV consist solely of the GF<sub>n</sub> type fructans while ITF I and II consist of both GF<sub>n</sub> and F<sub>n</sub> fructans. Glucose and fructose monomers, and GF and GF<sub>2</sub> (i.e. dimers of glucose and fructose subunits and trimers made up of one glucose subunit and two fructose subunits) are present in all four ITFs.

### $\beta$ 2→1-fructans Induce Cytokine Production in Human PBMCs

To gain insight into dose effects and to determine which cytokines are induced by  $\beta$ 2→1-fructans we first screened for a

**Table 1.** Cell densities and positive controls used in reporter cell stimulations.

Cell line	Cell density	Positive control
THP-1 MD2-CD14	1*10 <sup>6</sup> cells/ml	<i>E.coli</i> K12 lipopolysaccharide Ultrapure (LPS, 100 ng/ml)
THP-1 DefMyD	2*10 <sup>6</sup> cells/ml	L-Ala-gamma-D-Glu-mDAP (TriDAP, 10 µg/ml)
HEK-Blue TLR2	2.8*10 <sup>5</sup> cells/ml	Pam2CGDPKHPKSF (FSL-1, 1 µg/ml)
HEK-Blue TLR3	2.8*10 <sup>5</sup> cells/ml	Poly(I:C) high molecular weight (HMW, 1 mg/ml)
HEK-Blue TLR4	1.4*10 <sup>6</sup> cells/ml	<i>E.coli</i> K12 lipopolysaccharide Ultrapure (LPS, 100 ng/ml)
HEK-Blue TLR5	1.4*10 <sup>6</sup> cells/ml	Recombinant flagellin from <i>S.typhimurium</i> (Rec-FLA-ST,100 ng/ml)
HEK-Blue TLR7	2.2*10 <sup>6</sup> cells/ml	Imiquimod (5 µg/ml)
HEK-Blue TLR8	2.2*10 <sup>6</sup> cells/ml	Single stranded RNA (ssRNA40/LyoVec™, 5 µg/ml)
HEK-Blue TLR9	4.5*10 <sup>6</sup> cells/ml	Type B CpG oligonucleotide (ODN 2006, 100 mg/ml)
HEK-Blue NOD1	2.8*10 <sup>6</sup> cells/ml	L-Ala-gamma-D-Glu-mDAP (TriDAP, 10 µg/ml)
HEK-Blue NOD2	1.4*10 <sup>6</sup> cells/ml	MurNac-L-Ala-gamma-D-Glu-mDAP (M-TriDAP, 10 µg/ml)

doi:10.1371/journal.pone.0068367.t001

panel of pro- and anti-inflammatory cytokines, i.e. IL-1β, IL-6, IL-12, TNF-α, IL-1Ra, and IL-10, after stimulating human PBMCs for 24 hours with β2→1-fructans. This was done for ITF I, in doses of 1/2,5/5/10, and 100 µg/ml. As shown in Figure 2, IL-1Ra expression was decreased for lower doses (2,5 to 10 µg/ml β2→1-fructans) and increased at 100 µg/ml β2→1-fructans (p<0.05, panel A). The same pattern was observed for IL-6; expression was decreased for 2.5 µg/ml (p<0.05) and a substantial increase was observed at 100 µg/ml (p<0.01, panel C). IL-10 expression was increased, especially at 1 µg/ml (p<0.001) and 100 µg/ml (p<0.0001, panel D). A small increase for TNFα was observed at 100 µg/ml (12%, p<0.05, panel F). Since a dose as low as 1 µg/ml induced a significant increase for IL-10, and 100 µg/ml induced significant and substantial increases, these low end and high end doses were subsequently used for further cytokine measurements in PBMCs.

### Inulin-type Fructan-induced Cytokine Production by Human PBMCs is Chain Length Dependent

To gain insight into size-response relationships, we investigated whether the chain length of β2→1-fructans has an effect on the type and quantity of cytokines produced by PBMCs. ITF I to IV were tested at the concentrations of 1 and 100 µg/ml. IL-6 and IL-10 were mainly induced by ITF I and ITF II (Fig. 3) which can therefore be mainly attributed to short chain molecules enriched in these formulations, i.e. F<sub>2</sub>-F<sub>5</sub> and GF<sub>2</sub>-GF<sub>5</sub> respectively. Production of IL-12 was slightly increased by ITF II and strongly induced by ITF III and IV, indicating that longer chains (>DP8, GF<sub>n</sub> type) have to be held responsible for this effect. Similarly, the highest production of TNF-α was observed after incubation with ITF IV. These combined results showed that chain length profile of β2→1-fructan formulations is an important determinant of the cytokine profile which is induced upon stimulation.

Next, the IL-10/IL-12 ratios were calculated to quantify whether there is a correlation between chain length profile and cytokine balance (Fig.4). IL-10/IL12 ratios can be applied to calculate the regulatory effects of bioactive food components [36]. Controls were set to 100, and a ratio of more than 100 was considered to be regulatory or anti-inflammatory, whereas a ratio lower than 100 was considered to be proinflammatory. Strikingly, the IL-10/IL-12 ratios gradually decreased in the sequence of ITF I>ITF II>ITF III>ITF IV, indicating that shorter chain fructans induce a regulatory cytokine balance in human PBMCs compared


to longer chain fructans. When taking the structural differences into consideration (Fig.1) it can be concluded that the molecules F<sub>2</sub>-F<sub>5</sub> and/or GF<sub>2</sub>-GF<sub>5</sub>, which are enriched in ITF I and II, skew the IL-10/IL-12 ratio in PBMCs more towards IL-10, and thus induce a more anti-inflammatory balance.

### TLR Mediated Activation of NF-κB/AP-1 by β2→1-fructans is Dependent on the Presence of Functional TLR Adapter MyD88


Next, we investigated whether the activation of immune cells can be explained by activation of PRRs by the β2→1-fructans. To this end we applied the following strategy. First, β2→1-fructans were tested for the ability to induce NF-κB/AP-1 transcription in a THP-1 reporter cell line which endogenously expresses all TLRs, and carries extra inserts for co-signalling molecules CD14 and MD2 to increase the TLR-mediated responses. NF-κB and AP-1 are essential transcription factors in signalling for cytokine release [37].

ITF I and II induced a statistically significant elevation of NF-κB/AP-1 activation in THP-1 MD2-CD14 cells, while ITF III and ITF IV only induced activation at higher doses (Fig.5). The strongest activation was observed for ITF II, which is mainly short chain GF<sub>n</sub>. Next, we investigated whether the activation was TLR-dependent by testing the effect of β2→1-fructans on a specific THP-1 reporter cell line carrying the truncated TLR adapter molecule MyD88. These THP-1 DefMyD cells express only non-functional MyD88, whereas the functional MyD88 is an essential adapter molecule for TLR2, 4, 5, 7, 8, and 9 signalling. The β2→1-fructan doses which significantly activated THP-1 MD2-CD14 cells were subsequently tested in the THP-1 DefMyD cells.


β2→1-fructan mediated NF-κB/AP-1 activation appeared to be TLR dependent as the activation pattern as observed in THP-1 MD2-CD14, was virtually absent in the MyD88 deficient cell line. This indicated that β2→1-fructan mediated signalling to NF-κB/AP-1 was TLR2, 4, 5, 7, 8, and/or 9 dependent. Slight activation of THP-1 DefMyD cells by ITF IV suggests that other activation pathways independent of MyD88 may also have been induced by high concentrations of the longer chain β2→1-fructans which are enriched in this ITF.




**Figure 1. HPAEC profiles and HPSEC elution patterns of different β2→1-fructan formulations.** Figure A depicts the fructose (F) and glucose (G) monomers, dimers, and fructan oligomers present in the ITF formulations. GF<sub>n</sub> chains are terminated by a glucose molecule, and F<sub>n</sub> chains consist of only fructose moieties. In both cases, n represents the number of fructose moieties in the chain. Figure B depicts the elution patterns as a measure for molecular weight distribution profiles of the four different β2→1-fructan formulations in kDa. doi:10.1371/journal.pone.0068367.g001




**Figure 2. Induction of cytokines by ITF I, in a dose range of 0 to 100 μg/ml.** Statistical significance levels were determined with a parametric Student's t-test for unpaired observations (two-tailed). Mean and SEM of cytokine production is plotted as percentage of controls, which were set to 100% (n=4). Panels A to F show the results for IL-1Ra, IL-1β, IL-6, IL-10, IL-12, and TNF-α respectively.  
doi:10.1371/journal.pone.0068367.g002



**Figure 3. Induction of cytokines by  $\beta$ 2 $\rightarrow$ 1-fructan formulations with different chain length (DP range), at 0, 1, and 100  $\mu$ g/ml.** Statistical significance levels were determined with a non-parametric Mann-Whitney U-test for unpaired observations (two-tailed). Mean and SD of cytokine expression are plotted as percentage of controls (represented by 0  $\mu$ g/ml), which were set to 100% (n = 6). Panel A to D represent cytokine expression induced by ITF I to IV respectively. doi:10.1371/journal.pone.0068367.g003



**Figure 4. Ratio of IL-10/IL-12 upon incubation of PBMCs with different chain length  $\beta$ 2 $\rightarrow$ 1-fructans.** PBMCs (n = 6) were stimulated with 1  $\mu$ g/ml (panel A) and 100  $\mu$ g/ml (panel B)  $\beta$ 2 $\rightarrow$ 1-fructans for 24h. Statistical significance levels were determined with a non-parametric Mann-Whitney U-test for unpaired observations (two-tailed). Mean and SD of the IL-10/IL-12 ratios is plotted for the different  $\beta$ 2 $\rightarrow$ 1-fructans as percentage of controls, which were set to 100% (n = 6) and horizontal bars indicate the significant differences between  $\beta$ 2 $\rightarrow$ 1-fructan treatments. doi:10.1371/journal.pone.0068367.g004




**Figure 5. NF-κB/AP-1 activation in THP-1 MD2-CD14 and THP-1 defMyD reporter cells.** Statistical significance levels were determined with a non-parametric Mann-Whitney U-test for unpaired observations (two-tailed). Mean and SD of NF-κB/AP-1 activation by β2→1-fructans of different chain lengths (ITF I, II, III and IV) are plotted as percentage of negative controls (unstimulated cells), which were set to 100% for both cell lines. LPS stimulation was used as a common positive control for TLR4/MyD88 signalling to NF-κB/AP-1 in the functional THP-1 MD2-CD14 cell line. TriDAP was used as a positive control in THP-1 DefMyD cells, which induces MyD88-independent signalling to NF-κB/AP-1. Panel A to D represent cytokine expression induced by ITF I to IV respectively. Only concentrations which induced activation in the THP-1 MD1-CD14 cells are shown for comparison of these concentrations in the THP-1 DefMyD cells. doi:10.1371/journal.pone.0068367.g005

**β2→1-fructans Induce Strong Dose Dependent Activation of TLR2, Slight Activation of TLR4, 5, 7, 8, and NOD2, but No Activation of TLR3, TLR9, or NOD1**

The results from the THP-1 cell stimulations showed that TLRs are involved in β2→1-fructan signalling. Next, we investigated which TLRs are specifically activated by β2→1-fructans and whether the intracellular PRRs NODs were activated. To this end, HEK reporter cell lines, each carrying one construct for a specific TLR or NOD, were stimulated with β2→1-fructans in the same

way as described for the THP-1 cells (Fig.6). Strikingly, HEK cells with TLR2 construct were strongly and dose dependently activated by the β2→1-fructans. Moreover, this response was chain length dependent, as NF-κB/AP-1 activation in HEK TLR2 cells increased with increasing chain length in the sequence of ITF I<ITF II<ITF III<ITF IV. In these cells, long chain fructans induced the strongest activation, up to 6-fold induction as compared to control. To a lesser extent, HEK cells carrying either TLR4, 5, 7, 8, or NOD2 were activated upon β2→1-fructan stimulation. HEK cells expressing TLR3, 9 or NOD1 were



**Figure 6. NF-κB/AP-1 activation of HEK cell lines overexpressing separate TLRs or NODs.** Statistical significance levels were determined with a non-parametric Mann-Whitney U-test for unpaired observations (two-tailed). Mean and SD of NF-κB/AP-1 activation in HEK cell lines stimulated with  $\beta$ 2→1-fructans for 24 h are plotted as percentage of unstimulated controls, which were set to 100%. Dosages are plotted in  $\mu$ g/ml. Endotoxin free H<sub>2</sub>O was used as an additional negative control and per cell line the relevant positive controls were applied as mentioned in Table 1. doi:10.1371/journal.pone.0068367.g006

not significantly activated by  $\beta$ 2→1-fructan stimulation (data not shown). These data combined show that TLR binding was  $\beta$ 2→1-fructan type-, and thus chain length dependent.

**Discussion**

We hypothesised that  $\beta$ 2→1-fructans can affect the immune status through physical contact with pattern recognition receptors

(PRRs) on gut immune cells such as intra-epithelial lymphocytes [38] or DCs [27,28]. As TLR activation in PBMCs leads to cytokine production [31,39], we used these cells to test whether  $\beta$ 2→1-fructans would also induce production of cytokines. Both anti-inflammatory and proinflammatory cytokines were produced, demonstrating that  $\beta$ 2→1-fructans have the ability to activate human immune cells. Interestingly, chain length proved to be an important factor in skewing the cytokine balance. Anti-inflamma-

tory IL-10 was strongly induced by short chain  $\beta$ 2→1-fructans and a striking correlation was observed between the ratio of IL-10/IL-12 and the chain lengths of the fibers. The IL-10/IL-12 ratio has previously been used to evaluate the effects of probiotic bacteria on immune responses [40,41], and to describe anti-inflammatory effects of butyrate [42]. It is therefore useful to apply the same approach to study the effect of prebiotic fibers.

It can be concluded on the basis of our data that the short chain molecules which are more abundant in ITF I and II, skew the IL-10/IL-12 ratio in PBMCs more towards IL-10, and thus induce a more anti-inflammatory balance. When considering that the proportion of short chain fructans of the GF<sub>n</sub> type are also distributed as ITF I>ITF II>ITF III, the results also suggest that specifically the short chain fructans of the GF<sub>n</sub> type determine the outcome of this cytokine ratio. In spite of the fact that longer chains induced more IL-12 compared to shorter chains, the induced IL-10/IL-12 ratio is not significantly different from the ratio measured in the control cells, indicating that they are pro-inflammatory as compared to the shorter chains but not pro-inflammatory per se, when compared to controls.

IL-6 production was observed for higher concentrations of ITF I and II. Although IL-6 production is generally regarded as proinflammatory, IL-6 can exert several anti-inflammatory effects such as inhibition of TNF- $\alpha$  function [43], and activation of IL-1Ra and IL-10 [44]. Only few studies have addressed in vitro stimulation of immune cells by  $\beta$ 2→1-fructans and cytokine profiles. In a study of Eiwegger *et al.* [45] three types of prebiotic oligosaccharides, including a mixture of short chain galactooligosaccharides (scGOS) and long chain  $\beta$ 2→1-fructans were analysed for the induction of a selected panel of cytokines in cord blood mononuclear cells (CBMCs). The mixture containing long chain  $\beta$ 2→1-fructans did not induce IL-10 in these experiments which corroborates our findings. Long chain  $\beta$ 2→1-fructans alone, or short chain  $\beta$ 2→1-fructans were not tested in the Eiwegger study, which would allow for a more complete comparison with our results. In supplementation studies in experimental animals, the observed immune effects with  $\beta$ 2→1-fructans such as increased cytokine productions, increased serum and secretory immunoglobulins [13,46–49], and increased numbers of IL-10<sup>+</sup>, TLR2<sup>+</sup>, and TLR4<sup>+</sup> DCs [50] could be due to direct effects of  $\beta$ 2→1-fructan-mediated TLR activation in the intestine, be it on immune cells or intestinal epithelial cells [51]. However, the underlying mechanisms in these in vivo experiments are likely to be more complex and are probably the result of combined direct and indirect effects in the intestine.

Also other studies corroborate our findings on chain length effects and differences in induced immune parameters. In a study where rats were supplemented with shorter chain  $\beta$ 2→1-fructans (DP2-8), increased ex vivo secretions of IL-10 were observed in cells from the Peyer's patches [13,52–54] and mesenteric lymph nodes [53]. In another study by Ito *et al.* the effect of different chain length  $\beta$ 2→1-fructans on prebiotic and immunomodulatory parameters in rats was studied [55]. In this study,  $\beta$ 2→1-fructans supplementation increased the cecal lactobacilli and bifidobacteria counts and that IgA concentrations were increased in the order DP4> DP8> DP16. In addition, DP4, DP8, and DP16, but not DP23, increased IgA-producing plasma cells in the cecal mucosa. IFN- $\gamma$  and IL-10 production in cecal CD4(+) T cells was enhanced solely by DP4. These results confirm that in vivo, chain length of  $\beta$ 2→1-fructans is also of importance to the immunological response and the fact that these parameters were not correlated to any of the observed bacterial changes suggests that this is a direct effect on (immune) cells. This size exclusion effect for direct

signalling of  $\beta$ 2→1-fructans in vivo was previously suggested by Seifert and Watzl [8].

As there are many families of PRRs with typical carbohydrate binding properties and immunomodulatory capacities [56], we applied a strategy to target TLRs and NODs: as shown in this study, the activating capacity of  $\beta$ 2→1-fructans on THP-1 cells was MyD88 dependent, and activation of TLR2 expressing HEK cells was strongly induced, which implies that signalling through TLRs, and specifically TLR2, is important in the immunomodulatory capacity of  $\beta$ 2→1-fructans [11,27,28,57]. Other PRRs, i.e. C-type Lectin-like receptors or RIG-like receptors do not signal via MyD88, but utilise other mechanisms and molecules to transduce their signal [58,59].

When studying  $\beta$ 2→1-fructan mediated activation of HEK cells expressing individual TLRs or NODs, we observed that NF- $\kappa$ B/AP-1 was strongly induced via activation of TLR2, and to a lesser extent by TLR4, 5, 7, 8, and NOD2. No activation was observed for TLR3, 9, or NOD1. TLR3 does not signal to NF- $\kappa$ B/AP-1 via MyD88 but through its adapter Toll/interleukin-1 receptor homology-domain-containing adapter-inducing interferon- $\beta$  (TRIF) [60]. However, activation by  $\beta$ 2→1-fructans was virtually absent in the THP-1 DefMyD cells, indicating the importance of MyD88-related TLRs and no involvement of TLR3.

TLR2 has many natural ligands of bacterial, fungal, viral, and even endogenous nature [61]. One of these is zymosan, a yeast PAMP consisting of  $\beta$ -glucans [62], which bear similarity to  $\beta$ 2→1-fructans because both are prebiotic polysaccharides with  $\beta$ -glycosidic bonds. Although  $\beta$ -glucans are different molecules, it is tempting to speculate that the mechanism of TLR ligation by  $\beta$ 2→1-fructans could be similar to that of  $\beta$ -glucans, but this requires further biochemical studies.

Another way of inducing different signals might be mechanistic differences in receptor interactions at the cellular surface. It is possible that the shorter chains only activate a few receptors at a time, and the receptors may be located relatively distant from each other, whereas the longer chain  $\beta$ 2→1-fructans might show a property of clustering the relevant receptors on the membrane, thereby creating a molecular complex which enhances signal transduction or alters the downstream outcome. This clustering mechanism has been described for lipopolysaccharide (LPS), clustering substantial numbers of TLR4 [63] and may be a relevant mechanism for other TLRs as well.

To our best knowledge this is the first study addressing the direct mechanism behind the immunomodulating capacity of specific  $\beta$ 2→1-fructans, with the aim to get more insight into the protective associations attributed to dietary fibers in epidemiological studies [4]. We demonstrated the principle that  $\beta$ 2→1-fructans possess direct signalling capacity on human immune cells, mainly through TLR2. These results suggest that direct TLR2 signalling events on immune cells could be part of the mechanism by which IL-10 production is induced in *in vivo*  $\beta$ 2→1-fructan supplementation studies. Also we show structure-function relationships *in vitro* for  $\beta$ 2→1-fructans which illustrate that caution should be taken in ascribing beneficial health effects to families of molecules with similar structural features, as seemingly minor differences in chain length could induce opposite effects.

## Author Contributions

Conceived and designed the experiments: LV UR DM GP KV MMF HAS PdV. Performed the experiments: LV UR. Analyzed the data: LV UR MMF HAS PdV. Contributed reagents/materials/analysis tools: LV UR DM GP HAS PdV. Wrote the paper: LV MMF PdV.

## References

- acobs DR Jr, Andersen LF, Blomhoff R (2007) Whole-grain consumption is associated with a reduced risk of noncardiovascular, noncancer death attributed to inflammatory diseases in the iowa women's health study. *Am J Clin Nutr* 85: 1606–1614.
- Park Y, Subar AF, Hollenbeck A, Schatzkin A (2011) Dietary fiber intake and mortality in the NIH-AARP diet and health study. *Arch Intern Med* 171: 1061–1068.
- Chuang SC, Norat T, Murphy N, Olsen A, Tjønneland A, et al. (2012) Fiber intake and total and cause-specific mortality in the european prospective investigation into cancer and nutrition cohort. *Am J Clin Nutr* 96: 164–174.
- Landberg R (2012) Dietary fiber and mortality: Convincing observations that call for mechanistic investigations. *Am J Clin Nutr* 96: 3–4.
- Fardet A (2010) New hypotheses for the health-protective mechanisms of whole-grain cereals: What is beyond fibre? *Nutr Res Rev* 23: 65–134.
- Gibson GR, Roberfroid MB (1995) Dietary modulation of the human colonic microbiota: Introducing the concept of prebiotics. *J Nutr* 125: 1401–1412.
- Vos AP, M'Rabet L, Stahl B, Boehm G, Garssen J (2007) Immune-modulatory effects and potential working mechanisms of orally applied nondigestible carbohydrates. *Crit Rev Immunol* 27: 97–140.
- Seifert S, Watzl B (2007) Inulin and oligofructose: Review of experimental data on immune modulation. *J Nutr* 137: 2563S–2567S.
- Lomax AR, Calder PC (2009) Probiotics, immune function, infection and inflammation: A review of the evidence from studies conducted in humans. *Curr Pharm Des* 15: 1428–1518.
- Roberfroid M, Gibson GR, Hoyles L, McCartney AL, Rastall R, et al. (2010) Prebiotic effects: Metabolic and health benefits. *Br J Nutr* 104 Suppl 2: S1–63.
- Vogt LM, Meyer D, Pullens G, Faas MM, Smelt MJ, et al. Immunological properties of inulin-type fructans. Critical reviews in food science and nutrition (in press) DOI:10.1080/10408398.2012.656772.
- Swanson KS, Grieshop CM, Flickinger EA, Healy HP, Dawson KA, et al. (2002) Effects of supplemental fructooligosaccharides plus mannanoligosaccharides on immune function and ileal and fecal microbial populations in adult dogs. *Arch Tierernahr* 56: 309–318.
- Hosono A, Ozawa A, Kato R, Ohnishi Y, Nakanishi Y, et al. (2003) Dietary fructooligosaccharides induce immunoregulation of intestinal IgA secretion by murine peyer's patch cells. *Biosci Biotechnol Biochem* 67: 758–764.
- Nakamura Y, Nosaka S, Suzuki M, Nagafuchi S, Takahashi T, et al. (2004) Dietary fructooligosaccharides up-regulate immunoglobulin A response and polymeric immunoglobulin receptor expression in intestines of infant mice. *Clin Exp Immunol* 137: 52–58.
- Meijer K, de Vos P, Priebe MG (2010) Butyrate and other short-chain fatty acids as modulators of immunity: What relevance for health? *Curr Opin Clin Nutr Metab Care* 13: 715–721.
- Vinolo MA, Rodrigues HG, Nachbar RT, Curi R (2011) Regulation of inflammation by short chain fatty acids. *Nutrients* 3: 858–876.
- Maslowski KM, Vieira AT, Ng A, Kranich J, Sierro F, et al. (2009) Regulation of inflammatory responses by gut microbiota and chemoattractant receptor GPR43. *Nature* 461: 1282–1286.
- Vinolo MA, Rodrigues HG, Hatanaka E, Hebeda CB, Farsky SH, et al. (2009) Short-chain fatty acids stimulate the migration of neutrophils to inflammatory sites. *Clin Sci (Lond)* 117: 331–338.
- Park JS, Lee EJ, Lee JC, Kim WK, Kim HS (2007) Anti-inflammatory effects of short chain fatty acids in IFN-gamma-stimulated RAW 264.7 murine macrophage cells: Involvement of NF-kappaB and ERK signaling pathways. *Int Immunopharmacol* 7: 70–77.
- Vinolo MA, Rodrigues HG, Hatanaka E, Sato FT, Sampaio SC, et al. (2011) Suppressive effect of short-chain fatty acids on production of proinflammatory mediators by neutrophils. *J Nutr Biochem* 22: 849–855.
- McDole JR, Wheeler LW, McDonald KG, Wang B, Konjufca V, et al. (2012) Goblet cells deliver luminal antigen to CD103+ dendritic cells in the small intestine. *Nature* 483: 345–349.
- Lelouard H, Fallet M, de Bovis B, Meresse S, Gorvel JP (2012) Peyer's patch dendritic cells sample antigens by extending dendrites through M cell-specific transcellular pores. *Gastroenterology* 142: 592–601.e3.
- Brown GD, Taylor PR, Reid DM, Willment JA, Williams DL, et al. (2002) Dectin-1 is a major beta-glucan receptor on macrophages. *J Exp Med* 196: 407–412.
- Underhill DM, Rossnagle E, Lowell CA, Simmons RM (2005) Dectin-1 activates syk tyrosine kinase in a dynamic subset of macrophages for reactive oxygen production. *Blood* 106: 2543–2550.
- Underhill DM, Ozinsky A, Hajjar AM, Stevens A, Wilson CB, et al. (1999) The toll-like receptor 2 is recruited to macrophage phagosomes and discriminates between pathogens. *Nature* 401: 811–815.
- Janeway CA Jr (1989) Approaching the asymptote? evolution and revolution in immunology. *Cold Spring Harb Symp Quant Biol* 54 Pt 1: 1–13.
- Cambi A, Figdor CG (2003) Dual function of C-type lectin-like receptors in the immune system. *Curr Opin Cell Biol* 15: 539–546.
- Abreu MT (2010) Toll-like receptor signalling in the intestinal epithelium: How bacterial recognition shapes intestinal function. *Nat Rev Immunol* 10: 131–144.
- Kelly G (2008) Inulin-type prebiotics—a review: Part 1. *Altern Med Rev* 13: 315–329.
- Niness KR (1999) Inulin and oligofructose: What are they? *J Nutr* 129: 1402S–6S.
- Chen Z, Cheng Y, Xu Y, Liao J, Zhang X, et al. (2008) Expression profiles and function of toll-like receptors 2 and 4 in peripheral blood mononuclear cells of chronic hepatitis B patients. *Clin Immunol* 128: 400–408.
- Siednienko J, Miggin SM (2009) Expression analysis of the toll-like receptors in human peripheral blood mononuclear cells. *Methods Mol Biol* 517: 3–14.
- Joosten LA, Heinhuis B, Abdollahi-Roodsaz S, Ferwerda G, Lebourhis L, et al. (2008) Differential function of the NACHT-LRR (NLR) members Nod1 and Nod2 in arthritis. *Proc Natl Acad Sci U S A* 105: 9017–9022.
- Meijerink M, van Hemert S, Taverne N, Wels M, de Vos P, et al. (2010) Identification of genetic loci in *Lactobacillus plantarum* that modulate the immune response of dendritic cells using comparative genome hybridization. *PLoS One* 5: e10632.
- van Hemert S, Meijerink M, Molenaar D, Bron PA, de Vos P, et al. (2010) Identification of *Lactobacillus plantarum* genes modulating the cytokine response of human peripheral blood mononuclear cells. *BMC Microbiol* 10: 293–2180–10–293.
- Shalhoub J, Falck-Hansen MA, Davies AH, Monaco C (2011) Innate immunity and monocyte-macrophage activation in atherosclerosis. *J Inflamm (Lond)* 8: 9.
- Hayden MS, Ghosh S (2004) Signaling to NF-kappaB. *Genes Dev* 18: 2195–2224.
- Abadie V, Discepolo V, Jabri B (2012) Intraepithelial lymphocytes in celiac disease immunopathology. *Semin Immunopathol* 34: 551–566.
- Zheng H, Luo X, Segura M, Sun H, Ye C, et al. (2012) The role of toll-like receptors in the pathogenesis of streptococcus suis. *Vet Microbiol* 156: 147–156.
- Kwon HK, Lee CG, So JS, Chae CS, Hwang JS, et al. (2010) Generation of regulatory dendritic cells and CD4+Foxp3+ T cells by probiotics administration suppresses immune disorders. *Proc Natl Acad Sci U S A* 107: 2159–2164.
- Meijerink M, Wells JM, Taverne N, de Zeeuw Brouwer ML, Hilhorst B, et al. (2012) Immunomodulatory effects of potential probiotics in a mouse peanut sensitization model. *FEMS Immunol Med Microbiol* 65: 488–496.
- Hamer HM, Jonkers DM, Vanhoutvin SA, Troost FJ, Rijkers G, et al. (2010) Effect of butyrate enemas on inflammation and antioxidant status in the colonic mucosa of patients with ulcerative colitis in remission. *Clin Nutr* 29: 738–744.
- Starkie R, Ostrowski SR, Jauffred S, Febbraio M, Pedersen BK (2003) Exercise and IL-6 infusion inhibit endotoxin-induced TNF-alpha production in humans. *FASEB J* 17: 884–886.
- Petersen AM, Pedersen BK (2005) The anti-inflammatory effect of exercise. *J Appl Physiol* 98: 1154–1162.
- Eiwegger T, Stahl B, Haidl P, Schmitt J, Boehm G, et al. (2010) Prebiotic oligosaccharides: In vitro evidence for gastrointestinal epithelial transfer and immunomodulatory properties. *Pediatr Allergy Immunol* 21: 1179–1188.
- Janardhana V, Broadway MM, Bruce MP, Lowenthal JW, Geier MS, et al. (2009) Probiotics modulate immune responses in the gut-associated lymphoid tissue of chickens. *J Nutr* 139: 1404–1409.
- Benyacoub J, Rochat F, Saudan KY, Rochat I, Antille N, et al. (2008) Feeding a diet containing a fructooligosaccharide mix can enhance salmonella vaccine efficacy in mice. *J Nutr* 138: 123–129.
- Saavedra JM, Tschernia A (2002) Human studies with probiotics and prebiotics: Clinical implications. *Br J Nutr* 87 Suppl 2: S241–6.
- van Hoffen E, Ruter B, Faber J, M'Rabet L, Knol EF, et al. (2009) A specific mixture of short-chain galacto-oligosaccharides and long-chain fructo-oligosaccharides induces a beneficial immunoglobulin profile in infants at high risk for allergy. *Allergy* 64: 484–487.
- Lindsay JO, Whelan K, Stagg AJ, Gobin P, Al-Hassi HO, et al. (2006) Clinical, microbiological, and immunological effects of fructo-oligosaccharide in patients with crohn's disease. *Gut* 55: 348–355.
- Wells JM, Rossi O, Meijerink M, van Baaren P (2011) Epithelial crosstalk at the microbiota-mucosal interface. *Proc Natl Acad Sci U S A* 108 Suppl 1: 4607–4614.
- Roller M, Rechkemmer G, Watzl B (2004) Prebiotic inulin enriched with oligofructose in combination with the probiotics *Lactobacillus rhamnosus* and *Bifidobacterium lactis* modulates intestinal immune functions in rats. *J Nutr* 134: 153–156.
- Roller M, Pietro Femia A, Caderni G, Rechkemmer G, Watzl B (2004) Intestinal immunity of rats with colon cancer is modulated by oligofructose-enriched inulin combined with *Lactobacillus rhamnosus* and *Bifidobacterium lactis*. *Br J Nutr* 92: 931–938.
- Manhart N, Spittler A, Bergmeister H, Mittlbock M, Roth E (2003) Influence of fructooligosaccharides on peyer's patch lymphocyte numbers in healthy and dioxinemic mice. *Nutrition* 19: 657–660.
- Ito H, Takemura N, Sonoyama K, Kawagishi H, Topping DL, et al. (2011) Degree of polymerization of inulin-type fructans differentially affects number of lactic acid bacteria, intestinal immune functions, and immunoglobulin A secretion in the rat cecum. *J Agric Food Chem* 59: 5771–5778.
- Delgado MA, Deretic V (2009) Toll-like receptors in control of immunological autophagy. *Cell Death Differ* 16: 976–983.
- de Kivit S, Kraneveld AD, Garssen J, Willemsen LE (2011) Glycan recognition at the interface of the intestinal immune system: Target for immune modulation via dietary components. *Eur J Pharmacol*.

58. Kerrigan AM, Brown GD (2010) Syk-coupled C-type lectin receptors that mediate cellular activation via single tyrosine based activation motifs. *Immunol Rev* 234: 335–352.
59. Bruns AM, Horvath CM (2012) Activation of RIG-I-like receptor signal transduction. *Crit Rev Biochem Mol Biol* 47: 194–206.
60. Kawai T, Akira S (2010) The role of pattern-recognition receptors in innate immunity: Update on toll-like receptors. *Nat Immunol* 11: 373–384.
61. Oliveira-Nascimento L, Massari P, Wetzler LM (2012) The role of TLR2 in infection and immunity. *Front Immunol* 3: 79.
62. Sato M, Sano H, Iwaki D, Kudo K, Konishi M, et al. (2003) Direct binding of toll-like receptor 2 to zymosan, and zymosan-induced NF-kappa B activation and TNF-alpha secretion are down-regulated by lung collectin surfactant protein A. *J Immunol* 171: 417–425.
63. Visintin A, Latz E, Monks BG, Espevik T, Golenbock DT (2003) Lysines 128 and 132 enable lipopolysaccharide binding to MD-2, leading to toll-like receptor-4 aggregation and signal transduction. *J Biol Chem* 278: 48313–48320.