

1 Individualized on-line monitoring tool to analyse the complex physiological signal of 2 heart beat fluctuations in pigs

3
4 I. D. E. van Dixhoorn ^a, G. André ^{a,c}, B. Lambooij ^a, B. Kemp ^b, P. W. G. Groot Koerkamp ^{a,c}

5 ^a Wageningen UR Livestock Research, P.O. box 65, 8200 AB, Lelystad, The Netherlands

6 ^b Adaptation Physiology Group, Department of Animal Sciences, Wageningen University,
7 P.O. Box 338, 6700 AH, Wageningen, the Netherlands

8 ^c Farm Technology Group, Wageningen University, P.O. box 317, 6700 AH, Wageningen,
9 The Netherlands

10 ingrid.vandixhoorn@wur.nl

11 12 Abstract

13
14 Quantifying and understanding the complex fluctuations of physiological signals is the focus
15 of many research. The complexity of physiological signals reflect the ability of organisms to
16 adapt and function within an ever changing environment. The most studied physiological
17 signal to reflect the autonomic imbalance in cases of disease, chronic stress and impaired
18 welfare, in both human and animals is Heart Rate Variability (HRV). Given the limitations of
19 current HRV analyses, like spectral or total entropy analyses, it is suggested to improve our
20 understanding by decomposing the heart rate fluctuations with a dynamic model, based on a
21 Bayesian approach for time series analysis.

22 The model consists of three components describing the interbeat-intervals dynamics: (1) level
23 and trend, (2) autoregressive process of order 2 and (3) observation error (white noise). The
24 analysis also detects abrupt changes and slightly growing deteriorations. A dataset with
25 continuous ECG recordings from 2 pigs was analysed. The model decomposed total Heart
26 Rate dynamics successfully in the time domain. So, it was possible to detect abrupt changes
27 and slightly growing deteriorations in RR-interval in terms of the moments of occurrence, the
28 magnitude, the duration and recovery.

29
30 **Key words:** Dynamic modelling, Bayesian forecasting, inter-beat interval, heart rate
31 variability, autonomic nervous system

32 33 Introduction

34
35 Early detection of alteration in health status is of great importance in farm animals, as early
36 intervention after e.g. infection can minimize symptoms, may shorten the recovery period and
37 diminishes production losses due to disease. Most early warning systems in animals focus on
38 the early detection of disease symptoms or infectious agents. When a certain population of
39 pathogens is present, it is uncertain whether, when and to what extent animals will get ill or
40 how capable the animals are to cope with the present pathogenic load without showing
41 symptoms. This is caused by differences in the capability to maintain a certain health state
42 under varying conditions.

43 The complexity of cardiovascular dynamics has been studied as an indicator for autonomic
44 balance or health state. To describe the variations in both instantaneous heart rate and

45 fluctuations in RR-intervals over time, the term that has become conventionally accepted is
46 “Heart Rate Variability“ (HRV) (Malik 1996). In both humans and animals, high complexity
47 of HRV is associated with good health and high adaptive capacity. Low complexity of HRV
48 has become a popular marker for autonomic imbalance in cases of disease, chronic stress and
49 impaired welfare (Malik 1996; Goldberger, Challapalli et al. 2001; von Borell, Langbein et al.
50 2007) and has gained widespread acceptance as a clinical and investigational tool (Billman
51 2011).

52 HRV is usually measured under one of the two common settings: short term measurements
53 under controlled laboratory conditions, or long term measurements derived from 24 hour
54 ECG-recordings made while performing daily activities. The current methods used to describe
55 HRV deliver conventional time domain measures, spectral measures, geometric measures,
56 and a variety of nonlinear variables. Each single variable derived from the different methods
57 reflects a different aspect of HRV. They all show significant associations with physiological
58 outcome or state, but it is still unclear which is the best variable to describe and assess HRV
59 (Kleiger, Stein et al. 2005). Therefore, the clinical utility and predictive value of the existing
60 HRV measurements is under discussion, which focusses on (1) the relationship between the
61 autonomic nervous system (parasympathic and sympathetic effects) and the descriptive
62 variables of HRV and (2) the substantial variance in HRV within and between normal
63 individuals (Goldberger, Challapalli et al. 2001; Peng, Costa et al. 2009). It can be concluded
64 that we still lack an effective reconstruction method for the phenomenon of interbeat
65 fluctuations in heart rate, that is characterized by a non-stationary process and a degree of
66 stochasticity (Ghasemi, Sahimi et al. 2006).

67 The goal of this paper is to show the first experiences with an adaptive dynamic model that is
68 able to analyse the non-stationary and non-linear heart rate fluctuations in the time-domain in
69 terms of fast and slow variation. We used the model to analyse time series of ECG recordings
70 (RR-intervals) of 2 pigs to show that it reveals the non-stationary complexity of the cardiac
71 function both in terms of fast and slow variation simultaneously, as well as the detection of
72 the moment of occurrence and extent of changes in its dynamics. Results are presented and it
73 is discussed whether this model can be used in longitudinal on farm settings as an online
74 monitoring tool to detect process disturbances in early stage.

75

76 **Materials & Methods**

77

78 Experimental protocol

79 Four crossbred pigs (Yorkshire x Landrace) were kept together in a pen with rubber flooring
80 (150 cm wide and 120 cm deep) from 11 days until 86 days of age. Milk replacer (Sprayfo
81 Pork, Husdyr Systemer as, Mosby, Norge) was provided *ad libitum* with an automatic
82 wetfeeder (Mambo, Husdyr Systemer as, Mosby, Norge). Milk replacer was slowly reduced
83 until the Mambo was removed at 46 days of age. Solid piglet feed (pellets) were available *ad*
84 *libitum* from the start of the experiment. Water was provided *ad libitum* during the
85 experimental period. The pigs were exposed to a 12h light (6am until 6 pm) - 12 h dark (6pm
86 until 6 am) photoperiod. Temperature was kept at 28° C from start until 35 days, at 25° C until
87 64 days age and at 24° C until the end of the experiment. Average relative humidity of the air

88 was 55% during the experiment. The pigs had 2 weeks for acclimatization to the housing
89 facility and weaning regime. At the age of 26 days, two of the four pigs were surgically
90 equipped (intra-abdominal) with implantable telemetry transmitters (Data Science
91 International, DSI, St Paul, MN) to record continuously temperature, activity and cardiac
92 activity (ECG) until the end of the experiment. When the pigs were 35 days old, the two pigs
93 without implants were removed from the pen to enlarge individual space. The growth curve of
94 the two pigs with implants was similar to the curves of the animals that did not have surgical
95 treatment. One pig at the time was monitored during 24 hours (weekdays) and 72 hours
96 (weekends). Every day the pens were cleaned and the animals were inspected clinically.
97 During the experimental period no signs of illness appeared.¹

98

99 Data acquisition telemetry system & video recording

100 The telemetry system (Data Science International, DSI, St Paul, MN, USA) consisted of
101 implantable transmitters with two bio-potential leads (positive and negative) for measuring
102 ECG signal, a temperature sensor to measure body temperature and one blood pressure fluid
103 filled catheter (model TL11M2-D70-PCT; 49 grams, 33cc), a data exchange matrix and
104 receivers (DSI PhysioTel® Receivers - RMC-1 Model for Large Animals). Only the ECG
105 recordings were analysed in this paper. Signal strength between transmitter and receiver was
106 recorded and expressed in a physical meaningless number ranging from 0 to 51 units. No data
107 were recorded when signal strength dropped below 17 units. During the experimental period
108 video recordings were made with a Sanyo (type RC506CH) camera and Samsung SHR-2040
109 digital recording system of the complete pen during the light period (6am until 6pm).
110 Telemetry data of the implants were collected with DSI Dataquest A.R.T.TM version 4.31. The
111 ECG signal was measured and stored at 1 kHz.

112

113 Data pre-processing

114 The length (time in msec) of the RR-interval was derived from the raw ECG data with the
115 Ponemah Physiology Platform version 5.0 software (Data Science International, DSI, St Paul,
116 MN, USA). Due to weak radio or transfer signal or high noise some R-peaks were incorrectly
117 detected, resulting in extreme RR-intervals ($RR \leq 200$ msec or $RR \geq 700$ msec). These RR-
118 intervals (denoted as extreme RR-intervals), together with sections where no PQRS complex
119 could be detected (denoted as missing RR-intervals), were omitted from the data series.
120 Subsequently, equidistant time series were formed by averaging the RR-intervals per second
121 (further indicated as averaged or observed RR-intervals). Missing values in the time series
122 were replaced by interpolated data and weighted zero in the analysis.

123

¹ The established principles of laboratory animal use and care were followed as well as the Dutch law on animal experiments. The Wageningen University Animal Care and Use Committee (Lelystad Department) approved the experiment under number 2011040.f.

124 Data selection

125 From the experiments we randomly selected and analysed one hour of one pig (pig #231 from
126 15:00 until 16:00 on the 23d of January 2012). For detailed explanation of the Bayesian
127 model for analysis of HRV, two periods of 5 minutes were selected within this hour. The two
128 periods of 5 minutes were selected, one with high and one with low activity of the pig based
129 on the video recordings.

130


131 Modelling and analysis

132 Variation in interbeat intervals was decomposed in three parts:

133
$$\left\{ \begin{array}{l} \text{interbeat} \\ \text{interval} \end{array} \right\} = \left\{ \begin{array}{l} \text{level\&} \\ \text{trend} \end{array} \right\} + \left\{ \begin{array}{l} \text{autoregression} \\ \text{component} \end{array} \right\} + \left\{ \begin{array}{l} \text{white} \\ \text{noise} \end{array} \right\}$$

134

135 Level and trend represent slow dynamic variation, the autoregression component and the
136 white noise represent fast dynamic variation. The white noise represents the independent
137 random residual error. Figure 1 shows a simulated series of 5 min (=300 sec) illustrating the
138 3 components of the model.



139

140

141 Figure 1: Simulated time series of RR intervals of 5 minutes (300 sec.) showing the 3
142 components: level and trend (thick straight line); auto-regression component (thin fluctuating
143 line) and white noise scattered around the thin line (points).

144

145 A dynamic model was formulated for analysis of the time series of averaged RR-intervals per
146 second Y_t . The dynamic model consisted of an observation and a system equation. The
147 observation equation (1) described the relation between the observation and the parameters:

148
$$Y_t = \mu_t + Z_t + v_t \tag{1}$$

149 with the parameters: μ_t (level), Z_t (autoregression, AR2) and $v_t \sim N(0, k_t V_t)$ (observation
150 error or white noise, with unknown variance V_t and weight k_t equal to the number of beats per
151 second). The system equations (2) and (3) describe the evolution of the parameters over time:

152

$$\begin{aligned}\mu_t &= \mu_{t-1} + \beta_{t-1} + \omega_{1,t} \\ \beta_t &= \beta_{t-1} + \omega_{2,t}\end{aligned}\quad (2)$$

153

$$\begin{aligned}Z_t &= \phi_{1,t-1}Z_{t-1} + \phi_{2,t-1}Z_{t-2} + \omega_{3,t} \\ \phi_{1,t} &= \phi_{1,t-1} + \omega_{4,t} \\ \phi_{2,t} &= \phi_{2,t-1} + \omega_{5,t}\end{aligned}\quad (3)$$

154

155 with level μ_t , trend or incremental change β_t , two autoregression coefficients $\phi_{1,t}$ and $\phi_{2,t}$ 156 with system errors $\omega_{1...5,t}$ normally distributed with zero mean and variance matrix \mathbf{W}_t . The

157 dynamic parameters are locally constant and follow a random walk. Two kinds of parameter

158 estimates were achieved: online estimates, which were based on observations from the past

159 only, and retrospective estimates, which were based on all observations from the whole time

160 series. All parameters were recursively estimated following the Bayesian approach to the

161 analysis of time series according to (West and Harrison 1997) including a monitoring

162 procedure followed by automatic intervention to detect process disturbances.

163 The results describing variation in the time domain were linked to results in the frequency

164 domain. Level and trend corresponded to ultralow frequency variation, the pseudo (or

165 stochastic) cyclic behaviour, described by the autoregression AR(2) component corresponded

166 with low frequency variation and the white noise (or random error) corresponded to the high

167 frequency variation. From the parameter estimates at any point in time the correlogram,

168 spectrum and variances were calculated (Diggle 1990).

169 **Results and Discussion**

170

171 Within the complete analysed hour, 147 RR-intervals were indicated as extreme values and

172 left out of this analysis. Out of the 3600 averaged RR-intervals, 3480 intervals were classified

173 by the monitoring routine as normal, 51 as warnings and 69 as outliers. According to the

174 video recordings, between 15:00 and 15:30, the pigs were mostly resting/sleeping. After this

175 period they awoke, stood up and started eating and drinking, at 15:44 the pigs lied down

176 again.

177 The results of the analysis are shown in figures 2 to 5. Figures 2a and b show the observed


178 (averaged per second) and forecasted RR intervals for the two periods of 5 minutes within the

179 hour. Figure 2a shows the period of 5 minutes in which the pig was mostly resting and figure


180 2b shows a more active period where the pig was eating, drinking and playing with its pen

181 mate. Even during these short periods a substantial variation in RR-intervals was detected.

182



183
184




185


186 Figure 2a&b Observed and forecasted RR-interval; 2a: Selection of 5 min. period (resting):
187 above and 2b (active): below; observed values (grey points), forecasts (black line), upper and
188 lower level of 90% confidence interval (light grey lines).

189

190 Figure 3a and b show the estimated parameters for level (μ_t) and trend (β_t) for the selected
191 hour, representing the slow changing dynamics. Level was accurately estimated at any point
192 in time. More active periods were described by a shorter RR-interval and changes in level
193 were easily picked up, even sudden changes. Figure 4b shows the trend β_t . In the first period
194 of the hour, the trend was not significantly different from zero (the confidence intervals were
195 not above or below zero). It resulted in a more or less constant level, which is shown in figure
196 3a. Only at t=15:36 a significant negative trend was seen, followed by a significant positive
197 trend, which resulted in an temporary reduction of RR-interval. This is the period the pig was
198 active and was detected by a change in level and trend by the model. Thus, the moment of
199 occurrence of changes in trend of RR-intervals were properly detected by the analysis and
200 could directly be related to the behaviour of the animals. Rapid changes (e.g. at t=15:10 and
201 t=15:12) did not result in a change in level or trend. However, the variance of the process
202 increased, which resulted in a larger confidence interval.



203



204

205 Figure 3a: Estimated level μ_t (above) and 3b: Trend β_t or incremental growth (below);
 206 online estimates (grey points); retrospective estimates (black line) incl. 90% confidence
 207 interval (light grey lines)

208

209 In fig 4a and 4b the parameter estimates that describe the fast dynamics changes are shown.
 210 These are the autoregression component Z_t , the forecast variance Q_t , and observation variance
 211 S_t . The variances fluctuated, even during both 5 minute periods. The variances increased,
 212 especially after abrupt changes. The two AR-parameters $\phi_{1,t}$ and $\phi_{2,t}$ described the properties
 213 of the AR(2)-process, i.e. the shape and rate of decay of random deviations from the base
 214 level. In figure 5 $\phi_{1,t}$ and $\phi_{2,t}$ are plotted for both selected periods of 5 minutes.

215


216 The AR-process is stable when the AR-parameters lie within the triangle (figure 5). If they lie
 217 outside the triangle, the process is unstable and the heart rate will exponentially grow or
 218 decline which is physiologically not possible in living organisms. When the AR-parameters
 219 lie within the triangle and below the parabola the decay of the deviations follows a decaying
 220 cycle, otherwise they follow a decaying curve.

220

221 The stochastic cycle during the resting and active period showed different characteristics. It
 222 appeared that when the pig was resting, the AR-parameters lied clustered together and more
 223 to the left in the upper part above the parabola. In this area, the fluctuations around the base
 224 level are characterized by a relatively fast exponential decay (dark grey cluster in figure 5).
 When the pig was more active, the estimated AR-parameters lied in a strip near to the right

225 side of the triangle (light grey strip in figure 5) where the decay of fluctuations appear to be
 226 slower (light grey line in figure 5).

227




228

229

230 Figure 4a & b: Autoregression component Z_t and variances within the two selected periods of
 231 5 minutes; 4a sleep (above) 4b active (below): Online estimates (grey points); retrospective
 232 estimates (black line) incl. 90% confidence interval (light grey lines).

233 Forecast variance Q_t (-----) and observation variance S_t (- . -)

234



235

236

237 Figure 5 Retrospective estimated AR parameters $\phi_{1,t}$ and $\phi_{2,t}$ showing the two selected 5
 238 minute periods while sleeping (dark grey line 15:03-15:08) and active (light grey line 15:31-
 239 15:36).

240

241 The first two components level (μ_t) and trend (β_t) adjust the base level of heart rate to the
 242 required set point according the behaviour and physiological demands at that moment.

243 Significant trends appeared whenever an increase or decrease in base level is required to meet
244 the physiological needs. Changes in RR-interval at arousal were described by a decreasing
245 trend, followed by a slow increasing trend to adjust to the desired level again. Sudden fast
246 changes in level (mainly drops in RR-intervals) were not detected by trend since the process
247 variance increased simultaneously. This indicates that the level and trend components of the
248 model reflect the slow dynamic changes in RR-interval.

249 Within 5 minute periods substantial variation in interbeat interval can be seen. This actual,
250 more subtle, variation in RR-intervals cannot be described accurately by models based on the
251 assumption of a stationary process, like the Fast Fourier Transformation (FFT). They were,
252 however, captured by the decomposing model, which moment by moment follows every
253 fluctuation and any changes in dynamics. These fast dynamic fluctuations around the slowly
254 changing base level is described by the autoregression and error components, corresponding
255 to low and high frequency variation. The AR2-process ensured that the modelled RR-interval
256 oscillated around the base level that was set by level and trend. This process with more or less
257 decaying deviations can be seen as continuously fine-tuning towards the desired heart rate.

258 As expected, the random error component v_t could not be captured within any pattern or
259 related to any cause. This is the white noise component, which is shown as random scattering
260 around the AR oscillations as shown in figure 1. As shown in figure 4 the variance of this
261 white noise component was not constant within the 5 minute periods. So, the white noise
262 variance is dynamically changing even within short periods.

263 The parameters were estimated online at each moment of the time series and together
264 characterized the total dynamic variation in RR-interval and the dependency between
265 successive observations in the actual situation. The accuracy of the online estimates was
266 improved afterwards by the backward smoothing procedure, resulting in the retrospective
267 estimates.

268 269 **Conclusion**

270
271 Our model is based on a beat to beat calculation, which is necessary for timely detection of
272 the dynamical changes in HRV that apparently occur continuously over time. Techniques that
273 cannot calculate on a beat to beat basis, such as other time or frequency domain techniques,
274 overlook the dynamic changes when a mean value over a longer time period is calculated, and
275 therefore provide less (detailed) information.

276 We concluded that the adaptive dynamic model is able to analyse the non-stationary and non-
277 linear heart rate fluctuations in the time-domain in terms of fast and slow variation. The
278 model detects sudden changes as well as slightly growing deteriorations.

279 These characterisations of dynamics will differ depending on the physiological state of the
280 animal and may be individually different. Further research will focus on the relation between
281 estimated parameter values and the physiological state of animals. After the physiological
282 interpretation of the parameters with respect to the autonomic nervous system the model will
283 be further developed as an online monitoring tool that can be used to detect early alterations
284 in physiological state of animals in husbandry systems.

285
286

287 **References**

288

- 289 Billman, G. E. (2011). "Heart rate variability - a historical perspective." Frontiers in
290 Physiology **2**(86): 1-13.
- 291 Conte, E., A. Federici, et al. (2009). "A new method based on fractal variance function for
292 analysis and quantification of sympathetic and vagal activity in variability of R-R time
293 series in ECG signals." Chaos, Solitons and Fractals **41**(3): 1416-1426.
- 294 Diggle, P. J. (1990). Time series: a biostatistical introduction, Oxford University Press, New
295 York.
- 296 Ghasemi, F., M. Sahimi, et al. (2006). "Analysis of non-stationary data for heart-rate
297 fluctuations in terms of drift and diffusion coefficients." Journal of Biological Physics
298 **32**(2): 117-128.
- 299 Gleick, J. (1987). Chaos, making a new science, Penguin Group USA.
- 300 Goldberger, A. L. (1989). "Cardiac chaos [2]." Science **243**(4897): 1419.
- 301 Goldberger, A. L., D. R. Rigney, et al. (1990). "Chaos and fractals in human physiology."
302 Scientific American **262**(2): 42-49.
- 303 Goldberger, J. J., S. Challapalli, et al. (2001). "Relationship of heart rate variability to
304 parasympathetic effect." Circulation **103**(15): 1977-1983.
- 305 Kleiger, R. E., P. K. Stein, et al. (2005). "Heart rate variability: Measurement and clinical
306 utility." Annals of Noninvasive Electrocardiology **10**(1): 88-101.
- 307 Malik, M. (1996). "Heart rate variability: Standards of measurement, physiological
308 interpretation, and clinical use." Circulation **93**(5): 1043-1065.
- 309 Peng, C. K., S. V. Buldyrev, et al. (1994). "Non-equilibrium dynamics as an indispensable
310 characteristic of a healthy biological system." Integrative Physiological and
311 Behavioral Science **29**(3): 283-293.
- 312 Peng, C. K., M. Costa, et al. (2009). "Adaptive data analysis of complex fluctuations in
313 physiologic time series." Advances in Adaptive Data Analysis **1**(1): 61-70.
- 314 von Borell, E., J. Langbein, et al. (2007). "Heart rate variability as a measure of autonomic
315 regulation of cardiac activity for assessing stress and welfare in farm animals - A
316 review." Physiology and Behavior **92**(3): 293-316.
- 317 West, M. and J. Harrison (1997). Bayesian Forecasting and Dynamic Models. New
318 York. West, Springer.

319

320