

**The *Rubetum taxandriæ* ass. nov.
(*Lonicero-Rubion silvatici*, *Lonicero-Rubetea plicati*),
a new bramble association from the Belgian and Dutch Campine**

– Rense Haveman, Iris de Ronde & Eddy J. Weeda –

Abstract

Bramble scrubs are among the least known and understood vegetation types in Europe. In the Dutch National Vegetation Overview, three associations belonging to the *Lonicero-Rubion silvatici* were distinguished, viz. the *Rubetum grati*, *Rubetum silvatici*, and *Rubetum pedemontani*. During several vegetation mapping projects and *Rubus* excursions, a distinct type of bramble scrub was recorded repeatedly in the Campine in the province Noord-Brabant in the southern part of the Netherlands. In this paper, this scrub is described as a new association, the *Rubetum taxandriæ* Haveman, de Ronde & Weeda, with *R. taxandriæ*, *R. campaniensis*, and *R. baronicus* as character species, and *R. insectifolius* as regional character species. Variation, ecology and distribution of this new association are given and discussed, and two subassociations are distinguished. The differences with the *Rubetum silvatici*, to which this community was believed to belong, are discussed. Based on an earlier analysis of the centres of diversity of the genus *Rubus* in the Netherlands, it is supposed that the *Rubetum silvatici* in its circumscription in the Dutch National Vegetation Overview can be divided in more regionally distributed communities, partly as subassociations of the *Rubetum silvatici*, partly as independent associations.

**Zusammenfassung: Das *Rubetum taxandriæ* ass. nov.
(*Lonicero-Rubion silvatici*, *Lonicero-Rubetea plicati*), eine neue Brombeer-Assoziation
im Gebiet der belgischen und niederländischen „Kempen“**

Brombeergebüsche gehören zu den am wenigsten bekannten Vegetationstypen in Europa. Bisher wurden drei Assoziationen für die Niederlande beschrieben: *Rubetum grati*, *Rubetum silvatici* und *Rubetum pedemontani*. Bei verschiedenen Kartierungsarbeiten und Exkursionen wurde in „De Kempen“ in der Provinz Noord-Brabant im südlichen Teil der Niederlande mehrmals ein abweichendes Gebüsch beobachtet. Dieses Gebüsch wird hier als neue Assoziation: *Rubetum taxandriæ* Haveman, de Ronde & Weeda beschrieben, mit *R. taxandriæ*, *R. campaniensis* und *R. baronicus* als Kennarten und *R. insectifolius* als regionaler Kennart. Artenverbindung, Ökologie und Verbreitung werden erörtert, und zwei Subassoziationen werden beschrieben. Der Unterschied zum *Rubetum silvatici* wird diskutiert. Aufgrund einer früher publizierten Analyse von Diverzitätszentren der Gattung *Rubus* in den Niederlanden wird angenommen, dass das *Rubetum silvatici* dort in mehrere regional verbreitete Gebüschtypen aufgespalten werden kann, teilweise als Subassoziationen, teilweise vielleicht besser als eigene Assoziation.

Keywords: apomicts, bramble scrub, syntaxonomy, *Rubus*, woodland edge.

1. Introduction

Bramble scrubs are among the least known and understood vegetation types in Europe. In the comprehensive overview of the scrubs in the temperate and boreal parts of Europe given by WEBER (1997, 1998b), relevés in which apomictic *Rubus* species were adequately identified had almost only been made in Germany. The bramble scrubs of Northwestern Europe were placed in separate alliances and in two separate classes. The scrubs of relatively nutrient rich and/or base rich soils are united in the *Pruno-Rubion radulae* Weber 1974, an alliance of the *Rhamno-Prunetea* Rivas Goday & Borja Carbonell ex Tüxen 1962, whereas the *Lonicero-Rubion silvatici* Tüxen et Neumann ex Wittig 1977, containing scrubs of nutrient poor and acidic soils, were placed in the *Franguletea* Doing 1962 ex Westhoff in Westhoff & Den Held 1969 (POTT 1995, WEBER 1998a, 2003, 1999, 1990). In the Dutch national vegetation overview, this classification scheme was adopted for the greater part, and several *Rubus* associations were documented shortly after Weber published his overview (HAVEMAN et al. 1999a, 1999b). A remarkable difference was the description of a separate class

Lonicero-Rubetea plicati Haveman, Schaminée & Stortelder 1999 for the scrubs on nutrient poor acidic soils, as advocated by HAVEMAN (1997) some years before.

HAVEMAN et al. (1999a) distinguished three associations within the *Lonicero-Rubetea plicati* for the Netherlands: the *Rubetum grati* Tüxen et Neumann ex Weber 1976, the *Rubetum silvatici* Weber in Pott 1995, and the *Rubetum pedemontani* Weber in Pott 1995. According to WEBER (1998a), the character species of the *Rubetum silvatici* are *Rubus silvaticus*, *R. pyramidalis*, *R. flexuosa*, and *R. sprengelii* but HAVEMAN et al. (1999a) and WEEDA et al. (2005) considered many more species as such: *R. adspersus*, *R. campaniensis*, *R. drenthicus*, *R. erinulus*, *R. foliosus*, *R. glandithyrsos*, *R. lasiandrus*, *R. mucronulatus*, *R. rubercadaver*, *R. schlechtendalii* and *R. taxandriae*. However, the authors suggested that when more relevés would become available from the Netherlands, the *Rubetum silvatici* in the given circumscription would probably appear to be a ‘composite association’, comprising of a number of more clearly defined, regionally distributed vegetation types (HAVEMAN et al. 1999a, p. 96).

During the vegetation mapping projects of military training areas, and a study of the bramble scrubs along three landscape transects (HAVEMAN et al. in prep.) a distinct bramble scrub was repeatedly recorded in the Campine in the province Noord-Brabant, in the southern part of the Netherlands. The two most frequent *Rubus* species in this scrub type are *R. campaniensis* and *R. taxandriae*, which were among the species considered to be character species of the *Rubetum silvatici* by HAVEMAN et al. (1999a). Two other species occurring in these scrubs are *R. baronicus* and *R. insectifolius*. These four species are lacking in the North-German lowlands, which is the distribution centre of the *Rubetum silvatici*. At least the first two are frequently found in the southern part of the Netherlands and the adjacent part of Flanders/Belgium. Reversely, the character species of the *Rubetum silvatici* s.str. are virtually absent in the scrubs formed by *R. campaniensis* and *R. taxandriae*.

In this paper we describe this bramble scrub from the Campine as a new association, belonging to the *Lonicero-Rubion silvatici*.

2. Area

The Campine is the area between the rivers Scheldt and Meuse, roughly situated between Antwerpen and Hasselt in Belgium and Eindhoven in the Netherlands (fig. 1). The area was never covered by the ice caps during the Pleistocene, in contrast with more northern parts of the Netherlands. The area is largely covered by Pleniglacial aeolian cover sands from the Weichselian (116,000–11,500 BP), incised by many rivulets and brooks draining the area towards the two mentioned large rivers. Although the area is slightly undulating, the differences in altitude are small (0–50 m above sea level). The climate is subatlantic, with a precipitation surplus in all months (fig. 2), a total annual precipitation of 707 mm, and an annual average temperature of 9.8° C. (SLUIJTER 2011).

3. Material and methods

3.1 Relevés

For this study 38 relevés were used in which *Rubus taxandriae*, *R. campaniensis*, *R. baronicus*, and *R. insectifolius* reach a combined cover of at least 5%. They were collected between 1998 and 2011 in several mapping projects of military ranges, during several excursions in the region, and in a study of the species composition of bramble scrubs along three landscape transects (HAVEMAN et al. in prep.). The relevés are all stored in the Dutch National Vegetation Database (JANSEN et al. 2011) using TurboVeg (HENNEKENS & SCHAMINÉE 2001).

All phanerogam and cryptogam species were recorded in as many layers as necessary to describe the vegetation structure. Species names are according VAN DE BEEK et al. (in press) for the *Rubus* species, VAN DER MEIJDEN (2005) for the other phanerogams, and SIEBEL & DURING (2006) for the mosses. The abundance of the species was recorded using the modified scale of Braun-Blanquet (BARKMAN et al. 1964). Due to the diverse origin of the data, the area of the used relevés varies considerably (from 15 to 70 m²), but all relevés are considered to be homogeneous as to the species composition.

For a comparison with the *Rubetum silvatici*, we selected relevés from the Dutch National Vegetation Database (JANSEN et al. 2011, SCHAMINÉE et al. 2006) which met the following three conditions: 1.) the combined abundance of the four character species (*Rubus silvaticus*, *R. pyramidalis*, *R. flexuosa*, and *R. sprengelii*) >5%, 2.) relevé area 10–50 m², and 3.) the combined abundance of species of the *Rhamno-Prunetea* (e.g. *Crataegus monogyna*, *Prunus spinosa*) <5%.

3.2 Synthetic phase

Both the subdivision of the new association and the comparison with the *Rubetum silvatici* were performed in JUICE (TICHÝ 2002). For these analyses, we combined high, middle, and low tree layer into one tree layer, and we used two shrub layers (high for the scrubs excl. *Rubus* and lianas, low for the *Rubus* species and lianas), one herb layer (combining herbs, juveniles and saplings of trees and shrubs), and a moss layer.

To detect patterns in the species composition in the used dataset, we used the modified TWINSPAN algorithm as suggested by ROLEK et al. (2009); only one subdivision was recognised and the results given by TWINSPAN were refined subjectively on the basis of expert judgement. To compare differences in species composition, we made a combined synoptic table (in percentages, see the recommendations by DENGLER et al. 2006, p. 84) of the new association and the *Rubetum silvatici*. For the identi-




Fig. 1: The Benelux countries with some major towns and an indication of the location of the Campine.
Abb. 1: Die Benelux-Länder mit einigen Größeren Städten und der Lage der „Kempen“.




Fig. 2: Walter & Lieth climate diagram Gilze-Rijen (SLUIJTER 2011). Given in the title are the name and location of the station, height above sea level (m), mean annual temperature (°C), and annual precipitation (mm) for the 30 yrs normal period 1981–2010.

Abb. 2: Klimadiagramm nach Walter & Lieth von Gilze-Rijen (SLUIJTER 2011). Angegeben sind Name und Lage der Station, Höhenlage, mittlere Jahrestemperatur (°C), jährlicher Niederschlag (mm) für die 30-Jahresperiode 1981–2010.

fication of differential species of the *Rubetum taxandriæ* and the *Rubetum silvatici* frequency classes (DIERSCHKE 1994) were used: species are considered to be differential if they differ at least two classes between the columns.

4. *Rubetum taxandriæ* Haveman, de Ronde & Weeda ass. nova

Holotypus: relevé 31 (September 8th 2009, western border Ulvenhoutse Bos, authors R. Haveman 09-515 and J.H.J. Schaminée 09-120), table 1 of this publication.

4.1 Species composition, structure, and variation

Table 1 shows the species composition of the newly described *Rubetum taxandriæ*. *Rubus taxandriæ* and *R. campaniensis* are the character species, and most probably the same holds for the rare *R. baronicus*. *Rubus insectifolius* can only be considered as regional character species, since it has a much wider distribution than the aforementioned three species. Character species of the *Rubetum silvatici* rarely occur together with those of the *Rubetum taxandriæ*, and *vice versa* (table 2); until now only *R. pyramidalis* has sometimes been recorded in the *Rubetum taxandriæ*.

Frequent companions among other *Rubus* species are *R. gratus*, *R. plicatus*, *R. integrifolius*, *R. nessensis*, *R. frederici*, *R. calotemnus*, and *R. macrophyllus*. As is clear from table 2, *R. frederici*, *R. integrifolius*, *R. macrophyllus*, *R. calotemnus*, *R. geniculatus*, *R. bertramii*, and *R. planus* have a higher frequency in the *Rubetum taxandriæ* than in the *Rubetum silvatici*, and can be regarded as differential species of the new association. They either have their optimum in *Pruno-Rubion radulae* communites (*R. macrophyllus*, *R. geniculatus*) (HAVEMAN et al. 1999b, WEBER 2003, 1999, 1995), or have a broad sociological amplitude and may be regarded as character species of the *Lonicero-Rubion silvatici* occurring in the southern part of the area of this alliance (HAVEMAN et al. in prep., 1999a, WEBER 1998a,

2003, 1995). On the other hand, *R. drenthicus*, *R. glandithyrsos*, *R. erinulus*, *R. discors*, *R. idaeus*, *R. ammobius*, and *R. schlechtendalii* are more frequent in the *Rubetum silvatici*. While *R. ammobius*, *R. discors*, and *R. drenthicus* can probably be regarded as character species of the *Lonicero-Rubion silvatici*, the sociological optimum of the other species is not yet clear. Probably they have their optimum in the *Rubetum silvatici*, characterising regional subassociations (see also the last section of this paper).

Frequently accompanying species are *Quercus robur* and *Betula pendula* in the tree and shrub layers, and *Sorbus aucuparia*, *Lonicera periclymenum*, *Rhamnus frangula*, and *Amelanchier lamarckii* in the shrub layer. Typically, the herb and moss layer are not well developed because of the dense bramble layer. *Agrostis capillaris*, *Molinia caerulea*, and *Holcus mollis* are the most frequent species in the herb layer, while *Kindbergia praelonga*, *Brachythecium rutabulum*, *Hypnum cupressiforme*, and *Pseudoscleropodium purum* can be found more or less frequently in the moss layer. *Betula pendula* is a weak differential species of the *Rubetum taxandriæ* in comparison to the *Rubetum silvatici*. The opposite holds for *Sorbus aucuparia* and *Holcus mollis*, which occur more often in the *Rubetum silvatici* (table 2).

Typically, the *Rubetum taxandriæ* is a dense thicket of about 1 m height in which two or more *Rubus* species grow together under an open tree layer and higher shrub layer. Coenoses with only one of the character species without accompanying *Rubus* species are considered as association fragments. Such fragments can be found throughout the distribution area of the association, and also beyond the border of the main distribution area in the Campine, for instance in the Peel region (see under "Distribution").

Two subassociations can be distinguished:

***Rubetum taxandriæ typicum* subass. nova**

Holotypus: as association.

The *Rubetum taxandriæ typicum* is negatively characterised by the absence of the species of the *deschampsietosum* subassociation. Besides this, it has several rather weak differential species: *Lonicera periclymenum*, *Urtica dioica*, *Dactylis glomerata*, *Holcus lanatus*, *Calamagrostis epigejos*, and *Hedera helix*. Among the brambles, *Rubus idaeus*, *R. macrophyllus*, and *R. poliothyrsus* show a clear preference for this subassociation (table 1).

***Rubetum taxandriæ deschampsietosum flexuosae* subass. nova**

Holotypus: relevé 12 (May 30th 2011, military air field Woensdrecht, author R. Haveman 11-019 and I. de Ronde 11-015), table 1 of this publication.

The *Rubetum taxandriæ deschampsietosum flexuosae* is characterised by the occurrence of *Deschampsia flexuosa*, *Molinia caerulea*, *Dryopteris dilatata*, *Vaccinium myrtillus*, *Brachythecium rutabulum*, and *Dicranum scoparium*. *Rubus campaniensis* is more frequent in this subassociation than in the *Rubetum taxandriæ typicum* (table 1).

4.2 Ecology

The *Rubetum taxandriæ* is a mesophilous and silvicolous (i.e. confined to the direct influence of woodlands) bramble scrub in woodland edges and forest clearings on loamy to loam-poor, rather nutrient poor, sandy soils. Far less often it is found in hedges, wooded banks or between agricultural tracts. The *Rubetum taxandriæ deschampsietosum* is confined to heathland afforestations from the late 19th and early 20th century, and is mainly found along woodland paths and in woodland clearings ("Innensäume"). The *Rubetum taxandriæ typicum* grows on sites with a slightly better nutrient availability, especially at the outer borders of forests, and in old forest remnants. Under a dense canopy the brambles die back, since the light demands of characteristic *Rubus* species are no longer met. Only in woodlands with an open canopy, especially in *Pinus sylvestris* and *Quercus robur* forests, both *Rubus campaniensis* and *R. taxandriæ* can form a dense, species poor knee-high bramble layer in the undergrowth (cf. BIJLSMA 2004); the synsystematic position of such *Rubus*-rich

Table 2: Shortened frequency table of the *Rubetum taxandriae* (column 1) and the *Rubetum sylvatici* (column 2). Frequencies are given in percentages. Except for the *Rubus* species, species with a frequency < 20% in both columns are omitted. Character and differential species (with a frequency class difference of at least two classes) are boxed.

Tabelle 2: Gekürzte Stetigkeitstabelle des *Rubetum taxandriae* (Spalte 1) und des *Rubetum sylvatici* (Spalte 2). Stetigkeitswerte in Prozent. Außer *Rubus*-Arten sind alle Arten mit < 20 % weggelassen. Charakter- und Differentialarten (mit Prozentunterschieden von wenigstens zwei Klassen) sind eingerahmt.

Column Number of relevés	1	2
	39	67
ca Rubetum taxandriae		
<i>Rubus taxandriae</i>	73	.
<i>Rubus campaniensis</i>	45	.
<i>Rubus baronicus</i>	18	.
<i>Rubus insectifolius</i>	15	.
ca Rubetum sylvatici		
<i>Rubus flexuosa</i>	.	51
<i>Rubus pyramidalis</i>	5	39
<i>Rubus sylvaticus</i>	.	33
<i>Rubus sprengelii</i>	.	31
optimum in Rubetum taxandriae (partly da)		
<i>Rubus integrabasis</i>	33	3
<i>Rubus frederici</i>	33	.
<i>Rubus calotemnus</i>	18	1
<i>Rubus macrophyllus</i>	18	4
<i>Rubus geniculatus</i>	8	.
<i>Rubus planus</i>	8	.
<i>Rubus bertramii</i>	8	.
optimum in Rubetum sylvatici (partly da)		
<i>Rubus drenthicus</i>	.	21
<i>Rubus idaeus</i>	10	36
<i>Rubus glandithyrsos</i>	.	15
<i>Rubus erinulus</i>	.	10
<i>Rubus discors</i>	.	10
<i>Rubus ammobius</i>	.	4
<i>Rubus schlechtendalii</i>	.	4
da associations		
<i>Betula pendula</i>	60	19
<i>Sorbus aucuparia</i>	50	81
<i>Holcus mollis</i>	35	61
bramble layer		
<i>Rubus gratus</i>	68	79
<i>Rubus plicatus</i>	30	24
<i>Rubus nessensis</i>	23	16
<i>Rubus calvus</i>	3	6
<i>Rubus scissus</i>	3	4
<i>Rubus poliothyrsus</i>	5	1
<i>Rubus vigorosus</i>	.	4
<i>Rubus ferocior</i>	3	1
<i>Rubus laevicaulis</i>	.	3
<i>Rubus nemoralis</i>	.	3
<i>Rubus guestphalicus</i>	3	.
<i>Rubus glareosus</i>	3	.
<i>Rubus x pseudoidaeus</i>	3	.
<i>Rubus subsectio Subidaeus</i>	3	.
<i>Rubus contritidens</i>	.	1

Column	1	2
Number of relevés	39	67
<i>Rubus arrhenii</i>	.	1
<i>Rubus adspersus</i>	.	1
<i>Rubus trichanthus</i>	.	1
<i>Rubus camptostachys</i>	.	1
<i>Rubus calviformis</i>	.	1
<i>Rubus speculans</i>	.	1
other species		
tree and scrub layer		
<i>Quercus robur</i>	83	75
<i>Lonicera periclymenum</i>	28	52
<i>Rhamnus frangula</i>	33	30
<i>Betula pubescens</i>	25	19
<i>Quercus rubra</i>	25	4
<i>Pinus sylvestris</i>	20	3
herb layer		
grasses		
<i>Agrostis capillaris</i>	65	69
<i>Molinia caerulea</i>	30	18
<i>Deschampsia flexuosa</i>	25	21
<i>Elytrigia repens</i>	5	22
herbs		
<i>Urtica dioica</i>	30	49
<i>Galeopsis tetrahit</i>	3	24
<i>Ceratocapnos claviculata</i>	3	19
<i>Cirsium arvense</i>	10	1
<i>Teucrium scorodonia</i>	10	.
ferns		
<i>Dryopteris dilatata</i>	25	42
<i>Pteridium aquilinum</i>	13	4
<i>Dryopteris carthusiana</i>	18	1
moss layer		
<i>Hypnum cupressiforme</i>	13	27
<i>Eurhynchium praelongum</i>	20	6
<i>Pseudoscleropodium purum</i>	20	3

forests still has to be settled. On very nutrient-poor soils the *Rubetum taxandriæ* is replaced by the *Rubetum grati*.

4.3 Distribution

The distribution area of the *Rubetum taxandriæ* includes the Dutch part of the Campine and the Baronie (Western Noord-Brabant, eastwards to the Eindhoven area) and at least parts of the Belgian Campine; the latter area has not yet been explored extensively. Fig. 5 gives the distribution of the relevés used in this paper. Association fragments can also be found in the “Peel” region, northeast of the Campine, where *Rubus campaniensis* is the only character species of the association. Distribution maps of the character species of the association (KURTTO et al. 2010) suggest that it can be found south of the Campine, but relevés are lacking.

According to our observations, the *Rubetum taxandriæ* is much more common in the western part of the Campine, between the towns of Antwerpen and Breda, than in the eastern part, around Turnhout and Eindhoven. Probably this has its background in the landscape history of the region. The western part was developed much earlier than the eastern part. This is obvious from the military topographical map of around 1850 (ANONYMUS 1995): the development of the western part of the Campine has led to a small scale landscape pattern with fields, meadows, woodlands and hedgerows, whereas the region of Eindhoven



Fig. 3: The *Rubetum taxandriæ* is usually found along forest edges. Here with *Rubus taxandriæ* and *R. gratus*. (Photo R. Haveman)

Abb. 3: Das *Rubetum taxandriæ* wird gewöhnlich an Waldrändern gefunden, hier mit *Rubus taxandriæ* und *R. gratus*.



Fig. 4: Primocane with leafs of *Rubus campaniensis*, one of the character species of the *Rubetum taxandriæ*. (Photo R. Haveman)

Abb. 4: Schössling mit Blättern von *Rubus campaniensis*, einer der Charakterarten des *Rubetum taxandriæ*.




Fig. 5: Location of vegetation relevés of the *Rubetum taxandriæ* used in this study.

Abb. 5: Lage der Vegetationsaufnahmen des *Rubetum taxandriæ* für diese Arbeit.

is characterised mainly by large scale heathlands which are devoid of brambles to a large extent, with agricultural activity only along the brooks and rivulets. Heathland reclamation took place between 1875 and 1950 mainly, after which bramble species spread into the reclaimed area. Although both *Rubus taxandriæ* and *R. campaniensis* have a wide distribution in the eastern part of the Campine, they are much more abundant in the western part, which might lead to the conclusion that the colonisation of the eastern part of the Campine by these species has not yet come to an end: the distribution area seems to be unsaturated, as might be expected for most species in this agamic complex (WEBER 1987).

5. Some considerations on the classification of the *Lonicero-Rubion silvatici*

The syntaxonomic place of the bramble scrubs united in the *Lonicero-Rubion silvatici* is subject to discussion. After its publication it was placed in the *Epilobietea* Tüxen & Preisig in Tüxen 1950, but as WEBER (1977) argued there are strong objections concerning the vegetation structure: the *Epilobietea* are dominated by herbs and grasses, and the *Lonicero-Rubion* by scrub species. On the basis of the frequent occurrence of *Rhamnus frangula* and the occurrence of *Salix cinerea* and *S. aurita* on somewhat moister soils, WEBER (1998a, 2003) places this alliance in the *Franguletea*. As mentioned in the introduction, HAVEMAN et al. (1999a) placed the alliance in a new class, the *Lonicero-Rubetea plicati*. In the more (sub-)atlantic parts of Northwestern Europe, both mentioned *Salix* species play hardly any role in the bramble scrubs, and their occurrence in the German examples of the *Lonicero-Rubion* communities could be a first indication that the *Rubus* species tend to grow in more tempered, humid conditions in less atlantic climates, retracting in woodlands completely further to the east. This stresses the fact that the bramble scrubs of the *Lonicero-Rubion* have their optimum in the (sub-)atlantic parts of Northwestern Europe, and that alliance is less typical developed in Northern Germany, thereby intermingling with the *Franguletea*. Therefore, we prefer to place the bramble scrubs of the *Lonicero-Rubion silvatici* in a separate class, the *Lonicero-Rubetea plicati*.

The lowland sand areas in Northwest-Europe form a centre of diversity of *Rubus* subgen. *Rubus* (cf. KURTTO et al. 2010, pp. 42, MATZKE-HAJEK 1997). In an analysis of the centres of diversity of *Rubus* in the Netherlands (VEEKEN & HAVEMAN 2008), three regions could be distinguished in the major Dutch sand landscapes, which fall apart in seven sub -

regions. The distribution area of the *Rubetum taxandriae* in the Netherlands corresponds remarkably well with the subregion of western Noord-Brabant. This raises the question about a probable further subdivision of the *Rubetum silvatici* in the Netherlands: is it possible to distinguish further communities within the *Rubetum silvatici*, characteristic for the various regions and/or subregions? There are indications that this is indeed the case. In the northern region, lying on the Drenthian bolder-clay plateau, a bramble community occurs for which the combination of *Rubus glandithyrsos*, *R. erinulus* and *R. flexuosus* is very characteristic. In the central subregion, the first two species are practically absent in the *Rubetum silvatici*, and their place is taken by *R. lasiandrus* instead, while *R. pyramidalis* is much more prominently present. Unlike the *Rubetum taxandriae*, these communities show a rather high frequency of character species of the *Rubetum silvatici*, so that they can therefore probably better be distinguished as regional subassociations of this association. Further investigations must make clear whether more *Rubus* scrub types can be distinguished in the Northwest-European sand landscapes, and what the underlying mechanisms and causes of the regional variation are.

Acknowledgements

Our thanks goes to Henk Reitsema for checking and improving the English, and we want to thank Anja van der Berg and Joop Schaminée for accompanying us during numerous field trips. This paper benefited a lot from the stimulating discussions! We are obliged to prof. dr. H. Dierschke and two anonymous reviewers, who made it possible to improve this paper on the basis of their comments.

References

- ANONYMUS (1995): Grote historische atlas van Nederland 1:50.000. 4. Zuid-Nederland 1838–1857. – Wolters-Noordhoff bv., Groningen: 127 pp.
- BARKMAN, J. J., DOING, H. & SEGAL, S. (1964): Kritische Bemerkungen und Vorschläge zur quantitativen Vegetationsanalyse. – Acta Bot.Neerl. 13: 394–419.
- BIJLSMA, R. J. (2004): Verbraming: oorzaken en ecologische plaats. – Levende Nat. 105 (4): 138–144.
- DENGLER, J., EISENBERG, M. & SCHRÖDER, J. (2006): Die grundwasserfernen Saumgesellschaften Nordostniedersachsens im europäischen Kontext – Teil I: Säume magerer Standorte (*Trifolio-Geranietea sanguinei*). – Tuexenia 26: 51–93.
- DIERSCHKE, H. (1994). Pflanzensoziologie: Grundlagen und Methoden. – Ulmer, Stuttgart: 683 S.
- HAVEMAN, R. (1997). Het *Rubetum grati* (*Lonicero-Rubetea plicati* classis nova) in Nederland. – Stratiotes 14: 41–51.
- , DE RONDE, I., BIJLSMA, R. J. & SCHAMINÉE, J. H. J. (in prep.): Regional variation in species composition of *Rubus* scrubs along three landscape transects in the Netherlands. – Phytocoen.
- , SCHAMINÉE, J. H. J. & STORTELDER, A. H. F. (1999a): *Lonicero-Rubetea plicati*. – In: A. H. F. STORTELDER, J. H. J. SCHAMINÉE & P. W. F. M. HOMMEL (Eds.): De vegetatie van Nederland. Deel 5. Plantengemeenschappen van ruigten, struwelen en bossen: 89–104. Opulus Press, Uppsala, Leiden.
- , SCHAMINÉE, J. H. J. & WEEDA, E. J. (1999b): *Rhamno-Prunetea*. – In: A. H. F. STORTELDER, J. H. J. SCHAMINÉE & P. W. F. M. HOMMEL (Eds.): De vegetatie van Nederland. Deel 5. Plantengemeenschappen van ruigten, struwelen en bossen: 121–164. Opulus Press, Uppsala, Leiden.
- HENNEKENS, S. M. & SCHAMINÉE, J. H. J. (2001): TURBOVEG, a comprehensive data base management system for vegetation data. – J.Veg.Sci. 12: 589–591.
- JANSSEN, F., DENGLER, J., GLÖCKLER, F., CHYTRÝ, M., EWALD, J., OLDELAND, J. & SCHAMINÉE, J. H. J. (2011): Die mitteleuropäischen Datenbanken im Global Index of Vegetation-Plot Databases (GIVD). – Tuexenia 31: 351–367.
- KURTTO, A., WEBER, H. E., LAMPINEN, R. & SENNIKOV, A. N. (2010): Atlas Flora Europaea. Distribution of vascular plants in Europe. Rosaceae (*Rubus*). – The Committee for Mapping the Flora of Europe & Societas Biologica Fennica Vanamo, Helsinki: 362 pp.
- MATZKE-HAJEK, G. (1997): Zur Evolution und Ausbreitung apomiktischer *Rubus*-Arten (Rosaceae) in Offenland-Ökosystemen. – Bull.Geobot.Inst. ETH 63: 33–44.
- POTT, R. (1995). Die Pflanzengesellschaften Deutschlands. 2., überarb. Auflage. – Ulmer, Stuttgart: 622 S.
- ROLEČEK, J., TICHÝ, L., ZELENÝ, D. & CHYTRÝ, M. (2009): Modified TWINSPLAN classification in which the hierarchy respects cluster heterogeneity. – 20 (4): 596–602.

- SCHAMINÉE, J. H. J., JANSSEN, J. A. M., HAVEMAN, R., HENNEKENS, S. M., HEUVELINK, G. B. M., HUISKES, H. P. J. & WEEDA, E. J. (2006): Schatten voor de natuur. Achtergronden, inventaris en toepassingen van de Landelijke Vegetatie Databank. – KNNV Uitgeverij, Utrecht: 112 pp.
- SIEBEL, H. & DURING, H. (2006): Beknopte Mosflora van Nederland en België. – KNNV Uitgeverij, Utrecht: 559 pp.
- SLUIJTER, R. (2011): De Bosatlas van het klimaat. – KNMI & Noordhoff Atlasproducties, De Bilt: 112 pp.
- TICHÝ, L. (2002). JUICE, software for vegetation classification. – *J. Veg. Sci.* 13: 451–453.
- VAN DER BEEK, A., BIJLSMA, R. J., HAVEMAN, R., MEIJER, K., DE RONDE, I. & TROELSTRA, A. (in press): Naamlijst en verspreidingsgegevens van de Nederlandse bramen (*Rubus* L.). – Gorteria.
- VAN DER MEJDEN, R. (2005). Heukels' Flora van Nederland. – Wolters-Noordhoff, Groningen/Houten: 685 pp.
- VEEKEN, J. & HAVEMAN, R. (2008): Plantengeografische en ecologische aspecten van apomicten. – In: J. H. J. SCHAMINÉE & E. J. WEEDA (Eds.): Grenzen in Beweging. Beschouwingen over vegetatiegeografie: 73–88. KNNV Uitgeverij, Utrecht.
- WEBER, H. E. (1977): Beiträge zur Systematik der Brombeergebüsche auf potentiell natürlichen *Quercion robori-petraeae*-Standorten in Nordwestdeutschland. – *Mitt. Flor-soz. Arbeitsgem. N.F.* 19/20: 343–351.
- (1987): Typen ornithochorer Arealentwicklung, dargestellt an Beispielen der Gattung *Rubus* L. (Rosaceae) in Europa. – *Bot. Jahrb. Syst.* 108 (2/3): 525–535.
 - (1990). Übersicht über die Brombeergebüsche der *Pteridio-Rubetalia (Franguletea)* und *Prunetalia (Rhamno-Prunetea)* in Westdeutschland mit grundsätzlichen Bemerkungen zur Bedeutung der Vegetationsstruktur. – *Ber. Reinhold-Tüxen-Ges.* 2: 91–119.
 - (1995): *Rubus* L. – In: H. J. CONERT, E. J. JÄGER, J. W. KADEREIT, W. SCHULTZE-MOTEL, G. WAGENITZ & H. E. WEBER (Hrsg.): Gustav Hegi Illustrierte Flora von Mitteleuropa. Band IV. Teil 2A. Spermatophyta: Angiospermae: Dicotyledones 2(2): 284–595. Blackwell, Berlin.
 - (1997): Hecken und Gebüsche in den Kulturlandschaft Europas – Pflanzensoziologische Dokumentation als Basis für Schutzmassnahmen. – *Ber. Reinhold-Tüxen-Ges.* 9: 75–106.
 - (1998a). *Franguletea* (H1). Faulbaum-Gebüsche. – *Synopsis Pflanzenges. Deutschlands* 4: 3–86.
 - (1998b): Outline of the vegetation of scrubs and hedges in the temperate and boreal zone of Europe. – *Itiner. Geobot.* 11: 85–120.
 - (1999): *Rhamno-Prunetea* (H2A). Schlehen- und Traubenhölzer-Gebüsche. – *Synopsis Pflanzenges. Deutschlands* 5: 3–108.
 - (2003): Gebüsche, Hecken, Krautsäume. – Ulmer, Stuttgart: 229 S.
- WEEDA, E. J., SCHAMINÉE, J. H. J. & VAN DUUREN, L. (2005): Atlas van Plantengemeenschappen in Nederland deel 4: Bossen, struwelen en ruigten. – KNNV Uitgeverij, Utrecht: 282 pp.

Rense Haveman, Iris de Ronde
Dienst Vastgoed Defensie Noord
P.O. Box 47, 6700 AA Wageningen, The Netherlands

Eddy Weeda
Alterra-Wageningen University and Research
P.O. Box 47, 6700 AA Wageningen, The Netherlands

Manuscript received: 30.09.2011; accepted: 02.01.2012