

# Effects of shoot tipping on development and yield of the tuber crop *Plectranthus edulis*

M. TAYE<sup>1</sup>, W. J. M. LOMMEN<sup>2\*</sup> AND P. C. STRUIK<sup>2</sup>

<sup>1</sup>Hawassa University College of Agriculture, P. O. Box 5, Awassa, Ethiopia

<sup>2</sup>Centre for Crop Systems Analysis, Plant Sciences Group, Wageningen University, Droevendaalsesteeg 1, 6708 PB Wageningen, The Netherlands

(Received 2 December 2010; revised 31 May 2011; accepted 29 August 2011)

## SUMMARY

*Plectranthus edulis* (Vatke) Agnew is one of the tuber crops of the genus *Plectranthus* that is widely cultivated in Africa and Asia. *P. edulis* produces below-ground tubers on stolons originating from the stems, comparable to the potato (*Solanum tuberosum* L.). Farmers apply several laborious cultural practices to enhance shoot growth and yield, among which shoot tipping is very common. Tipping (pinching) is the removal of the shoot apex with one or two pairs of leaves from the main stems and branches. The rationale of this practice, especially when repeated more than once during one cropping season, is not fully understood. One similar experiment with two cultivars was carried out at two locations (Awassa and Wondogenet) in Ethiopia to assess and analyse the effects of shoot tipping and its frequency on crop development and tuber production. Tipping treatments included zero tipping, tipping once, tipping twice and tipping thrice, with the first tipping taking place 68 days after planting (DAP), a stage at which most of the stems reached a height of about 0.15 m, and the remainder following at intervals of 44–46 days. Tipping stimulated stem branching; it significantly increased the number of primary, secondary and tertiary stems in both experiments. Soil cover increased with an increase in the frequency of the tipping in Awassa, because of the tipping effects on the different canopy development variables. Tipping also enhanced the soil cover in Wondogenet, but the crop did not gain any extra benefit from a third tipping. Tipping enhanced early stolon formation, but did not consistently affect the number of stolons per hole later in the growing season. The number of tubers increased with an increase in the frequency of tipping in both cultivars in Wondogenet and in one cultivar in Awassa. Tuber dry matter yield increased with an increase in the frequency of tipping at both sites. Fresh tuber yield in the final harvest at 208 DAP was c. 1.9 kg/m<sup>2</sup>. Tipping on average increased fresh tuber yield by 17% in Wondogenet, whereas the difference was not detectable in Awassa. Because senescence was delayed slightly by tipping, yield effects of tipping might be larger when harvesting later. In general, there was a positive effect of tipping on canopy development and tuber yield.

## INTRODUCTION

Tuber-forming species of the genus *Plectranthus* (Lamiaceae) are widely cultivated in Africa and Asia (e.g. Jansen 1996), but agronomic information on their cultivation is scarce. *Plectranthus edulis* (Vatke) Agnew (syn. *Coleus edulis* Vatke), locally called Wolaita dono, Dinicha Oromo, Gamo dinich, Agaw dinich or Ethiopian potato, can grow up to a height of 1.5 m and produces edible tubers on below-ground stolons. The below-ground habit of the plant resembles that of the potato (*Solanum tuberosum* L.). *P. edulis* is

grown at mid to high altitudes (Greenway 1944; Asfaw & Woldu 1997) in the southern, western and northern parts of Ethiopia (Westphal 1975; Taye *et al.* 2007).

Tipping (pinching), i.e. the removal of the shoot apices with one or two pairs of leaves is a common element of the crop husbandry applied by farmers. Some farmers claim that tipping increases tuber yield (Taye *et al.* 2007). In a survey carried out in the regions of Chencha and Wolaita in Ethiopia, Taye *et al.* (2007) found the number of tippings vary between one and three, depending on the availability of time and labour. Most growers carried out the first tipping when the plants reached a height of 0.15 m, which is 1–1.5 months after planting. If farmers tipped more

\* To whom all correspondence should be addressed. Email: Willemien.Lommen@wur.nl

frequently, the second and third tippings were carried out about 1 and 2 months after the first tipping, respectively.

Shoot tipping or decapitation is well known to stimulate branching in many plant species (e.g. Dun *et al.* 2006) by breaking apical dominance. Tipping is carried out in various ornamentals, including *Plectranthus*, *Angelonia* and *Calibrachoa* species, and ornamental sweet potato (Smith 2003) to make the plant bushy, and in potato to stimulate branching for production of stem cuttings (e.g. Bryan *et al.* 1981; Lommen & Struik 2007). For the same reason, tipping may also increase the number of stolons, and thus the number of potential tuber sites. However, the practice can only be justified if the size of the canopy or the number of potential tuber sites is indeed a factor limiting production.

Although tipping has been practised in *P. edulis* for years, it is not known how it affects crop development and tuber yield. Therefore, field experiments were conducted to assess and analyse the effects of tipping and tipping frequency on canopy branching and soil cover, yield components and tuber production of *P. edulis*.

## MATERIALS AND METHODS

### Experimental sites

A similar experiment was carried out at two sites in southern Ethiopia, namely Awassa and Wondogenet, between 19 April and 13 November 2004 (Awassa) and between 20 April and 14 November 2004 (Wondogenet). The Awassa field (7°03'N and 38°30' E, 1650 m asl) was a sandy loam soil with an organic matter (OM) concentration of 24 g/kg, pH-H<sub>2</sub>O 6.9, 1 g N/kg and 14.4 mg P/kg; the Wondogenet field (7°06'N 38°37'E, 1850 m asl) was a sandy loam soil with 29 g OM/kg, pH-H<sub>2</sub>O 6.1, 1.1 g N/kg and 15 mg P/kg. Wondogenet is relatively cooler and more humid than Awassa (average temperatures of 18.4 and 20.2 °C, respectively). Daily average temperatures during the experiments ranged from 17.4 to 23.7 °C and from 14.9 to 21.6 °C in Awassa and Wondogenet, respectively. Rainfall was not limiting.

### Experimental design and treatments

The experiments had a complete split-split-plot design with three factors, namely cultivar, number of tippings and harvesting time, replicated in six blocks. Four

harvesting dates were randomized within four tipping practices and the latter within two cultivars.

The two cultivars were Lofuwa and Chankua. The leaf and stem colours of Lofuwa are green and its tuber skin is dull white, whereas the leaf, branch and tuber skin of Chankua are red or reddish. Farmers consider Lofuwa as relatively early maturing and Chankua as late maturing. The four tipping treatments included zero tipping, tipping once (carried out 68 days after planting (DAP), when most shoots attained a height of about 0.15 m), tipping twice (in which the first tipping was followed by a second tipping at 114 DAP, i.e. 46 days after the first tipping) and tipping thrice (in which the third tipping was carried out at 158 DAP, i.e. 44 days after the second tipping). While tipping, the top parts (i.e. the parts above the first clear internode, containing the apex with one or two pairs of visible leaves) were removed from those main stems and branches that were developed to such an extent that at least one pair of leaves remained on the stem or branch. In the second and third tipping, only outgrowths were tipped from stems that had been tipped in the previous tipping. To study the crop development at the moment(s) of tipping and the development with time thereafter, the crops from different tipping treatments were harvested on four dates. The 1st harvest was on 26 and 27 June (68 DAP), the 2nd on 11 and 12 August (114 DAP), the 3rd on 25 and 26 September (159 DAP) and the final harvest on 13 and 14 November (208 DAP) in Awassa and Wondogenet, respectively. The first three harvest dates coincided with the days of the 1st and 2nd tipping of the respective treatments (immediately after tipping), and one day after the 3rd tipping, which was the moment maximum haulm development was measured. The final harvest was 50 days after the 3rd tipping, when the haulm was senescing.

Gross plot size was 3.60 × 3.75 m, comprising plants from 20 holes in four rows of five holes. Hole spacing was 0.90 m between rows and 0.75 m within a row, consistent with common practice. Only plants from the inner six positions (two rows × three holes, i.e. 1.80 × 2.25 m net plot size) were harvested, those from the remaining 14 positions were kept as guard plants.

### Crop management

Before planting, the experimental field was ploughed, disc harrowed and furrowed with a width of 0.50 m between ridges and the furrows 0.20 m below the

ridges. The crops were planted on 19 April 2004 (Awassa) and 20 April 2004 (Wondogenet). Planting holes 0.05 m deep were made in furrows. Three desprouted, broken tuber pieces (prepared by breaking a mother tuber of 0.10–0.15 m length into three) were planted per hole in a triangle with sides of about 0.05 m, and covered with soil. During growth the soil was cultivated two to three times in order to control weeds, and plants were earthed up (i.e. soil was piled up around the stems) three times, in order to support the stems and also cover the stolons. These are farmer practices as indicated in Taye *et al.* (2007). Manure or chemical fertilizer was not applied.

## Observations and calculations

### *Ground cover*

Ground cover was estimated at the harvest dates (1st and 2nd harvest) or one day before the harvest dates (3rd and 4th harvest) on the plots to be harvested in the final harvest, using a frame of 0.90 m × 0.75 m that was divided into 100 cells of 90 mm × 75 mm each. The number of cells filled to at least 0.5 with green leaves was counted, the observer looking down from above to avoid parallax. Two measurements were taken per net plot, one from each inner row. This grid method provides a non-destructive measurement, which correlates well with the proportion of intercepted photosynthetically active radiation as measured with a tube solarimeter (Haverkort *et al.* 1991).

### *Crop analysis and related measurements*

Data from above-ground and below-ground parts were recorded during each harvest. Whole plants from all six holes of the net plots were dug out carefully and cleaned. The number of main stems from all six holes and the number of branches (primary, secondary and tertiary) from a subsample of three holes were recorded. Main stem refers to each stem that originated from the bud of a mother tuber piece, primary branches originated from main stems, secondary branches from primary branches and tertiary branches from secondary branches. Stolon number was recorded from main stems and primary branches from three holes. Secondary and tertiary branches produced no stolons. Tuber yield was determined from all six holes.

## Statistical analysis

GenStat release 13.3 (VSN International Ltd, Hemel Hempstead, UK) was used to conduct separate ANOVA on the data from each harvest date. For branch numbers and yield components, tests were conducted to determine whether the response to the frequency of tipping was linear or quadratic. Correlations across tipping treatments within sites and cultivars between number of secondary branches and the total number of green leaves and between the number of stolons per hole or the tuber:stolon ratio and the number of tubers per hole were determined by Excel 2003.

## RESULTS

### Canopy development

#### *Number of main stems*

At 159 and 208 DAP, the number of main stems per hole was increased significantly by the tipping treatments in Awassa (Table 1), with no difference in the number of main stems between the frequencies of tipping. In Wondogenet, the number of main stems was not affected by tipping at 159 DAP in either cultivar, whereas at 208 DAP the number of main stems per hole was not affected by tipping in cvar Lofuwa, but was lower for higher frequencies of tipping in cvar Chankua (Table 1). There were no significant differences between cultivars in the number of main stems in Awassa, whereas in Wondogenet Chankua had fewer stems than Lofuwa at 159 DAP, but still had more stems than Lofuwa for the zero tipping and lower frequencies of tipping at 208 DAP (Table 1).

#### *Number of branches*


For all types of branches, the number of branches per hole was highest at 159 DAP and had decreased at the final harvest due to crop senescence. Primary branches were already present in the first harvest at 68 DAP (data not shown). At 159 and 208 DAP, tipping had significantly increased the number of primary branches per hole as compared with zero tipping at both sites (Table 1). The first secondary branches were observed in the harvest at 114 DAP (data not shown). Tipping also increased the number of secondary branches at 159 DAP, but had no consistent effect on the number of secondary branches remaining at 208 DAP (Table 1). Tertiary branches were not observed at

Table 1. Number of main stems and different order branches per hole of the *P. edulis* cultivars Lofuwa and Chankua as affected by zero (T0), one (T1), two (T2) and three tipplings (T3) for two harvests in Awassa and Wondogenet

Cultivar (CV) and tipping (T) treatment	Awassa								Wondogenet							
	Main stems		Primary branches		Secondary branches		Tertiary branches		Main stems		Primary branches		Secondary branches		Tertiary branches	
	3rd harvest (159 DAP)	Final harvest (208 DAP)	3rd harvest (159 DAP)	Final harvest (208 DAP)	3rd harvest (159 DAP)	Final harvest (208 DAP)	3rd harvest (159 DAP)	Final harvest (208 DAP)	3rd harvest (159 DAP)	Final harvest (208 DAP)	3rd harvest (159 DAP)	Final harvest (208 DAP)	3rd harvest (159 DAP)	Final harvest (208 DAP)	3rd harvest (159 DAP)	Final harvest (208 DAP)
Cultivar Lofuwa																
T0	1.5	1.3	22	4.4	50	10	35	23	1.6	1.2	22	3.4	56	8	39	21
T1 (68 DAP)	2.0	2.1	24	8.3	95	5	36	17	1.9	1.3	27	6.4	89	9	36	16
T2 (68+114 DAP)	2.2	1.9	25	7.0	123	10	85	26	1.8	1.0	28	6.6	126	8	83	25
T3 (68+114+ 158 DAP)	2.0	2.1	32	7.4	117	23	104	20	1.7	1.2	29	5.9	125	22	82	20
Average	1.92	1.83	25.6	6.81	96.1	12.1	64.9	21.4	1.74	1.17	26.5	5.58	99.1	11.8	60.0	20.4
Cultivar Chankua																
T0	1.4	1.3	26	4.6	47	13	35	19	1.6	1.9	22	3.9	44	8	37	19
T1 (68 DAP)	2.0	1.8	31	8.0	100	7	33	14	1.6	1.7	29	4.9	113	10	57	22
T2 (68+114 DAP)	2.2	1.9	32	7.1	149	16	76	25	1.6	1.4	28	5.6	128	9	119	26
T3 (68+114+ 158 DAP)	2.0	2.3	35	6.6	173	10	78	27	1.6	1.4	29	5.4	128	8	86	22
Average	1.92	1.84	31.0	6.56	117.1	11.4	55.5	21.2	1.62	1.58	26.8	4.94	103.2	8.7	74.8	22.2
Average over cultivars																
T0	1.5	1.3	24	4.5	48	12	35	21	1.6	1.5	22	3.7	50	8	38	20
T1 (68 DAP)	2.0	1.9	27	8.2	97	6	34	15	1.8	1.5	28	5.7	101	9	47	19
T2 (68+114 DAP)	2.2	1.9	29	7.0	136	13	81	26	1.7	1.2	28	6.1	127	9	101	25
T3 (68+114+ 158 DAP)	2.0	2.2	33	7.0	145	17	91	23	1.6	1.3	29	5.7	127	15	84	21
Average	1.92	1.83	28.3	6.68	106.6	11.8	60.2	21.3	1.68	1.37	26.6	5.26	101.1	10.3	67.4	21.3
S.E.D.																
CV (5 D.F.)	ns	ns	1.71	ns	4.15	ns	ns	ns	0.047	0.101	ns	ns	ns	0.62	3.00	ns
Tlinear (30 D.F.)	0.135	0.146	2.00	0.631	6.84	1.15	ns	2.23	ns	0.877	1.15	0.479	6.15	0.92	4.75	ns
Tquadratic (30 D.F.)	0.135	ns	ns	0.631	6.84	1.15	ns	ns	ns	ns	1.15	0.479	6.15	0.92	4.75	ns
CV × Tlinear (30 D.F.)*	ns	ns	ns	ns	9.68	1.62	ns	3.15	ns	0.124	ns	ns	ns	1.30	ns	ns
CV × Tquadratic (30 D.F.)*	ns	ns	ns	ns	ns	1.62	ns	ns	ns	ns	ns	ns	ns	1.30	6.72	ns

DAP, days after planting.

\* For comparisons of tipping treatments within a cultivar only.



**Fig. 1.** Effect of zero, one, two and three tipplings on development of green leaf number per hole (a, b) and fraction ground cover (c, d) of *P. edulis* with time in Awassa (a, c) and Wondogenet (b, d). Average values over two cultivars. Bars: s.e.d. (30 D.F.).

68 and 114 DAP. The number of tertiary branches at 159 DAP was increased by tipping twice or thrice in Awassa and in cvar Lofuwa in Wondogenet, and by all tipping frequencies in cvar Chankua in Wondogenet (Table 1), but there were no consistent effects at 208 DAP.

#### Number of green leaves

The number of green leaves per hole was highest at 159 DAP (Fig. 1a, b). At this harvest date, the number of leaves per hole both in Awassa and Wondogenet showed significant interaction between cultivar and tipping. The more frequent the tipping was, the more green leaves per hole, especially for Chankua. There were significant correlations between the number of secondary branches at 159 DAP and the total number of leaves within individual cultivar  $\times$  location combinations ( $r = 0.822\text{--}0.889$ ,  $n = 24$ ).

#### Ground cover

Significant differences due to tipping only appeared later in the growing season (Fig. 1). This was true for both sites. Tipping increased both the maximum ground cover recorded and the ground cover in the period during which leaves senesced.

#### Below-ground development

##### Number of stolons

Stolons generally did not appear until after the 1st harvest (Table 2). At 114 DAP, a small number of stolons had appeared and the number was significantly higher in tipped plants than in control plants (Table 2). Stolon numbers were maximum at 159 DAP. Tipping no longer had any effect on the number of stolons at 159 DAP in Wondogenet and in cvar Chankua in Awassa. For Lofuwa at Awassa, higher

Table 2. Number of stolons and tubers per hole of the *P. edulis* cultivars Lofuwa and Chankua as affected by zero (T0), one (T1), two (T2) and three (T3) tipplings in four harvests in Awassa and Wondogenet

Cultivar (CV) and tipping (T) treatment	Number of stolons per hole								Number of tubers per hole							
	Awassa				Wondogenet				Awassa				Wondogenet			
	1st harvest (68 DAP)	2nd harvest (114 DAP)	3rd harvest (159 DAP)	Final harvest (208 DAP)	1st harvest (68 DAP)	2nd harvest (114 DAP)	3rd harvest (159 DAP)	Final harvest (208 DAP)	1st harvest (68 DAP)	2nd harvest (114 DAP)	3rd harvest (159 DAP)	Final harvest (208 DAP)	1st harvest (68 DAP)	2nd harvest (114 DAP)	3rd harvest (159 DAP)	Final harvest (208 DAP)
Cultivar Lofuwa																
T0	0.056	5.1	26.6	3.2	0.111	9.7	35.9	13.4	0	0	6.5	112	0	0	5.5	122
T1 (68 DAP)	0	6.1	32.7	2.7	0	10.2	39.9	7.1	0	0	7.7	135	0	0	6.6	109
T2 (68+114 DAP)	0	6.2	20.8	2.7	0	10.4	34.4	9.2	0	0	8.7	137	0	0	5.6	132
T3 (68+114+ 158 DAP)	0	7.3	20.7	3.1	0	14.6	30.0	9.1	0	0	7.2	147	0	0	5.1	144
Average	0.0139	6.18	25.18	2.93	0.0278	11.06	35.05	9.69	0	0	7.52	132.6	0	0	5.69	126.8
Cultivar Chankua																
T0	0	4.7	18.3	3.3	0	8.8	28.6	11.7	0	0	5.9	136	0	0	5.1	117
T1 (68 DAP)	0	5.8	17.4	3.1	0	15.4	28.6	14.0	0	0	7.4	122	0	0	5.6	145
T2 (68+114 DAP)	0	7.8	21.0	3.3	0	13.0	35.0	5.6	0	0	6.9	132	0	0	5.1	141
T3 (68+114+ 158 DAP)	0	6.6	19.7	3.2	0	12.9	29.0	12.7	0	0	7.9	135	0	0	6.0	145
Average	0	6.19	19.11	3.22	0	12.51	30.30	10.99	0	0	7.03	131.1	0	0	5.46	136.8
Average over cultivars																
T0	0.028	4.9	22.4	3.2	0.056	9.2	32.2	12.6	0	0	6.2	124	0	0	5.3	120
T1 (68 DAP)	0	5.9	25.1	2.9	0	12.8	34.2	10.5	0	0	7.6	128	0	0	6.1	127
T2 (68+114 DAP)	0	7.0	20.9	3.0	0	11.7	34.7	7.4	0	0	7.8	134	0	0	5.4	136
T3 (68+114+ 158 DAP)	0	6.9	20.2	3.2	0	13.7	29.5	10.9	0	0	7.5	141	0	0	5.5	144
Average	0.0069	6.19	22.14	3.08	0.0139	11.78	32.67	10.34	0	0	7.28	131.8	0	0	5.57	131.8
S.E.D.																
CV (5 D.F.)	ns	ns	1.04	ns	ns	ns	ns	ns			ns	ns			ns	2.7
Tlinear (30 D.F.)	ns	0.57	ns	ns	ns	1.22	ns	ns			ns	7.77			ns	11.4
Tquadratic (30 D.F.)	ns	ns	ns	ns	ns	ns	ns	ns			ns	ns			ns	ns
CV × Tlinear (30 D.F.)*	ns	ns	2.73†	ns	ns	ns	ns	ns			ns	10.99			ns	ns
CV × Tquadratic (30 D.F.)*	ns	ns	ns	ns	ns	1.72	ns	ns			ns	ns			ns	ns

DAP, days after planting.

\* For comparisons of tipping treatments within a cultivar only.

† 28 D.F.

tipping frequencies resulted in fewer stolons at 159 DAP (Table 2).

#### *Number of tubers*

Tubers did not appear until after 114 DAP. Only a few tubers had appeared at 159 DAP and tipping had no effect on that number. The number of tubers was highest at 208 DAP, when there was a linear increase in the number of tubers with an increase in the frequency of tipping in both cultivars in Wondogenet and in cvar Lofuwa in Awassa (Table 2). At the final harvest in Wondogenet, the number of tubers was significantly higher for Chankua than for Lofuwa (Table 2).

#### *Yield components and dry matter production*

In Awassa, tipping increased the number of main stems per hole, decreased the number of stolons per main stem in cvar Lofuwa, and had no effect on the number of stolons per hole in cvar Chankua and an inconsistent effect in cvar Lofuwa (Table 3). In Wondogenet, tipping had no effect on the number of main stems per hole, and did not affect the number of stolons per main stem or per hole (Table 4). Effects of tipping on the tuber:stolon ratio were inconsistent over cultivars in Awassa and absent in Wondogenet (Tables 3 and 4). Nevertheless, there was a linear increase in the number of tubers with an increase in the frequency of tipping for cvar Lofuwa in Awassa and for both cultivars in Wondogenet (Tables 3 and 4). Within individual combinations of cultivar and location, the number of tubers per hole was correlated with the tuber:stolon ratio for both cultivars in Awassa and cvar Lofuwa in Wondogenet ( $r=0.446-0.719$ ,  $n=22-24$ ), but not with the number of stolons per hole ( $r=0.087-0.260$ ,  $n=24$ ). No effect of tipping on the average weight per tuber was found (Tables 3 and 4). There was no detectable effect of tipping on fresh tuber yield in Awassa (Table 3), but tipping increased fresh tuber yield in Wondogenet. The quadratic nature of the effect showed that optimum tipping frequencies were one or two, with two tippings leading to only limited yield increase compared with one (Table 4). Also the tuber dry matter concentration was not affected significantly by tipping in Awassa, but increased significantly with increase in the tipping frequency in Wondogenet (Tables 3 and 4). The tuber dry weight per hole increased with increase in tipping frequency at both sites (Tables 3 and 4). Within individual combinations of cultivar and location, tuber dry

weight was correlated with tuber dry matter concentration ( $r=0.427-0.644$ ,  $n=23-24$ ), and with tuber fresh weight per hole ( $r=0.702-0.920$ ,  $n=23-24$ ).

In Awassa, the number of stolons produced per main stem was significantly higher for cvar Lofuwa than for cvar Chankua in the zero tipping and tipping once treatments, and so was the number of stolons per hole (Tables 3 and 4). The tuber:stolon ratio, however, was significantly higher for cvar Chankua than for cvar Lofuwa in the zero tipping and tipping once treatments. There were no differences between cultivars in any of the other production components in Awassa. At Wondogenet, cvar Chankua produced fewer main stems, but more tubers per hole than Lofuwa, had a smaller average tuber size, and had a higher tuber dry matter concentration and a higher tuber dry matter yield (Table 4).

## DISCUSSION

### Canopy development

Tipping enhanced the number of main stems in Awassa, but not in Wondogenet (Table 1). Tipping, however, clearly enhanced branching in both experiments (Table 1), most likely due to the breaking of apical dominance, which stimulates branching in many crops (e.g. Bryan *et al.* 1981; Dun *et al.* 2006; Lommen & Struik 2007). Due to increased branching there were also more leaves (Fig. 1a, b). This enhanced branching and leaf production increased the canopy cover and probably delayed crop senescence (Fig. 1). This is consistent with findings in okra (*Abelmoschus esculentus* (L.) Moench), where apical bud removal is applied in combination with leaf removal to prolong the growing period (Olasantan & Salau 2008).

### Below-ground development

Below-ground development was affected less by tipping than above-ground development. Tipping increased stolon numbers in the earlier phases of growth, whereas by the time of the 3rd harvest (when stolon numbers were maximum and the first tubers were initiated) effects of tipping on stolon numbers were no longer visible (Table 2). More main stems – as found in Awassa – tended to reduce the number of stolons per stem, thus counterbalancing the positive effect of tipping on the number of main stems.

The number of tubers obtained from these stolons increased with increase in the frequency of tipping in

Table 3. Yield characteristics of the *P. edulis* cultivars Lofuwa and Chankua as affected by zero (T0), one (T1), two (T2) and three (T3) tipplings in Awassa. Stem and stolon data from the harvest at 159 DAP, tuber data are from the final harvest at 208 DAP

Cultivar (CV) and tipping (T) treatment	Number of main stems per hole (#/hole)	×	Number of stolons per main stem (#/main stem)	=	Number of stolons per hole (#/hole)	×	Tuber: stolon ratio (#/#)	=	Number of tubers per hole (#/hole)	×	Average weight per tuber (g/tuber)	=	Tuber fresh weight per hole (g/hole)	×	Number of holes per m <sup>2</sup> (#/m <sup>2</sup> )	=	Tuber fresh yield per m <sup>2</sup> (g/m <sup>2</sup> )	×	Tuber dry matter concentration (g/g)	=	Tuber dry weight per m <sup>2</sup> (g/m <sup>2</sup> )
Cultivar Lofuwa																					
T0	1.5	×	18	=	27	×	4.3	=	112	×	12	=	1288	×	1.48	=	1906	×	0.16	=	302
T1 (68 DAP)	2.0	×	15	=	33	×	4.5	=	135	×	11	=	1406	×	1.48	=	2081	×	0.16	=	335
T2 (68+114 DAP)	2.2	×	10	=	21	×	6.7	=	137	×	10	=	1401	×	1.48	=	2074	×	0.16	=	321
T3 (68+114+158 DAP)	2.0	×	11	=	21	×	7.2	=	147	×	10	=	1439	×	1.48	=	2130	×	0.16	=	340
Average	1.92	×	13.6	=	25.2	×	5.66	=	132.6	×	10.7	=	1383.5	×	1.48	=	2049.6	×	0.159	=	324.3
Cultivar Chankua																					
T0	1.4	×	13	=	18	×	8.9	=	136	×	10	=	1342	×	1.48	=	1986	×	0.16	=	316
T1 (68 DAP)	2.0	×	9	=	17	×	7.7	=	122	×	11	=	1293	×	1.48	=	1914	×	0.16	=	298
T2 (68+114 DAP)	2.2	×	9	=	21	×	6.5	=	132	×	11	=	1468	×	1.48	=	2173	×	0.17	=	368
T3 (68+114+158 DAP)	2.0	×	10	=	20	×	7.1	=	135	×	11	=	1453	×	1.48	=	2150	×	0.16	=	352
Average	1.92	×	10.4	=	19.1	×	7.54	=	131.1	×	10.7	=	1389.0	×	1.48	=	2057.8	×	0.162	=	333.6
Average over cultivars																					
T0	1.5	×	16	=	22	×	6.6	=	124	×	11	=	1315	×	1.48	=	1946	×	0.16	=	309
T1 (68 DAP)	2.0	×	12	=	25	×	6.1	=	128	×	11	=	1350	×	1.48	=	1997	×	0.16	=	317
T2 (68+114 DAP)	2.2	×	10	=	21	×	6.6	=	134	×	11	=	1435	×	1.48	=	2123	×	0.16	=	344
T3 (68+114+158 DAP)	2.0	×	11	=	20	×	7.1	=	141	×	10	=	1445	×	1.48	=	2140	×	0.16	=	346
Average	1.92	×	12.0	=	22.2	×	6.60	=	131.8	×	10.7	=	1386.2	×	1.48	=	2053.7	×	0.160	=	328.9
S.E.D.																					
CV (5 D.F.)	ns		1.05		1.04		0.597		ns		ns		ns		ns		ns		ns		ns
Tlinear (30 D.F.)	0.135		1.44†		ns		ns†		7.8		ns		ns		ns		ns		ns		20.0
Tquadratic (30 D.F.)	0.135		1.44†		ns		ns†		ns		ns		ns		ns		ns		ns		ns
CV × Tlinear (30 D.F.)*	ns		2.03†		2.73†		1.391†		11.0		ns		ns		ns		ns		ns		ns
CV × Tquadratic (30 D.F.)*	ns		ns†		ns		ns†		ns		ns		ns		ns		ns		ns		ns

DAP, days after planting.

\* For comparisons of tipping treatments within a cultivar only.

† 28 D.F.

Given product values are not the same as the ones found after re-calculation based on the averages in this and the following table, because  $(A1 \times B1 + \dots + An \times Bn)/n$  is mathematically unequal to  $[(A1 + \dots + An)/n] \times [(B1 + \dots + Bn)/n]$ .



Table 4. Yield characteristics of the *P. edulis* cultivars Lofuwa and Chankua as affected by zero (T0), one (T1), two (T2) and three (T3) tipplings in Wondogenet. Stem and stolon data from the harvest at 159 DAP, tuber data are from the final harvest at 208 DAP

Cultivar (CV) and tipping (T) treatment	Number of main stems per hole (#/hole)	×	Number of stolons per main stem (#/main stem)	=	Number of stolons per hole (#/hole)	×	Tuber: stolon ratio (#/#)	=	Number of tubers per hole (#/hole)	×	Average weight per tuber (g/tuber)	=	Tuber fresh weight per hole (g/hole)	×	Number of holes per m <sup>2</sup> (#/m <sup>2</sup> )	=	Tuber fresh yield per m <sup>2</sup> (g/m <sup>2</sup> )	×	Tuber dry matter concentration (g/g)	=	Tuber dry weight per m <sup>2</sup> (g/m <sup>2</sup> )
Cultivar Lofuwa																					
T0	1.6	×	23	=	36	×	3.4	=	122	×	11	=	1258	×	1.48	=	1864	×	0.14	=	268
T1 (68 DAP)	1.9	×	23	=	40	×	3.6	=	109	×	14	=	1465	×	1.48	=	2170	×	0.14	=	298
T2 (68+114 DAP)	1.8	×	19	=	34	×	3.8	=	132	×	11	=	1374	×	1.48	=	2036	×	0.15	=	298
T3 (68+114+158 DAP)	1.7	×	19	=	30	×	5.2	=	144	×	11	=	1329	×	1.48	=	1969	×	0.17	=	330
Average	1.74	×	20.9	=	35.0	×	4.03	=	126.8	×	11.4	=	1356.5	×	1.48	=	2009.8	×	0.148	=	298.3
Cultivar Chankua																					
T0	1.6	×	18	=	29	×	4.5	=	117	×	10	=	1195	×	1.48	=	1771	×	0.16	=	291
T1 (68 DAP)	1.6	×	18	=	29	×	5.4	=	145	×	10	=	1462	×	1.48	=	2167	×	0.17	=	368
T2 (68+114 DAP)	1.6	×	22	=	35	×	4.2	=	141	×	11	=	1561	×	1.48	=	2313	×	0.17	=	393
T3 (68+114+158 DAP)	1.6	×	18	=	29	×	5.7	=	145	×	10	=	1455	×	1.48	=	2155	×	0.17	=	366
Average	1.62	×	19.3	=	30.3	×	4.95	=	136.8	×	10.4	=	1418.5	×	1.48	=	2101.4	×	0.168	=	354.5
Average over cultivars																					
T0	1.6	×	21	=	32	×	4.0	=	120	×	10	=	1227	×	1.48	=	1818	×	0.15	=	280
T1 (68 DAP)	1.8	×	21	=	34	×	4.5	=	127	×	12	=	1463	×	1.48	=	2168	×	0.15	=	333
T2 (68+114 DAP)	1.7	×	21	=	35	×	4.0	=	136	×	11	=	1468	×	1.48	=	2174	×	0.16	=	345
T3 (68+114+158 DAP)	1.6	×	18	=	30	×	5.5	=	144	×	10	=	1392	×	1.48	=	2062	×	0.17	=	348
Average	1.68	×	20.1	=	32.7	×	4.49	=	131.8	×	10.9	=	1387.5	×	1.48	=	2055.6	×	0.158	=	326.4
S.E.D.																					
CV (5 D.F.)	0.047		ns		ns		ns		2.7		0.27		ns				ns		0.0042		11.8
Tlinear (30 D.F.)	ns		ns		ns		ns		11.4		ns†		69.2†				102.5†		0.0058		20.4†
Tquadratic (30 D.F.)	ns		ns		ns		ns		ns		ns†		69.2†				102.5†		ns		ns†
CV × Tlinear (30 D.F.)*	ns		ns		ns		ns		ns		ns†		ns†				ns†		ns		ns†
CV × Tquadratic (30 D.F.)*	ns		ns		ns		ns		ns		ns†		ns†				ns†		ns		ns†

DAP, days after planting.

\* For comparisons of tipping treatments within a cultivar only.

† 29 D.F.

cvar Lofuwa in Awassa (Table 3) and in both cultivars in Wondogenet (Table 4). This increase was not related to an increase in the number of stolons: the positive effects of tipping on stolon numbers early in the growing season were no longer obvious by the time the first tubers were initiated, and across all tipping treatments, there were no significant correlations between the numbers of tubers and stolons per hole. This is different from what is commonly found in potato, where there is generally a close relationship between the number of stolons and the number of tubers (e.g. Haverkort *et al.* 1990). An increase in tuber number per hole was more likely to be associated with more tubers per stolon. Although direct effects of tipping on the tuber:stolon ratios were inconsistent in Awassa and not significant in Wondogenet, significant correlations between the tuber:stolon ratio and the number of tubers per hole were found over tipping treatments in Awassa and one cultivar in Wondogenet. It seems likely that more tubers were initiated in the tipped treatments because of the higher light interception by the individual plants. This might be because of a higher rate of tuber initiation because of the increased canopy cover or because of a longer duration of tuber initiation because of the delayed senescence. In Irish potato, a higher light interception at the moment of tuber initiation – because of a larger canopy (e.g. as a result of a lower plant density) or because of less shading in shading experiments – was thought to increase the number of tubers initiated (Burstall *et al.* 1987; O'Brien *et al.* 1998; Van der Veen & Lommen 2009). The increase in number of tubers is consistent with the positive effect found by Menzel (1981) on number of tubers in potato after decapitating potato plants growing at high temperatures. This effect was attributed by Menzel (1981) to lowering of the gibberellin concentrations in the stolon, where they inhibit tuberization.

#### Intercepted radiation and yield

More leaves were produced on tipped plants than on non-tipped plants (Fig. 1a, b) and, as a result, more light was intercepted by tipped plants (Fig. 1c, d). Also, the stem parts of the extra branches (Table 1) may have contributed to the greater light interception, explaining the relatively small differences in green leaf numbers at the final harvest compared with the still larger difference in ground cover by the green canopy (Fig. 1). The heavy foliage formed by the

tipping treatments increased tuber dry matter yield at both sites by an accumulation of small effects on tuber fresh yield and tuber dry matter concentration (Tables 3 and 4).

Besides tipping, other cultural techniques might also enhance canopy cover and therefore radiation interception and yield. Alternatives could be to increase planting density (e.g. Bremner & Taha 1966; Van der Veen & Lommen 2009) or changing to a more evenly distributed planting pattern than the conventional planting of three tuber pieces per hole with a between-hole spacing of 0.75 m within a row and 0.90 m between rows. Also, changing the weight and pre-treatment of the seed tuber (pieces) are candidates for further investigation, because they are known to affect the number of main stems and/or canopy cover in potato (e.g. Bremner & Taha 1966; Allen *et al.* 1979; Lommen & Struik 1994).

#### Practical implications

Tipping stimulated the production of stems and branches and thus contributed to leaf area development and radiation interception and may have reduced weed problems. Tuber dry matter production and tuber numbers increased at both sites with an increase in the frequency of tipping, whereas the average tuber size was not affected. Fresh tuber yields obtained at 208 DAP without tipping were 1.90–1.98 kg/m<sup>2</sup> in Awassa and 1.86–1.77 kg/m<sup>2</sup> in Wondogenet. Tipping on average increased the fresh tuber yields by 17% in Wondogenet, but had no significant effect in Awassa ( $P=0.094$ ). Slightly later harvesting might have increased the differences in tuber yield between the tipped treatments and the control, because the ground cover of tipped plants was still higher than that of control plants at the final harvest. The results obtained in the present experiments confirm the farmers' perception that tipping will not only stimulate canopy development but also increase tuber yield. The latter, however, was only shown for tuber dry matter yield. For fresh tuber yield, limiting the tipping frequency to one early tipping when plants are 0.15 m tall might suffice, as more frequent tipping did not result in higher yields and would require more time and labour.

We gratefully acknowledge the financial support from the Norwegian Agency for Development and Corporation (NORAD).

## REFERENCES

- ALLEN, E. J., BEAN, J. N., GRIFFITH, R. L. & O'BRIEN, P. J. (1979). Effects of length of sprouting period on growth and yield of contrasting early potato varieties. *Journal of Agricultural Science, Cambridge* **92**, 151–163.
- ASFAW, Z. & WOLDU, Z. (1997). Crop associations of home-gardens in Welayta and Gurage in southern Ethiopia. *SINET, An Ethiopian Journal of Science* **20**, 73–90.
- BREMNER, P. M. & TAHA, M. A. (1966). Studies in potato agronomy I. The effects of variety, seed size and spacing on growth, development and yield. *Journal of Agricultural Science, Cambridge* **66**, 241–252.
- BRYAN, J. E., MELÉNDEZ, G. N. & JACKSON, M. T. (1981). *Stem Cuttings, a Rapid Multiplication Technique for Potatoes. CIP Slide Training Series: Series I: Rapid Multiplication Techniques. Guide-Book 1/3*. Lima, Peru: International Potato Center (CIP).
- BURSTALL, L., THOMAS, M. N. & ALLEN, E. J. (1987). Some effects of shading on tuber set and development. *Potato Research* **30**, 142–143.
- DUN, E. A., FERGUSON, B. J. & BEVERIDGE, C. A. (2006). Apical dominance and shoot branching: divergent opinions or divergent mechanisms. *Plant Physiology* **142**, 812–819.
- GREENWAY, P. J. (1944). Origins of some of East African food plants. Part 1. *East African Agricultural Journal* **10**, 34–39.
- HAVERKORT, A. J., VAN DE WAART, M. & BODLAENDER, K. B. A. (1990). Interrelationships of the number of initial sprouts, stems, stolons and tubers per potato plant. *Potato Research* **33**, 269–274.
- HAVERKORT, A. J., UENK, D., VEROUDE, H. & VAN DE WAART, M. (1991). Relationships between ground cover, intercepted solar radiation, leaf area index and infrared reflectance of potato crops. *Potato Research* **34**, 113–121.
- JANSEN, P. C. M. (1996). *Plectranthus rotundifolius* (Poir.) Sprengel. In *Plant Resources of South-East Asia No. 9. Plants Yielding Non-Seed Carbohydrates* (Eds M. Flach & F. Rumawas), pp. 141–143. Leiden: Backhuys Publishers.
- LOMMEN, W. J. M. & STRUIK, P. C. (1994). Field performance of potato minitubers with different fresh weights and conventional seed tubers: crop establishment and yield formation. *Potato Research* **37**, 301–313.
- LOMMEN, W. J. M. & STRUIK, P. C. (2007). The canon of potato science: 26. *In vivo* cuttings. *Potato Research* **50**, 309–313.
- MENZEL, C. M. (1981). Tuberization in potato at high temperatures: promotion by disbudding. *Annals of Botany* **47**, 727–733.
- O'BRIEN, P. J., FIRMAN, D. M. & ALLEN, E. J. (1998). Effects of shading and seed tuber spacing on initiation and number of tubers in potato crops (*Solanum tuberosum*). *Journal of Agricultural Science, Cambridge* **130**, 431–449.
- OLASANTAN, F. O. & SALAU, A. W. (2008). Effect of pruning on growth, leaf yield and pod yields of okra (*Abelmoschus esculentus* (L.) Moench). *Journal of Agricultural Science, Cambridge* **146**, 93–102.
- SMITH, T. M. (2003). *Production Guidelines for Four Crops: Osteospermum, Angelonia, Calibrachoa and Ornamental Sweet Potato (Ipomoea batatas)*. Fact Sheet. Department of Plant and Soil Sciences, University of Massachusetts, Amherst, USA. Available online [http://www.umass.edu/umext/floriculture/fact\\_sheets/specific\\_crops/newcrops.html](http://www.umass.edu/umext/floriculture/fact_sheets/specific_crops/newcrops.html) (verified 9 August 2011).
- TAYE, M., LOMMEN, W. J. M. & STRUIK, P. C. (2007). Indigenous multiplication and production practices for the tuber crop *Plectranthus edulis* in Chenchu and Wolaita, southern Ethiopia. *Experimental Agriculture* **43**, 381–400.
- VAN DER VEEKEN, A. J. H. & LOMMEN, W. J. M. (2009). How planting density affects number and yield of potato minitubers in a commercial glasshouse production system. *Potato Research* **52**, 105–119.
- WESTPHAL, E. (1975). *Agricultural Systems in Ethiopia. Agricultural Research Reports 826*. Wageningen, The Netherlands: Centre for Agricultural Publishing and Documentation (PUDOC).