

Euphytica (2011) 179:427–438
DOI 10.1007/s10681-010-0340-7

Identification and mapping of quantitative resistance to late blight (*Phytophthora infestans*) in *Solanum habrochaites* LA1777

Junming Li · Lei Liu · Yuling Bai · Richard Finkers ·
Feng Wang · Yongchen Du · Yuhong Yang · Bingyan Xie ·
Richard G. F. Visser · Adriaan W. van Heusden

Received: 19 November 2009 / Accepted: 28 December 2010 / Published online: 19 February 2011
© The Author(s) 2011. This article is published with open access at Springerlink.com

Abstract Late blight (*Phytophthora infestans*) can have devastating effects on tomato production over the whole world. Most of the commercial cultivars of tomato, *Solanum lycopersicum*, are susceptible. Qualitative and quantitative resistance has been described in wild relatives of tomato. In general qualitative resistance can more easily be overcome by newly evolved isolates. Screening of three *S. habrochaites* accessions (LA1033, LA2099 and LA1777) through a whole plant assay showed that accession LA1777 had a good level of resistance to several isolates of *P. infestans*. To explore the potential in this wild species, an introgression line (IL) population of *S. habrochaites* LA1777 was used to screen individual chromosome regions of the wild species by a detached leaf assay. Two major isolates (T_{1,2} and T_{1,2,4}) were used and two parameters were measured: lesion size (LS), and disease incidence (DI). Substantial variation was observed between the individual lines. QTLs were identified for LS but not for DI.

The presence of five QTLs derived from LA1777 (*Rlbq4a*, *Rlbq4b*, *Rlbq7*, *Rlbq8* and *Rlbq12*) results in unambiguous higher levels of resistance. All QTLs co-localized with previously described QTLs from *S. habrochaites* LA2099 except QTL *Rlbq4b*, which is therefore a novel QTL.

Keywords Tomato · Late blight · *Phytophthora infestans* · Quantitative resistance · *Solanum habrochaites* · Introgression lines

Introduction

The oomycete *Phytophthora infestans* (Mont.) de Bary, the causal agent of late blight, is one of the most destructive pathogens of potato and tomato. Late blight causes serious yield and economic losses especially under favorable conditions for the pathogen (wet and cool temperatures) both in the open field as well as in non-heated greenhouses. The responsible pathogen is heterothallic and forms oospores with A1 and A2 mating types and has been found in different areas of the world (Gotoh et al. 2005). The co-existence of mating types and the sexual reproduction increase the chance of developing resistance to fungicides such as metalaxyl (Goodwin et al. 1998; Gotoh et al. 2005). In addition, the spread of the disease may be also initiated from spores present in the soil (Widmark et al. 2007). The genetic diversity,

J. Li (✉) · L. Liu · F. Wang · Y. Du · Y. Yang · B. Xie
Institute of Vegetables and Flowers, Chinese Academy of
Agricultural Sciences, No. 12 Zhongguancun Nandajie,
Haidian District, Beijing 100081, China
e-mail: junmingli@mail.caas.net.cn

J. Li · Y. Bai · R. Finkers · R. G. F. Visser ·
A. W. van Heusden
Graduate School Experimental Plant Sciences,
Wageningen UR Plant Breeding, Wageningen University
and Research Center, PO box 386, 6700 AJ Wageningen,
The Netherlands

rapid evolution and the broader range of virulence factors have made this pathogen more and more aggressive (Drenth et al. 1995; Gotoh et al. 2005). Tomato plants can be completely destroyed in a few weeks despite the use of chemicals to control the infections (Jones et al. 1991).

The most effective and environmentally favorable way to prevent devastation of tomato plants by this pathogen is to incorporate natural resistance into cultivars. Two kinds of host plant resistance to *P. infestans* have been described in tomato (Labate et al. 2007). Firstly there is the qualitative or race-specific resistance. This resistance is based on “R” genes, examples are *Ph-1*, *Ph-2*, *Ph-3*, *Ph-4* and *Ph-5* which originate from the wild species *S. pimpinellifolium* and the position of these R-genes has been determined on chromosomes 7, 10, 9, 2 and 1, respectively (Chunwongse et al. 2002; Peirce 1971; Foolad et al. 2008; Kole et al. 2006; Moreau et al. 1998). However, these qualitative resistances are not durable due to the rapid evolution of compatible races of the pathogen. The first four genes have already been broken by newly evolved races of *P. infestans* (Kole et al. 2006; Foolad et al. 2008) and the durability of the *Ph-5* gene is still a question. It is expected that combining or pyramiding several genes could provide a more durable resistance than deploying just a single one (Foolad et al. 2008).

The second type of resistance is quantitative resistance that is controlled by quantitative trait loci (QTLs). Quantitative resistance is often non race-specific and more durable than qualitative resistance governed by R-genes (Brun et al. 2010; Palloix et al. 2009; Robert et al. 2009). For resistance to late blight, research in potato has been since 1970 focusing on introducing quantitative resistance (Wastie 1991), in spite of the fact that already eleven single R-genes were known. In tomato, *S. habrochaites* is believed to be a potential donor for high levels of quantitative resistance (Brouwer et al. 2004). Five to six consistent QTLs have been identified in two BC₁ populations derived from *S. habrochaites* LA2099 and lines with all four major QTLs introgressed were more resistant to *P. infestans* in different environments. Recently, one QTL was identified from the wild species *S. pennellii* LA716 (Smart et al. 2007). These identified QTLs could be explored in tomato breeding for quantitative and non race-specific resistance to *P. infestans*.

An introgression line population (IL) has several advantages over segregating populations such as F₂ and BC₁. Such a population is advantageous for QTL mapping because it can be phenotyped with many replicates and in different environments, which makes it possible to detect QTLs with smaller effects and allows also an estimation of the Genotype × Environment (G × E) interaction (Chaïb et al. 2006; Eshed and Zamir 1996; Gur and Zamir 2004; Lecomte et al. 2004; Monforte et al. 2001). At least five IL populations have been developed in tomato; they are derived from *S. pennellii* LA716 (Eshed and Zamir 1994), *S. habrochaites* LA1777 (Monforte and Tanksley 2000), *S. habrochaites* LA407 (Francis et al. 2001), *S. habrochaites* LYC4 (Finkers et al. 2007) and *S. lycopersicoides* LA2951 (Canady et al. 2005). The *S. pennellii* IL library has been extensively explored to identify QTLs for several traits including disease resistances (Astua-Monge et al. 2000; Smart et al. 2007), fruit quality (Rousseaux et al. 2005; Tieman et al. 2006) and yield (Eshed et al. 1996). Recently, an introgression line of *S. habrochaites* LA1777 has been identified with a significant contribution to marketable fruit yield (Hanson et al. 2007).

In this paper we describe the screening of three *S. habrochaites* accessions (LA1777, LA2099 and LA1033), which have shown high levels of resistance to *P. infestans* (Brouwer et al. 2004), and we found that of the three accessions LA1777 gave the highest resistance levels to several races of late blight originating from China, especially to *P. infestans* race T_{1,2,3,4} which already overcame the *Ph-1*, *Ph-2*, *Ph-3* and *Ph-4* genes. Since the genetic distance between LA2099 and LA1777 is substantial, LA1777 might harbor other QTLs for late blight resistance as LA2099. In this paper, the IL population derived from *S. habrochaites* LA1777 (Monforte and Tanksley 2000) has been screened. Results and comparisons with earlier studies are presented and discussed.

Materials and methods

Plant material

Three accessions of *S. habrochaites* (LA1777, LA2099 and LA1033) were tested together with control lines. As susceptible control, we used

S. lycopersicum 99165 (tomato inbred line derived from the progeny of OH211 × NS217, USA) and *S. lycopersicum* HZ14 and HZ18 (two commercial hybrids widely used in China for processing tomato production). As resistance control, we included the inbred line CLN2037B (provided by AVRDC–The World Vegetable Centre) harboring the *Ph-3* gene. Disease testing was always performed after the sixth true leaf had developed.

The introgression lines (IL) used in this study were derived from *S. habrochaites* accession LA1777 (a self-incompatible, homozygous green fruited, indeterminate accession) in the background of *S. lycopersicum* E6203 (a red fruited, determinate, processing-type tomato). In total this IL population has 98 ILs that cover at least 85% of the wild species genome (Monforte and Tanksley 2000). In this study, all these ILs were screened except for LA3973, LA3974, LA3982, LA3987 and LA3992. Since introgressions in different lines overlap and the introgressions in the 93 lines used in this study are located on all the 12 tomato chromosomes, we expect that the 93 lines could still cover about 85% of the wild species genome. Seeds were kindly provided by the Tomato Genetic Resource Center (TGRC, Davis USA). In order to get new seeds fruits of the individual introgression lines were collected after self pollination in the greenhouse and the accession LA1777 was maintained by pollination with a pollen mixture from different plants.

The seeds were germinated in an incubator at 25°C and then transferred to 10 cm pots containing a medium of peat–vermiculite with organic fertilizer. Greenhouse temperature ranged from 15 to 18°C at night and from 20 to 25°C at day time.

Inoculum preparation

Two *P. infestans* isolates from China were used in the resistance assays: T_{1,2} and T_{1,2,4} (Chen et al. 2008). Among them, race T_{1,2} is the most epidemic isolate in China and present in eighteen provinces (Dr. Feng IVF, CAAS Beijing, China and Dr. Tian AVRDC, Taiwan; personal communication). Isolates T_{1,2}, and T_{1,2,4} are virulent on tomato genotypes containing the resistance genes *Ph-1* and *Ph-2* or *Ph-1*, *Ph-2* and *Ph-4*, respectively. For the three *S. habrochaites* accessions one extra *P. infestans* race was used. This isolate, T_{1,2,3,4}; is the most virulent race found in

China and was collected in a greenhouse in Beijing, China by the department of Pathology, IVF, CAAS.

Cultures of *P. infestans* were grown at 17°C on Rye B agar (Caten and Jinks 1968) and transferred to new plates monthly. Isolates were periodically grown on leaves of susceptible control tomato cv Zaofeng no.2 to maintain pathogenicity and profuse sporulation. Inoculum for disease assays was prepared by washing 8-day-old sporulating lesions with sterile distilled water. Spore concentrations were determined using a hemocytometer and diluted to the desired concentration (1×10^4 spores ml⁻¹).

Detached-leaflet assay

The *S. habrochaites* IL population was evaluated by a droplet method using T_{1,2} race in five independent experiments and T_{1,2,4} race in one independent experiment. Five to fifteen plants of each genotype were used for each experiment. From each individual plant the sixth true leaf was detached with a razor blade and immediately inserted in moist florist foam. The abaxial surface of three of the top leaflets was inoculated with a drop of 20 µl of sporangial suspension (1×10^4 spores ml⁻¹). Leaves were transferred to transparent plastic boxes, sealed with a transparent plastic membrane, covered by the lids and randomly placed in a growth cabinet at 16°C without light. After 24 h, the regime was changed to 16°C with 12 h light and 12 h dark. Late blight resistance was assessed six days post inoculation (dpi). The largest length and width (perpendicular to the length) of each lesion was measured resulting in the lesion size (LS) and the ellipse area was calculated following the formula $LS = (\text{length} \times \text{width} \times \pi)/4$. No lesion or a lesion remaining within the size of the inoculum droplet ($\leq 0.3 \text{ cm}^2$) was considered as no infection or as arrested lesion (Vleeshouwers et al. 1999). For each genotype, the percentage of successfully infected leaflets was calculated as disease incidence (DI).

Whole plant assay in growth cabinets

Wild species were tested for resistance using a whole plant assay with races T_{1,2}, T_{1,2,4} and T_{1,2,3,4}. Greenhouse-grown plants with fully stretched six true leaves were moved to cabinets. Thirty plants of each genotype were evaluated in three blocks using a randomized complete block design.

For whole plant inoculations, each plant was spray inoculated until the water started to drip off. The dew cabinets had an air temperature of 17–18°C. The first 24 h after inoculation no light was used, after this a regime of 12 h light (18°C): 12 h dark (16°C) was used. After seven days the plants were scored individually for disease severity on a scale of 0–6, where 0 = no symptoms; 1 = < 5% leaf area affected and small (<2 mm) lesions; 2 = 6–15% leaf area affected and restricted (<4 mm) lesions; 3 = 16–30% leaf area affected and/or few superficial small stem lesions; 4 = 31–60% leaf area affected and/or few small penetrating stem lesions; 5 = 61–90% leaf area affected and/or deep expanding stem lesions; 6 = 91–100% leaf area affected, extensive stem damage, or plant death (Chunwongse et al. 2002; Brouwer et al. 2004). Percentage disease index (PDI) was calculated with the following formula for each genotype: $PDI = \text{sum of all ratings} \times 100 / \text{total no. of observations} \times \text{maximum rating grade}$ (Chaerani et al. 2007).

Statistical analysis

All statistical analyses were performed using SPSS 13.0. Differences in *P. infestans* resistance in the *S. habrochaites* IL population were analyzed using the procedure of general linear model (GLM). LS data was transformed by square root to meet a normal distribution. Mean estimates for each IL line were calculated using the following models: $PDI = \text{constant} + \text{genotype} + \text{block} + \text{genotype} \times \text{block}$. $LS = \text{constant} + \text{genotype} + \text{experiment} + \text{genotype} \times \text{experiment}$. $DI = \text{constant} + \text{genotype} + \text{experiment} + \text{genotype} \times \text{experiment}$. The correlation between traits was calculated by Pearson correlation coefficients. Trait data for experiments were tested for homogeneity of variance using a Levene test in order to evaluate the normal distribution. Significance of QTL was determined by comparing mean values of individual ILs to the control *S. lycopersicum* E6203 at the 0.05 level by Dunnett test.

Results

Comparison of three accessions of *S. habrochaites* with whole plant assay

From an earlier experiment it was clear that *S. habrochaites* accession LA1777 was resistant to

P. infestans race $T_{1,2}$ (Fig. 1). Analysis of different accessions and controls with race $T_{1,2,4}$ of *P. infestans* (Fig. 2) show that all three accessions of the wild species *S. habrochaites* gave a good resistance level to that particular race. However, the two accessions LA1777 and LA2099 with a mean PDI of 20.0 and 21.7 were more resistant than the accession LA1033 (mean PDI ~ 34), but all were significantly more resistant than the susceptible controls (mean PDI ~ 86.1). The same two accessions of *S. habrochaites* (LA1777 and LA2099) also showed enhanced resistance levels to the most virulent race: $T_{1,2,3,4}$ (Fig. 2).

Detached leaflet assay in LA1777 IL population with race $T_{1,2}$

The IL population was analyzed with the detached leaflet assay using isolate $T_{1,2}$ in five independent experiments over two years. Two traits were evaluated for each individual ILs: lesion size (LS) expressed as the mean size of *P. infestans* lesions of infected leaves, and disease incidence (DI) expressed as the percentage of inoculated leaves that were successfully infected.

Between experiments, the mean LS varied from 2.34 ± 0.06 to 5.00 ± 0.05 cm², while the mean DI varied from 75.6 ± 0.9 to $84.0 \pm 0.6\%$ for the IL population. The disease scorings of LS were higher in 2006 than in 2007. However, the mean DI remained more or less the same over all five experiments. Significant correlations were observed for LS between experiments 1, 2, 4 and 5 ($r = 0.21$ – 0.37 , $P < 0.05$ or $P < 0.01$) but no significant correlation was present with the LS of experiment 3 (Table 2). Experiment 3 was excluded for the final analysis of the results, but this only influenced the level of significance for each IL. Significant correlations for DI were only observed between experiments 1 and 4 ($r = 0.357$, $P < 0.01$), and between experiments 3 and 5 ($r = 0.283$, $P < 0.05$). The data of the experiments with significant correlation were analyzed. There is a significant difference between *S. habrochaites* LA1777 (2.33 ± 0.31 cm²) and *S. lycopersicum* E6203 (4.05 ± 0.16 cm²) and the mean LS ranged from 2.89 ± 0.19 to 6.28 ± 0.48 cm² among the ILs. A total of fifty-four lines showed smaller LS (0.30–28.5%) and this was significant in thirty-one lines. Thirty-two lines identified herein can harbor a number of QTLs conferring resistance to *P. infestans* as determined by LS (Table 1). We designated the identified QTLs as

Fig. 1 Screening of tomato wild species for resistance to *P. infestans* race T_{1,2}. *Top left* *S. habrochaites* LA1777. *Top right* the susceptible control *S. lycopersicum* 99165. *Bottom left* the susceptible control *S. lycopersicum* HZ14. *Bottom right* the resistant control *S. lycopersicum* CLN2037B with *Ph-3* gene (provided by AVRDC)





Fig. 2 Comparison of three accessions of wild species *S. habrochaites* for resistance to isolates T_{1,2,4} and T_{1,2,3,4}. The bar indicates the standard error. ** indicates values in the lines that are significantly different from the susceptible control (99165) at the 0.01 level

Resistance to Late Blight QTL (*Rlbq*) followed by the number of the chromosome on which they are located.

Based on the IL map constructed by Monforte and Tanksley (2000), the introgressions of these thirty-one lines derived from wild species *S. habrochaites* LA1777 were distributed on 11 of the 12 chromosomes. We will focus in this paper only on the five

most significant and substantial QTLs, which were located on chromosomes 4 (2 QTLs), 7, 8 and 12.

Rlbq4a and *Rlbq4b*. Sixteen of the 93 lines contain an introgression of chromosome 4. Ten of these lines were significantly more resistant than the control (Table 2 and Fig. 3). The significant effects in lines LA3931, LA3959, LA3979 and LA4006 show the presence of a QTL (*Rlbq4a*) at the top of Chromosome 4. More markers will have to be determined to pinpoint the QTL more precisely. The fact that lines LA3930 and LA4000 are not resistant will make it likely that the position can be determined rather precisely. However, also some lines with other introgressions on Chromosome 4 had higher resistance levels, the significant higher resistance levels of ILs LA3934, LA3935, LA3937, LA3976 and LA4007 shows that there must be a second QTL (*Rlbq4b*) towards the bottom of Chromosome 4. Again additional markers are needed to explain why lines such as LA3936, LA3977 and LA3978 do not show resistance.

In the same manner, we identified three other QTLs for smaller LS: on top of Chromosome 7: *Rlbq7*, and

Table 1 Estimated mean of lesion size (LS) and disease incidence (DI) in introgression lines (IL) and their two parents, resistant *S. habrochaites* LA1777 and susceptible *S. lycopersicum* cv. E6203

ILs	LS (cm ²)	N ^a	DI (%)	N ^b
LA3915	3.91 ± 0.21*	98	83.8 ± 4.6	117
LA3916	3.76 ± 0.19*	107	86.3 ± 3.9	124
LA3918	3.95 ± 0.21*	97	82.9 ± 4.2	117
LA3919	3.72 ± 0.24**	86	72.9 ± 4.5	118
LA3921	3.75 ± 0.20**	101	82.1 ± 4.2	123
LA3922	3.92 ± 0.20**	97	85.1 ± 4.3	114
LA3923	3.66 ± 0.22**	83	76.6 ± 4.4	109
LA3925	3.52 ± 0.21**	88	82.2 ± 4.3	107
LA3929	3.48 ± 0.19**	101	82.8 ± 4.0	122
LA3931	3.82 ± 0.19*	106	82.2 ± 3.9	129
LA3932	3.61 ± 0.22**	82	73.9 ± 4.0	111
LA3934	3.49 ± 0.21**	95	76.6 ± 4.3	124
LA3935	3.59 ± 0.22**	93	76.9 ± 4.6	121
LA3937	3.20 ± 0.21**	85	72.0 ± 3.9	118
LA3941	3.19 ± 0.26**	64	66.0 ± 4.8*	97
LA3948	3.67 ± 0.21*	92	74.8 ± 3.9	123
LA3949	2.89 ± 0.19**	102	77.3 ± 3.7	132
LA3959	3.39 ± 0.19**	102	82.3 ± 3.8	124
LA3961	3.72 ± 0.21*	96	76.2 ± 3.9	126
LA3963	3.67 ± 0.19*	100	84.0 ± 3.8	119
LA3964	3.92 ± 0.21*	84	83.2 ± 4.4	101
LA3965	3.56 ± 0.18**	107	80.9 ± 3.6	132
LA3967	3.40 ± 0.20*	101	82.8 ± 3.9	122
LA3969	3.46 ± 0.19**	111	88.8 ± 4.0	125
LA3976	3.45 ± 0.19**	108	87.1 ± 3.9	124
LA3979	3.75 ± 0.22**	86	76.1 ± 4.3	113
LA3988	3.73 ± 0.22**	88	80.7 ± 4.7	109
LA3989	3.75 ± 0.19**	113	88.3 ± 3.9	128
LA3990	3.95 ± 0.20*	85	70.8 ± 3.8	120
LA4006	3.59 ± 0.17**	119	86.2 ± 3.6	138
LA4007	3.74 ± 0.19*	104	83.9 ± 3.9	124
LA3954	3.78 ± 0.21	84	65.6 ± 3.7*	128
E6203	4.05 ± 0.16	141	83.4 ± 3.3	169
LA1777	2.33 ± 0.31**	51	38.9 ± 5.7**	131

Means of each trait for each IL were compared to the mean of the susceptible parent using a Dunnett test by GLM mode and significant differences are marked with * ($P < 0.05$) or ** ($P < 0.01$)

^a Number of leaflets that had lesion growth

^b Number of leaflets that were tested in four experiments

Fig. 3 Rows indicate introgression line (IL) and columns indicate chromosomes. For each chromosome, the top and bottom is from the left to right. Solid segments show introgressions in the IL based on Monforte and Tanksley (2000). Following each IL, the asterisk showed the significant differences as compared to the susceptible control *S. lycopersicum* E6203 for leaf LS (* $P < 0.05$ and ** $P < 0.01$). The arrows indicate the most likely location of the identified QTLs through their overlapped ILs


bottom of Chromosome 8: *Rlbq8*, and the middle of Chromosome 12: *Rlbq12* (Table 2 and Fig. 3).

The mean DI of *S. habrochaites* LA1777 was 55.9%, a significant lower value than in the susceptible parent *S. lycopersicum* E6203 with 83.4%. The values of the individual lines of the IL population varied from 65.6 to 90.8%. Two lines, LA3941 (Chr. 5) and LA3954 (Chr. 8), showed a significantly decreased DI. However, this was not confirmed in other lines with overlapping introgressions. Hence, the identified QTLs must be further confirmed in additional experiments.

Detached leaflet assay in LA1777 IL population with race $T_{1,2,4}$

In order to see whether the identified resistance is real and consistent, the more virulent race $T_{1,2,4}$ was used in one experiment. The difference in LS between the parental lines *S. habrochaites* LA1777 (0.91 ± 0.26 cm²) and *S. lycopersicum* E6203 (3.20 ± 0.29 cm²) was highly significant. For the individuals of the IL population LS varied from 1.22 ± 0.26 cm² (line LA3918) to 6.39 ± 0.25 cm² (LA3969). Of the 93 IL lines, 53 had a reduced LS ranging from 0.67 to 61.85%. Twelve lines gave a significant difference; these were LA3914, LA3918, LA3920, LA3922, LA3923, LA3928, LA3937, LA3948, LA3981, LA3999, LA4004 and LA4005. Five lines, LA3918, LA3922, LA3923, LA3937 and LA3948 were also identified after inoculation with race $T_{1,2}$. All lines containing QTL *Rlbq4b* or *Rlbq7* showed lower LS than the control.

The mean DI of *S. habrochaites* LA1777 (63.4%) was significantly different from the DI of *S. lycopersicum* E6203 (83.4%). The DI ranged among the individuals of the IL population from 66.0 to 92.7% (data not shown), but were not significantly different from the control.



Discussion

Wild species conferring resistance to *P. infestans*

Three wild species, *S. pimpinellifolium*, *S. pennellii* and *S. habrochaites*, have been reported to give

qualitative and quantitative resistance to *P. infestans* (Chunwongse et al. 2002; Conner and Walter 1953; Moreau et al. 1998; Smart et al. 2007; Turkensteen 1973). Quantitative resistance in plants has been shown to be more durable (Brun et al. 2010; Palloix et al. 2009). In *S. habrochaites* a high level of

Table 2 Leaf lesion size (LS) assayed in four independent experiments, and means of LS and disease incidence (DI) of ILs containing the introgressions and their overlapped ILs on chromosome 4, 7, 8 and 12, respectively and two parent lines

IL	QTL	LS (cm ²) in four experiments				Mean LS (cm ²)	Mean DI (%)
		1	2	4	5		
LA3930	<i>Rlbq4a</i>	4.62	5.84	2.28	3.49	4.06 ± 0.20	76.8 ± 3.8
LA3931		4.81	5.11	2.28	3.09	3.82 ± 0.19*	83.4 ± 3.9
LA3932		4.56	4.40	2.43	3.04	3.61 ± 0.23**	72.0 ± 4.0
LA3933		5.19	5.50	3.06	4.27	4.50 ± 0.18	86.7 ± 3.9
LA3959		4.66	3.84	2.41	2.64	3.39 ± 0.19**	81.1 ± 3.8
LA4000		5.85	5.29	2.70	3.34	4.30 ± 0.20	76.5 ± 4.0
LA4006		4.38	4.19	2.58	3.20	3.59 ± 0.17**	85.3 ± 3.6
LA3934	<i>Rlbq4b</i>	4.62	4.75	2.18	2.39	3.49 ± 0.21**	78.9 ± 4.3
LA3935		5.51	3.73	2.58	2.54	3.59 ± 0.22**	81.0 ± 4.6
LA3936		4.53	4.33	2.84	3.38	3.78 ± 0.19	83.7 ± 3.9
LA3937		3.83	3.29	2.34	3.34	3.20 ± 0.21**	70.5 ± 3.9
LA3976		4.33	3.96	2.77	2.73	3.45 ± 0.19**	86.1 ± 3.9
LA3977		6.38	4.80	2.76	3.16	4.27 ± 0.24	71.9 ± 4.7
LA3978		4.77	3.65	3.18	3.84	3.86 ± 0.21	71.0 ± 4.1
LA3979		3.71	5.13	2.70	3.46	3.75 ± 0.22**	73.8 ± 4.3
LA4007		4.96	3.79	2.58	3.57	3.73 ± 0.19*	82.9 ± 3.9
LA3948	<i>Rlbq7</i>	4.36	4.24	2.74	3.35	3.67 ± 0.21*	73.1 ± 3.9
LA3949		3.66	3.39	1.90	2.62	2.89 ± 0.19**	77.4 ± 3.7
LA3950		5.24	6.34	–	7.28	6.28 ± 0.48	65.8 ± 8.9
LA3951		5.93	5.06	2.60	3.98	4.39 ± 0.18	82.1 ± 3.7
LA3985		4.42	5.85	3.39	4.96	4.66 ± 0.21	87.3 ± 4.6
LA3986		4.16	2.19	2.99	4.41	3.44 ± 0.21	78.0 ± 4.3
LA4001		5.46	4.21	3.18	3.45	4.08 ± 0.22	85.5 ± 4.6
LA4009		5.72	4.65	–	–	5.19 ± 0.42	69.0 ± 7.9
LA3928	<i>Rlbq8</i>	3.66	5.31	–	–	4.49 ± 0.34	76.2 ± 6.4
LA3929		4.43	4.61	2.06	2.80	3.48 ± 0.19**	82.3 ± 4.0
LA3942		4.39	3.81	3.22	3.61	3.76 ± 0.20	70.6 ± 3.8
LA3952		5.24	4.26	2.59	3.595	3.92 ± 0.18	78.8 ± 3.7
LA3953		4.65	4.57	2.79	3.10	3.78 ± 0.20	84.1 ± 4.0
LA3954		4.97	4.15	2.85	3.16	3.78 ± 0.20	65.6 ± 3.7**
LA3955		4.19	5.09	3.54	2.71	3.88 ± 0.19	80.6 ± 3.8
LA3988		5.52	3.55	3.19	2.66	3.73 ± 0.22**	85.2 ± 4.7
LA3989		5.36	4.06	2.31	3.28	3.75 ± 0.19**	86.0 ± 3.9
LA3990		5.14	5.09	2.32	3.27	3.95 ± 0.20*	69.9 ± 3.8**
LA4002		5.09	4.38	3.64	4.05	4.29 ± 0.22	78.2 ± 4.4
LA4010		6.57	4.67	2.82	3.88	4.48 ± 0.19	84.2 ± 3.9

Table 2 continued

IL	QTL	LS (cm ²) in four experiments				Mean LS (cm ²)	Mean DI (%)
		1	2	4	5		
LA3917	<i>Rlbq12</i>	5.55	5.53	2.92	3.38	4.35 ± 0.28	67.6 ± 5.5*
LA3960		4.79	4.52	2.45	3.47	3.81 ± 0.19	81.3 ± 3.5
LA3962		5.40	5.56	2.74	3.48	4.29 ± 0.18	81.0 ± 3.5
LA3966		6.23	4.90	3.18	3.57	4.47 ± 0.22	88.8 ± 3.8
LA3968		5.45	5.75	3.28	3.24	4.43 ± 0.19	72.5 ± 3.5
LA3969		4.28	4.09	2.36	3.12	3.46 ± 0.19**	88.4 ± 3.7
LA3995		4.61	5.19	3.03	3.56	4.10 ± 0.20	86.9 ± 3.6
LA3998		5.11	6.16	3.76	4.02	4.76 ± 0.20	81.8 ± 3.8
LA3999		3.97	6.24	3.64	2.55	4.10 ± 0.18	79.2 ± 3.3
LA4001		5.46	4.21	3.18	3.45	4.08 ± 0.22	83.1 ± 4.1
LA4002		5.09	4.38	3.64	4.05	4.29 ± 0.22	78.6 ± 4.0
LA4007		4.96	3.79	2.58	3.77	3.73 ± 0.29*	83.6 ± 3.6
LA4010		6.57	4.67	2.82	3.88	4.48 ± 0.29	84.7 ± 3.6
E6203		6.23	4.05	3.10	2.94	4.05 ± 0.16	83.4 ± 3.3
LA1777		2.75	2.17	2.25	2.51	2.33 ± 0.31**	55.9 ± 5.7

Means of LS for each IL were compared to the mean of the susceptible parent using a Dunnett test by GLM mode and significant differences are marked with * ($P < 0.05$) or ** ($P < 0.01$)

quantitative resistance to several isolates has been found (Brouwer et al. 2004). In our study, three wild accessions of *S. habrochaites* were evaluated for resistance to different races of *P. infestans* and the resistance level of the three accessions is quite different for two races of *P. infestans*. Accessions LA1777 and LA2099 had a higher level of resistance to race T_{1,2,4} than accession LA1033. To the most virulent race T_{1,2,3,4} accession LA1033 was almost completely susceptible in contrast to a very good level of resistance in both LA1777 and LA2099. The virulence of race T_{1,2,3,4} on accession LA1033 maybe caused by the fact that this accession has allelic variants of the *Ph-3* gene, a gene which can be overcome by race T_{1,2,3} (Chunwongse et al. 2002). Brouwer et al. (2004) have shown that LA2099 has a very good resistance to USA isolates 7629 and 9175, which are virulent on tomato genotypes containing the *Ph-1* and *Ph-2* gene. We found in this study that LA1777 and LA2099 have potential resistances to several other races of *P. infestans* with LA1777 as the most resistant. Recently, two new resistance genes, *Ph-4* and *Ph-5*, have been identified from *S. pimpinellifolium* (Foolad et al. 2008; Kole et al. 2006) and another gene located on chromosome 6 of *S. pennellii*

was described by Smart et al. (2007). We also screened a nightshade *S. lycopersicoides* LA2951 IL population and it seems that this population also harbors resistance to *P. infestans* (data not shown). Hence, it might be that more wild species can serve as potential sources for resistance genes to *P. infestans*. These wild species provide a rich resource for breeding tomatoes with resistance to late blight.

QTLs identified in different experiments

We have made an effort to explore introgression lines of *S. habrochaites* LA1777 for resistance loci against *P. infestans* by a detached leaflet assay. As a permanent population, these ILs allowed us to test them with a major race (T_{1,2}) of *P. infestans* in five independent detached leaf experiments over two years. The mean LS in these two years varied greatly which might be caused by inoculum quality or differences in individual lines under different experimental conditions (Vleeshouwers et al. 1999) and each genotype from an IL population might adapt in a different way to the environment. Low to moderately low correlations were also reported among different assay methods (Brouwer et al. 2004). To identify introgressions with

resistance genes only the experiments with significant correlation between experiments were used in this study. While analyzing the data of IL populations, the five QTLs could only be identified after combining the data of four independent experiments because not all of them were significant in single experiments. Some of the identified QTLs were not detected when another inoculum (race T_{1,2,4}) was used. In conclusion independent experiments are needed in search for quantitative resistance to *P. infestans*. The whole-plant assay was also used to screen the *S. habrochaites* LA1777 IL population in three replications (data not shown). However, only small differences could be observed between ILs and the susceptible control. With the detached leaf assay substantial differences were observed. Brouwer et al. (2004) reported that neither a detached leaflet nor a whole-plant assay can entirely substitute *P. infestans* screenings in tomato. Hence, field or greenhouse tests should add more evidence to prove the true nature of the identified QTLs (Brouwer et al. 2004).

DI as a measure for infection efficiency was also evaluated in our study. For DI, low or no correlation between experiments was found and QTLs responsible for DI could not be identified in this study. Vleeshouwers et al. (1999) showed that a highly constant humidity in closed trays apparently enhances infection by the zoospores causing a very high disease pressure. In our experiments, the plastic box is always covered by the lid, which would enhance the chance for the successful infection as proved by Vleeshouwers et al. (1999) and Brouwer et al. (2004). In the detached leaf assay some well known resistant wild *Solanum* genotypes were partially infected. A high amount of successful infections reduces the change to find QTLs for disease incidence but makes the chance higher to find QTLs for LS (more data points). Vleeshouwers et al. (1999) suggested when the DI is to be used as a parameter for resistance, a different screening methodology must be chosen, e.g., incubation of detached leaves in open trays, or intact plants in climate chamber or field. Other complicating factors in a bioassay can be the significant negative correlation between percentage infection and plant height (Brouwer et al. 2004) and rotting of leaves during the evaluation. Brouwer et al. (2004) found that determinate genotypes were more susceptible at least to late blight than indeterminate genotypes because the indeterminate type is higher

than the determinate type. In our paper, all of ILs are determinate type, which would be more susceptible. Also, we tried not to include these rotten leaves in the data analysis, but it is sometimes difficult to distinguish between infected and rotten leaves. This can be the reason that introgression lines appear to be more susceptible than the control as the leaves of *S. habrochaites* LA1777 rot more easily than the leaves of the ILs.

In this study we only focused on lines with larger effect on resistance to *P. infestans*. Only QTLs identified in several lines with overlapping introgressions are considered as reliable. Since the IL population we used covered about 85% of the genome of LA1777 accession, some QTLs for resistance to *P. infestans* might be missed therefore. Moreover, the exact location of each QTL still needs to be confirmed due to lack of precise flanking markers in each introgression line.

Comparative analysis of QTLs from different populations

On all 12 tomato chromosomes many QTLs for resistance to *P. infestans* have been detected in another *S. habrochaites* accession namely LA2099 with only eight QTLs showed consistent resistance over experiments (Brouwer et al. 2004). We found that *Rlbq4a*, *Rlbq7*, *Rlbq8b* and *Rlbhq12* co-localize with previously identified *lb4a*, *lb7a*, *lb8b* and *lb12b* respectively on chromosome 4, 7, 8 and 12 (Brouwer et al. 2004). Because *Rlbq4b* was not detected in the previous study and the fact that it showed a good resistance against two different races of late blight, we think that this novel QTL from LA1777 is important and worthwhile introgressing in tomato varieties. A next step could be to clone the genes underlying these QTLs.

Potential of pyramiding QTLs

Up to now, both qualitative and quantitative genes have been identified in several different wild tomato species. The interaction between these QTLs and single genes is still unknown but lines with all four QTLs, from *S. habrochaites* LA2099, have shown a high level of resistance to *P. infestans* under different environments (Brouwer et al. 2004; Brouwer and St Clair 2004). In our study, five QTLs have been

identified and two of them give also resistance to the most virulent race. Not all QTLs were identified in each experiment and in all lines with overlapping introgressions, indicating that most QTLs have a limited effect and interactions between QTLs as well as between QTLs and environment play a role in obtaining a high level of resistance. Combining QTLs and single resistance genes can be beneficial for durability of the resistance to several fungi (Brun et al. 2010; Palloix et al. 2009; Stall et al. 2009). Therefore we suggest to develop combinations of several QTLs or QTLs and single genes, not only from one wild species but also from different wild species. For example a combination of QTL *Rlbq4b* derived from *S. habrochaites* LA1777, one QTL derived from *S. pennellii* LA716 (Smart et al. 2007) and some QTLs derived from *S. habrochaites* LA2099 (Brouwer et al. 2004). In this way pyramiding of these or similar effective QTLs might pave the way for durable resistance to *P. infestans*.

Acknowledgments This project was sponsored by National Basic Research and Development Program (2009CB119000), Hi-Tech Research and Development Program (2006AA10Z1A6), Key Laboratory of Horticultural Crops Genetic Improvement (Ministry of Agriculture of the People's Republic of China), the Royal Dutch Academy of Sciences (KNAW) and the Asian Facility (project AF01/CH/8 “Sino-Dutch Genomic Lab and Vegetable Research Center”). The authors would like to thank Dr. Guusje Bonnema for the experimental suggestions. Zhang Pujuan, Wang Baike, Wu Yuhui, Su Caixia and Fan Weiqiang are acknowledged for their kind help during the artificial inoculation.

Open Access This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

References

- Astua-Monge G, Minsavage GV, Stall RE, Vallejos CE, Davis MJ, Jones JB (2000) *Xv4-vrxv4*: a new gene-for-gene interaction identified between *Xanthomonas campestris* pv. *Vesicatoria* race T3 and the wild tomato relative *Lycopersicon pennellii*. *Mol Plant Microbe Interact* 13:1346–1355
- Brouwer DJ, St Clair DA (2004) Fine mapping of three quantitative trait loci for late blight resistance in tomato using near isogenic lines (NILs) and sub-NILs. *Theor Appl Genet* 108:628–638
- Brouwer DJ, Jones ES, St Clair DA (2004) QTL analysis of quantitative resistance to *Phytophthora infestans* (late blight) in tomato and comparisons with potato. *Genome* 47:475–492
- Brun H, Chevre AM, Fitt BDL, Powers S, Besnard AL, Ermel M, Huteau V, Marquer B, Eber F, Renard M, Andrivon D (2010) Quantitative resistance increases the durability of qualitative resistance to *Leptosphaeria maculans* in *Brassica napus*. *New Phytol* 185:285–299
- Canady MA, Meglic V, Chetelat RT (2005) A library of *Solanum lycopersicoides* introgression lines in cultivated tomato. *Genome* 48:685–697
- Caten C, Jinks J (1968) Variation spontaneous variability of single isolates of *Phytophthora infestans* I. cultural. *Can J Bot* 46:329
- Chaerani R, Smulders MJ, van der Linden CG, Vosman B, Stam P, Voorrips RE (2007) QTL identification for early blight resistance (*Alternaria solani*) in a *Solanum lycopersicum* x *S. arcanum* cross. *Theor Appl Genet* 114:439–450
- Chaïb J, Lecomte L, Buret M, Causse M (2006) Stability over genetic backgrounds, generations and years of quantitative trait locus (QTLs) for organoleptic quality in tomato. *Theor Appl Genet* 112:934–944
- Chen CH, Sheu ZM, Wang TC (2008) Host specificity and tomato-related race composition of *Phytophthora infestans* isolates in Taiwan during 2004 and 2005. *Plant Dis* 92:751–755
- Chunwongse J, Chunwongse C, Black L, Hanson P (2002) Molecular mapping of the *Ph-3* gene for late blight resistance in tomato. *J Hortic Sci Biotech* 77:281–286
- Drenth A, Janssen EM, Govers F (1995) Formation and survival of oospores of *Phytophthora infestans* under natural conditions. *Plant Pathol* 44:86–94
- Eshed Y, Zamir D (1994) A Genomic library of *Lycopersicon pennellii* in *Lycopersicon esculentum*—a tool for fine mapping of genes. *Euphytica* 79:175–179
- Eshed Y, Zamir D (1996) Less-than-additive epistatic interactions of quantitative trait loci in tomato. *Genetics* 143:1807–1817
- Eshed Y, Gera G, Zamir D (1996) A genome-wide search for wild-species alleles that increase horticultural yield of processing tomatoes. *Theor Appl Genet* 93:877–886
- Finkers R, van Heusden A, Meijer-Dekens F, van Kan J, Maris P, Lindhout P (2007) The construction of a *Solanum habrochaites* LYC4 introgression line population and the identification of QTLs for resistance to *Botrytis cinerea*. *Theor Appl Genet* 114:1071–1080
- Foolad MR, Merk HL, Ashrafi H (2008) Genetics, genomics and breeding of late blight and early blight resistance in tomato. *Crit Rev Plant Sci* 27:75–107
- Francis DM, Kabelka E, Bell J, Franchino B, St. Clair D (2001) Resistance to bacterial canker in tomato (*Lycopersicon hirsutum* LA407) and its progeny derived from crosses to *L. esculentum*. *Plant Dis* 85:1171–1176
- Goodwin SB, Smart CD, Sandrock RW, Deahl KL, Punja ZK, Fry WE (1998) Genetic change within populations of *Phytophthora infestans* in the United States and Canada during 1994 to 1996: role of migration and recombination. *Phytopathology* 88:939–949

- Gotoh K, Akino S, Maeda A, Kondo N, Naito S, Kato M, Ogoshi A (2005) Characterization of some Asian isolates of *Phytophthora infestans*. *Plant Pathol* 54:733–739
- Gur A, Zamir D (2004) Unused natural variation can lift yield barriers in plant breeding. *PLoS Biol* 2:e245
- Hanson P, Sitathani K, Sadashiva A, Yang R-y, Graham E, Ledesma D (2007) Performance of *Solanum habrochaites* LA1777 introgression line hybrids for marketable tomato fruit yield in Asia. *Euphytica* 158:167–178
- Jones J, Jones J, Stall R, Zitter TA (1991) Compendium of tomato disease. American Phytopathological Society, St. Paul, Minnesota
- Kole C, Ashrafi H, Lin G, Foolad M (2006) Identification and molecular mapping of a new R gene, *Ph-4*, conferring resistance to late blight in tomato. Solanaceae Conference, University of Wisconsin, Madison, Abstract 449
- Labate JA, Grandillo S, Fulton T, Munos S, Caicedo AL, Peralta I, Ji Y, Chetelat RT, Scott JW, Gonzalo MJ, Francis D, Yang W, van der Knaap E, Baldo AM, Smith-White B, Mueller LA, Prince JP, Blanchard NE, Storey DB, Stevens MR, Robbins MD, Wang JF, Liedl BE, O'Connell MA, Stommel JR, Aoki K, Iijima Y, Slade AJ, Hurst SR, Loeffler D, Steine MN, Vafeados D, McGuire C, Freeman C, Amen A, Goodstal J, Facciotti D, Van Eck J, Causse M (2007) Tomato. In: Labate CR et al (eds) Genome mapping, molecular breeding in plants, Vegetables, vol 5. Springer-Verlag, Berlin, Heidelberg, pp 1–125
- Lecomte L, Duffé P, Buret M, Servin B, Hospital F, Causse M (2004) Marker-assisted introgression of five QTLs controlling fruit quality traits into three tomato lines revealed interactions between QTLs and genetic backgrounds. *Theor Appl Genet* 109:658–668
- Monforte AJ, Tanksley SD (2000) Development of a set of near isogenic and backcross recombinant inbred lines containing most of the *Lycopersicon hirsutum* genome in a *L. esculentum* genetic background: a tool for gene mapping and gene discovery. *Genome* 43:803–813
- Monforte AJ, Friedman E, Zamir D, Tanksley SD (2001) Comparison of a set of allelic QTL-NILs for chromosome 4 of tomato: deductions about natural variation and implications for germplasm utilization. *Theor Appl Genet* 102:572–590
- Moreau P, Thoquet P, Olivier J, Laterrot H, Grimsley N (1998) Genetic mapping of *Ph-2*, a single locus controlling partial resistance to *Phytophthora infestans* in tomato. *Mol Plant Microbe Interact* 11:259–269
- Palloix A, Ayme V, Moury B (2009) Durability of plant major resistance genes to pathogens depends on the genetic background, experimental evidence and consequences for breeding strategies. *New Phytol* 183:190–199
- Peirce LC (1971) Linkage tests with Ph conditioning resistance to race 0, *Phytophthora infestans*. *Tomato Genet Coop Rep* 21:30
- Rousseaux M, Jones C, Adams D, Chetelat R, Bennett A, Powell A (2005) QTL analysis of fruit antioxidants in tomato using *Lycopersicon pennellii* introgression lines. *Theor Appl Genet* 111:1396–1408
- Smart CD, Tanksley SD, Mayton H, Fry WE (2007) Resistance to *Phytophthora infestans* in *Lycopersicon pennellii*. *Plant Dis* 91:1045–1049
- Stall RE, Jones JB, Minsavage GV (2009) Durability of resistance in tomato and pepper to *Xanthomonads* causing bacterial spot. *Annu Rev Phytopathol* 47:265–284
- Tieman DM, Zeigler M, Schmelz EA, Taylor MG, Bliss P, Kirst M, Klee HJ (2006) Identification of loci affecting flavour volatile emissions in tomato fruits. *J Exp Bot* 57: 887–896
- Turkensteen L (1973) Partial resistance of tomatoes against *Phytophthora infestans*, the late blight fungus. Dissertation, Wageningen University, The Netherlands
- Vleeshouwers VGAA, van Dooijeweert W, Paul Keizer LC, Sijpkens L, Govers F, Colon LT (1999) A laboratory assay for *Phytophthora infestans* resistance in various *Solanum* species reflects the field situation. *Eur J Plant Pathol* 105: 241–250
- Wastie R (1991) Resistance to powdery scab of seedling progenies of *Solanum tuberosum*. *Potato Res* 34:249–252
- Widmark A-K, Andersson B, Cassel-Lundhagen A, Yuen MSJE (2007) *Phytophthora infestans* in a single field in southwest Sweden early in spring: symptoms, spatial distribution and genotypic variation. *Plant Pathol* 56:573–579