

# Heavy metals in paddy fields in Taiwan: chemical behavior in soil and uptake by brown rice

C.L. Chu  
P. F.A.M. Römken  
H.Y. Guo



Alterra-rapport 1823, ISSN 1566-7197





Heavy metals in paddy fields in Taiwan: chemical behavior in soil and uptake by brown rice



# Heavy metals in paddy fields in Taiwan: chemical behavior in soil and uptake by brown rice

C.L. Chu†  
P.F.A.M. Römkens  
H.Y. Guo

Alterra-rapport 1823

Alterra, Wageningen, 2009

## ABSTRACT

Chu, C.L., P.F.A.M. Römken & H.Y. Guo, 2009. *Heavy metals in paddy fields in Taiwan: chemical behavior in soil and uptake by brown rice*. TARI – Taichung and Alterra, Wageningen, Alterra-rapport 1823. 112 blz.; 25 figs.; 41 tables; 119 refs.

Levels of cadmium (Cd), chromium (Cr), copper (Cu), nickel (Ni), lead (Pb) and zinc (Zn) were measured in 19 individual paddy fields in Taiwan. Total, reactive, and available metal levels were measured using Aqua Regia, 0.43 N HNO<sub>3</sub>, 0.1 M HCl, 0.05 M EDTA and 0.01 M CaCl<sub>2</sub>. Total metal levels ranged from below background levels to polluted and were highly heterogeneous across most fields. In general levels of metals in the soil decreased with an increase in distance from the water inlet which suggests that most metals originate from the irrigation water. Availability as measured by 0.01 M CaCl<sub>2</sub> could be predicted well (Cd, Ni, Zn) by a Freundlich model similar to the one used in non-tropical soils. The fit of models for Cu and Pb was poor due to the lack of data on dissolved organic carbon (DOC). For Cr no fit was obtained at all. Uptake of Cd by rice was highly correlated to the availability as measured by CaCl<sub>2</sub>. Uptake models based on either the CaCl<sub>2</sub> extractable Cd and Zn in soil solution, or a combination of the reactive Cd content in combination with pH and CEC proved equally suitable to predict Cd in rice. The impact of pH and, to a lesser extent, CEC urges the need to consider both properties when deriving soil quality standards (SQS). Uptake by rice by Indica species was markedly higher than that of Japonica although uptake by roots proved to be similar between both species. This suggests that differences between Japonica and Indica are more related to internal redistribution rather than differences in root uptake processes. Using the models, user friendly tools are designed allowing farmers and policy makers alike to evaluate the quality of the soil for a specific cultivar. This allows for a more accurate assessment of the suitability of the soil to be used for rice cropping compared to present soil standards based on Aqua Regia or HCl. As such the approach can be transferred easily to other countries as well based on a limited number of field tests.

Keywords: bioavailability, cadmium, food safety, paddy fields, risk assessment, soil pollution, Taiwan, rice

ISSN 1566-7197

The pdf file is free of charge and can be downloaded via the website [www.alterra.wur.nl](http://www.alterra.wur.nl) (go to Alterra reports). Alterra does not deliver printed versions of the Alterra reports. Printed versions can be ordered via the external distributor. For ordering have a look at [www.boomblad.nl/rapportenservice](http://www.boomblad.nl/rapportenservice).

© 2009 Alterra

P.O. Box 47; 6700 AA Wageningen; The Netherlands

Phone: + 31 317 484700; fax: +31 317 419000; e-mail: [info.alterra@wur.nl](mailto:info.alterra@wur.nl)

No part of this publication may be reproduced or published in any form or by any means, or stored in a database or retrieval system without the written permission of Alterra.

Alterra assumes no liability for any losses resulting from the use of the research results or recommendations in this report.

## Contents

Preface	7
Preface (continued)	9
Summary	13
1 Introduction	15
2 Description of sites and methods used in the field study	17
2.1 Location of sampling plots and cultivars used in the study	17
2.2 Soil pretreatment and chemical analyses	18
2.3 Plant pretreatment and chemical plant analyses	19
2.4 Conceptual framework to describe metal pools in soil according to their chemical availability.	19
2.5 Statistical analysis: soil – solution equilibrium	20
2.6 Statistical analysis: soil - plant equilibrium	21
3 Heavy metals in Paddy fields: overview of soil data	23
3.1 Soil properties of the fields included in the study	23
3.2 Heavy metal levels in soil	24
3.2.1 Total heavy metal content	27
3.2.2 Reactive metal pools	29
3.2.3 Actual available metal pools and solid-solution partitioning models	33
3.2.4 Effect of sampling time on the directly available metal pools and on solid-solution partitioning models	38
3.3 Mechanistic modeling of the availability of rice in soils	41
3.4 Conclusions	42
4 Uptake of metals by rice	45
4.1 General quality assessment of rice grown on study locations	45
4.2 Transfer of Cd from soil to roots and shoot-root ratios	48
4.3 Models to describe Cd uptake by rice: why?	49
4.4 Regression models to predict Cd levels in rice grain	50
4.5 Effect of time of harvest on the uptake of Cd by rice	53
5 Assessment of soil tests and soil-plant models to evaluate soil quality of paddy fields in relation to rice grain quality	57
5.1 Evaluation of the suitability of standards and models to test soils	57
5.2 Alternative soil standards based on soil tests with AR and HCl	60
5.3 Conclusions	61
6 Uncertainty of models to predict Cd uptake by rice	63
6.1 Introduction	63
6.2 Materials and Methods	64
6.3 Statistical approach to derive SQS based on soil to plant transfer	

models	65
6.4 Results	65
6.4.1 Derivation of soil plant transfer models for different extracts	65
6.4.2 Soil quality standards (SQS) for individual cultivars	68
6.4.3 Assessment of soil quality based on the P50, P90 or P95 percentile SQS	72
7 Tools to assist farmers and policy makers to evaluate soil quality for rice cropping	77
7.1 Introduction	77
7.2 Look-up tables	77
7.3 Soil specific evaluation	80
7.4 Issues to be considered when using average field data	82
8 Conclusions and Recommendations	83
References	85

### ***Appendices***

1 Overview of model coefficients used to calculate Cd in brown rice from soil data (soil model)	93
2 Look-up tables for individual cultivars	95
3 Overview of soil data per site	97
4 Overview of plant data	105
5 Theory behind the derivation of P50, P90 and P95 confidence levels of soil quality standards	111



## Preface

This report is dedicated to the memory of Chien-Liang Chu who tragically lost his life due to a fatal traffic accident on May 3<sup>rd</sup> 2009.

Without his help this study would not have been possible.



C.L. Chu



## Preface (continued)

Rice is an important food crop in Taiwan. Historically, a large part of the rice consumed in the country has been produced locally in Taiwan. However, due to industrial activities across Taiwan, soil heavy metal levels (especially cadmium -Cd- and lead -Pb-) have increased which results in an increasing number of samples that do not meet the food quality standard for rice from the EU/WHO<sup>1</sup> (0.2 mg kg<sup>-1</sup>) or even the Japanese/Taiwan standard of 0.4 mg kg<sup>-1</sup>. This increased uptake by rice is largely due to the increased level of metals in the soil.

In Taiwan atmospheric deposition (Pb) and the use of contaminated irrigation water (Cd) are important sources which have resulted in the observed increase of metal levels in soil.

This stresses the need for specific soil quality standards that can be used to identify soils where the quality is insufficient for being used for rice cropping. Present soil quality standards appear not to be protective enough since in many cases Cd levels in rice exceed food quality standards although the soil Cd level is below the standard of 5 mg kg<sup>-1</sup> which is used for arable soils in Taiwan.

Ideally, soil quality standards (SQS) for agriculture are based on the relationship between the quality of soil and that of the product grown on the soil. To establish such relationships, data on both soil and crop are needed as well as information on the availability of metals in the soil.

To address these issues a large field study has been performed at 19 sites. In total 12 varieties of rice have been grown on a range of soils which reflect the range in heavy metals currently present in soils in Taiwan. The reason for studying different varieties, is that uptake of Cd strongly depends on the variety. Varieties that accumulate Cd are, therefore not suitable for being used on Cd rich soils, while species that do not accumulated can be grown safely.

The aims of this study are to:

1. Establish the availability of metals in soil for plant uptake. This will be accomplished by measuring the metal content in different extracts including strong acid (Aqua Regia, AR), dilute hydrochloric or nitric acid (0.1 M HCl, 0.43 M HNO<sub>3</sub>), synthetic chelates (EDTA) and neutral salt solutions (0.01 M CaCl<sub>2</sub>). These data will be used to evaluate whether the availability of metals can be estimated using one or more of the extracts mentioned here. It also can be used to assess whether methods used in different countries are comparable.

---

<sup>1</sup> Recently the WHO/FAO food standard of 0.2 mg kg<sup>-1</sup> is raised to 0.4 mg kg<sup>-1</sup> but with reservations from representatives from the EU and other countries. However, in the remainder of this report we will often refer to the (old) WHO standard of 0.2 mg kg<sup>-1</sup>

2. Establish the relationship between the heavy metal content in the soil and the uptake by 12 different varieties of rice. Based on the results the varieties will be grouped relative to their suitability for being used on polluted soils.
3. Develop tools for end-users (farmers, policy makers) that enable them to evaluate the quality of local fields. At present existing soil quality guidelines are insufficient to make an accurate assessment of the soil quality and rice crops often do not meet the food quality standards used in Taiwan.

Ultimately the data will be used to construct models to predict uptake of Cd from soil by the cultivars tested here. These models will be used to derive more meaningful soil quality standards to be used across Taiwan. The approaches tested here are similar to the ones currently applied in the Netherlands. As such, the concept allows for the derivation of standards to be used on a national level (generic standards) as well as standards or soil testing values on a local scale.

Also, easy to use tools (so-called 'look-up' tables) will be developed to help farmers to decide which cultivars can be grown safely (if any) and if soil management (including lime application) can be used to improve soil conditions to such an extent that Cd levels in rice will remain below the food quality standard used in Taiwan.

The approach is designed such that the set-up of the fieldwork can be applied in other areas of Asia as well which yields model parameters for cultivars used in other countries. Ideally, the methodology can be used to derive a framework for soil policy in the light of food safety which has a common basis but allows for regional standards.

### ***How to read this report***

Chapter 2 contains information on the location of the sites included in the study, analytical procedures and the models developed to predict uptake of metals by rice. Chapter 3 contains all results of the soil and soil solution data. This includes results from all metals which were analyzed initially (Cu, Cr, Ni, Pb and Zn). Data on the total metal content in the fields as well as the reactive and available metal content are described. Models to predict the availability are discussed with an emphasis on Cd. Chapter 4 discusses the major findings regarding the levels of Cd in rice and models to predict the transfer of Cd from soil into the rice grain. Chapter 5 contains the results from an analysis on the accuracy of soil tests and models to predict whether or not a soil is suitable for rice cropping. Also alternatives for the standards based on Aqua Regia and HCl are discussed. In chapter 6 a discussion on the reliability of the models to predict Cd in rice is presented.

Chapters 3 to 6 contain all experimental results and reflect the scientific knowledge gained by this study. Aim of this study was to help policy makers and farmers alike make better decisions on the soil use. Therefore chapter 7 is dedicated to the practical application of the results. Here we will present applicable tools derived from the data and models to assist non-scientists in decision making on whether or not a soil can be used safely.

All soil, plant and model data are summarized in 4 appendices in the back of the report.

The results presented in chapters 3 to 7 reflect the contents of three scientific papers which have been published or submitted recently in the following papers as well:

Römkens, P.F.A.M., Guo, H.Y., Chu, C.L., Liu, T.S., Chiang, C.F., Koopmans, G.F. 2009. *Characterization of soil metal pools in paddy fields in Taiwan: chemical extraction and solid-solution partitioning*. J Soils Sediments (2009) 9:216–228.

Römkens, P.F.A.M., H.Y. Guo, C.L. Chu, T.S.Liu, C.F. Chiang, and G.F. Koopmans. 2009. *Prediction of Cadmium uptake by brown rice and derivation of soil-plant transfer models to improve soil protection guidelines*. Environmental Pollution 157 (2009) 2435–2444.

Römkens, P.F.A.M. D.J. Brus, G.F. Koopmans, H.Y. Guo, C.L. Chu, T.S.Liu, and C.F. Chiang. *Derivation of probabilistic local soil quality standards for paddy fields using soil to plant transfer models and 0.01 M CaCl<sub>2</sub> extracts*. Submitted for publication in Environmental Pollution.

### ***Acknowledgements***

The authors thank Dr T.S. Liu, Mr C.F. Chiang, Dr. D. Brus and Dr. G.F Koopmans for their contributions to this report. Also we thank EPA for the financial support to perform this study and financially supporting the field visits by P. Römkens to Taiwan. The financial means to complete this report have been supplied by the Dutch Ministry of Agriculture, Nature Conservation and Food Protection (LNV), BO Program ‘Internationaal’, BO-010-006-119.



## Summary

A large field study was conducted in 2005 – 2006 to measure the degree of contamination with heavy metals (HM) in paddy fields across the western plains of Taiwan. The second goal was to establish relationships between soil and the levels of HM in rice. The aim is to use such relationships to derive more meaningful soil protection guidelines for paddy fields and to construct easy to use tools for farmers and local policy makers to assess the suitability of paddy fields for rice cropping. Below the main findings are summarized, first results from the soil inventory are presented followed by results on rice and the relationship between soil and rice.

### ***Summary for major findings on HM in soils:***

- HM levels in paddy fields are highly variable within a single field and between fields;
- The in-field variation is related to the distance of the irrigation water inlet and levels of most HM decrease with an increasing distance from the inlet;
- The between-field variation is related to the vicinity of different types of industry and even within one community clear differences between the source of the HM can be found based on the ratio of HM in the soil;
- Total HM levels extracted by Aqua Regia exceed current limits for arable soils;
- The amount of HM extracted by dilute acids (0.43 N HNO<sub>3</sub> or 0.1 M HCl) or EDTA are comparable, difference between HNO<sub>3</sub> and HCl are small and both acids can be used to obtain an estimate of the reactive Cd content of the soil;
- The ratio of reactive (as determined by dilute acids) and total HM content is comparable to those found in soils from the EU;
- The availability of HM as measured by 0.01 M CaCl<sub>2</sub> can be modeled quite well (except for Cr) using the reactive metal content, soil pH and CEC.
- Only for Cu and Pb the models seem to be less accurate probably because no information on the levels of Dissolved Organic Matter (DOM) is available.
- The models to describe the 0.01 M CaCl<sub>2</sub> extractable metal content in paddy soils are comparable to those derived for soils from Belgium and the Netherlands. This suggests that the chemical behavior of metals in paddy fields during the drained periods is comparable to that in well-drained soils from moderate climates.

### ***Summary of major findings on HM in rice with emphasis on Cd:***

- Uptake of Cd by rice strongly depends on the cultivar: levels of Cd in Indica type cultivars exceed that of Japonica type species;
- Even at low soil Cd levels (below 0.5 mg kg<sup>-1</sup>) levels of Cd in Indica rice species exceed the food quality standard of 0.4 mg kg<sup>-1</sup>;
- Levels of Cd in brown rice are highly correlated to those in the roots;
- The amount of Cd in roots can be explained largely by the variation in the amount of Cd and Zn in the 0.01 M CaCl<sub>2</sub> extract;
- For the species tested in this study Zn reduces the sorption of Cd to the root leading to lower levels of Cd in the rice grain;

- The significant relation between Cd (and Zn) in the 0.01 M CaCl<sub>2</sub> extract and levels in roots and rice grain, suggests that the total amount taken up by the rice plant largely depends on the chemical availability of Cd in the soil which can be predicted well by the Cd content of the soil, pH and CEC.

***Summary of major findings on soil quality guidelines and tools developed in this study***

- Current soil quality guidelines used in Taiwan (5 mg kg<sup>-1</sup> for Aqua Regia or 2 mg kg<sup>-1</sup> for 0.1 M HCl) are too high to safeguard the quality of rice; a large number of soil samples with Cd below the (soil) standard have high (> food quality standard) levels of Cd in rice;
- The use of pH and CEC when evaluating the suitability of soils for the production of rice leads to accurate assessments of the quality of the rice;
- For soils with a pH of more than 6, fixed standards (1 or 2 mg kg<sup>-1</sup> based on HCl) can be used for Japonica but these are not suitable for Indica;
- Alternatively, soil tests based on a single extract with 0.01 M CaCl<sub>2</sub> are also suitable to evaluate the quality of the soil for rice cropping;
- The models developed and calibrated in this field study can be used to construct look-up tables which show the suitability of a specific soil for individual cultivars depending on the level of Cd in soil and pH
- Differences in uptake of Cd by various cultivars can be used as a tool by farmers to select those cultivars that can be grown safely on a specific soil.
- To obtain an accurate prediction of the suitability of a specific paddy field as whole for rice cropping, careful sampling schemes are essential due to the large observed variation in both Cd in soil and in the crop. Despite the fact that the models are good compared to previously published models from the literature, the uncertainty is still too large to obtain accurate predictions for individual samples.

***Recommendations for policy regarding Cd in paddy fields***

This study clearly reveals that the levels of Cd in rice exceed current food quality standards even though in many cases the soil Cd content is below the soil quality standard. This indicates that the present soil standard of 5 mg kg<sup>-1</sup> is too high. For Japonica which is less sensitive regarding uptake of Cd from soil, a standard of 2 mg kg<sup>-1</sup> in clayey soils with a pH of 6 or higher seems protective enough. This is however true only for the food quality standard of 0.4 mg kg<sup>-1</sup> which is currently being used in Taiwan. If the food quality standard is reduced to 0.2 mg kg<sup>-1</sup> (current level used in the EU), the allowed Cd content in the soil obviously will be lower as well.

For soils with a lower pH or lower clay content, as well as for soils used for Indica species, fixed soil standards are not suitable since they do not account for the strong impact of pH on the level of Cd in grains of brown rice. For those soils or rice varieties, the pH and CEC have to be considered. The use of crop specific look-up tables can be of help to assess whether or not the combination of the pH and the Cd content leads to Cd in rice below or above the food quality standard.



## 1 Introduction

Soil pollution in industrialized countries is of growing concern in relation to human health. Emission by industry (atmospheric, waste water and solid waste) as well as agriculture (manure disposal, hoof disinfection solutions, and medicinal waste material) has resulted in an increase in the content of many contaminants in soil. Especially metals like Cd pose a significant threat since this metal not only is toxic for animals and plants alike, it can also be taken up by various important food crops such as rice and wheat. To protect human health food quality standards are in place, which for rice vary between  $0.2 \text{ mg kg}^{-1}$  (EU standard as well as former WHO/FAO standard) and  $0.4 \text{ mg kg}^{-1}$  (Japanese/Taiwan standard and revised WHO standard).

Due to soil pollution, soil Cd levels have increased all over the world which can result in an increase in uptake of Cd by arable crops as well. Especially in areas with high atmospheric deposition from industry (especially non-Ferro industry like Zn smelters) levels in soil have reached excessive levels. Usually Cd background levels in soil are low (less than  $0.5 \text{ mg kg}^{-1}$  depending on the parent material) but all over the world levels of more than  $500 \text{ mg kg}^{-1}$  have been reported although these levels are confined to hot spots. Estimates of the background levels for Cd in Taiwan are as low as  $0.16 \text{ mg kg}^{-1}$  (Yang and Chang, 2005) and soils with Cd levels higher than  $0.64 \text{ mg kg}^{-1}$  can be considered as polluted (Yang and Chang, 2005). However, levels between 1 and  $20 \text{ mg kg}^{-1}$  have become more and more common near industrial areas. At these levels, uptake of Cd by rice can result in Cd grain levels beyond the food quality standard.

Major sources of heavy metals in Taiwan include direct deposition from the atmosphere near industrial zones, but more important is the use of polluted irrigation water. Untreated waste water from cities and industries has been used for decades to irrigate rice fields. A large part of the metals in the water is bound to some form of organic colloidal material which, after being led onto the field, deposits at the soil. Due to the limited solubility, metals are retained in the topsoil and can be taken up by plants depending on the geochemical availability and the plants selected.

In many countries including Taiwan, soil standards have been developed to protect the ecosystem and human health. One of the problems with soil quality standards is that they do not, or only to a limited extent address the specific protection of the quality of arable products. In many cases these soil protection levels are related to direct effects on human health and not so much related to the quality of arable crops. Hence, to assess whether or not a soil is suitable for the production of specific crops like rice it is necessary to study the relationship between the soil and the crop of interest. Based on the results from such studies (which until now are rather scarce) specific target levels for agricultural soils can be developed.

At present soil quality standards are in place. Soil Cd levels of more than  $5 \text{ mg kg}^{-1}$  (farmland, other soil use  $20 \text{ mg kg}^{-1}$ ) are considered dangerous and have to be reported (EPA, 2006). Aside from the standard based on the total amount in soils

(Aqua Regia, AR) also a proposal for a standard of  $2 \text{ mg kg}^{-1}$  based on  $0.1 \text{ M HCl}$  was put forward (Chen, 2000). It has become clear however, that the measurement of the total amount in soil is not always relevant since some part of the heavy metal content is not available for plant uptake. The use of dilute acids to measure the fraction of metals which can be taken up by plants instead of the total metal content is an option to correct for this. However, differences between plant species or varieties cause a large range in uptake by crops from similar soils.

To derive meaningful standards for metals in soils, therefore, it is imperative to study the uptake in a wide range of soil including different varieties of crops. Measurements of soil properties such as pH and organic matter in addition to the metal content in the soil is important as well since differences in uptake are often related to differences in soil type. For several crops it has been documented that Cd uptake from a sandy soil with a low organic matter content at pH 4.5 exceeds that from soils at pH levels above 5 or 6.

The aim of this study was to include those variables that affect the level of Cd in the crop which includes both differences in soil type and differences in uptake (species). Aside from this, different extracts are tested to evaluate which ones are most suitable to predict Cd levels in crops. Here, 3 extracts are tested (apart from the total metal content measured by Aqua Regia) that are commonly used to measure the availability of metals in soil:  $0.1 \text{ M HCl}$  (currently used in Taiwan),  $0.43 \text{ N HNO}_3$  (currently used in the Netherlands) en  $0.05 \text{ M EDTA}$  which is often used in plant uptake studies. Apart from these strong extracts also a salt solution ( $0.01 \text{ M CaCl}_2$ ) is used to determine the amount of Cd that is believed to be available for plant uptake in a specific year.

Apart from soil data, levels of Cd in various parts of rice plants (root, stem, leaf, husk and rice grain) are measured to assess the relation between soil quality and uptake by the crop. The data will allow for the development of models that predict both the chemical availability of Cd in paddy soils as well as models that predict the uptake of Cd by rice. In this report we focus on the quality of the rice grains since this is the plant part which is used for human consumption.

Ultimately the results of this study will be used to derive soil quality standards that are protective enough to ensure the farmer that the rice grown on the soil indeed meets the food quality standard. A different approach to avoid excess uptake of Cd by rice is to select those rice cultivars which do not accumulate Cd. Especially in moderately polluted soils, crop (or cultivar) selection seems a good way to be able to use the soil in a safe way. Both approaches (soil tests and crop selection) will be discussed here.

## 2 Description of sites and methods used in the field study

### 2.1 Location of sampling plots and cultivars used in the study

In 2005 and 2006, soil samples were taken from 19 different paddy fields across the western plains in Taiwan. The fields were located in the towns of Chang Hua (CH, 3 fields), Ho Mei (HM, 3 fields), Lu Kang (LK, 2 fields), Hsin Chu (HC, 3 fields) and Pa-Deh (PD, 8 fields). In figure 2.1, an overview of the communities where the fields are located is shown. These plots were selected based on the known history of soil pollution and include both sites known to be heavily polluted (e.g. the PD fields) but also relatively clean sites with low metal levels in soil (such as the LK sites).




Figure 2.1 Map of Taiwan and the location of the communities where the paddy fields used in this study are located (PD: Pa-Deh city, Tao-Yuan county; HC: Hsin-Chu city, Hsin-Chu county; CH: Chang-Hua city; HM: Ho-Mei city, LK: Lu-Kang city; the latter three communities are all located in Chang-Hua county)

Soils in these parts of Taiwan are mostly developed in unconsolidated river sediments in flat to undulating areas. All paddy fields were used for rice cropping prior to the fieldwork done described in this report. At each field, 12 cultivars of Indica and Japonica rice plant species were planted on plots with 5 to 9 replicates for each cultivar (depending on the size of the field, schematically shown in figure 2.2) to account for the high spatial variability of the metal levels in soil across the paddy field. In table 2.1 an overview of the cultivars used is shown.

Table 2.1 List of rice cultivars and number of samples for each cultivar.

Cultivar	Cultivar name	Family	Number of samples
1	Tainung no.70	Japonica	277
2	Taiken no.8	Japonica	278
3	Tainung no.72	Japonica	278
4	Kaohsiung no.143	Japonica	278
5	Taitung no.30	Japonica	277
6	Tainung Sen no.20	Indica	277
7	Tainung no.71	Japonica	278
8	Tainung no.67	Japonica	278
9	Kaohsiung Sen Yu no.1151	Indica	210
10 (A)	Taichung Sen Waxy no.1	Indica	278
11 (B)	Taichung Sen no.10	Indica	278
12 (C)	Kaohsiung no.144	Japonica	211

The total number of plots per field thus ranged from 60 (i.e., 5 replicates times 12 cultivars) to 108 (i.e., 9 replicates times 12 cultivars). Since the fields are different in size and shape, the size of the individual plots ranged from 1 to 2 m (width) by 9 to 11 m (length) depending on the total size of the field. Each plot was sampled twice during the year in May and November. In 2005, the CH, LK and HM fields were sampled while the HC and PD fields were sampled in 2006, which results in a database containing 3265 individual sampling data. Soil samples were taken from the topsoil (0–25 cm) from each plot using a hand auger at time of harvest when the paddy field was drained. Within each plot, up to 25 soil samples were taken at random across the plot and mixed manually resulting in a mixed sample of approximately 3 to 4 kg of field-moist soil.




Figure 2.2 Schematical lay-out of the fields included in the 2005 and 2006 fieldwork. Note that not all fields include 9 sections (ranging from 5 to 9 depending on the size of the field).

## 2.2 Soil pretreatment and chemical analyses

After collection of field-moist samples, soil samples were air-dried in air-conditioned rooms with a dehumidifier during 96 hr at 25 to 30°C. Dried soil samples were passed through a 2-mm sieve. After drying, the remaining moisture content was measured at 105°C to calculate the dry matter content. Total metal pools in air-dried

samples were determined by extraction with AR using a microwave digestion device (Lamothe et al., 1986). After destruction, extracts were filtered through a filter with a pore size of 2.5  $\mu\text{m}$  (Whatman no.42), and analyzed for Cd, Cr, Cu, Ni, Pb, and Zn by an inductively coupled plasma-atomic emission spectrometer (ICP-AES). The reactive metal pool was measured by three reagents: (i) 0.43 M  $\text{HNO}_3$  (Houba et al., 1997), (ii) 0.1 M  $\text{HCl}$  (Nelson et al., 1959), and (iii) 0.05 M EDTA in the Na form ( $\text{Na}_2\text{-EDTA}\cdot 2\text{H}_2\text{O}$ ; Tiwari and Kumar 1982). All extractions were performed at a soil to solution ratio of 1:10 (w:v) for 1 hr. After shaking, all extracts were filtered (Whatman no.42), and metal concentrations were measured by ICP-AES. The directly available metal pool was measured by extraction with 0.01 M  $\text{CaCl}_2$  according to Houba et al. (2000). After shaking for 2 hr on a table-top shaker, extracts were filtered (Whatman No.42 filters), and metal concentrations were measured by ICP-AES. The pH was measured directly in  $\text{CaCl}_2$  extracts prior to filtration. SOM was determined by loss-on-ignition after heating at 400°C for 8 hr (US-EPA, 2002). CEC was determined using 1 M  $\text{NH}_4\text{OAc}$  buffered at pH 7.0 (USDA 1996). Clay content was measured by laser spectroscopy (Xu 2000). SOM, pH and CEC were analyzed in all soil samples but clay content was only measured on a limited number of samples from each field.

### **2.3 Plant pretreatment and chemical plant analyses**

Root tissues were washed with tap water and subsequently rinsed with de-ionized water. Large root and plant (stem, leaf) tissue samples were shortened using a ceramic knife. Root, stem, leaves, husk and unpolished rice samples were dried at 70°C for about 96 hr and ground to powder using a titanium-coated blade mill. Dried plant (except rice) material was digested with an acid mixture containing concentrated  $\text{HNO}_3$  and  $\text{HClO}_4$  mixed at a ratio of 5:1 (v:v) at 140°C for 2 hr followed by a second step at 170°C for 1 hr. After filtration (Whatman no. 42), Cd concentration in plant digests were measured by ICP-AES. Chemical analysis of rice grains was done on unpolished or brown rice grain samples. Dried rice grain material was digested using concentrated  $\text{HNO}_3$  at room temperature overnight. Subsequently,  $\text{H}_2\text{O}_2$  was added followed by a second digestion in a microwave oven at 180°C for 20 min. After filtration (Whatman no. 42), Cd concentrations in the extracts were determined by ICP-MS. Quality of the crop and soil analyses was monitored by including blanks and extracts obtained from soil or plant reference materials in each batch.

### **2.4 Conceptual framework to describe metal pools in soil according to their chemical availability.**

The concept of availability of metals in soils has gained considerable attention during the last few decades. It is clear that not all metals (and organic contaminants alike) are equally available for uptake by plants or leaching to ground- and surface waters. To improve risk assessment approaches it is therefore imperative to find methods that are able to more specifically measure the available fractions in soil. In this report

we want to assess to what extent the behavior of heavy metals in soils can be characterized according to the scheme illustrated in figure 2.3. Ultimately the goal is to link plant uptake to the pools characterized by the methods as mentioned in figure 2.3




Figure 2.3 methods used in this report to measure different fractions of metals in soil ranging from inert (not available) to directly available for plant uptake.

Please note that the size of the individual pools is not according to scale and may differ depending on the metal of interest. A rather fast (hours to days) equilibrium is assumed to exist between the reactive metal pool and the directly available pool. The release of metals from the inert fraction on the other hand is very slow (years to decades) and is mainly due to weathering of soil minerals. Total metal content can be measured by extraction of soil with Aqua Regia (AR) or hydrofluoric acid (HF) or by X-ray fluorescence (XRF). Reactive metal pool can be determined by extraction of soil with 0.43 M HNO<sub>3</sub>, 0.1 M HCl or 0.05 M EDTA, whereas the directly available metal pool can be determined by extraction with 0.01 M CaCl<sub>2</sub>. The methods actually used in the study are included because they are either commonly used in current legislation (like Aqua Regia for the total metal content and 0.1 M HCl for the reactive part) or because it has been shown experimentally that the methods appear suitable (like 0.01 M CaCl<sub>2</sub> for the directly available pool and 0.43 M HNO<sub>3</sub> for the reactive fraction).

## 2.5 Statistical analysis: soil – solution equilibrium

Multiple linear regression analysis was performed to derive the solid-solution partition models of metals. All data were log-transformed (except for pH) prior to the regression. An extended Freundlich equation was used to relate the directly available metal pool to the reactive metal pool in combination with soil properties like pH and CEC. This approach has been used previously by several authors and was able to explain a large part of the observed variability in the solubility of metals like Cd, Zn, Cu and Pb (Römken et al., 2004; Ingwersen and Streck 2006; Meers et al., 2007; Koopmans et al., 2008a).

$$\log[\text{Me-CaCl}_2] = \text{intercept} + a \cdot \text{pH} + b \cdot \log[\text{CEC}] + c \cdot \log[\text{Me-reactive pool}] \quad [2.1]$$

With	
[Me-CaCl <sub>2</sub> ]	directly available metal pool extracted with 0.01 M CaCl <sub>2</sub> in mg kg <sup>-1</sup> ;
pH	pH in the 0.01 M CaCl <sub>2</sub> extracts;
[CEC]	cation exchange capacity in cmol(+) kg <sup>-1</sup> ;
[Me-reactive pool]	reactive soil metal content determined by extraction with either HCl, HNO <sub>3</sub> or EDTA in mg kg <sup>-1</sup> .

SOM was not included in Eq. 2.1, because its contribution to the regression in addition to CEC appeared to be not significant. The CEC alone is able to represent the exchange capacity of the soil and as such includes the contribution from different soil components such as SOM, clay and amorphous Al and Fe oxides.

All regression analyses and further statistical tests were carried out using Genstat (version 11.1)

## 2.6 Statistical analysis: soil - plant equilibrium

Accumulation of Cd in rice grains can be related to the available Cd and Zn pool in soil (0.01M CaCl<sub>2</sub>):

$$^{10}\log[\text{Cd-rice grains}] = \text{intercept} + a \cdot ^{10}\log[\text{Cd-CaCl}_2] + b \cdot ^{10}\log[\text{Zn-CaCl}_2] \quad [2.2]$$

Combining eqs. [2.1] and [2.2] results in a soil-plant transfer function (McGrath et al., 2000; Efrogmson et al., 2001; Krauss et al., 2002; Brus et al., 2005; Japenga et al., 2007):

$$^{10}\log[\text{Cd-grain}] = \text{intercept} + f \cdot ^{10}\log[\text{Cd-HNO}_3] + g \cdot \text{pH} + h \cdot ^{10}\log[\text{CEC}] \quad [2.3]$$

In this report, we use eqs. [2.2] and [2.3] to predict Cd levels in rice grains. We use the terms 'CaCl<sub>2</sub>-model' or 'soil solution model' for eq. [2.2] and 'HNO<sub>3</sub>-model' or 'soil to plant relationship' for eq. [2.3]. Relationships were derived for each cultivar separately and for the combined data for Japonica or Indica cultivars.

All regression analyses and further statistical tests were carried out using Genstat (version 11.1)





### 3 Heavy metals in Paddy fields: overview of soil data

In this chapter, data on the level of soil pollution will be presented. Aside from Cd also other metals including Cu, Pb, Zn, Ni and Cr were measured and will be included here. The remainder of the report will focus on Cd since this element is of prime importance in relation to food safety.

#### 3.1 Soil properties of the fields included in the study

In table 3.1 an overview of the soil properties measured in the study is shown. The different fields from each location (e.g. CH, HC etc) are numbered. Hence, data in each row correspond to an individual field. On average the PD fields (PDC through PDH) are smaller than the CH, HM, and LK fields which results in a smaller number of samples per field.

*Table 3.1 Soil properties (mean and standard deviation) from the fields included in this study (average values from all individual plots within a field; data from all harvests combined)*

Field	n	Clay (%)		SOM (%)		pH		CEC (cmol <sup>[+]</sup> kg <sup>-1</sup> )	
		Mean	s.d.	Mean	s.d.	Mean	s.d.	Mean	s.d.
CH6	216	70.7	5.0	6.7	0.5	4.7	0.4	16.9	1.5
CH7	215	59.1	9.1	4.6	0.5	6.1	0.5	11.1	1.4
CH8	217	75.0	3.2	6.3	0.5	6.3	0.3	19.0	2.1
HC3	138	15.7	4.7 <sup>1</sup>	2.7	0.4	5.4	0.2	5.5	1.1
HC4	115	15.7	4.7 <sup>1</sup>	4.5	0.5	5.6	0.2	8.7	1.5
HC5	184	15.7	4.7 <sup>1</sup>	3.5	0.4	5.1	0.3	6.9	1.3
HM3	215	70.1	3.0	7.5	0.6	6.2	0.4	19.1	1.8
HM4	215	69.7	2.8	6.2	0.6	5.9	0.4	16.2	1.6
HM5	215	66.6	3.0	5.9	0.5	5.3	0.3	15.1	1.4
LK1	216	3.8	0.4	2.7	0.7	5.3	0.4	5.9	1.9
LK2	216	51.9	9.3	4.4	0.5	5.8	0.3	10.6	1.5
PDA	229	33.9	1.6 <sup>1</sup>	7.1	0.4	5.3	0.2	13.1	1.5
PDB	230	33.9	1.6 <sup>1</sup>	7.5	0.4	5.2	0.2	14.4	1.9
PDC	92	33.9	1.6 <sup>1</sup>	6.8	0.4	5.2	0.2	13.3	1.7
PDD	92	33.9	1.6 <sup>1</sup>	7.8	0.4	5.0	0.2	14.6	1.3
PDE	115	33.9	1.6 <sup>1</sup>	7.5	0.6	5.1	0.2	14.1	1.5
PDF	92	33.9	1.6 <sup>1</sup>	7.8	0.4	5.2	0.2	15.2	1.7
PDG	92	33.9	1.6 <sup>1</sup>	7.7	0.4	5.1	0.1	14.9	1.7
PDH	161	33.9	1.6 <sup>1</sup>	8.7	0.5	5.0	0.1	16.4	1.9

<sup>1</sup>The clay content in the HC and PD fields was measured in 69 (PD) and 12 (HC) individual samples taken from the PDA field through the PDH field and from the HC3 field through the HC5 field respectively, but no distinction between fields was made.

The soils included in the study possess a rather wide range in texture, ranging from sandy soils (LK1) to heavy clay soils (HM plots and CH plots). The pH varies widely from acid (CH6) to near neutral (CH8), although most of the soils in this study tend to be slightly acid with pH values ranging between 5 and 6 with 25% of the samples having a pH value below 5.1. SOM levels are fairly comparable and range from less than 3% in the LK1 and HC3 plot to almost 9% in the PDH plot. CEC values can be classified broadly into two groups, soils with a rather low CEC around 5 cmol(+) kg<sup>-1</sup> (HC and LK1 plots) and soils with CEC values around 15 cmol(+) kg<sup>-1</sup>. Differences in CEC were closely linked those in the SOM content as is shown in figure 3.1 which reflects the impact of SOM on the capacity of the soil to bind cations.




Figure 3.1. Relation between organic matter (median value per site) and CEC (median value per site)

Soil pH had little effect on CEC which is not surprising since CEC was measured at pH 7 in all soils using a buffered NH<sub>4</sub>OAc solution. The range in soil properties within each field is limited as is indicated by the low standard deviation. As such, the ranges found for these soil properties are comparable to those measured by others within paddy rice fields (e.g., He et al., 2006; Simmons et al., 2008) and reflect the ranges in soil properties commonly observed.

### 3.2 Heavy metal levels in soil

An overview of the amount of metals present in the soils studied is shown in table 3.2 and graphically in figure 3.2.

Table 3.2 Overview of the total, reactive and directly available metal pools in all 19 fields as measured by extraction with Aqua Regia (AR; total metal content), 0.1 M HCl, 0.43 M HNO<sub>3</sub> or 0.05 M EDTA (reactive metal pool), and 0.01 M CaCl<sub>2</sub> (directly available pool)

Metal	Extract	n	Percentiles of HM content in various extracts (mg kg <sup>-1</sup> )								Soil quality standard <sup>1</sup> (mg kg <sup>-1</sup> )
			Min.	5%	25%	50%	Mean <sup>2</sup>	75%	95%	Max.	
Cd	AR	3265	0.06	0.21	0.39	0.64	3.8	5.3	14.9	29.4	5.0
	HCl	3250	0.01	0.11	0.22	0.41	3.0	4.3	12.1	25.7	
	HNO <sub>3</sub>	3266	0.08	0.13	0.22	0.41	3.1	4.6	12.3	20.6	
	EDTA	3265	0.05	0.11	0.21	0.39	2.3	3.6	9.1	16.4	
	CaCl <sub>2</sub>	3255	0.001	0.004	0.01	0.04	0.29	0.46	1.3	3.2	
Cr	AR	3265	38.9	47.2	54.8	80.0	94.4	112.4	203.9	483.0	250
	HCl	2560	0.01	0.13	0.89	2.1	2.9	3.5	9.1	31.6	
	HNO <sub>3</sub>	3265	0.93	1.8	2.6	4.9	7.2	8.3	22.2	82.4	
	EDTA	2072	0.004	0.1	0.63	1.1	1.5	1.9	4.0	9.2	
	CaCl <sub>2</sub>	2567	< 0.001	0.002	0.01	0.02	0.02	0.02	0.04	0.2	
Cu	AR	3265	22.9	29.9	55.0	101.5	123.9	150.4	312.1	903.5	200
	HCl	3265	6.7	11.3	22.6	50.9	64.7	73.3	180.9	640.3	
	HNO <sub>3</sub>	3265	9.8	14.5	27.8	60.3	74.4	86.5	199.8	635.8	
	EDTA	3265	5.0	12.9	24.0	56.4	70.5	86.5	193.2	539.3	
	CaCl <sub>2</sub>	2776	< 0.001	0.008	0.03	0.07	0.18	0.18	0.73	9.2	

<sup>1</sup>Soil monitoring value for cropped soils used at present in Taiwan (EPA 2006).

<sup>2</sup>All elements have a log-normal distribution, which results in a large difference between the median and the average value.

Table 3.2 Continued

Metal	Extract	n	Percentiles of HM content in various extracts (mg kg <sup>-1</sup> )								Soil quality standard <sup>1</sup> (mg kg <sup>-1</sup> )
			Min.	5%	25%	50%	Mean <sup>2</sup>	75%	95%	Max.	
Ni	AR	3265	16.8	26.1	32.5	72.9	97.9	147.1	245.6	461.4	200
	HCl	3265	0.83	2.1	3.9	23.9	23.9	39.5	59.4	101.7	
	HNO <sub>3</sub>	3265	1.7	2.7	4.9	27.3	26.8	44.7	64.8	102.2	
	EDTA	3265	0.80	2.1	4.1	22.7	23.9	40.1	60.3	89.1	
	CaCl <sub>2</sub>	3185	0.002	0.082	0.257	0.954	3.1	4.8	12.0	29.4	
Pb	AR	3265	10.8	21.2	30.9	48.8	164.4	194.1	665.5	1171.4	500
	HCl	3262	0.30	4.7	7.8	13.6	84.6	83.7	389.1	821.7	
	HNO <sub>3</sub>	3265	6.7	12.0	17.5	26.2	110.4	127.3	476.7	755.6	
	EDTA	3265	2.0	7.8	11.9	19.8	92.3	109.0	400.7	624.9	
	CaCl <sub>2</sub>	2197	< 0.001	0.01	0.06	0.15	0.20	0.309	0.543	1.05	
Zn	AR	3265	63.1	99.1	197.1	277.2	311.7	386.3	666.5	1225.3	600
	HCl	3265	11.9	22.8	41.7	60.8	77.7	97.0	179.5	537.3	
	HNO <sub>3</sub>	3265	17.3	25.7	49.8	70.8	90.4	111.8	209.5	574.2	
	EDTA	3265	10.7	20.1	36.0	54.5	68.6	86.4	160.6	383.6	
	CaCl <sub>2</sub>	3244	0.009	0.13	1.9	5.5	8.0	10.5	25.0	96.9	

<sup>1</sup>Soil monitoring value for cropped soils used at present in Taiwan (EPA 2006).

<sup>2</sup>All elements have a log-normal distribution, which results in a large difference between the median and the average value.




Figure 3.2 Heavy metal contents as measured by extraction with Aqua Regia in the 19 paddy fields used in this study. The boxes represent the 25 and 75 percentile levels whereas the solid Y-bars represent the minimum and maximum level found within each field. The dark bar within the box is equal to the median value (50 percentile). The dashed red lines represent the soil monitoring value for cropped land in Taiwan (EPA 2006).

### 3.2.1 Total heavy metal content

Total metal levels vary considerably between and within fields, and they range from background levels to levels beyond current soil quality standards used in Taiwan (EPA 2006; table 3.2). For the metals studied here, the number of soil samples with total metal contents in excess of the present soil monitoring values for cropped soils increases from 2.9% for Cr, 6.3% for Zn, 10.1% for Ni, 13.0% for Pb, 13.3% for Cu to 26.7% for Cd. Not all fields are equally polluted as is shown in figure 3.2. Total Cd and Pb contents are especially high in the PD fields whereas total Ni and Zn contents in the HM and LK fields exceed those of the other locations. Figure 3.2 not only illustrates the broad range in the total metal contents between fields but it also shows the broad range observed within each field. Within the HC3 field, for example, the total Cd content in soil ranges from 0.19 mg kg<sup>-1</sup> to 6.8 mg kg<sup>-1</sup>. Likewise, the total Cd contents in the PDD and PDB fields, which range from 4.0 to 21.6 mg kg<sup>-1</sup> and from 6.2 to 29.4 mg kg<sup>-1</sup>, respectively, are highly variable. Even within 50 m from the inlet, total Cd content decreases up to a factor of 35 (HC3 plot). This is illustrated as well in figure 3.3 which shows the maps of HCl extractable

Cd in the fields HC3 and HM3. Highest Cd levels are found close to the inlet of the irrigation water and Cd levels gradually decrease across the field. Similar trends in metal levels in soil within paddy fields have been observed by Simmons et al. (2005). Hence, the distance within a paddy field to the main inlet of irrigation water is crucial in relation to the metal levels observed in soil.




Figure 3.3 Spatial distribution of Cd as measured by extraction with 0.1 M HCl in the soil across field HM3 (left) and HC3 (right). The inlets for irrigation water are marked with a blue arrow. The scale is different in both fields (units in mg kg<sup>-1</sup>). Spatial distribution patterns were obtained by kriging of the individual data points from each field.

In contrast to the HM3 and HC3 fields, total Cd contents in the PDF, PDG, LK1, and CH6 fields contain much less variation (figure 3.2). In the CH, HM, and LK fields on the other hand, total Cu contents are highly variable, reflecting the apparent specific Cu emission of a nearby industry. In table 3.3, a correlation matrix between total metal contents is presented for all PD fields together. In these fields, Cd, Pb, Cu and Zn are likely to originate from the same source, because of the close correlation found between the total contents of these metals.

Table 3.3 Cross-correlation matrix of metals in soil in the Pab-Deb fields based on total metal contents as extracted with Aqua Regia

	Cd	Cr	Cu	Ni	Pb	Zn
Cd	1.000	-	-	-	-	-
Cr	0.1759	1.000	-	-	-	-
Cu	0.6432	0.1721	1.000	-	-	-
Ni	0.1445	0.5240	0.1223	1.000	-	-
Pb	0.9109	0.1218	0.8115	0.1051	1.000	-
Zn	0.8754	0.2179	0.9105	0.1578	0.9432	1.000

In the CH fields, however, Cu and Ni seem to be closely related. Indeed, a close correlation for Cu and Ni was found between the total contents of these metals within the CH fields, but it is different for the three individual fields within the Chang-Hua community (figure 3.4). Apparently, the polluted irrigation water entering the fields originates from different sources with Cu:Ni ratios decreasing in the order CH8 > CH7 > CH6. Hence, soil monitoring programs have to consider the heterogeneous nature of metal distribution between and within paddy fields in order to be able to accurately assess the biological risks related to metals in soil in industrialized areas like in Taiwan.




Figure 3.4 Relationship between total Cu and Ni contents as measured by extraction with Aqua Regia in soil samples from the three Chang-Hua fields. The dotted lines represent a best-fit line.

### 3.2.2 Reactive metal pools

Amounts of metals extracted by HNO<sub>3</sub>, HCl, and EDTA are markedly smaller than those extracted by AR (table 3.2). Figure 3.5 and table 3.4 show the median ratio for the reactive metal pool extracted with HNO<sub>3</sub>, HCl or EDTA relative to the total metal content.




Figure 3.5 Mean median ratio of reactive metal pool relative to the total metal content as found for 0.43 M HNO<sub>3</sub>, 0.1 M HCl, and 0.05 M EDTA. The error bars represent the standard deviation.

The size of the reactive metal pool relative to the total metal content decreases in the order Cd > Cu > Pb > Zn ≈ Ni >> Cr. Differences between HNO<sub>3</sub>, HCl, and EDTA are different for the metals measured in this study, although the amounts extracted with HNO<sub>3</sub>, HCl and EDTA are closely correlated (figure 3.6 and table 3.5)

Table 3.4. Ratio of 0.43 N HNO<sub>3</sub> and Aqua Regia extractable metals in soil

	Metal (0.43 N HNO <sub>3</sub> /AR)					
	Cu	Zn	Cd	Cr	Ni	Pb
minimum	0.08	0.05	0.08	0.01	0.04	0.16
1%	0.44	0.14	0.31	0.02	0.12	0.32
average	0.61	0.24	0.68	0.07	0.27	0.58
median	0.60	0.23	0.68	0.07	0.27	0.57
99%	0.81	0.42	1.18	0.17	0.42	0.84
maximum	6.42 <sup>1</sup>	1.07	6.66 <sup>1</sup>	0.94	1.70	2.061 <sup>1</sup>

<sup>1</sup> Theoretically, the ratio of 0.43 HNO<sub>3</sub>/AR cannot exceed 1 since 0.43 N HNO<sub>3</sub> only extracts a part of what is extracted by Aqua Regia. The number of samples for which the ratio 0.43 HNO<sub>3</sub>/AR exceeds 1 is however limited as appears also from figure 1 (few data are positioned above the 1:1 line).



Table 3.5 Correlation matrix of metals in soil as measured by extraction with Aqua Regia (AR), 0.43 M HNO<sub>3</sub>, 0.05 M EDTA, and 0.1 M HCl

	MeAR	MeHNO <sub>3</sub>	MeEDTA		MeAR	MeHNO <sub>3</sub>	MeEDTA
Cd-AR	1.00	-	-	Zn-AR	1.00	-	-
Cd-HNO <sub>3</sub>	0.99	1.00	-	Zn-HNO <sub>3</sub>	0.84	1.00	-
Cd-EDTA	0.98	0.98	1.00	Zn-HCl	0.85	0.98	1.00
Cd-HCl	0.99	0.98	0.98	Zn-EDTA	0.83	0.97	0.97
Pb-AR	1.00	-	-	Ni-AR	1.00	-	-
Pb-HNO <sub>3</sub>	0.98	1.00	-	Ni-HNO <sub>3</sub>	0.90	1.00	-
Pb-EDTA	0.98	0.98	1.00	Ni-EDTA	0.86	0.98	1.00
Pb-HCl	0.95	0.94	0.95	Ni-HCl	0.89	0.99	0.98
Cu-AR	1.00	-	-	Cr-AR	1.00	-	-
Cu-HNO <sub>3</sub>	0.96	1.00	-	Cr-HNO <sub>3</sub>	0.88	1.00	-
Cu-EDTA	0.95	0.98	1.00	Cr-EDTA	0.69	0.82	1.00
Cu-HCl	0.96	0.98	0.98	Cr-HCl	0.85	0.92	0.84

Only for Cr, the amounts of this metal extracted with AR and EDTA are less strongly correlated. For Cd, Zn, and Ni, the amounts extracted with HNO<sub>3</sub> are slightly, but significantly ( $p < 0.001$ ; based on two-sample paired T-test) larger than those extracted with HCl. For Pb, however, the difference is substantial and the amounts extracted with HNO<sub>3</sub> are almost twice those extracted with HCl. Apparently, the stronger acidity of the HNO<sub>3</sub> solution is able to release a significant part of Pb which cannot be extracted with HCl. The amounts of metals extracted by EDTA are always smallest as compared to those extracted with the other two extracts. Only for Cu and Pb, the amounts extracted with EDTA exceed those extracted with HCl. This difference may reflect the amount of these heavy metals adsorbed or bound to SOM which can be released by EDTA but not by HCl. Both Cu and Pb are known for their high affinity binding to organic ligands present in SOM as well as in dissolved organic carbon (DOC) in the soil solution (Weng et al., 2001; Qin et al., 2006; Koopmans et al., 2008b). For Cr, none of the three extracts tested here extracts significant amounts of this metal as compared to its total content. The maximum value of the amount of Cr extracted with HNO<sub>3</sub>, HCl or EDTA relative to the total content of this metal was 12%. Hence, Cr in the soils studied here is probably present in rather immobile and chemically inert fractions which can be dissolved only to a slight extent by extraction with HNO<sub>3</sub>, HCl or EDTA. The reactive pool of Pb, which is commonly considered a rather immobile metal (Tipping et al., 2006), is high relative to its total content with a median ratio equaling 61% while the relative sizes of the reactive Zn and Ni pools are much smaller. For Zn, a median ratio of only 23% of its reactive pool relative to its total content could be extracted with HNO<sub>3</sub>, which, for Zn, is the strongest of the three extracts tested here. Apparently, Zn and also Ni are mostly present in rather immobile and chemically inert fractions forms in the soils studied here. Both Zn and Ni are known for their tendency to become immobile due to fixation and surface precipitation of these metals on oxihydroxides (Singhal and Gupta 1978; Bruemmer et al., 1981; Elzinga and Sparks 1999; Buekers 2007). In contrast to Cr, Ni, and Zn, Cd remains highly available in the soils studied here which is in agreement with data from soils for moderate climate zones (McBride et al., 2006). For both HNO<sub>3</sub> and HCl, more than 70% can be extracted from soil compared to its total content. In 5% of all soil

samples, the ratio of the reactive Cd pool relative to its total content increases to even more than 90%. Hence, Cd in these soils remains in a potentially available pool which can become available for uptake by rice plants on the long-term. The median value of 74% for Cd as measured by HNO<sub>3</sub> is in close agreement with data from non-polluted soils from the Netherlands and Belgium (Römken et al., 2004; Meers et al., 2007). Apparently, the chemical processes controlling the reactivity or binding of Cd to soil components like SOM and clay are comparable regardless the obvious differences between well-drained soils from moderate climate zones and the paddy field soils from Taiwan studied here. Also, the ratio of the reactive Pb, Zn, Cu, and Ni pools relative to the total contents of these metals was the same for the soils studied here and non-polluted non-tropical soils (Römken et al., 2004; Meers et al., 2007).




Figure 3.6. Relation between total (Aqua Regia) and reactive metal fraction (0.43 N HNO<sub>3</sub>), the solid line is the 1:1 line.




Figure 3.7. Comparison between the metals extracted by 0.1 M HCl (Y-axis) and 0.43 N HNO<sub>3</sub> (X-axis). The dotted line is the 1:1 line for comparison, the dashed line marks the regression line

### 3.2.3 Actual available metal pools and solid-solution partitioning models

The amounts of metals extracted with CaCl<sub>2</sub> are much lower compared to the total contents or the reactive pools (table 3.2). Table 3.6 contains an overview of the average ratio of the directly available metal pool relative to the reactive metal pool as extracted with HNO<sub>3</sub>. Especially for Cd, Zn, and Ni, the directly available pool of metals is rather high relative to the reactive metal pool; approximately 10% of these metals can be considered to be directly available for uptake by rice plants. In contrast to this, the relative size of the directly available metal pool for Cu, Cr, and Pb is less than 0.3%, while the maximum value remains below 4%.

Table 3.6 Overview of the ratio of the directly available metal pool as measured by extraction with 0.01 M CaCl<sub>2</sub> relative to the reactive metal pool as measured by extraction with 0.43 M HNO<sub>3</sub>

	Ratio directly available metal pool/reactive metal pool (%)					
	Cu	Zn	Cd	Cr	Ni	Pb
Minimum	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1
Median	0.1	7.9	8.8	0.2	7.1	0.1
Average	0.2	9.3	10.4	0.3	9.4	0.2
Maximum	2.7	75.0	65.8	3.3	77.1	3.8

Table 3.7 shows the results of the multiple regression analysis using equation [2.1]. Here, only the reactive metal pool as measured by extraction with HNO<sub>3</sub> and soil properties like pH and CEC were used as input parameters.

Table 3.7 Regression coefficients of the Freundlich equation to predict the size of the 0.01 M CaCl<sub>2</sub>-extractable metal pool using 0.43 M HNO<sub>3</sub> and soil properties like pH and CEC as input parameters

Metal	n	Regression coefficients				R <sup>2</sup>	se(Y-est)
		Intercept	a (pH)	b (CEC)	c (HNO <sub>3</sub> )		
Cd	3255	2.64	-0.53	-0.80	0.96	92.9	0.22
Cu	2776	0.16	-0.38	-1.18	1.17	58.7	0.38
Cr	2567	-1.12	-0.05	-0.59	0.17	9.9	0.41
Ni	3185	2.73	-0.58	-0.95	1.22	86.0	0.27
Pb	2197	-1.69	-0.02	-0.28	0.65	46.7	0.40
Zn	3244	4.57	-0.88	-1.25	1.16	80.3	0.30

For Cd, Zn, and Ni, the main part of the observed variation in the directly available metal pool as measured by extraction with CaCl<sub>2</sub> can be explained by these parameters as is shown in figure 3.8 as well.




Figure 3.8 Measured data of metals in the 0.01 M CaCl<sub>2</sub> extracts versus model fit based on Eq. 2.1. The solid line indicates the 1:1 line, while the dotted lines are equal to  $Y\text{-est} \pm 2 \cdot se(Y\text{-est})$  which is approximately equal to the 95%-confidence interval of the predicted mean levels

For Cr, Cu, and Pb, much less of the variation in the directly available metal pool was explained by the reactive metal pool, pH and CEC. Apparently, the parameters included here are not the only ones controlling the solid-solution partitioning of Cu, Pb and especially of Cr. For Cu and Pb, the role of DOC is important in the prediction of the solubility of these elements (Weng et al., 2002; Römken et al., 2004; Koopmans et al., 2008b). In this study, however, DOC was not measured which partly explains the rather low explained variability of the directly available Cu and Pb pools. For Cr, the model as such is not an appropriate one since Cr behaves rather different being an anion and prone to precipitation and dissolution processes even at low concentrations. Neither the reverse effect of pH on the solid-solution partitioning for anions compared to cations nor the precipitation phenomena are accounted for in the Freundlich equation (Eq. 2.1).

A further analysis of the data revealed that Zn had a significant influence on the amounts of Cd, Ni and to a lesser extent also of Cu extracted with CaCl<sub>2</sub>. Since Zn and Cd have similar affinity for sorption sites in soils (McBride 1994), an increase in the available pool of Zn should induce competition between these metals for sorption leading to higher directly available Cd levels in soil as measured by extraction with CaCl<sub>2</sub>. Including Zn in the regression indeed improved the model even further ( $R^2 = 0.95$ ;  $se(Y\text{-est}) = 0.19$ ) as is shown for Cd in Eq. [3.1]:

$$\begin{aligned} \log[\text{Cd-CaCl}_2] = & 0.83 \\ & - 0.29 \cdot \text{pH} \\ & - 0.50 \cdot \log[\text{CEC}] \\ & + 0.91 \cdot \log[\text{Cd-HNO}_3] \\ & + 0.29 \cdot \log[\text{Zn-CaCl}_2] \end{aligned} \quad [3.1]$$

With  
[Zn-CaCl<sub>2</sub>] directly available metal pool extracted with 0.01 M CaCl<sub>2</sub> in mg kg<sup>-1</sup>

For Ni and Cu, similar improvements were obtained upon inclusion of Zn in the regression (results not shown). For Pb and Cr, however, no effect of Zn was observed. Apparently, Pb and Zn do not compete for the same sorption sites in paddy soils, whereas a strong competition effect was found to exist between Cd, Ni and to a lesser extent for Cu on the one hand and Zn on the other for sorption to the soil solid phase.

In table 3.8, results of the multiple regression analysis are presented when using all soil tests (i.e., AR, HNO<sub>3</sub>, HCl, and EDTA) only for Cd, Ni, and Zn. For Pb and Cr, and to a lesser extent also for Cu, the error of prediction using HNO<sub>3</sub> was high (table 3.8), but the use of other soil tests like AR, HCl, and EDTA did not improve this (results not shown). The data presented in table 3.8 indicate that the use of HCl or HNO<sub>3</sub> and, for Ni even EDTA, result in comparable regression coefficients for the prediction of the directly available metal pool in soil. This is not surprising considering the similarity between these soil tests in extracting metals from soil. Results from an all-subset analysis (Genstat 11.1) nevertheless showed that the combination of pH/CEC/HNO<sub>3</sub> was the most significant model based on 3 terms.

After including Zn in the regression terms (equation 3.1) the best model still was the one using the 0.43 M HNO<sub>3</sub> extractable pool. The use of AR results in both a lower R<sup>2</sup> and a higher standard error of the predicted values which indicates that the prediction is less accurate. This is in line with the fact that AR extracts the inert fraction as well, as is shown in figure 3.6 and this inert fraction is thought not to be in equilibrium with the reactive or directly available metal pools. This effect is more pronounced for Ni and Zn for which also the difference between the reactive and total metal content was much larger than for Cd. For Cd, differences between model coefficients based on HNO<sub>3</sub>, HCl or EDTA are comparable to those for AR which shows that the reactive metal pool is closely related to the total metal content in the soil. Also for Cd, however, the model fit based on AR is less accurate compared to that based on HNO<sub>3</sub>, HCl or EDTA.

Table 3.8 Regression coefficients of the Freundlich equation to predict the size of the 0.01 M CaCl<sub>2</sub> extractable metal pool based on Aqua Regia (AR), 0.43 M HNO<sub>3</sub>, 0.1 M HCl or 0.05 M EDTA and soil properties like pH and CEC (Eq. 2.1)

Metal and extract	Regression coefficients				R <sup>2</sup>	se(Y-est)
	Intercept	a (pH)	b (CEC)	c (Reactive metal pool)		
Cd						
HNO <sub>3</sub>	2.64	-0.53	-0.80	0.96	92.9	0.22
HCl	2.69	-0.53	-0.83	0.96	92.8	0.23
EDTA	2.81	-0.53	-0.87	1.02	93.0	0.22
AR	2.58	-0.55	-0.79	1.04	91.2	0.25
Zn						
HNO <sub>3</sub>	4.57	-0.88	-1.25	1.16	80.3	0.30
HCl	4.54	-0.87	-1.21	1.16	80.7	0.30
EDTA	4.34	-0.83	-1.18	1.17	82.1	0.29
AR	4.65	-1.01	-1.18	1.12	74.1	0.35
Ni						
HNO <sub>3</sub>	2.73	-0.58	-0.95	1.22	86.0	0.27
HCl	2.88	-0.57	-0.98	1.16	86.2	0.27
EDTA	2.78	-0.55	-0.99	1.17	86.4	0.27
AR	2.01	-0.61	-1.63	1.67	77.3	0.35

For Cd, a similar regression analysis performed by Meers et al. (2007) resulted in remarkably similar model coefficients. Meers et al. (2007) used the total Cd content as extracted with AR instead of a reactive Cd pool to predict the directly available pool extracted with 0.01 M CaCl<sub>2</sub> and obtained the following regression model for Cd:

$$\begin{aligned} \log[\text{Cd-CaCl}_2] = & 2.91 (\pm 0.26) \\ & - 0.55 (\pm 0.04) \cdot \text{pH} \\ & - 0.94 (\pm 0.14) \cdot \log(\text{CEC}) \\ & + 1.13 (\pm 0.06) \cdot [\text{Cd-AR}] \end{aligned} \quad [3.2]$$

With  
[Cd-CaCl<sub>2</sub>] directly available metal pool extracted with 0.01 M CaCl<sub>2</sub> in mg kg<sup>-1</sup>

pH	pH in the 0.01 M CaCl <sub>2</sub> extracts
[CEC]	cation exchange capacity in cmol(+) kg <sup>-1</sup>
[Cd-AR]	total Cd determined by extraction with AR in mg kg <sup>-1</sup>

All coefficients related to soil properties are remarkably close with exactly the same effect of pH and only minor differences in the coefficients for the total Cd content and CEC (table 3.8). The higher model intercept of 2.91 derived by Meers et al. (2007) compared to the intercept of 2.58 from our data results in a significant overestimation of approximately a factor of 2 when applied to the data from this study. The slope of log-transformed data from this study, however, versus predicted log-transformed values using Eq. [2.1] is 0.93 with an R<sup>2</sup> of 0.91 (results not shown). This suggests that the response of Cd in the CaCl<sub>2</sub> extracts to changes in pH, CEC and even the total Cd content is the same in our data as in those obtained by Meers et al. (2007).

### 3.2.4 Effect of sampling time on the directly available metal pools and on solid-solution partitioning models

Despite similar weather conditions in 2005 and 2006 (Central Weather Bureau 2008), significant differences in the directly available pools of most metals between the first harvest obtained in May and the second harvest obtained in November were obtained (table 3.9). For all metals in the PD fields, an increase in the directly available metal pools at the second harvest ranging from 0.05 (Pb) to 0.58 (Cu) log units. In the CH, HM, LK and HC fields, however, the reverse trend was observed and directly available metal pools at the first harvest exceeded those extracted at the second harvest, although differences were less pronounced compared to those obtained in the PD fields (table 3.9). Only for Pb, CaCl<sub>2</sub>-extractable metal levels at the second harvest exceeded those from the first harvest in all fields. Also, pH levels in all fields were higher at the second harvest compared to the first harvest which explains the larger size of the directly available metal pools in the CH, HM and LK fields at the first harvest but not in the PD fields. The higher pH at the second harvest fails to explain the higher CaCl<sub>2</sub>-extractable metal levels in the PD fields since the size of the directly available metal pool for metals like Cd, Zn and Ni usually decreases with an increase in pH (McBride, 1994). In addition to this, there were no statistically significant differences in either CEC or the reactive soil metal pools as measured by extraction with HNO<sub>3</sub>. Possibly dynamics in other soil parameters including DOC, salinity or redox potential which were not included in this study play an important role in the temporal variability of the directly available metal pools.



Table 3.9 Mean difference and 95% confidence interval of the directly available metal pool measured with 0.01 M CaCl<sub>2</sub> in soil samples from harvest 1 (May) and harvest 2 (November). This difference was calculated as the size of the metal pool at harvest 1 minus the size of the metal pool at harvest 2.

	CH, HM, and LK fields			PD fields		
	Mean	95%- confidence interval	p	Mean	95%- confidence interval	p
Cd	0.09 <sup>1</sup>	0.08 - 0.11	<0.001	-0.20	-0.22 - -0.17	<0.001
Cu	0.11	0.08 - 0.14	<0.001	-0.58	-0.64 - -0.52	<0.001
Cr	0.06	0.02 - 0.11	<0.01	-0.34	-0.39 - -0.29	<0.001
Ni	0.05	0.03 - 0.08	<0.001	-0.12	-0.14 - -0.09	<0.001
Pb	-0.18	-0.36 - -0.1	<0.001	-0.05	-0.07 - -0.02	<0.001
Zn	0.06	0.03 - 0.09	<0.001	-0.12	-0.13 - -0.10	<0.001
pH	-0.09	-0.12 - -0.06	<0.001	-0.14	-0.16 - -0.12	<0.001

<sup>1</sup> All data are based on differences between log transformed values. The significance of the differences between harvest 1 and 2 were calculated based on a paired two sample T-test.

The differences between the directly available metal pool at the first and the second harvest became evident after deriving separate solid-solution partitioning models for the data from these harvests as shown in table 3.10.

Table 3.10 Regression coefficients of the Freundlich equation to predict the size of the 0.01 M CaCl<sub>2</sub>-extractable metal pool using 0.43 M HNO<sub>3</sub> and soil properties like pH and CEC as input parameters (Eq. 1) at harvest 1 (H1) and at harvest 2 (H2)

Metal	Regression coefficients				R <sup>2</sup>	se(Y-est)
	Intercept	a (pH)	b (CEC)	c (HNO <sub>3</sub> )		
Cd-H1&H2	2.64	-0.53	-0.80	0.96	92.9	0.22
Cd-H1	2.20	-0.47	-0.71	0.88		
Cd-H2	3.17	-0.61	-0.88	1.05		
Zn-H1&H2	4.57	-0.88	-1.25	1.16	80.3	0.30
Zn-H1	3.96	-0.77	-1.28	1.17		
Zn-H2	5.32	-1.00	-1.20	1.14		
Cu-H1&H2	0.16	-0.38	-1.18	1.17	58.7	0.38
Cu-H1	-1.22	-0.25	-1.26	1.55		
Cu-H2	1.35	-0.52	-1.23	1.01		
Ni-H1&H2	2.73	-0.58	-0.95	1.22	86.0	0.27
Ni-H1	2.07	-0.50	-0.87	1.31		
Ni-H2	3.49	-0.67	-1.03	1.13		
Pb-H1&H2	-1.69	-0.02	-0.28	0.65	46.7	0.40
Pb-H1	-1.41	-0.07	-0.48	0.73		
Pb-H2	-1.79	0.00 (ns)	-0.05 <sup>1</sup>	0.55		

<sup>1</sup>Not significant

The data were split solely based on the time of sampling but no further distinction was made between fields. For all metals except Pb, the intercept increased between 0.97 (Cd) and 2.57 (Cu) log units. The ultimate effect of the time of harvest on the solubility of metals appeared to be most pronounced in the more polluted paddy fields where, for all metals, a clear increase in the size of the directly available pool was observed at the second harvest. This is schematically shown for Cd in figure 3.9

where the ratio of the directly available Cd pool at the second harvest relative to that obtained at the first harvest is plotted. For each combination of the reactive Cd pool and pH, the directly available Cd pool was calculated for both the first and second harvest, and plotted as a function of the reactive Cd pool and pH. This figure illustrates the overall effect of the time of harvest across all fields which is difficult to assess from the data points for each field alone. The increase in the directly available Cd pool at the second harvest is most pronounced in polluted soils with a reactive Cd pool  $> 1 \text{ mg kg}^{-1}$  and a pH  $< 5.5$ . In contrast, whereas the opposite effect was observed at a higher pH or at a smaller reactive Cd pool.

At present, it remains unclear as to what causes these changes in both pH and  $\text{CaCl}_2$ -extractable metal levels but the observations clearly show that the dynamics of the directly available metal pools in paddy fields throughout the year can be considerable. Obviously, the change from reducing to oxidizing conditions and vice versa can result in long-lasting changes in the directly available pool of metals in soil. Further investigation on the controls of metal solubility under drained and water-logged conditions and its dynamics after changing from the anaerobic state of the soil to the aerobic state seems essential and can be the clue to understanding some of the contrasting results that have been obtained not only in this study but by others as well. Simmons et al. (2008), for example, showed that the availability of Cd in extracts from air-dried soils differed from that from field-moist samples. However, the direction of this change was different and depended on the type of extract used. Dissolved Cd concentrations in a 0.01 M  $\text{CaCl}_2$  extract decreased upon drying, but those in 0.05 M  $\text{CaCl}_2$  extracts increased upon drying. This shows that the time of sampling, conditions at the time of soil sampling, soil pretreatment and extraction procedure all affect the amounts of metals extracted by standard soil tests. Obviously, the usefulness of the available fraction of metals in soils in assessment of soil quality and biological risks is evident but these results suggest that the underlying mechanisms that control the amount of metals present in such extracts still require ample attention.




Figure 3.9 Difference in the size of the 0.01 M  $\text{CaCl}_2$ -extractable Cd pool as measured at harvest 1 (H1) and at harvest 2 (H2)

### 3.3 Mechanistic modeling of the availability of rice in soils

The partition equations described in the previous sections suggest that the availability of Cd in the soil samples studied here is mainly controlled by sorption processes. To verify this hypothesis a mechanistic model has been applied where the solubility of Cd as measured in the  $\text{CaCl}_2$  extract was predicted by a previously calibrated model (Bonten et al., 2008). In this model sorption of Cd to different soil components like organic matter, iron oxides and clay is described taking into account sorption and precipitation processes. Also speciation in solution is considered (for example Cd binding to chlorine and dissolved organic carbon). Some data were not available, such as dissolved organic carbon and clay and these were estimated from either organic matter (DOC) or CEC (clay). pH  $\text{CaCl}_2$  was used as well as the 0.43 N  $\text{HNO}_3$  extractable mild content as the total Cd pool available for sorption processes. More details on the model used can be found in Weng et al. (2001) and Bonten et al. (2008).

In figure 3.10 the predicted Cd concentration in  $\text{CaCl}_2$  is plotted against the measured concentration. Here, the amount in  $\text{mg kg}^{-1}$  has been converted to mol per liter based on the 1:10 soil-solution ratio of the extraction.




Figure 3.10. Measured versus predicted Cd concentration in all samples

Although there is a considerable scatter around the 1:1 line, the overall prediction range is in close agreement with measured data. Considering the fact that this model was not calibrated on the data from Taiwan, the fit is rather good.

Also, the relation with predictions from the partitioning model is quite good as is shown in figure 3.11. The dotted line represents the 1:1 line, and most predicted values center around this line although there is a small tilt between both predictions. Nevertheless it confirms that the mechanistic model is able to predict the measured variability quite well. On the other hand it also shows that partitioning models which

can be easily derived from data from field studies are able to predict the availability of Cd in soils.




Figure 3.11. Correlation between model predicted values from the partitioning model versus the mechanistic model

### 3.4 Conclusions

Due to the use of polluted irrigation water, metals in paddy fields in Taiwan have reached levels that exceed soil quality standards above which soil cleanup is required. The levels of all metals in the 19 field studied here are quite heterogeneous across the field and are always highest near the inlet of irrigation water. To assess the availability of metals in soil, different extraction methods were tested. Results indicated that methods used in the EU (HNO<sub>3</sub>), Australia (EDTA), and Taiwan (HCl) give comparable results, although HNO<sub>3</sub> consistently extracts more metals from soil than HCl or EDTA. As such, all three soil tests can be used to predict the size of the directly available pool of Cd, Zn, and Ni and to a lesser extent also of Cu and Pb. The regression analysis nevertheless indicated that the combination of HNO<sub>3</sub>-extractable metals with CEC and pH is the most suitable one to explain the observed variation in the directly available metal pool. For Cr, no such relation could be obtained since the solubility of Cr is controlled by other processes as compared to cationic metals like Cd or Zn. For Cd, Ni, and Cu, a competition effect of Zn was observed with the size of the directly available Cd, Ni, and Cu pools increasing with an increase of the directly available Zn pool. The soils studied here have been developed under very different conditions compared to soils in moderate climate zones, but the variability in the directly available Cd, Zn, and Ni pools and to a lesser extent in the directly available Cu and Pb pools can be described well by solid-solution partitioning models similar to those published for non-paddy soils. There is a close agreement in the solid-solution partitioning model for Cd obtained here and those published by Römkens et al. (2004) and Meers et al. (2007). This suggests that processes that control the variability in the directly available Cd pool as measured by extraction with CaCl<sub>2</sub> in drained paddy fields are comparable to those in soils from

moderate climate zones like Western Europe. This is an important finding since it suggests that knowledge and model concepts for metals which have been developed elsewhere can also be of use to predict the availability of metals in paddy fields. In this study, we focused on the prediction of the directly available metal pool under drained, oxidizing conditions since it has been suggested that the uptake of metals by rice is controlled largely during the period that the soil is drained (Simmons et al., 2008). It is known, however, that both the solubility and availability of metals decrease under reducing conditions when the soil is water-logged (Kashem and Singh 2004; Contin et al., 2007). The return to oxidizing conditions can lead to an increase in the availability and uptake of metals (Daum et al., 2001; Kelderman and Osman 2007), although the degree to which metals are bound or released by sulfides has been shown to be quite variable. Barret and McBride (2007) showed that an increase in the Cd:Zn ratio in soil reduced the metal scavenging effect of sulphides for Cd considerably. Charlatchka and Cambier (2000) even observed an initial increase in the solubility of metals after flooding of a soil due to dissolution of ferric and manganese oxides. These contrasting observations, together with the observed differences in the size of the directly available metal pool between the first and the second harvest, suggest that careful monitoring of soil conditions, metal ratio's and time period of drainage is essential in order to better understand the dynamics of the availability and uptake of metals in paddy soils. The data collected here nevertheless illustrate that the use of a dilute acid extracts like 0.43 M HNO<sub>3</sub> or 0.1 M HCl in combination with generally available soil properties like pH and CEC can be used to obtain a robust estimate of the availability of some of the most critical elements in relation to food safety and environmental effects (Cd, Zn, and Ni). To better explain the observed differences in the solubility of Cu and Pb, the influence of DOC needs to be considered.

The final step is to link model concepts such as the ones described here to those that predict uptake by arable crops and/or leaching to ground- and surface waters on a regional scale as was recently accomplished for the Netherlands (de Vries et al., 2008). The data presented in this study will therefore be used to establish a link between the availability of metals in paddy fields and uptake by rice (see Chapter 4). The aim is to use such a concept to assess the suitability of contaminated land for rice cropping on a local and regional scale.



## 4 Uptake of metals by rice

### 4.1 General quality assessment of rice grown on study locations

Table 4.1 gives an overview of soil and rice data used in this study including ranges and minimum and maximum values. The data were sorted by rice cultivar because our aim is to relate the availability of Cd in soil to uptake by different cultivars and not so much to derive relationships for specific fields.

*Table 4.1 Overview of pH, cation exchange capacity (CEC), soil organic matter (SOM), reactive (0.43 M HNO<sub>3</sub>) and total (Aqua Regia) soil Cd sorted by rice cultivar. Minimum and maximum values and median (in brackets) are presented*

Cult.	Family	pH	CEC (cmol[+] kg <sup>-1</sup> )	SOM (%)	Reactive Cd (mg kg <sup>-1</sup> )	Total Cd (mg kg <sup>-1</sup> )
1	Japonica	4.0 - 6.8 (5.4)	2.6 - 21.3 (14.4)	1.7 - 9.4 (6.5)	0.09 - 19.5 (0.39)	0.13 - 27.8 (0.60)
2	Japonica	4.1 - 7.2 (5.4)	3.1 - 21.1 (14.2)	1.6 - 8.9 (6.3)	0.09 - 16.4 (0.37)	0.09 - 18.8 (0.61)
3	Japonica	4.2 - 7.2 (5.4)	3.8 - 20.9 (14.1)	1.6 - 9.2 (6.5)	0.09 - 17.5 (0.41)	0.11 - 18.2 (0.65)
4	Japonica	3.8 - 7.2 (5.4)	4.1 - 21.8 (13.7)	2.0 - 9.5 (6.5)	0.09 - 15.0 (0.43)	0.07 - 17.4 (0.64)
5	Japonica	3.9 - 6.9 (5.4)	3.2 - 21.6 (14.2)	1.7 - 9.2 (6.5)	0.09 - 17.3 (0.40)	0.06 - 23.9 (0.59)
6	Indica	4.3 - 7.0 (5.4)	3.2 - 25.1 (14.5)	1.3 - 9.6 (6.4)	0.08 - 18.1 (0.43)	0.08 - 21.2 (0.71)
7	Japonica	4.2 - 6.8 (5.3)	3.3 - 24.2 (14.4)	2.2 - 9.3 (6.6)	0.09 - 19.1 (0.43)	0.13 - 25.9 (0.60)
8	Japonica	3.9 - 7.2 (5.4)	4.2 - 21.6 (14.0)	1.4 - 9.3 (6.6)	0.09 - 20.6 (0.44)	0.08 - 26.6 (0.66)
9	Indica	4.1 - 6.8 (5.4)	2.8 - 21.7 (14.8)	1.9 - 9.9 (6.1)	0.08 - 19.9 (0.33)	0.09 - 25.9 (0.57)
10	Indica	4.1 - 7.0 (5.3)	2.6 - 21.7 (14.0)	1.9 - 10.2 (6.5)	0.10 - 20.2 (0.43)	0.14 - 25.8 (0.71)
11	Indica	4.2 - 6.9 (5.4)	2.9 - 22.5 (14.5)	1.7 - 8.9 (6.3)	0.08 - 19.8 (0.41)	0.14 - 22.7 (0.70)
12	Japonica	4.0 - 6.8 (5.4)	3.0 - 20.5 (14.8)	1.9 - 9.0 (6.3)	0.08 - 12.6 (0.36)	0.13 - 16.8 (0.59)

Observed total Cd concentrations in soil range from less than 0.1 mg kg<sup>-1</sup> to almost 30 mg kg<sup>-1</sup> which covers the range from background levels to heavily polluted soils. In Taiwan, total Cd concentrations in soil below 0.4 mg kg<sup>-1</sup> are considered background levels without a clear indication of anthropogenic pollution (Lin et al., 2002). As mentioned in chapter 3, a SQS of 5 mg kg<sup>-1</sup> as determined by extraction with AR is used (EPA, 2006) to identify the suitability of land for agricultural purposes. According to this definition, almost 27% of the plots is unsuitable to serve as arable land.

Although the ranges in soil properties including total Cd concentration in soil are rather similar for all rice cultivars, Cd levels in rice grains are quite different among the cultivars used in this study as is shown in table 4.2.

Table 4.2 Overview of Cd concentrations in roots and grains of rice cultivars used in this study. For Cd levels in roots, minimum and maximum values and median are presented (in brackets), while for Cd levels in rice grains the 25%- and 75%-percentiles are presented as well. For ratio of Cd levels in rice grains over roots, mean is presented. Numbers in Italics indicate Cd levels exceeding the food quality standard (FQS) of 0.2 mg kg<sup>-1</sup> from the WHO while numbers in bold indicate levels exceeding the Japanese and Taiwanese FQS of 0.4 mg kg<sup>-1</sup>

Cultivar	Family	Cd roots (mg kg <sup>-1</sup> )	Cd rice grains (mg kg <sup>-1</sup> )					Ratio Cd rice grains/roots
			minimum	25%	median	75%	maximum	
1	Japonica	0.8 - 373.4 (8.0)	0.02	0.11	<i>0.21</i>	<b>1.06</b>	<b>4.57</b>	0.029
2	Japonica	0.5 - 181.3 (6.6)	0.01	0.11	<i>0.23</i>	<b>1.15</b>	<b>6.00</b>	0.037
3	Japonica	0.4 - 403.9 (9.7)	0.02	0.08	0.19	<b>1.09</b>	<b>2.98</b>	0.026
4	Japonica	0.5 - 198.5 (8.0)	0.01	0.09	0.19	<b>0.97</b>	<b>4.47</b>	0.029
5	Japonica	0.7 - 213.6 (8.8)	0.02	0.08	0.18	<b>1.13</b>	<b>3.32</b>	0.025
6	Indica	0.6 - 247.7 (6.5)	0.02	<i>0.22</i>	<b>0.43</b>	<b>1.69</b>	<b>12.57</b>	0.096
7	Japonica	0.6 - 175.2 (7.2)	0.02	0.10	<i>0.20</i>	<b>1.31</b>	<b>3.71</b>	0.030
8	Japonica	0.6 - 139.2 (5.4)	0.01	0.09	0.19	<b>1.16</b>	<b>3.39</b>	0.032
9	Indica	0.5 - 107.0 (6.2)	0.03	<i>0.23</i>	<b>0.44</b>	<b>1.39</b>	<b>7.64</b>	0.075
10 (A)	Indica	0.5 - 161.8 (6.4)	0.04	<i>0.25</i>	<b>0.60</b>	<b>3.65</b>	<b>25.32</b>	0.092
11 (B)	Indica	0.7 - 266.6 (9.6)	0.03	0.19	<i>0.37</i>	<b>2.85</b>	<b>29.08</b>	0.061
12 (C)	Japonica	0.6 - 107.7 (4.6)	0.01	0.08	0.16	<b>0.49</b>	<b>3.72</b>	0.034



Cd accumulation in the rice grains of the Indica species tested here proved to be very high. For all Indica species tested here, median levels of Cd exceed the FQS of the WHO (i.e., 0.2 mg kg<sup>-1</sup>) and the FQS used in Japan and Taiwan (i.e., 0.4 mg kg<sup>-1</sup>). Hence, such Indica species are not suitable for cropping on soils affected by Cd pollution as has been previously reported by He et al. (2006). In contrast to the Indica species, Cd accumulation in rice grains in Japonica rice species is lower, although median levels found in the grain are close to or in excess of the FQS from the WHO as well. Apparently, the combination of elevated total Cd concentrations in soil and soil properties like pH and SOM results in an increased availability of Cd in soil, and, in turn, a higher uptake of Cd by rice plants. This is further illustrated in figures 4.1 and 4.2 which show the percentage of samples where Cd levels in rice grains exceed the FQS of 0.2 or 0.4 mg kg<sup>-1</sup> at a given total Cd concentration in soil (figure 4.1) and the distribution of the Cd levels per cultivar (figure 4.2).




Figure 4.1. Percentage of samples where Cd concentration in rice grains exceeds the food quality standard (FQS) of 0.2 mg kg<sup>-1</sup> from the WHO (left) and the Japanese and Taiwanese FQS of 0.4 mg kg<sup>-1</sup> (right) at a given total Cd concentration in soil as measured by extraction with Aqua Regia for Japonica and Indica rice cultivars.




Figure 4.2. Levels of Cd in rice grains of the 12 cultivars grown in the study (level in mg kg<sup>-1</sup>)

Even at total Cd levels in soil below 0.3 mg kg<sup>-1</sup>, still a large number of rice grain samples of both families does not meet the FQS. For Indica cultivars, for example,

this percentage ranges from 11.3% for the FQS of 0.4 mg kg<sup>-1</sup> to 51.1% for the FQS of 0.2 mg kg<sup>-1</sup>. Apparently, Indica species cannot be grown safely in Cd-enriched soils without a considerable risk of exceeding the FQS of the WHO. Furthermore, a SQS of 5 mg kg<sup>-1</sup> based on extraction of soil with AR used in Taiwan for arable land is insufficient as a tool to assess the suitability of soils for rice cropping. Even for Japonica cultivars, which accumulated significantly lower amounts of Cd, between 17% for a FQS of 0.4 mg kg<sup>-1</sup> and 30% for a FQS of 0.2 mg kg<sup>-1</sup> of all rice grain samples grown on soils with total Cd levels below 5 mg kg<sup>-1</sup> do not meet the FQS. For Indica cultivars, these percentages increase even to 41% for a FQS of 0.4 mg kg<sup>-1</sup> and to 73% for a FQS 0.2 mg kg<sup>-1</sup>. Clearly, SQS based on AR fails to assess the suitability of soils for rice cropping. These results clearly stress the need to develop alternative soil tests or soil-plant models to predict Cd uptake by rice plants in order to identify those soils where rice can be grown safely without a considerable risk of Cd levels in rice grains exceeding the FQS.

#### 4.2 Transfer of Cd from soil to roots and shoot-root ratios

Uptake of contaminants from soil has to occur through root action and hence can only be realized through uptake from the soil solution. The first step is the transfer of metals from the soil solution onto the root surface followed by a translocation into the root. Despite the observed difference in Cd levels in Indica versus Japonica, uptake of Cd from soil by rice roots from all fields appeared to be equal for all cultivars tested here (table 4.2). No significant differences in Cd root levels between Indica and Japonica were observed. Mean Cd levels in roots from Indica species equaled 30.4 mg kg<sup>-1</sup> whereas those for Japonica species equaled 30.6 mg kg<sup>-1</sup> (LSD at the 5% level equals 3.1, taking into account differences across the field due to heterogeneity in soil Cd levels). Similar results were obtained by Liu et al. (2003) despite the fact that they used a spiked soil with up to 100 mg kg<sup>-1</sup> Cd added in a highly mobile form.

For all rice cultivars, the available Zn pool as determined by extraction with CaCl<sub>2</sub> suppressed the transfer of Cd into the root (regression based on all data, regardless the cultivar):

$$^{10}\log[\text{Cd-root}] = 2.31 + 0.88 \cdot ^{10}\log[\text{Cd-CaCl}_2] - 0.38 \cdot ^{10}\log[\text{Zn-CaCl}_2]; \quad [4.1]$$

$$R^2 = 0.85; \text{se-}Y_{\text{est}} = 0.25$$

where [Cd-root] is the measured Cd content in the root in mg kg<sup>-1</sup>. This suggests that an increase of Zn reduces the Cd binding to the root surface of rice plants which is probably due to competition with Zn for similar sorption sites on the root surface. A similar effect of Zn on Cd root levels was observed by Liu et al. (2007). This finding suggests that Indica and Japonica cultivars tested in this study mainly differ in their ability to transfer Cd from the root into the shoot.

In contrast to the similarity in the Cd-root levels, the rice-root distribution coefficients, obviously, differ substantially between Japonica (mean rice - root ratio of 0.030 which is similar to reported values by Kukier and Chaney, 2002) and Indica (mean rice - root ratio of 0.082, LSD at 1% is equal to 0.007). Differences between individual Japonica or Indica species are also significant and range from 0.025 (C5) to 0.037 (C2) for Japonica cultivars and from 0.061 (C11) to 0.096 (C6.)

The effect of Zn on uptake of Cd has been reported previously (Girling and Peterson, 1981; Hassan et al., 2006) although contrasting effects have been demonstrated as well depending on the levels of Zn and Cd in the soil and the Cd:Zn ratio for other crops as well (Kukier and Chaney, 2002; Dunbar, 2004). Cui et al. (2008) found a similar interaction between Cu and Cd, with Cd uptake decreasing at higher Cu addition levels. Liu et al. (2007) found that Zn in solution suppressed Cd in roots but increased Cd levels in rice. Several of the studies mentioned here, however, have disadvantages which hamper a clear evaluation of the results or a translation to field conditions. In most cases the available Cd pool was not actually measured (e.g. Liu et al., 2007). Likewise, soils are spiked with considerable amounts of Cd (e.g. Zhou et al., 2003; Diao et al., 2005; Li et al., 2005) to mimic uptake of Cd from polluted soils. The availability of Cd and Zn in freshly spiked soils, however, decreases over time and can thus be different from that in non-spiked soils (Naidu et al., 2003, Ma and Uren, 2006). Such changes over time clearly will affect the uptake of Cd by rice. Also hydroponic studies are often performed (Cui et al., 2008) where conditions clearly are different from those in the soil. It is therefore not surprising that results from different studies can differ substantially. In order to be able to translate experimental results to the field level, we therefore suggest that the experiments or monitoring studies performed should mimic conditions that prevail in the field.

### **4.3 Models to describe Cd uptake by rice: why?**

Ultimately one of the major goals of this study is to assess whether it is possible to predict the quality of rice (grains) based on available soil data. At present, rice is tested after the harvest and depending on the quality it cannot be sold on the market if Cd in rice grains exceed the food quality standard. In the previous paragraph it became clear that current soil quality standards fail to identify those soils where rice will not meet the quality demanded by law. Obviously a suitable soil test or model based on easily available soil data would be much more efficient since this would allow to assess the quality of rice prior to seeding the plants. In paragraph 4.4 results from the model derivation are shown. Two types of models are tested as described in the materials and methods section. One is the direct relation between Cd in rice and Cd (and Zn) measured in the 0.01 M CaCl<sub>2</sub> extract and the second model is to relate the soil properties (reactive Cd content, pH and CEC) to the measured Cd levels in rice (illustrated in figure 4.3). The advantage of the first approach (CaCl<sub>2</sub> model) is that only one soil test is sufficient to predict the quality of the rice. The disadvantage is that at present, limited data on the CaCl<sub>2</sub> extractable metals levels in paddy fields are available. This means that this approach cannot be applied on a regional scale

without first having to measure the  $\text{CaCl}_2$  extractable metal content. The second method based on soil properties does allow for a regional approach since many countries do have regional data on soil pH, CEC and some form of the reactive Cd content in soil. In Taiwan for example many soils are tested using 0.1 M HCl which can be used as a suitable indicator of the reactive soil Cd content as was shown in chapter 3.




Figure 4.3 Schematic representation of the models used to predict Cd levels in rice. At the bottom the current approach based on AR is shown.

Both model 1 and 2 are based on the Freundlich equation which has been used previously for other crops as well:

$$\text{Model 1: } {}^{10}\log[\text{Cd-rice grains}] = \text{intercept} + a \cdot {}^{10}\log[\text{Cd-CaCl}_2] + b \cdot {}^{10}\log[\text{Zn-CaCl}_2]$$

$$\text{Model 2: } {}^{10}\log[\text{Cd-grain}] = \text{intercept} + f \cdot {}^{10}\log[\text{Cd-HNO}_3] + g \cdot \text{pH} + h \cdot {}^{10}\log[\text{CEC}]$$

Ultimately these models can be used to calculate critical levels in soil beyond which Cd in rice exceeds the food quality standard

#### 4.4 Regression models to predict Cd levels in rice grain

In table 4.3, regression coefficients of the  $\text{CaCl}_2$ -model (i.e., model 1) and the  $\text{HNO}_3$ -model (i.e., model 2) are shown for all individual rice cultivars as well for all Japonica species or all Indica species grouped together. The overall model performance for the individual rice cultivars for both models was rather good with small but significant differences between model coefficients for different cultivars. Apart from  $\text{HNO}_3$ , HCl and EDTA were used as well to estimate the reactive metal pool in the  $\text{HNO}_3$ -model. However, the combination of the reactive Cd pool as measured by extraction with  $\text{HNO}_3$  and pH and CEC proved to be the most significant model (results not shown). In the remainder of the study, therefore, the results of the soil test based on  $\text{HNO}_3$  will be used for the reactive soil Cd pool. Both the  $\text{CaCl}_2$ -model and the  $\text{HNO}_3$ -model for individual Japonica cultivars outperformed those of the individual Indica cultivars as indicated by a higher  $R^2$  value and lower standard errors of the predicted Y-values. The model for all Japonica cultivars or for all Indica

cultivars grouped together has a good performance as well. Model coefficients for both the reactive Cd pool and pH in the HNO<sub>3</sub>-model are surprisingly close for all rice cultivars with marginal, non-significant differences between the Japonica and Indica cultivars. The significance of the CEC is lower than for pH and for the reactive Cd pool, (results not shown), although CEC is significant at the 1%-level for most cultivars except for the rice cultivars 4, 9 and 10.

Addition of pH in the CaCl<sub>2</sub>-model did not improve the model performance (results not shown). Obviously, pH has a profound impact on the availability of Cd in soil solution in paddy fields (Römkens et al., 2009b), and the solubility of Cd and Zn clearly increase with a decrease in pH. Indeed, the HNO<sub>3</sub>-model shows increased levels of Cd in rice grains with a decrease in pH (table 4.3). When using the available Cd pool as measured by extraction with CaCl<sub>2</sub> as in the CaCl<sub>2</sub>-model, however, this effect of pH is already accounted for since the Cd concentration in CaCl<sub>2</sub> extracts is higher in samples from soils with a low pH. For this reason, pH does not contribute significantly to the regression of the CaCl<sub>2</sub>-model.

According to the HNO<sub>3</sub>-model, the predicted Cd concentration in Indica cultivars is higher at similar HNO<sub>3</sub>-extractable Cd levels and pH and CEC as due to the higher intercept in the equation for all Indica cultivars grouped together as compared to the intercept found for all Japonica species. This similarity was also observed in the solution model which performs better compared to the soil model, more so for Japonica species than for Indica species (table 4.3). Again, model performance for Japonica species is slightly better than for Indica. Model coefficients for both Cd and Zn in solution are rather similar for Japonica and Indica and predicted Cd levels in rice differ mostly due to the higher off-set. In figure 4.4, all predicted values for Cd are shown for harvest 1 and harvest 2 based on the CaCl<sub>2</sub> model for the main cultivars (Indica and Japonica) using Cd and Zn in the 0.01 M CaCl<sub>2</sub> extract (equations from table 4.3). Data from both harvests can be explained by the model based on the amounts of Cd and Zn extracted by 0.01 M CaCl<sub>2</sub>. The 95% confidence interval roughly corresponds to a relative error of a factor of 2 on a linear scale.

Table 4.3. Regression coefficients of the CaCl<sub>2</sub>-model and the HNO<sub>3</sub>-model describing Cd accumulation in the grains of 12 rice cultivars used in this study.

Cultivar	Family	CaCl <sub>2</sub> -model (model 1)					HNO <sub>3</sub> -model (model 2)					
		Intercept	Cd-CaCl <sub>2</sub>	Zn-CaCl <sub>2</sub>	R <sup>2</sup>	Se (Y-est)	Intercept	Cd- HNO <sub>3</sub>	pH	CEC	R <sup>2</sup>	Se (Y-est)
1	J	0.54	0.78	-0.18	0.86	0.23	1.44	0.71	-0.24	-0.51	0.80	0.28
2	J	0.61	0.78	-0.28	0.88	0.21	0.74	0.70	-0.13	-0.43	0.81	0.25
3	J	0.58	0.82	-0.27	0.87	0.23	1.06	0.75	-0.22	-0.37	0.83	0.26
4	J	0.67	0.88	-0.32	0.88	0.22	0.94	0.79	-0.22	-0.26 <sup>1</sup>	0.82	0.28
5	J	0.51	0.79	-0.24	0.89	0.21	1.36	0.74	-0.22	-0.60	0.85	0.24
6	I	0.84	0.73	-0.27	0.73	0.32	1.32	0.71	-0.18	-0.44	0.69	0.34
7	J	0.72	0.86	-0.32	0.89	0.21	0.63 <sup>1</sup>	0.79	-0.13	-0.37	0.85	0.24
8	J	0.59	0.83	-0.36	0.87	0.22	0.85	0.72	-0.17	-0.41	0.79	0.28
9	I	0.75	0.65	-0.24	0.66	0.30	0.94	0.62	-0.15	-0.24 <sup>1</sup>	0.68	0.30
10	I	1.03	0.80	-0.29	0.77	0.32	1.33	0.78	-0.19	-0.25 <sup>1</sup>	0.79	0.30
11	I	1.06	0.89	-0.37	0.76	0.34	1.11	0.86	-0.16	-0.31	0.78	0.33
12	J	0.69	0.85	-0.28	0.82	0.23	0.75	0.74	-0.13	-0.53	0.79	0.25
All	J	0.60	0.82	-0.28	0.86	0.23	0.97	0.74	-0.18	-0.43	0.81	0.27
All	I	0.94	0.78	-0.30	0.73	0.33	1.20	0.76	-0.17	-0.32	0.74	0.33

<sup>1</sup>Significance of the coefficient exceeds 5%, all other coefficients are significant at p < 0.001.




Figure 4.4. Measured versus predicted levels of Cd in rice for the first harvest and the second harvest. The dotted lines are equal to the predicted value  $\pm 2 \cdot se(Y\text{-est})$ , which is approximately equal to the 95% confidence interval of the mean.

#### 4.5 Effect of time of harvest on the uptake of Cd by rice

The rice grain samples from this study were collected during four consecutive harvests in 2005 and 2006. Differences between years proved to be non-significant, but significant ( $P < 0.001$ ) differences between the two harvests obtained within one year were observed as is illustrated by the regression coefficients for each harvest (table 4.4).

Table 4.4 Regression coefficients of the  $\text{CaCl}_2$ -model and the  $\text{HNO}_3$ -model describing Cd accumulation in the grains of all Japonica cultivars or of all Indica cultivars grouped together at the first (May) and at the second harvest (November)

Family	Harvest (month)	$\text{CaCl}_2$ -model					
		Intercept	Cd- $\text{CaCl}_2$	Zn- $\text{CaCl}_2$	$R^2$	$se(Y\text{-est})$	
Japonica	5	0.55	0.83	-0.27	85.4	0.22	
Japonica	11	0.68	0.81	-0.30	89.2	0.21	
Indica	5	0.78	0.79	-0.25	85.3	0.23	
Indica	11	1.24	0.85	-0.33	74.5	0.34	
		$\text{HNO}_3$ model					
		Intercept	Cd- $\text{HNO}_3$	pH	CEC <sup>1</sup>	$R^2$	$se(Y\text{-est})$
Japonica	5	1.04	0.70	-0.16	-0.71	81.2	0.25
Japonica	11	1.04	0.80	-0.24	-0.13	87.3	0.23
Indica	5	1.71	0.69	-0.24	-0.57	80.8	0.27
Indica	11	1.32	0.89	-0.23	ns	81.2	0.29

<sup>1</sup>CEC = cation exchange capacity measured at pH 7

Cd concentrations in rice grains were, on average, higher in at the second harvest compared to those of the first harvest ( $P < 0.001$ ), despite the wide range in Cd levels in rice grains among the rice cultivars.

The differences between the first and second harvest are illustrated in figure 4.5 for paddy fields LK2 and PDE; the LK2 field can be considered as non-polluted because total Cd concentration is lower than the Taiwanese background level of  $0.4 \text{ mg kg}^{-1}$ , while the PDE field is heavily polluted with total Cd concentration up to  $30 \text{ mg kg}^{-1}$ . For both fields, average Cd concentration in rice grains of Japonica cultivars was 1.6 to 2.4 times higher at the second harvest compared to those at the first harvest. For the Indica cultivars, average Cd levels were 1.8 to 2.5 times higher at harvest 2 compared to those at harvest 1. However, this trend was not observed for all rice cultivars. At the LK2 field, for example, Cd uptake by cultivars 2 and 5 were not different between both harvests. In the PDE field, Cd accumulation in rice grains of cultivar 11 was even lower at the second harvest compared to Cd accumulation of this cultivar at the first harvest.




Figure 4.5. Difference in Cd concentration in rice grains between the first harvest (May) and the second harvest (November) at two paddy fields with contrasting total Cd concentration. The LK2 field has a relatively low total Cd concentration in soil whereas the PDE field has a relatively high total Cd concentration. Note the difference in the scale on the Y-axis. No results are shown for rice cultivars 9 and 12 since they were not cropped twice each year.

Although it is difficult to explain differences between harvests without additional data on factors like weather and irrigation data, the observed increase in Cd-uptake at time of harvest two largely coincides with an increase in the available Cd pool which was observed as well. The increase in the available Cd-pool could not be explained from the soil data, but it is likely that differences in temperature, rainfall or the duration of the drainage period of the soil prior to the harvest contribute to this difference. It does show, however, that the available pool for Cd throughout the year can vary which has an effect on the quality of rice originating from the same location. To evaluate to what extent the quality of rice fails to meet the FQS at either H1 or H2, data from H1 and H2 from each sampling plot (1442 combinations) were paired. Rice samples from 133 plots (9.2%) exceeded the Taiwan food standard at H2 but not at H1. Since the available Cd levels at H2 generally exceeded those in samples taken at H1, the reverse effect was almost not observed; only 18 samples out of all



1442 pairs (1.2%) exceeded the 0.4 mg kg<sup>-1</sup> standard for rice at H1 where they remained below the 0.4 mg kg<sup>-1</sup> level at H2. The remainder of the samples either was always below (786 plots, 54.5%) or above (505 plots, 35%) the 0.4 mg kg<sup>-1</sup> standard for rice.

Our findings reported here both support and contradict recent findings from a comparable field study conducted by Simmons et al. (2008) who found that, in agreement with our data, uptake of Cd by rice could be predicted well by the amount of Cd measured in a dilute CaCl<sub>2</sub> extract (0.1 M). However, the relation between Cd in rice (2 cultivars only) and Cd in the CaCl<sub>2</sub> extract was found only using field-moist soil samples. After drying, Simmons et al. (2008) did not find significant relationships between Cd-CaCl<sub>2</sub> and Cd in rice grains. Apparently, air-drying altered conditions in the soil samples to such an extent that the availability of Cd was affected when compared to field-moist soils in the study by Simmons et al. (2008). Whether this difference between air-dried and field moist soils is due to the higher ionic strength used by Simmons et al. (2008) or other factors (including soil type and the length of the draining period previous to harvest) remains unclear.



## 5 Assessment of soil tests and soil-plant models to evaluate soil quality of paddy fields in relation to rice grain quality

### 5.1 Evaluation of the suitability of standards and models to test soils

The main goal of SQS is to assess whether a soil can be used safely for the production of specific food crops. Here we evaluate whether present SQS in Taiwan are protective enough to ensure that soils that meet the SQS indeed are suitable for rice cropping. Both the SQS of  $5 \text{ mg kg}^{-1}$  (AR) and the proposed SQS of  $2 \text{ mg kg}^{-1}$  (0.1 M HCl) are used for this evaluation. For each soil sample, measured Cd concentrations in soil with AR or 0.1 M HCl are compared to the aforementioned SQS and based on the outcome of this comparison, the soil is either classified as suitable (soil-Cd < SQS; category A and C) or unsuitable (soil-Cd > SQS; category B and D). Subsequently, measured Cd rice concentrations grown on the same plot are compared to the FQS. This determines whether the soil test was correct (category A and D) or incorrect (category B and C). The FQS used here include the WHO standard ( $0.2 \text{ mg kg}^{-1}$ ) and the standard used in Japan and Taiwan ( $0.4 \text{ mg kg}^{-1}$ ). The outcome of this evaluation can be summarized as:

IF(soil-Cd < SQS)AND(Cd-rice grains < FQS):	Category A (correct, soil = suitable)
IF(soil-Cd > SQS)AND(Cd-rice grains < FQS):	Category B (false negative)
IF(soil-Cd < SQS)AND(Cd-rice grains > FQS):	Category C (false positive)
IF(soil-Cd > SQS)AND(Cd-rice grains > FQS):	Category D (correct, soil $\neq$ suitable)

The  $\text{CaCl}_2$ -model and the  $\text{HNO}_3$ -model are used to perform the same evaluation. Both models are applied to all paired soil – rice samples to predict the Cd level in rice grains. The predicted Cd rice levels are subsequently compared to the FQS. Predicted Cd rice levels are either lower or higher than the FQS and a combination of model predictions and measured Cd rice levels are used to classify all samples as A, B, C, or D:

IF(Cd-rice grains model < FQS) AND(Cd-rice grains < FQS):	Category A (correct, soil = suitable)
IF(Cd-rice grains model > FQS)AND(Cd-rice grains < FQS):	Category B (false negative)
IF(Cd-rice grains model < FQS)AND(Cd-rice grains > FQS):	Category C (false positive)
IF(Cd-rice grains model > FQS)AND(Cd-rice grains > FQS):	Category D (correct, soil $\neq$ suitable)

In summary, samples in category A and D are predicted correctly, whereas samples in category B and C are predicted incorrectly. False positives (C) are particularly undesirable because soil tests or soil-plant models suggest that rice can be grown safely whereas measured Cd rice levels exceed the FQS. False negatives are wrongly classified as unsuitable for rice cropping but rice samples do meet the FQS after all.

Sample classifications (A, B, C, and D) of measured Cd concentration in rice grains and those based on soil tests or soil-plant models are summarized in table 5.1.

*Table 5.1 Overview of the assessment of soil quality using either soil tests or the CaCl<sub>2</sub>-model and the HNO<sub>3</sub>-model to predict the quality of rice grains. Soil tests used are Aqua Regia (AR) and HCl with soil quality standards of 5 and 2 mg kg<sup>-1</sup>, respectively. Food quality standard (FQS) of 0.2 mg kg<sup>-1</sup> from the WHO and the Japanese and Taiwanese FQS of 0.4 mg kg<sup>-1</sup> are used. Model-I or model-A refers to whether the model was derived for individual (I) cultivars or for all (A) cultivars of Japonica or Indica species grouped together.*

Method	% classified as A,B,C,D FQS: 0.2 mg kg <sup>-1</sup>				% classified as A,B,C,D FQS: 0.4 mg kg <sup>-1</sup>			
	A	B	C	D	A	B	C	D
All cultivars together								
Rice grain data	41.2	-	-	58.8	55.8	-	-	44.2
AR	41.2	0.0	33.3	25.5	55.6	0.3	18.9	25.2
HCl	40.7	0.5	23.7	35.1	55.0	0.9	9.4	34.7
CaCl <sub>2</sub> -model-I	35.5	5.7	7.8	51.0	53.7	2.1	4.3	39.8
CaCl <sub>2</sub> -model-A	35.2	6.0	7.9	50.9	53.5	2.4	4.2	39.9
HNO <sub>3</sub> -model-I	35.2	6.1	7.6	51.2	53.1	2.8	4.4	39.7
HNO <sub>3</sub> -model-A	35.6	5.6	7.5	51.3	52.6	3.3	4.7	39.4
Data Japonica species only								
Rice grain data	51.4	-	-	48.6	61.7	-	-	38.3
AR	51.4	0.0	22.5	26.1	61.4	0.4	12.5	25.8
HCl	50.7	0.7	12.7	35.9	60.5	1.2	2.8	35.4
CaCl <sub>2</sub> -model-I	48.3	3.1	8.7	39.9	61.1	0.7	1.2	37.0
CaCl <sub>2</sub> -model-A	48.3	3.1	8.8	39.8	61.1	0.6	1.3	37.0
HNO <sub>3</sub> -model-I	48.7	2.7	8.1	40.5	61.2	0.6	1.3	37.0
HNO <sub>3</sub> -model-A	49.0	2.4	8.2	40.4	61.2	0.6	1.3	37.0
Data Indica species only								
Rice grain data	21.4	-	-	78.6	46.0	-	-	54.0
AR	21.4	0.0	56.5	22.1	46.0	0.0	31.9	22.1
HCl	21.4	0.0	47.8	30.7	45.8	0.2	23.5	30.5
CaCl <sub>2</sub> -model-I	10.1	11.3	6.5	72.1	40.7	5.3	11.2	42.8
CaCl <sub>2</sub> -model-A	9.3	12.1	6.4	72.2	39.8	6.3	10.8	43.2
HNO <sub>3</sub> -model-I	8.2	13.2	7.0	71.6	38.6	7.4	11.5	42.5
HNO <sub>3</sub> -model-A	8.9	12.5	6.6	72.0	36.9	9.1	12.4	41.6

Between 41% (FQS 0.2 mg kg<sup>-1</sup>) and 56% (FQS 0.4 mg kg<sup>-1</sup>) met the FQS. An ideal soil test would predict these percentages, with 41% and 56% of all samples classified as A versus and 44% and 59% as D. This is not the case as is shown in table 5.1 for both the FQS of 0.2 and 0.4 mg kg<sup>-1</sup>. The SQS of 5 mg kg<sup>-1</sup> for AR is the worst method to test the quality of paddy soils. Between 19% for a FQS of 0.4 mg kg<sup>-1</sup> and 33% for a FQS of 0.2 mg kg<sup>-1</sup> are categorized as C. These soils are classified as suitable for rice cropping whereas in fact Cd concentrations in rice grains exceed the FQS. This confirms that accumulation of Cd in grains of rice grown on soils with a Cd concentration below 5 mg kg<sup>-1</sup> often leads to Cd levels in rice grains that exceed the FQS.

The SQS of 2 mg kg<sup>-1</sup> (0.1 M HCl) performs rather good compared to the SQS of 5 mg kg<sup>-1</sup> based on AR; for the FQS of 0.4 mg kg<sup>-1</sup>, only 9% of the samples is classified as C. However, using the 0.2 mg kg<sup>-1</sup> WHO standard the HCl soil test results in a significant underestimation of Cd in rice grains, and 24% of the samples are classified as C.

Data in table 5.1 illustrate the considerable differences in accumulation of Cd in rice grains between cultivars of Japonica and Indica, because the number of rice grain samples with Cd levels higher than the FQS is much higher for the latter. For both soil tests used here, the number of samples classified into category C is much higher for Indica than for Japonica. The CaCl<sub>2</sub>-model as well as the HNO<sub>3</sub>-model give a close to perfect assessment of the suitability of the soil for rice cropping with only 1% of the samples classified in category C at the FQS of 0.4 mg kg<sup>-1</sup>. At a FQS of 0.2 mg kg<sup>-1</sup>, the number of samples classified into this category increases to 8 to 9%. For Indica cultivars, the number of samples classified into category C increases as compared to Japonica cultivars for both FQS values used here, but it still is much lower than for the soil tests based on AR and HCl. Furthermore, only minor differences were found between the number of samples classified into category C when using the CaCl<sub>2</sub>-model and HNO<sub>3</sub>-model based on individual Japonica and Indica cultivars or when based on all Japonica cultivars or all Indica cultivars grouped together. This demonstrates again the minor differences in the behavior of different rice cultivars of Japonica and of Indica varieties with regard to Cd accumulation in rice grains (table 5.1). Obviously, differences between both varieties are much larger, as has been demonstrated previously.

In summary, the CaCl<sub>2</sub>-model and the HNO<sub>3</sub>-model perform equally well and clearly outperform the soil tests based on AR and HCl test where the number of erroneously classified samples is roughly 50% higher compared to both soil-plant models. When considering the number of chemical soil analyses needed to use the CaCl<sub>2</sub>-model and the HNO<sub>3</sub>-model, the use of the CaCl<sub>2</sub> model seems most attractive because only a single extraction with CaCl<sub>2</sub> is needed to assess the quality of paddy soils for rice cropping whereas the HNO<sub>3</sub>-model requires both an extraction of soil with HNO<sub>3</sub> and measurement of pH and CEC. On the other hand, in many countries data on the reactive metal pool and pH and CEC may be available which then can be used directly without the need for additional analyses.

## 5.2 Alternative soil standards based on soil tests with AR and HCl

The SQS currently used in Taiwan based on AR and as proposed for HCl are insufficient to assess the quality of paddy soils for rice cropping compared to the performance of the soil-plant models (table 5.1). Nevertheless, the use of fixed SQS values in combination with routinely applicable soil tests is very practical. A possible solution to test whether this would be feasible is to lower the SQS based on AR or HCl, and to assess the performance of these soil tests with respect to the number of samples classified into category C. Hence, the number of samples in category C, being the most serious error, was calculated at 'new' SQS based on AR and HCl ranging from 0.5 to 5 mg kg<sup>-1</sup>. In table 5.2, the results of this analysis are shown for both the FQS of 0.2 and 0.4 mg kg<sup>-1</sup>.

Table 5.2 Effect of alternative soil quality standards (SQS in mg kg<sup>-1</sup>) for soil tests like Aqua Regia (AR) and HCl on the number of soil samples incorrectly classified as safe (category C). Current SQS standards are 5 mg kg<sup>-1</sup> for AR and 2 mg kg<sup>-1</sup> for HCl.

% classified in category C at FQS 0.2 mg kg <sup>-1</sup>						
SQS	AR			HCl		
	All	Japonica	Indica	All	Japonica	Indica
0.5	10.2	4.6	21.8	17.2	8.2	36.8
1	19.0	9.5	39.9	20.5	10.1	43.2
2	21.7	11.1	44.9	23.7	12.7	47.8
3	24.7	13.8	48.7	26.4	15.7	49.8
5	33.3	22.5	56.5	38.9	27.9	62.0
% classified in category C at FQS 0.4 mg kg <sup>-1</sup>						
SQS	AR			HCl		
	All	Japonica	Indica	All	Japonica	Indica
0.5	2.5	0.4	6.4	5.3	0.9	14.3
1	6.1	1.1	16.8	7.0	1.1	19.5
2	7.6	1.4	20.6	9.4	2.8	23.5
3	10.5	3.9	24.2	12.2	5.9	25.4
5	18.9	12.5	31.9	24.5	18.0	37.5

Reducing the SQS for AR from 5 to 0.5 mg kg<sup>-1</sup> would decrease the number of soil samples for all rice cultivars in category C from 19 to 2.5% at a FQS of 0.4 mg kg<sup>-1</sup> and from 33 to 10% at a FQS of 0.2 mg kg<sup>-1</sup>. This performance is close to the performance of the CaCl<sub>2</sub>-model and the HNO<sub>3</sub>-model (table 5.1). However, reducing the SQS for AR to 0.5 mg kg<sup>-1</sup> has the side-effect of increasing the number of soil samples incorrectly classified as unsuitable for rice cropping whereas the Cd concentration in rice grains in meets the FQS (category B). The percentage of samples in category B increases to 20% at the SQS of 0.5 mg kg<sup>-1</sup> based on AR. A possible compromise resulting in a clear reduction of samples classified in category C without being too protective is a SQS of 2 mg kg<sup>-1</sup>. At least for the FQS of 0.4 mg kg<sup>-1</sup>, this seems an appropriate level for the Japonica cultivars tested here, because the

number of soil samples classified into category C amounts to 1.1% whereas the number of soil samples classified into category B amounts to 1% as well. For the Indica cultivars, however, there seems to be no suitable level based on AR resulting in an acceptable assessment of soil quality. At the FQS level of  $0.2 \text{ mg kg}^{-1}$  neither AR nor HCl SQS are accurate whereas both the  $\text{CaCl}_2$ -model (or soil test) or the  $\text{HNO}_3$ -model provide reliable alternatives to assess the suitability of the soil for rice cropping.

### 5.3 Conclusions

Uptake of Cd by 12 different rice varieties grown on non- to moderately polluted soils in Taiwan results in a significant number of rice samples that do not meet current food quality standards. Even at soil Cd levels far below current soil quality standards, Cd levels in Japonica and especially Indica species exceed both the WHO and Japanese food quality standard. This stresses the need for alternative testing methods that do consider the availability of Cd rather than the total Cd content of the soil. Multiple regression analysis indeed confirmed that uptake of Cd was highly correlated to the available Cd and Zn pool in soils which was measured in a 0.01 M  $\text{CaCl}_2$  extract. A soil-to-plant regression model accounting for differences in pH and CEC of the soil was equally good at simulating the range of Cd measured in rice. Apparently, the 0.01 M  $\text{CaCl}_2$  extractable Cd pool measured at the time of harvest, when the soil is drained, is a good indicator of the amount of Cd taken up by the plant. This has been suggested by others as well who call this period of drainage the *critical stage* because the filling of the rice grain takes place during this period (Simmons et al., 2008). Despite the fact that the samples in our study were air-dried, there was a highly significant relationship between Cd in the  $\text{CaCl}_2$  extract and Cd levels in both roots and rice which seems to contradict results by Simmons et al. (2008). In the study by Simmons et al. (2008) relations were only obtained when using field-moist soil. Whether this is related to the different behavior or even toxic effects of Cd in extremely polluted soils studied by Simmons et al. (2008) or differences between cultivars, remains unclear. In-situ measurements of the changes in the solubility and speciation of Cd in paddy soils and its effects on Cd uptake by rice plants are needed to further address this discrepancy.

In contrast to fixed soil standards, soil-to-plant models and the solution model were able to accurately identify those soils where food quality standards will be exceeded in rice. Especially at the WHO standard of  $0.2 \text{ mg kg}^{-1}$  both the soil-to-plant and the solution model were still able to discriminate between soils suitable and those not suitable for rice cropping whereas the soil standards based on AR or HCl failed to do so in a large number of samples.

In this paper the model concepts were used merely to assess whether or not rice can be grown on a specific soil. The approach can be used as well to derive soil specific target levels and regional risk maps which is another obvious advantage compared to single standards based on the analysis of the total metal content. Uptake models also allow for the region-specific calculation of exposure of human beings feeding from

specific areas as was shown by Thornton et al. (2008) or Brus et al. (2009). Obviously model parameters applied in these studies and those obtained here are not necessarily valid for all cultivars grown in paddy fields elsewhere, but as such, the approach can be adapted rather easily based on a limited number of soil and plant samples taken from specific sites or regions. This study nevertheless also confirms that differences between cultivars or even during the year can be significant and have to be considered when assessing the suitability of soils for rice cropping.



## 6 Uncertainty of models to predict Cd uptake by rice

### 6.1 Introduction

Soil pollution is a world-wide problem affecting the quality of both the terrestrial and aquatic environment. Transfer of contaminants from soil into the food chain is of growing concern due to rising levels of pollutants in food crops (Zhen et al., 2007; Grant and Sheppard, 2008). Uptake of Cd from soil by vegetables or rice is the starting point of an important exposure pathway for human beings (Muchuweti et al., 2006; Khan et al., 2008; Kobayashi et al., 2008; Franz et al., 2008). To protect the general public from elevated exposure to Cd through ingestion of food, soil quality standards (SQS) have been developed. Ideally, SQS represent relevant upper limits for metals and organic compounds in soils to ensure the production of crops that meet food quality standards set by WHO or individual countries. At present, most SQS are still based on the total contaminant content in soil (Carlson, 2007) although plant uptake depends on the availability in soil rather than the total content (Simmons et al., 2008; Römkens et al., 2009a). SQS based on the total content often do not account for differences in the availability of contaminants in soil and it is, therefore, not surprising to observe that SQS vary greatly across the world. SQS for Cd for example range from less than 1 mg kg<sup>-1</sup> (e.g. China: 0.6 mg kg<sup>-1</sup>; NEPAC, 1996) to more than 10 mg kg<sup>-1</sup> in some EU countries (Carlson, 2007). Across the EU alone, Cd SQS related to intermediate 'warning' levels range from 0.5 mg kg<sup>-1</sup> in Denmark to 20 mg kg<sup>-1</sup> in Germany (Carlson, 2007). This range is mainly due to both different concepts used to link an undesired effect (exposure, ecotoxicology, leaching etc.) to a level in soil, as well as differences in soil type across the EU.

Revisions of frameworks for risk assessment hence should consider the concept of (bio)availability rather than the total metal content (Fairbrother et al., 2007). (Bio)availability of metals like Cd depends on soil properties like acidity (pH) and the content of metal sorbing soil compounds including organic matter and clay content (Brus et al., 2005; Chaudri et al., 2007, Römkens et al., 2009b). The key for improving SQS therefore, is to develop tools that link the chemical availability in soil to levels of Cd in those parts of plants that are consumed by human beings. Both empirical as well as mechanistic models are currently being developed to bridge the gap from chemical speciation in the soil to levels of contaminants in plants. Mechanistic models like the Free Ionic Activity Model (FIAM) applied for Cu (Sauvé et al., 1996), or the Biotic Ligand Model applied to predict toxicity for Cu and Ni (Thakali et al., 2006) need to be improved before application on a field scale. Their main limitation is that both types of models focus on uptake processes by roots and transfer into above ground parts is not yet accounted for. François et al. (2009) demonstrated that empirical models outperformed models based on the predicted free metal activity in the soil solution to predict the Cd content in wheat. Also in studies by Efraymson (2001), Chaudri et al. (2007), Brus et al. (2009), Römkens et al. (2009b) empirical soil to plant transfer models using pH, CEC and the reactive soil Cd content were able to predict the metal content in arable crops including rice rather accurately. Aside from

such transfer models, the chemical availability of Cd in soils measured by dilute salt solutions (e.g. 0.01 M CaCl<sub>2</sub>) appears to be correlated well to Cd levels in rice grain (Simmons et al., 2008, Römkens et al., 2009*b*). Both soil-plant transfer models and the CaCl<sub>2</sub> model were able to identify with sufficient accuracy (> 90%) those soils where Cd levels in rice did not meet the food quality standard (FQS) applied in Taiwan (0.4 mg kg<sup>-1</sup> for brown rice). In contrast to these modeling approaches, classic SQS based on the total metal content failed to identify 40 to 50% of all soils where measured levels of Cd in rice exceeded the FQS (Römkens et al., 2009*a*). Despite the fact that soil-plant relationships have been derived for several crops across the world, few examples exist where such relationships are actually used to derive national or local SQS. Existing examples include the calculation of soil specific standards for Cd in wheat in the Netherlands and Australia (Brus et al., 2005; McLaughlin et al., 2006) and for Pb in wheat in the UK (Chaudri et al., 2007) but applications for important food crops like rice are still lacking.

The aim of this paper is apply soil to plant transfer models as well as the CaCl<sub>2</sub> model to derive SQS for 12 different rice cultivars grown under field conditions in Taiwan (Römkens et al., 2009*a* and *b*). Based on a method described by Brus et al. (2005), model uncertainty is taken into account which allows for the calculation of a 90% or 95% confidence level. To illustrate the possibilities of the approach, soil data from Taiwan are used to construct regional SQS maps. Based on soil pH and CEC, a critical Cd content can be calculated for each cell on the map resulting in a regional risk assessment tool. Such maps can be used by regional planners and farmers to identify potential risk areas related to Cd levels in rice. Obviously the model results presented here are not necessarily valid for all rice growing areas across Asia, but we show that this concept can be adapted easily to be applied elsewhere to obtain more realistic SQS.

## 6.2 Materials and Methods

In 2005 and 2006, more than 3000 samples of soil and rice were taken from 19 different paddy fields across Taiwan. A detailed description of the location of the fields, soil types, methods used and levels of Cd in soil and rice is given by Römkens et al. (2009*a*). Based on the results from the field study, soil to plant relationships for 11 different Japonica or Indica type cultivars were derived according to:

$$^{10}\log[\text{Cd}_{\text{rice grain}}] = \text{Constant} + a \cdot ^{10}\log[\text{Cd-soil}] + b \cdot \text{pH}_{0.01\text{M Ca}} + c \cdot ^{10}\log[\text{CEC}] \quad [6.1]$$

With:

- Cd<sub>rice grain</sub> = measured Cd content in rice grain (mg kg<sup>-1</sup>)
- Cd-soil = Cd content in soil based on extraction by Aqua Regia (AR), 0.43 M HNO<sub>3</sub>, and 0.1 M HCl; all extracts expressed in mg kg<sup>-1</sup>.
- pH<sub>0.01M Ca</sub> = pH measured in an 0.01 M CaCl<sub>2</sub> extract
- CEC = cation exchange capacity (cmol (+) kg<sup>-1</sup>)
- Constant, a, b, c = regression coefficients

For all cultivars results of eq. [6.1] were calculated using Aqua Regia (AR), 0.43 N HNO<sub>3</sub>, and 0.1 M HCl.

Estimates of the Cd content in rice grains were also obtained using a 0.01 M CaCl<sub>2</sub> extract (Römken et al., 2009b):

$$^{10}\log[\text{Cd}_{\text{rice grain}}] = \text{Constant} + a \cdot ^{10}\log[\text{Cd}_{\text{CaCl}_2}] + b \cdot ^{10}\log[\text{Zn}_{\text{CaCl}_2}] \quad [6.2]$$

Where

$\text{Cd}_{\text{CaCl}_2}$  = Cd content in soil measured in 0.01 M CaCl<sub>2</sub> in mg kg<sup>-1</sup>

$\text{Zn}_{\text{CaCl}_2}$  = Zn content in soil measured in 0.01 M CaCl<sub>2</sub> in mg kg<sup>-1</sup>

Coefficients for eq. [6.1] (for HNO<sub>3</sub>) and eq. [6.2] are listed in Römken et al. (2009b) and in appendix 2.

### 6.3 Statistical approach to derive SQS based on soil to plant transfer models

In this study eq. [6.1] and eq. [6.2] are used to derive SQS for Cd in soil based on the relation between Cd in soil and Cd in rice grains. The first model is a Freundlich equation similar to the ones used by Adams et al. (2004), Simmons et al. (2008), and Brus et al. (2008) although the choice of soil properties included can vary. Both models are derived by OLS (Ordinary Least Square fitting) in Genstat version 11.1. The equations are derived for 12 individual cultivars included in the field study as well as for all Japonica (C1, C2, C3, C4, C5, C7, C8, C12) or Indica (C6, C9, C10, C11) combined. Using the appropriate FQS for Cd in rice (WHO: 0.2; Japan/Taiwan: 0.4 mg kg<sup>-1</sup>), a SQS for Cd (depending on pH and CEC) or in the CaCl<sub>2</sub> extract can be derived after rearranging equation [6.1] and [6.2] and back transformation to a linear scale. Obviously, models that relate the Cd content in rice to that in soil or a soil extract are not perfect. In fact, the SQS derived from equation [6.1] or [6.2] by definition equals the P50 which means that the predicted Cd content in rice at the calculated SQS has a chance of 50% to exceed the FQS. Depending on the quality of the model, the distribution of the predicted levels can be narrow (good model) or wide. To correct for model uncertainty in order to reduce errors in the prediction of the suitability of soils, probabilistic SQS can be derived based on eq. [6.1] (Brus and Jansen, 2004; Brus et al., 2005). Based on the model uncertainty, the P90 and P95 can be calculated. At the P90 (or P95) the chance that the rice still exceeds the food quality criteria is less than 10% (P90) or 5% (P95).

## 6.4 Results

### 6.4.1 Derivation of soil plant transfer models for different extracts

Both AR and HCl are, at present, used in various countries including Taiwan to

evaluate the quality of soils in relation to land use. Elsewhere, also 0.43 N HNO<sub>3</sub> is used as a measure of the reactive metal content in soils. Coefficients of soil plant transfer models (Constant, *a*, *b*, and *c*) for HNO<sub>3</sub> were listed in Römken et al (2009a); in tables 6.1 and 6.2, coefficients of eq. [6.1] for AR and HCl are listed. The coefficients for the HNO<sub>3</sub> model are included for comparison. The coefficients in table 6.1 are for either Japonica or Indica species together, whereas table 6.2 lists the coefficients for the 12 individual cultivars.

Table 6.1. Coefficients of the soil to plant transfer model (eq. 6.1) for Japonica and Indica for models using Aqua Regia (AR), 0.43 N HNO<sub>3</sub> and 0.1 M HCl.

		Model coefficients					
	extract	Constant	Cd	pH	CEC	R <sup>2</sup>	Se-Yest
Japonica	AR	0.93 ± 0.07	0.80 ± 0.01	-0.19 ± 0.01	-0.44 ± 0.04	79.8	0.276
	HNO <sub>3</sub>	0.98 ± 0.07	0.74 ± 0.01	-0.18 ± 0.01	-0.43 ± 0.04	81.4	0.265
	HCl	1.01 ± 0.07	0.73 ± 0.01	-0.18 ± 0.01	-0.46 ± 0.04	80.8	0.269
Indica	AR	1.17 ± 0.10	0.83 ± 0.02	-0.20 ± 0.02	-0.30 ± 0.06	72.9	0.333
	HNO <sub>3</sub>	1.20 ± 0.10	0.76 ± 0.02	-0.17 ± 0.02	-0.32 ± 0.05	73.9	0.326
	HCl	1.25 ± 0.10	0.75 ± 0.02	-0.18 ± 0.02	-0.33 ± 0.05	73.6	0.328

As was shown for HNO<sub>3</sub> previously by Römken et al. (2009a) results for Japonica and Indica for AR and HCl are quite good when using in equation [6.1]. An extraction using HNO<sub>3</sub> proved to be the most accurate but differences between HNO<sub>3</sub> and HCl are small as is indicated by the coefficients of eq. [6.1] in table 6.1. This is not surprising since both HNO<sub>3</sub> and HCl extract almost identical amounts of Cd from soil (Römken et al., 2009b). Differences in model coefficients between individual cultivars are somewhat more pronounced especially those for CEC (table 6.2) although the difference between Japonica and Indica exceeds that of difference between individual Japonica or Indica species.

Table 6.2. Coefficients of the soil to plant transfer model (eq.6.1) for individual Japonica (J) and Indica (I) cultivars for Aqua Regia (AR), and 0.1 M HCl.

Cult.	Family	Extracts used to determine the metal content											
		AR						0.1 M HCl					
		Constant	<i>a</i> AR	<i>b</i> pH	<i>c</i> CEC	R <sup>2</sup>	Se	Constant	<i>a</i> HCl	<i>b</i> pH	<i>c</i> CEC	R <sup>2</sup>	se (Y-est)
1	J	1.37	0.76	-0.25	-0.53	79.4	0.2	1.38	0.67	-0.23	-0.50	77.9	0.29
2	J	0.63	0.75	-0.13	-0.42	77.1	0.2	0.75	0.70	-0.12	-0.46	80.0	0.26
3	J	0.92	0.82	-0.20	-0.43	81.4	0.2	1.04	0.74	-0.20	-0.39	82.6	0.26
4	J	0.89	0.84	-0.23	-0.26*	80.0	0.2	0.94	0.79	-0.21	-0.30	81.9	0.27
5	J	1.32	0.77	-0.24	-0.57	82.7	0.2	1.44	0.74	-0.23	-0.63	84.6	0.25
6	I	1.33	0.76	-0.21	-0.38	66.3	0.3	1.44	0.70	-0.20	-0.42	69.1	0.34
7	J	0.54*	0.84	-0.14	-0.32	83.2	0.2	0.73	0.79	-0.13	-0.41	84.8	0.24
8	J	0.94	0.78	-0.20	-0.46	78.5	0.2	0.92	0.72	-0.17	-0.46	79.1	0.28
9	I	0.95	0.68	-0.18	-0.22**	68.1	0.3	0.98	0.61	-0.15	-0.27*	67.6	0.30
10	I	1.34	0.86	-0.23	-0.23*	77.8	0.3	1.36	0.79	-0.20	-0.26**	78.9	0.30
11	I	0.97	0.96	-0.17	-0.32	78.3	0.3	1.11	0.86	-0.16	-0.33	77.9	0.33
12	J	0.76	0.81	-0.15	-0.54	76.8	0.2	0.86	0.71	-0.13	-0.58	77.7	0.25

t value < 0.05; \*\* t value < 0.1; all other parameters t < 0.001

### 6.4.2 Soil quality standards (SQS) for individual cultivars

One of the key features using soil to crop models is the derivation of cultivar specific critical SQS for Cd. Figure 6.1 illustrates the differences in P50 levels obtained for all 12 cultivars included in this study at pH 5 (left panel) and pH 7 (right panel) for a sandy soil (open bars) and clay soil (closed bars). Figure 6.1 clearly reveals the elevated uptake of Cd by Indica-type cultivars. SQS for Indica range from less than  $0.3 \text{ mg kg}^{-1}$  in sandy soils at pH 5 to approximately  $1.0 \text{ mg kg}^{-1}$  at pH 7 in clay soils. Figure 6.1 also illustrates that the use of the models based on all data from Japonica or Indica as presented in table 6.1 gives a good approximation of the average SQS for either Japonica or Indica. For nationwide applications it seems justified to use the overall Japonica or Indica model rather than models for individual cultivars. For local applications on the other hand, specific models can be used if the cultivars match the ones included in this study.




Figure 6.1. P50 SQS for individual Japonica and Indica-type cultivars at pH 5 (left hand panel) and pH7 (right hand panel) in a sandy ( $\text{CEC} = 5 \text{ cmol kg}^{-1}$ ) and clay soil ( $\text{CEC} = 20 \text{ cmol kg}^{-1}$ ).

#### **Impact of model uncertainty on the SQS**

In table 6.3, calculated values of the P50, P90, and P95 SQS for Cd are listed for a sandy soil ( $\text{CEC} = 5 \text{ cmol kg}^{-1}$ ) and clay soil ( $\text{CEC} = 20 \text{ cmol kg}^{-1}$ ). Here, eq. [6.1] was used based on Aqua Regia extractable metals in soil. Soil pH ranged from 4.5 to 7. The results are based on the combined equations for either Japonica or Indica species.

Table 6.3. Calculated P50, P90 and P95 SQS (*Aqua Regia* in mg kg<sup>-1</sup>) based on the average models for either Japonica or Indica for a sandy soil (CEC = 5 cmol<sup>+</sup> kg<sup>-1</sup>) and a clay soil (CEC = 20 cmol<sup>+</sup> kg<sup>-1</sup>)

<i>Japonica</i>						
pH	Sand			Clay		
	P50	P90	P95	P50	P90	P95
4.5	0.73	0.37	0.26	1.46	0.75	0.53
5	0.97	0.50	0.35	1.94	0.99	0.70
5.5	1.28	0.65	0.46	2.56	1.31	0.92
6	1.69	0.87	0.61	3.39	1.73	1.22
6.5	2.24	1.15	0.81	4.49	2.29	1.61
7	2.97	1.51	1.06	5.94	3.03	2.13

<i>Indica</i>						
pH	Sand			Clay		
	P50	P90	P95	P50	P90	P95
4.5	0.28	0.13	0.09	0.44	0.20	0.13
5	0.37	0.17	0.11	0.58	0.26	0.18
5.5	0.49	0.22	0.15	0.76	0.35	0.23
6	0.64	0.29	0.19	1.00	0.46	0.30
6.5	0.84	0.38	0.25	1.31	0.60	0.40
7	1.10	0.50	0.33	1.73	0.79	0.52

Critical levels (P50) for Cd in soil range from more than 5 mg kg<sup>-1</sup> at pH 7 for Japonica to less than 0.3 mg kg<sup>-1</sup> for Indica in sandy soils at pH 4.5. This large range clearly shows that the impact of soil acidity and, to a lesser extent, soil texture has a profound impact on the quality of soil regarding the Cd content of rice. As such it confirms conclusions from previous studies (Simmons et al., 2008; Römken et al., 2009a) suggesting that fixed soil standards for a wide range of soils are inappropriate. Also, it clearly shows that the SQS of 5 mg kg<sup>-1</sup> currently applied in Taiwan is not protective enough and only suitable for clay soils at near neutral pH. Results from other field and pot trials indeed confirm that for vegetables, wheat and rice alike, SQS can be below 0.5 mg kg<sup>-1</sup>. Shentu et al. (2008) obtained critical soil Cd levels ranging from 0.1 to 0.5 mg kg<sup>-1</sup> for vegetables using the total soil Cd content. Brus et al. (2005) obtained critical soil levels for Cd ranging from 0.5 to 1.5 mg kg<sup>-1</sup> for wheat grown under moderate climate conditions. For field-grown wheat, critical Cd levels in soil ranged from 0.3 mg kg<sup>-1</sup> at pH 4.5 in a sandy soil to 1.0 mg kg<sup>-1</sup> in clay soils at pH 7 (McLaughlin et al., 2006). These studies all suggest that differences between soils which can be characterized by differences in pH and CEC need to be considered to obtain realistic SQS.

For Japonica a reduction of the SQS to 2 mg kg<sup>-1</sup> seems appropriate in most soils as long as the soil pH is higher than 6. For Indica species on the other hand a fixed SQS of 0.5 to 1.0 is needed in soils with pH 6 or higher. For both Japonica and Indica, both CEC and pH have a pronounced effect on the level of the SQS although the impact of pH exceeds that of CEC as is illustrated in figure 6.2.




Figure 6.2. Effect of pH and CEC on the SQS (P50) for Japonica (left) and Indica (right) based on eq. 1 using AR.

The effect of CEC is more pronounced at high pH where SQS at pH 6.5 increase from 1.0 in soils with a CEC below 5  $\text{cmol kg}^{-1}$  to 4.0 in soils with a CEC of 30  $\text{cmol kg}^{-1}$ .

Table 6.3 clearly illustrates the impact of model uncertainty on the P90 and P95 values. Based on the soil extraction by AR, P90 values for the combined models are approximately half the value of P50 values. Levels of the P95 SQS range from 0.3 to 0.4 times the value of the P50. For Indica this means that reducing the error in the SQS to 5% results in an SQS value of less than 0.1  $\text{mg kg}^{-1}$  at pH 4.5 to about 0.3  $\text{mg kg}^{-1}$  at pH 7. This is equivalent to the natural background level of Cd in most soils around the world and illustrates the fact that for some crops including brown rice, uptake of Cd can be high even at low Cd levels in soil. This is in line with findings on Cd uptake by vegetables in non-tropical areas like Belgium and the Netherlands where Cd levels in leek and scorzonera exceeded current EU food quality standards (Rietra and Römken, 2007) in soils below generally accepted background levels (Brus et al., 2009). Although the low levels of the P95 SQS illustrate that levels of Cd in crops can exceed the FQS, P50 SQS are most commonly used in relation to crop protection (e.g. Shintu et al., 2008). Although the use of a P95, SQS level warrants the production of crops that meet the FQS in 19 out of 20 cases, application on a national level will result in a large number of sites that are incorrectly qualified as unsuitable. SQS at or below generally accepted background levels imply that even clean soils are not suitable for crop production. The fact that there is a risk that crops exceed FQS at low soil Cd levels seems unavoidable and is related to specific properties of crops like brown rice and vegetables including leek and scorzonera. P50 SQS levels on the other hand seem more suitable as a first screening level and have proven to be rather successful in identifying those soils where rice indeed did not meet FQS (Römken et al., 2009b).



### ***SQS based on CaCl<sub>2</sub> extracts***

The link between CaCl<sub>2</sub> extractable Cd and Zn on one hand and the levels of Cd in rice grain on the other has been established by various authors (Simmons et al., 2005; Römkens et al., 2009b). This link can be used to derive critical levels in such extracts beyond which the Cd level in rice exceeds the food quality standard. The obvious advantage of such critical limits is that only one soil test has to be performed that serves as indicator for the quality of rice. The study by Römkens et al. (2009b) clearly showed that the available Zn fraction measured in the CaCl<sub>2</sub> extract suppressed the level of Cd in rice. Hence, the critical Cd level depends on the available Zn fraction. In table 6.4 critical limits for Cd in the 0.01 M CaCl<sub>2</sub> extract are listed for Japonica and Indica as a function of the available Zn content. In the database used to derive the soil plant relationships based on CaCl<sub>2</sub>, the 10 to 90 percentile extractable Zn content ranged from 0.3 to 20 mg kg<sup>-1</sup>. This range was used to derive critical Cd levels in soil as listed in table 6.4. Here, average models for all Japonica or Indica species were used. Figure 6.3 shows the effect of Zn on the critical (P50) level of Cd for all individual 12 cultivars. Critical levels for Japonica exceed those of Indica by a factor of approximately 4 (table 6.4) and range from less than 0.05 mg kg<sup>-1</sup> at very low Zn levels in soil to more than 0.15 mg kg<sup>-1</sup> in soils with 20 mg kg<sup>-1</sup> extractable Zn. This clearly illustrates the considerable influence of Zn on the uptake of Cd by rice studied here. If the model without considering the impact of Zn is used to calculate critical Cd levels in 0.01 M CaCl<sub>2</sub>, the fixed critical Cd content is equal to 0.10 mg kg<sup>-1</sup> for Japonica and 0.027 mg kg<sup>-1</sup> for Indica. This is very close to the values reported in table 6.4 based on the average Zn content in all samples from the database which ranges from 3.6 mg kg<sup>-1</sup> for the plots used to grow Japonica rice and 3.9 mg kg<sup>-1</sup> for Indica. When used to evaluate the suitability of soils for rice cropping, the accuracy of the assessment obviously increases when using the extractable Zn content as well. After all, the local critical limit for Cd including Zn, varies up to 50% of the level of the P50 value compared to the model without Zn. Due to model uncertainty the P90 level for Japonica type rice species is roughly equal to 0.58 times the P50 level. For Indica type cultivars, most models proved to be somewhat less accurate and P90 levels are approximately 0.4 to 0.45 times the P50 level. P95 levels are approximately 0.45 times the P50 level for Japonica and 0.28 times the P50 level for Indica.

Differences between individual Japonica or Indica species (figure 6.4) are less pronounced. Only for cultivar 8 (Japonica) the critical levels seem higher compared to the other Japonica species. For Indica, all species tested here behave more or less similar with only minor differences between individual cultivars. Again, these results suggest that the use of a single model based on either Japonica or Indica will give a rather accurate assessment of the critical limit of Cd in soil as measured by CaCl<sub>2</sub>. As such the use of a single extract has its merits compared to the soil plant relationships, which requires the input of both soil pH and CEC. Especially the latter is more laborious to measure whereas the 0.01 M CaCl<sub>2</sub> is a rather robust method (Houba et al., 2000) which also has been promoted as an appropriate technique to assess the availability of other metals as well (Brand et al., 2009).

Table 6.4 Overview of P50, P90 and P95 levels for the critical Cd content in 0.01 M CaCl<sub>2</sub> (in mg kg<sup>-1</sup>) for Japonica and Indica species.

Zn (mg kg <sup>-1</sup> )	Critical Cd content in 0.01 M CaCl <sub>2</sub> extract					
	Japonica			Indica		
	P50	P90	P95	P50	P90	P95
0.3	0.040	0.024	0.018	0.012	0.005	0.004
0.5	0.047	0.028	0.021	0.015	0.006	0.004
0.8	0.055	0.032	0.025	0.018	0.008	0.005
1.3	0.065	0.038	0.029	0.021	0.009	0.006
2.0	0.076	0.045	0.034	0.025	0.011	0.007
3.2	0.089	0.052	0.040	0.030	0.013	0.009
5.0	0.104	0.061	0.046	0.036	0.016	0.010
7.9	0.122	0.072	0.054	0.042	0.019	0.012
12.6	0.143	0.084	0.064	0.051	0.022	0.014
20.0	0.168	0.099	0.075	0.060	0.027	0.017
No Zn	0.101	0.046	0.031	0.027	0.008	0.005




Figure 6.4. Critical Cd content in the 0.01 M CaCl<sub>2</sub> extract for 8 Japonica type cultivars (left) and 4 Indica type cultivars within the 10-90% range of the extractable Zn content (0.3 to 20 mg kg<sup>-1</sup>). The figures have the same scale to facilitate comparison between Japonica and Indica.

### 6.4.3 Assessment of soil quality based on the P50, P90 or P95 percentile SQS

As shown in tables 6.3 and 6.4, P90 or even P95 SQS levels are substantially lower compared to the P50 value. One of the aims of the research was to minimize the erroneous assessment of soils using fixed standards. The results shown in Römken et al (2009a) clearly show that soil plant relationships are far more accurate in determining whether a soil is suitable for rice cropping or not. Here we assess whether the P90 or even the P95 should be used in assessing soil quality or that a P50 is suitable as well. To do so, all samples were classified according to the method described by Römken et al. (2009a). In short, soil plant models are used to assess whether the soil is suitable or not comparing the predicted outcome of the model with the actual quality of rice from the same sample. To do so, the calculated P50,

P90, and P95 SQS values for the CaCl<sub>2</sub> and soil model were used to derive regression functions that calculate the P50, P90 and P95 SQS based on either the Zn measured in CaCl<sub>2</sub> or a combination of pH and CEC for the soil –plant model. In table 6.5 the coefficients of these equations are listed. The only difference between the three uncertainty levels is the constant in the equation; all coefficients of variables included are similar.

Table 6.5 Regression functions to calculate P50, P90 and P95 SQS from CEC and pH (soil model) or CaCl<sub>2</sub> extractable Zn (CaCl<sub>2</sub> model).

	Soil plant model					
	<i>Japonica</i>			<i>Indica</i>		
	Constant	CEC	pH	Constant	CEC	pH
P50	-1.85	0.585	0.246	-2.11	0.422	0.226
P90	-2.15	0.585	0.246	-2.47	0.423	0.226
P95	-2.30	0.586	0.246	-2.66	0.423	0.227
	CaCl <sub>2</sub> model					
	<i>Japonica</i>		<i>Indica</i>			
	Constant	Zn	Constant	Zn		
P50	-1.22	0.346	-1.72	0.383		
P90	-1.45	0.346	-2.07	0.383		
P95	-1.58	0.346	-2.26	0.383		

For each sample from the database, the P50, P90 and P95 SQS of Cd in soil (HNO<sub>3</sub> model) or CaCl<sub>2</sub> was calculated. Subsequently the SQS value thus calculated was compared to the actual Cd content of the soil or the Cd content in the CaCl<sub>2</sub> extract. This results in an assessment of the soil quality which then can be validated using the measured Cd content in rice grains. In table 6.6 the results of this assessment are shown listing the number of samples that are qualified correctly, which means that the soil test correctly predicts whether the rice is above or below the FQS, and the samples which are qualified incorrectly. The latter samples include those where the soil test indicates that the rice is of insufficient quality whereas in fact the Cd content in rice meets the FQS and the samples where the rice is predicted to meet the FQS whereas the actual Cd content in rice exceeds the FQS. For comparison this table also includes the same assessment based on the HCl and AR soil test used at present in Taiwan.

The results in table 6.6 show that models are clearly better compared to fixed soils tests although 2 mg kg<sup>-1</sup> HCl test for Japonica seems acceptable. However, using the P90 or even P95 level up to 42% of all Indica samples is incorrectly classified as not suitable which renders large areas as seemingly at risk whereas in fact the quality of rice will meet the FQS.

Table 6.6 Assessment of the accuracy of the P50, P90 and P95 SQS to identify soils unsuitable for rice cropping

		SOLUTION MODEL				SOIL MODEL			
		A	B	C	D	A	B	C	D
P50	All	53.6%	3.3%	4.5%	38.6%	54.5%	2.4%	4.2%	39.0%
	Japonica	61.3%	0.5%	1.1%	37.0%	61.2%	0.6%	1.1%	37.0%
	Indica	36.8%	9.2%	11.9%	42.1%	39.8%	6.2%	10.8%	43.2%
P90	All	46.5%	10.4%	1.4%	41.8%	45.9%	11.0%	1.0%	42.1%
	Japonica	61.1%	0.8%	1.1%	37.0%	59.2%	2.7%	1.0%	37.2%
	Indica	14.7%	31.3%	2.0%	52.1%	17.0%	29.0%	1.0%	53.0%
P95	All	40.6%	16.3%	0.9%	42.2%	38.0%	18.8%	0.6%	42.5%
	Japonica	57.1%	4.7%	1.0%	37.1%	53.7%	8.2%	0.7%	37.4%
	Indica	4.6%	41.4%	0.6%	53.4%	4.1%	41.9%	0.4%	53.6%
		AR soil test				HCl soil test			
		A	B	C	D	A	B	C	D
All		56.7%	0.3%	18.5%	24.6%	56.6%	0.4%	9.3%	33.8%
Japonica		61.5%	0.4%	12.3%	25.7%	61.4%	0.5%	2.7%	35.3%
Indica		46.1%	0.0%	31.8%	22.0%	46.1%	0.0%	23.5%	30.4%

Although both the soil model and the  $\text{CaCl}_2$  model at the P50 level perform much better compared to fixed soil standards, there is still a number of samples (11 to 12% for Indica) where the Cd content in rice exceeds the P50 FQS although models predict it to fall below this standard (category C). Using a P90 or even P95 SQS substantially reduces the samples in category C to less than 2% which means that only one in 50 samples will fail the FQS despite having a soil Cd level below the calculated SQS. However, the obvious disadvantage of P90 and P95 SQS for field applications is the considerable increase in the number of samples incorrectly classified as non-suitable whereas in fact the rice will meet the FQS. At the P95 level, up to 42% of Indica rice samples and 8% of Japonica rice samples is predicted not to meet the FQS at the given soil Cd level or the amount of Cd in the  $\text{CaCl}_2$  extract whereas in fact the data indicate that the rice meets the criteria. Reducing the number of soils that fail to be identified by models as not-safe by increasing the confidence level of the SQS (P90 or P95) invariably leads to an increase in the number of soils qualified as not-suitable whereas in fact they are suitable for rice production. As such P50 SQS levels seem to represent an acceptable compromise to identify soils not suitable for crop production without being too strict. The use of P90 and P95 SQS levels on one hand identifies those soils which are suitable for crop production but it also would erroneously identify a large number of fields as potentially not safe for use.

### Conclusions

- Soil specific SQS vary widely due differences in pH and CEC and range from less than 0.3 for Indica species in sandy soils to more than 5 in clay soils at neutral pH values;

- Model uncertainty leads to 50% (P90) or 65% (P95) reduction of SQS compared to P50;
- On average SQS for Japonica should be reduced at least to 2 mg kg<sup>-1</sup> for clay soils and 0.5 mg kg<sup>-1</sup> for Indica at pH 7 if no correction for pH or CEC is applied;
- Differences between individual cultivars are limited, large differences however exist between Japonica and Indica;
- Application of soil-plant models on a regional scale is easy and can be used as first assessment for potential risks. High variability of Cd in soil however makes regional maps not reliable for application on a local scale (unless local data are available obviously)
- Derivation of a single standard based on CaCl<sub>2</sub> seems promising but few data are available as of now (in contrast to AR/pH/CEC which are available on regional scale);
- Effect of Zn on uptake of Cd is pronounced and should be considered, which is easy due to the fact that both Cd and Zn can be measured in one extract (0.01 M CaCl<sub>2</sub>)
- P90 and especially P95 SQS lead to unacceptable high numbers of samples which are unnecessarily disqualified as not suitable. P50 levels seem to represent an acceptable compromise.



## 7 Tools to assist farmers and policy makers to evaluate soil quality for rice cropping

### 7.1 Introduction

At present, few tools are available that can be used by either farmers or policy makers to evaluate the quality of local fields used for agricultural purposes. As such, the models described in chapter 4 can be used to construct easy to use tools that enable the user to decide on the quality of the soil. This concept which has been developed for similar problems with Cd polluted soils in the Netherlands is based on the relationship between soil properties (including the Cd content) and the Cd content in rice. The user only has to fill in some basic properties including the pH, CEC and Cd content of the soil. Often these data are available and if not, are rather easy to obtain.

Here we present two types of output, the so-called 'look-up table' which shows the suitability of the soil for a specific cultivar in the form of a color-table, and output generated by an Excel file for specific evaluation of various cultivars.

### 7.2 Look-up tables

Look-up tables present an overview of the suitability of a range of soils (that is, a range in the Cd-level in the soil and pH). This concept is based on the soil (HNO<sub>3</sub>) model but can be used for Aqua Regia or HCl as well. Obviously, the parameters used in the model have to be derived for the correct soil analysis (i.e. HNO<sub>3</sub>, HCl or Aqua Regia). For every combination of pH, CEC and Cd content in the soil, the model calculates the level of Cd in the rice, either for the mixed models (Japonica or Indica) or for individual cultivars as was discussed in chapter 4 (table 4.3):

$$\text{Cd-rice} = \text{Intercept} + a \cdot \text{pH} + b \cdot \log \text{CEC} + c \cdot \log[\text{Cd-soil}] \quad [7.1]$$

In Appendix 1 all model coefficients (intercepts, values for a, b and c) are listed for all extracts (Aqua Regia, HNO<sub>3</sub> and HCl). Obviously, care should be taken to use the appropriate set of coefficients when dealing with a specific extract. To make the table flexible, the user can define the range of interest in pH and Cd-level in the soil which the table should reflect.

The results are then plotted in a square matrix as is shown in figure 7.1. Here the following soil properties were used:

pH range: 5 – 7  
Cd-soil range: 0.2 – 2 mg kg<sup>-1</sup> based on HCl

Obviously the table looks different when using other combinations of ranges in pH and Cd-soil. In figure 7.1 the models for Japonica and Indica are used, tables can be constructed for individual cultivars as well. The color scheme in figure 7.1 is chosen as follows to reflect the predicted quality of rice:

- Green fields: combination of pH and Cd-soil will result in rice of sufficient quality, the Cd level will be below the food quality standard (in this case we used a critical level of Cd in rice of 0.4 mg kg<sup>-1</sup> which is currently used in Taiwan)
- Red fields: combination of pH and Cd-soil will results in rice of insufficient quality, the rice will contain more than 0.4 mg kg<sup>-1</sup>
- Yellow fields: this combination results in a Cd level in rice close to the food quality standard used (in this case 0.4 mg kg<sup>-1</sup>). The model uncertainty does not allow for a specific evaluation.

		Japonica											Indica										
		pH range											pH range										
		5	5.2	5.4	5.6	5.8	6	6.2	6.4	6.6	6.8	7	5	5.2	5.4	5.6	5.8	6	6.2	6.4	6.6	6.8	7
Cd-soil	0.2	0.13	0.12	0.11	0.10	0.09	0.08	0.08	0.07	0.07	0.06	0.06	0.21	0.29	0.27	0.24	0.23	0.21	0.19	0.18	0.17	0.15	0.14
	0.4	0.21	0.19	0.17	0.16	0.15	0.14	0.12	0.11	0.11	0.10	0.09	0.50	0.47	0.43	0.40	0.37	0.34	0.31	0.29	0.27	0.25	0.23
	0.6	0.28	0.25	0.23	0.21	0.20	0.18	0.17	0.15	0.14	0.13	0.12	0.68	0.63	0.58	0.53	0.49	0.46	0.42	0.39	0.36	0.33	0.31
	0.7	0.34	0.31	0.29	0.26	0.24	0.22	0.20	0.19	0.17	0.16	0.15	0.84	0.77	0.71	0.66	0.61	0.56	0.52	0.48	0.44	0.41	0.38
	0.9	0.40	0.37	0.34	0.31	0.28	0.26	0.24	0.22	0.20	0.19	0.17	0.99	0.91	0.84	0.78	0.72	0.66	0.61	0.57	0.52	0.48	0.45
	1.1	0.46	0.42	0.39	0.35	0.33	0.30	0.27	0.25	0.23	0.21	0.20	1.13	1.04	0.96	0.89	0.82	0.76	0.70	0.65	0.60	0.56	0.51
	1.3	0.51	0.47	0.43	0.40	0.36	0.33	0.31	0.28	0.26	0.24	0.22	1.27	1.17	1.08	1.00	0.92	0.85	0.79	0.73	0.67	0.62	0.58
	1.5	0.56	0.52	0.48	0.44	0.40	0.37	0.34	0.31	0.29	0.26	0.24	1.40	1.29	1.20	1.11	1.02	0.94	0.87	0.81	0.74	0.69	0.64
	1.6	0.61	0.56	0.52	0.48	0.44	0.40	0.37	0.34	0.31	0.29	0.26	1.53	1.41	1.31	1.21	1.12	1.03	0.95	0.88	0.81	0.75	0.69
	1.8	0.66	0.61	0.56	0.51	0.47	0.43	0.40	0.37	0.34	0.31	0.29	1.66	1.53	1.41	1.31	1.21	1.12	1.03	0.95	0.88	0.81	0.75
	2.0	0.71	0.65	0.60	0.55	0.51	0.47	0.43	0.39	0.36	0.33	0.31	1.78	1.64	1.52	1.40	1.30	1.20	1.11	1.02	0.95	0.87	0.81

Figure 7.1 'Look-up table' for Japonica (left) and Indica (right) for soils with pH levels from 5 – 7 and HCl extractable Cd soil from 0.2 to 2.0 mg kg<sup>-1</sup> (CEC of the soil was set at 10 cmol<sup>+</sup> kg<sup>-1</sup>).

Figure 7.1 clearly illustrates the considerable difference between Japonica and Indica in terms of Cd accumulation and the importance of pH when dealing with risk assessment. Japonica species can be cultivated safely up to 1 mg kg<sup>-1</sup> in near neutral soil. Below pH 6 the risk that Cd levels in rice exceed the 0.4 food quality limit increases. Indica species on the other hand can be cultivated safely only in soils with a Cd content of less than 0.4 mg kg<sup>-1</sup> or even 0.2 below pH 6. This shows that, in line with the results presented in chapter 4 that in polluted soils (Cd-soil > 1 or 2 mg kg<sup>-1</sup>) Indica species are not suitable. This again also confirms that existing soil quality guidelines used at present in Taiwan (5 mg kg<sup>-1</sup> for Aqua Regia and 2 mg kg<sup>-1</sup> for HCl) are too high to safeguard the quality of rice.

To illustrate this, look-up tables for soils with a Cd content ranging from 1 to 5 mg kg<sup>-1</sup> are shown (based on extraction with Aqua Regia).






Figure 7.2 'Look-up table' for Japonica (left) and Indica (right) for soils with pH levels from 5 – 7 and Aqua Regia extractable Cd soil from 1 to 5 mg kg<sup>-1</sup> (CEC of the soil was set at 10 cmol+ kg<sup>-1</sup>).

Figure 7.2 confirms that Indica species are unsuitable for use on these soils, all combinations of Cd-soil and pH will most likely result in rice which contains more than 0.4 mg kg<sup>-1</sup>. For Japonica on the other hand, the look-up table shows that at pH levels higher than 6, the quality of Japonica rice probably is sufficient in soils with Cd levels below 2 mg kg<sup>-1</sup>. Between pH 5 and 6, the soils should contain less than 1.5 mg kg<sup>-1</sup> (approximately) to be suitable for rice cropping. Again, the influence of pH is pronounced and should be taken into account to make safe decisions on whether or not to grow rice on this soil.

The model used here also includes the CEC of the soil. In soils with a low CEC such as sandy soils, Cd uptake is higher compared to uptake from clay soils with a much higher CEC. This is illustrated in figure 7.3 for Japonica species. Here, the effect of CEC is illustrated for a soil with a CEC of 7 cmol(+) kg<sup>-1</sup> which is roughly equivalent to the soils from the HC fields versus a soil with a CEC of 20 cmol(+) kg<sup>-1</sup> which was found in the HM3 soil (table 2.1)




Figure 7.3 'Look-up table' for Japonica for soils with pH levels from 5 – 7 and Aqua Regia extractable Cd soil from 1 to 5 mg kg<sup>-1</sup>. CEC of the soil was set at 7 (left) and 20 (right) cmol+ kg<sup>-1</sup>.

The data in figure 7.3 illustrate that in clay soils (CEC = 20), Japonica species can be grown safely up to Cd levels of 3 mg kg<sup>-1</sup> provided that the soil pH is close to 7. In sandy soils on the other hand an upper limit of 2 mg kg<sup>-1</sup> at pH 7 should not be exceeded.

Figure 7.3 also shows a second use of look-up table which is the evaluation of soil management. Lime application for example to increase soil pH can be an option to reduce the uptake of Cd from soil. Increasing soil pH from values around 5 to 6 or higher in a clay soil (right table in figure 7.3) that contains  $2 \text{ mg kg}^{-1}$  Cd would reduce the uptake from soil to such an extent that rice will meet the food quality criteria. This on the other hand also indicates that small increases in soil pH are usually not sufficient to improve the quality of rice. Especially in acid soils, an increase in soil pH from levels below 5 to more than 6 would require a large amount of lime which is not a suitable option. Look-up tables nevertheless can be used to assess whether or not correction of soil pH is an option to reduce Cd uptake by the cultivar of choice.

The examples included here are merely meant to illustrate some of the differences between cultivars and impact of soil properties (and changes therein) on the Cd uptake by rice. The user can define his or her boundary conditions in terms of cultivars used, pH, soil Cd levels and CEC. Here we presented look-up tables based on the models for Japonica and Indica (all data combined for the two species) but such tables can be constructed for individual cultivars as well using the coefficients listed in Appendix 1. In Appendix 2 an example is shown for all 12 cultivars studied here. As such differences between the individual Japonica (or Indica) cultivars are small compared to the large difference between Japonica and Indica cultivars.

### 7.3 Soil specific evaluation

Instead of look-up tables, a farmer is merely interested if his or her soil is suitable for a specific cultivar. In this case, there is no need to construct a look-up table from which the soil of interest can be evaluated. In this case a soil-specific evaluation can be performed indicating the suitability of the local soil for one or more varieties available in the database.

The farmer has to fill in the soil properties required including:

1. soil metal content in  $\text{mg kg}^{-1}$
2. extraction method (choice of Aqua Regia,  $\text{HNO}_3$ , or HCl)
3. soil pH
4. CEC in  $\text{cmol}(+)\text{kg}^{-1}$

Using the appropriate model coefficients listed in Annex 1, the predicted level of Cd in rice is then reported for each cultivar separately as well as for the Japonica and Indica model as is illustrated in table 7.1

Table 7.1 Site specific evaluation of soil quality

<b>Soil Characteristics (to be filled in)</b>			
Cd-soil		<b>0.63</b>	mg kg <sup>-1</sup>
extract		<b>HNO<sub>3</sub></b>	
pH		<b>5</b>	
CEC		<b>10</b>	cmol(+)kg <sup>-1</sup>
<b>Suitability Assessment of soil for</b>		<b>Cd-rice</b>	
Japonica		0.30	mg kg <sup>-1</sup>
Indica		0.74	mg kg <sup>-1</sup>
<b>Cultivar</b>			
Tainung No.70	Japonica	0.37	mg kg <sup>-1</sup>
Taiken No.8	Japonica	0.33	mg kg <sup>-1</sup>
Tainung No.72	Japonica	0.29	mg kg <sup>-1</sup>
Kaohsiung No.143	Japonica	0.28	mg kg <sup>-1</sup>
Taitung No.30	Japonica	0.31	mg kg <sup>-1</sup>
Tainung Sen No.20	Indica	0.72	mg kg <sup>-1</sup>
Tainung No.71	Japonica	0.30	mg kg <sup>-1</sup>
Tainung No.67	Japonica	0.27	mg kg <sup>-1</sup>
Kaohsiung Sen Yu No.1151	Indica	0.67	mg kg <sup>-1</sup>
Taichung Sen Waxy No.1	Indica	0.89	mg kg <sup>-1</sup>
Taichung Sen No.10	Indica	0.66	mg kg <sup>-1</sup>
Kaohsiung No.144	Japonica	0.27	mg kg <sup>-1</sup>

Table 7.1 illustrates the differences between Japonica and Indica; for the soil given here, Indica species clearly are not suitable. For Japonica species a mixed result can be obtained. Overall, Cd levels in Japonica will be close to the food quality (average 0.3 mg kg<sup>-1</sup>) but in this case cultivar 8 and 12 (Tainung no. 67 and Kaoshiung 144) appear to be more suitable compared to for example cultivar 1 (Tainung no. 70). Again, a safety margin was considered and it can be seen from the coloring of the table that there is a reasonable chance that all Japonica species will meet the 0.4 mg kg<sup>-1</sup> food quality standard. If the farmer has a choice however, it seems best to choose those cultivars which accumulate the least Cd for the soil under consideration.

#### **7.4 Issues to be considered when using average field data**

An important issue that needs to be considered is the heterogeneity of the Cd content of the fields (illustrated in figure 3.3). In the majority of the fields the Cd content decreases across the field with an increasing distance from the inlet. Soil sampling procedures to assess the average quality of the field need to take into account this heterogeneity to avoid erroneous results. Samples taken close to the inlet probably contain (much) more Cd compared to those taken at the opposite end of the field. In contrast to the samples included in this study where each individual sample was treated as a soil sample that represents a soil, the farmer is interested merely in the quality of the rice from the entire field (mixed). This means that the soil sample used to determine the soil quality should reflect this average quality. Considering the large variability encountered here, this requires that samples have to be taken from the entire field before mixing it into one sample that represents the average quality across the field.

## 8 Conclusions and Recommendations

### *Revision of current soil quality standards for agricultural soils*

The data presented in chapter 3 and 4 clearly show that current soil quality standards used to distinguish clean from polluted soils are not protective enough. The soil monitoring value of 5 mg kg<sup>-1</sup> based on Aqua Regia leads to a large number of cases where Cd levels in rice are (far) higher than the allowed 0.4 mg kg<sup>-1</sup> currently used as the food quality standard in Taiwan.

The 2 mg kg<sup>-1</sup> standard based on 0.1M HCl works better for Japonica species but still results in a large number of samples for Indica that contain too much Cd. Alternative soil standards based on either Aqua Regia or HCl should not exceed 2 mg kg<sup>-1</sup> as was shown in chapter 5 (table 5.1 and 5.2). Such tests then are only suitable for Japonica species and not for Indica. Even a soil standard of 0.5 mg kg<sup>-1</sup> still results in a considerable number of samples of Indica rice samples that do not meet the food quality standard.

### *Application of models to derive soil-specific testing values.*

Results in chapter 4 clearly show that soil properties like pH and CEC have a large influence on the availability of Cd in the soil. Based on the metal content alone, the uptake of Cd by rice cannot be predicted accurately enough as was discussed previously. However, when using soil pH and CEC in combination with the amount of metals extracted by either HCl or HNO<sub>3</sub>, the availability of Cd can be assessed rather easily (Chapter 4). This results in a significant improvement of the prediction of Cd in rice and a clear reduction of the number of samples that were incorrectly classified as either suitable or unsuitable. The advantage of the application of such models is that they require relatively little information (pH and CEC) which can be obtained from soil maps or local soil investigations. The approach is very suitable for application on a regional or even national scale to identify areas where rice cropping should be monitored more closely.

### *Use of CaCl<sub>2</sub> for soil testing: advantage and disadvantages*

Instead of a soil test based on Aqua Regia or HCl, the use of 0.01 M CaCl<sub>2</sub> seems much more appropriate for either Japonica or Indica species, at least for the ones tested here. The use of CaCl<sub>2</sub> does account for differences in the availability of heavy metals in soil which is not the case when using acid soil extracts. Based on CaCl<sub>2</sub> extractions no additional soil properties need to be measured since CaCl<sub>2</sub> does reflect differences between soil types (as expressed by differences in pH, CEC, organic matter etc.). The results presented in this report are based on CaCl<sub>2</sub> extracts performed on dried soil samples (air dried). Results obtained in other field studies seem to indicate that there can be an effect of drying of soil samples on the relationship between soil extractable Cd and uptake by rice. In this study the soil in the immediate vicinity of the rice root (rhizosphere) soil was sampled which may explain why there is a close match despite drying the samples. If possible, soil samples should be taken from the rhizosphere.

The  $\text{CaCl}_2$  test is not so much suitable for regional studies as is the case of the soil models but it is especially suitable for local testing of soils where no information is available yet or to focus on areas that have been identified (by the risk maps constructed using the soil-rice model) as potentially not suitable. Careful sampling of local soils from selected fields is then crucial due to the large variability of the Cd content across many fields.

### ***Design of user-friendly tools to assess local soil quality for rice cropping***

Based on the soil-rice models, simple easy-to-use look-up tables can be constructed which enables the owner of a site to assess quickly whether or not a specific soil or field is suitable for one or more varieties to be grown. Based on the list of cultivars studied so far, farmers can choose to use those cultivars which accumulate the least amount of Cd.

### ***Extension of the method to other regions in South East Asia***

At present different approaches are used to determine the suitability of soils for agriculture. In most countries, either in Asia, the EU or elsewhere, the total metal content is still most commonly used to evaluate the suitability of soils for agriculture, housing or industry. This report shows, if anything, that considering the availability of metals (in this case Cd) in soil results in a considerable improvement of soil testing. The approach described here based on soil-plant models currently is being implemented in soil policy in several EU countries (Germany, the Netherlands) and seems quite appropriate for rice cropping as well as indicated by the results in this study. In fact the similarity in the behavior of Cd in paddy fields compared to that in soils from the EU was striking. Nevertheless, soils will be different in countries across Asia, but the approach nevertheless can be implemented rather easily based on a limited number of carefully selected field studies. Selection criteria include soil properties (range in soil types), degree of contamination (clean to polluted) and the rice-varieties used in each country.

### ***Research questions that need to be addressed***

Although it seems that the chemical behavior of Cd in paddy fields as well as the uptake of Cd by rice is rather similar compared to soils and crops from non-tropical countries, several aspects still need to be considered. This includes:

1. The observed difference in the availability of Cd between harvest 1 (May) and 2 (November) and the resulting increase in Cd uptake at harvest 2. Several reasons (difference in temperature, rainfall, and fertilization) can be mentioned but proof is lacking.
2. The apparent difference between results from this study and the one performed by Simmons et al (2008) where extracts with  $\text{CaCl}_2$  only showed a significant relationship with Cd in rice when using field moist soil. In our study rhizosphere soil was used which may explain the better fit. This nevertheless suggests that in order to obtain suitable relationships between soil extractable Cd and Cd in rice, clear sampling protocols need to be established.

## References

- Arao, T. and Ishikawa, S., 2006. Genotypic differences in Cd concentration and distribution of soybean and rice. *Japan Agricultural Research Quarterly* 40, 21–30.
- Adams, M.L., Zhao, F.J., McGrath, S.P., Nicholson, F.A. and Chambers, B.J. Predicting Cadmium Concentrations in Wheat and Barley Grain Using Soil Properties (2004) *Journal of Environmental Quality*, 33 (2), pp. 532-541.
- Barrett K.A. and McBride M.B., 2007. Dissolution of zinc-cadmium sulfide solid solutions in aerated aqueous suspension. *Soil Sci Soc Am J* 71:322–328.
- Bonten, L.T.C., Groenenberg J.E., Weng L. and Riemsdijk W.H. van, 2008. Use of speciation and complexation models to estimate heavy metal sorption in soils. *Geoderma*, 146:303-310.
- Brand, E., Peijnenburg W., Groenenberg B., Vink J., Lijzen J., Hulscher D. ten, Jonker C., Römkens P. and Roex E., 2009. Towards implementation of bioavailability measurements in the Dutch regulatory framework. RIVM report 711701084/2009, RIVM, Bilthoven, the Netherlands, 80 pp.
- Bruemmer G.W., Gerth J. and Tiller K.G., 1981. Reaction kinetics of the adsorption and desorption of nickel, zinc and cadmium by goethite. I. Adsorption and diffusion of metals. *J Soil Sci* 39:37-52.
- Brus D.J. and M.J.W. Jansen, 2004 Uncertainty and Sensitivity Analysis of Spatial Predictions of Heavy Metals in Wheat. *J. Environ. Qual.* 33:882-890.
- Brus, D.J., Z. Li, J. Song, G.F. Koopmans, E.J.M. Temminghoff, X. Yin, C. Yao, H. Zhang, Y. Luo and J. Japenga, 2009. Predictions of Spatially Averaged Cadmium Contents in Rice Grains in the Fuyang Valley, P.R. China. *J. Environ. Qual.* 38: 1126-1136.
- Brus, D.J., Gruijter J.J. de and Römkens P.F.A.M., 2005. Probabilistic quality standards for heavy metals in soil derived from quality standards in crops. *Geoderma* 128:301– 311.
- Buekers J., 2007. Fixation of cadmium, copper, nickel and zinc in soil: kinetics, mechanisms and its effect on metal bioavailability. PhD thesis no. 754, Catholic University Leuven, Leuven, Belgium.
- Cai, S., Hu Z., Zhong X., Ye Z., Xu H., Liu Y., Rong J., Zhang W. and Zhang F., 1990. Cd exposure and health effects among residents in an irrigation area with ore dressing wastewater. *Science of the Total Environment* 90, 67-73.
- Carlou, C. (Ed.), 2007. Derivation methods of soil screening values in Europe. A review and evaluation of national procedures towards harmonization. European Commission, Joint Research Centre, Ispra, EUR 22805-EN, 306 pp.
- Central Weather Bureau, 2008. Climate statistics. Central Weather Bureau of Taiwan, No. 64, Gongyuan Road, Taipei, Taiwan 10048, R.O.C.  
<http://www.cwb.gov.tw/eng/statistics/statistics.htm>.
- Charlatchka R. and Cambier P., 1999. Influence of reducing conditions on solubility of trace metals in contaminated soils. *Water Air Soil Pollut* 118:143–167.

- Chaudri A, McGrath S., Gibbs P., Chambers B., Carlton-Smith C., Godley A., Bacon J., Campbell C. and Aitken M., 2007. Cadmium availability to wheat grain in soils treated with sewage sludge or metal salts. *Chemosphere* 66 (2007) 1415–1423.
- Chen, Z.S., 1991. Cadmium and lead contamination of soils near plastic stabilizing materials producing plants in Northern Taiwan. *Water, Air, and Soil Pollution* 57-58, 745-754.
- Chen, Z.S., Tsai, C.C. and Tsui, C.C., 1999. Proposal regulation of soil pollutants in Taiwan soils. In: Chen, Z.S. (Ed.), *Proceedings of 6<sup>th</sup> Workshop on Soil Pollution and Prevention: Soil Remediation Techniques on Soils Contaminated by Organic Pollutants*, Taipei, Taiwan, ROC, pp. 169-207. (In Chinese, with English abstract).
- Choudhary, M., Bailey, L.D., Grant C.A. and Leisle D., 1995. Effect of Zn on the concentration of Cd and Zn in plant tissue of two Durum wheat lines. *Canadian Journal of Plant Science* 75, 445-448.
- Clayton P.M. and Tiller K.G., 1979. A chemical method for the determination of heavy metal content of soils in environmental studies. Technical paper no. 41, Division of Soils, CSIRO Australia.
- COA, 2008. Food supply and utilization 2005. Council of Agriculture, Taipei, Taiwan. Available at <http://eng.coa.gov.tw> (checked October 30, 2008).
- Colombo C. and Van Den Berg C.M.G., 1998. Determination of trace metals (Cu, Pb, Zn and Ni) in soil extracts by flow analysis with voltammetric detection. *Int J Environ Anal Chem* 71:1-17.
- Contín M., Mondini C., Leita L. and De Nobili, M., 2007. Enhanced soil toxic metal fixation in iron (hydr)oxides by redox cycles. *Geoderma* 140:164–175.
- Cui, Y., Zhang, X., Zhu, Y., 2008. Does copper reduce Cd uptake by different rice genotypes? *Journal of Environmental Sciences* 20, 332-338.
- Das, P., Samantaray, S., Rout, G.R., 1997. Studies on Cd toxicity in plants. A Review. *Environmental Pollution* 98, 29-36.
- Daum D., Bogdan K., Schenk M.K., Merkel D., 2001. Influence of the field water management on accumulation of arsenic and cadmium in paddy rice. In: Horst et al. (Eds.): *Plant nutrition - food security and sustainability of agro-ecosystems through basic and applied research*, Kluwer, pp 290-291.
- De Vries W., Curlík J., Murányi A., Alloway B., J.E Groenenberg, 2005. Assessment of relationships between total and reactive concentrations of Cd, Cu, Pb and Zn in Hungarian and Slovakian soils. *Ekológia* 24:152-169
- De Vries W., Römkens P.F.A.M., Bonten L.T.C., 2008. Spatially explicit integrated risk assessment of present soil concentrations of Cd, Pb, Cu and Zn in the Netherlands. *Water Air Soil Pollut* 191:199-215.
- Degryse, F., Broos, K., Smolders, E., Merckx, R., 2003. Soil solution concentration of Cd and Zn can be predicted with a CaCl<sub>2</sub> soil extract. *European Journal of Soil Science* 54, 149-157.
- Diao, W.P., Ni, W.Z., Ma, H.Y., Yang, X.E., 2005. Cd Pollution in paddy soil as affected by different rice (*Oryza sativa* L.) cultivars. *Bulletin of Environmental Contamination and Toxicology* 75, 731-738.
- Dudka S., Miller W.P., 1999. Accumulation of potentially toxic elements in plants and their transfer to human food chain. *J Environ Sci Health B* 34:681–708.



- Dunbar, K.R., 2004. Uptake and partitioning of Cd in two cultivars of potato (*Solanum tuberosum* L.). Ph.D. thesis, University of Adelaide, School of Earth and Environmental Sciences, Adelaide, Australia, 123 p.
- Efroymsen, R. A., B. E. Sample, and G. W. Suter II, 2001. Uptake of inorganic chemicals from soil by plant leaves: regressions of field data. *Environ. Toxicol. Chem.* 20:2561-2571.
- Elzinga E.J., Sparks D.L., 1999. Nickel sorption mechanisms in a pyrophyllite-montmorillonite mixture. *J Colloid Interf Sci* 213:506–512.
- EPA, 2006. Regulations governing the preliminary assessment of soil and groundwater pollution control sites. Environmental Protection Administration order Huan-Shu-Tu-Tzu no.0950023629, March 29, 2006.
- Fairbrother, A. R. Wenstel, K. Sappington, and W. Wood, 2007. Framework for Metals Risk Assessment. *Ecotoxicology and Environmental Safety* 68 (2007) 145–227.
- François, M., Grant, C., Lambert, R. Sauvé, S., 2009. Prediction of cadmium and zinc concentration in wheat grain from soils affected by the application of phosphate fertilizers varying in Cd concentration. *Nutr Cycl Agroecosyst* 83:125–133.
- Franz, E., Römken, P.F.A.M., van Raamsdonk, L., van der Fels-Klerx, I., 2008. A chain modeling approach to estimate the impact of soil cadmium pollution on human dietary exposure. *Journal Food Protection* 71, 2504-2513.
- Fu J, Zhou Q., Liu J., Liu W., Wang T., Zhang Q., Jiang G., 2008. High levels of heavy metals in rice (*Oryza sativa* L.) from a typical E-waste recycling area in southeast China and its potential risk to human health. *Chemosphere* 71:1269-1275.
- Girling, C.A., Peterson, P.J., 1981. The significance of the Cd species in uptake and metabolism of cd in crop plants. *Journal of Plant Nutrition* 1981, 707-720.
- Grant C.A., Sheppard S.C., 2008. Fertilizer Impacts on Cadmium Availability in Agricultural Soils and Crops. *Journal Human and Ecological Risk Assessment* 14: 210 – 228.
- Guo, H.Y., Liu, T.S., Chu, C.L., Chiang, C.F., 2007. Prediction of heavy metals uptake by different rice species in paddy soils near contaminated sites of Taiwan. In: *Proceedings of the International Workshop of ESAFS 8: New Solutions to Soil Pollution and Distribution, Bioavailability and Management of Heavy Metals*, October 22-23, 2007, Tsukuba, Japan, pp. 95-116.
- Hassan, M.J., Zhu, Z., Ahmad, B., Mahmood, Q., 2006. Influence of Cd toxicity on rice genotypes as affected by Zn, sulfur and nitrogen fertilizers. *Caspian Journal of Environmental Sciences* 4, 1-8.
- He J, Zhu C., Ren Y., Yan Y., Jiang D., 2006. Genotypic variation in grain Cd concentration of lowland rice. *J Plant Nutr Soil Sci* 169:711–716.
- Hochi, Y., Kido, T., Nogawa, K., Kito, H., Shaikh, Z.A., 1995. Dose-response relationship between total Cd intake and prevalence of renal dysfunction using general linear models. *Journal of Applied Toxicology* 15, 109-116.
- Houba V.J.G., Van Der Lee J.J., Novozamsky I., 1997. *Soil and Plant Analysis. Part 1: Soil Analysis Procedures*. Wageningen University, Wageningen, the Netherlands.

- Houba, V.J.G., Temminghoff, E.J.M., Gaikhorst, G.A., Van Vark, W., 2000. Soil analysis procedures using 0.01 M calcium chloride as extraction reagent. *Commun. Soil Sci. Plant Anal.* 31:1299–1396.
- Hough R.L., Tye A.M., Crout N.M.J., McGrath S.P., Zhang H., Young S.D., 2005. Evaluating a 'Free Ion Activity Model' applied to metal uptake by *Lolium perenne* L. grown in contaminated soils. *Plant Soil* 270:1-12.
- Hseh, Y.H., Wang, M.K., 1990. Soil survey on the potential contaminated sites of Taichung City, Taiwan, ROC. Project Report of Environmental Protection Bureau of Taichung City, Taiwan, 121 p.
- Huang, Y.M., Huang, J.H., 1993. Monitoring studies of heavy metals in potential contaminated soils of Changhwa City. Taiwan, ROC. Project report of Environmental Protection Bureau of Chang Hwa County, Taiwan. (Unpub. mimeograph).
- Ingwersen J, Streck T., 2006. Modeling the environmental fate of Cd in a large wastewater irrigation area. *J Environ Qual* 35:1702-1714.
- Japenga, J., Koopmans, G.F., Song, J., Römkens, P.F.A.M., 2007. A feasibility test to estimate the duration of phytoextraction of heavy metals from polluted soils. *International Journal of Phytoremediation* 9, 115–132.
- Kalis, E.J.J., Temminghoff, E.J.M., Visser, A., Van Riemsdijk, W.H., 2007. Metal uptake by *Lolium perenne* in contaminated soils using a four-step approach. *Environmental Toxicology and Chemistry* 26, 335-345.
- Kashem M.A., Singh B.R., 2004. Transformations in solid phase species of metals as affected by flooding and organic matter. *Commun Soil Sci Plant Anal* 35:1435–1456.
- Kelderman P., Osman A.A., 2007. Effect of redox potential on heavy metal binding forms in polluted canal sediments in Delft (The Netherlands). *Wat Res* 41:4251–4261.
- Khan, S., Q. Cao, Y.M. Zheng, Y.Z. Huang, Y.G. Zhu, 2008. Health risks of heavy metals in contaminated soils and food crops irrigated with wastewater in Beijing, China. *Environmental Pollution* 152:686-692.
- Kikuchi T., Okazaki M., Kimura S.D., Motobayashi T., Baasansuren J., Hattori T., Abe T., 2008. Suppressive effects of magnesium oxide materials on Cd uptake and accumulation into rice grains II: suppression of Cd uptake and accumulation into rice grains due to application of magnesium oxide materials. *J Hazard Mat* 154:294–299.
- Kirkham, M.B., 2006. Cd in plants on polluted soils: Effects of soil factors, hyperaccumulation, and amendments. *Geoderma* 137, 19-32.
- Kobayashi, E., Y. Suwazono, R. Hondab, M. Dochi, M. Nishijo, T. Kido, H. Nakagawa, 2008. Serial changes in urinary cadmium concentrations and degree of renal tubular injury after soil replacement in cadmium-polluted rice paddies. *Toxicology Letters* 176: 124–130.
- Koopmans G.F., Schenkeveld W.D.C., Song J., Luo Y.M., Japenga J., Temminghoff E.J.M., 2008b. Influence of EDDS on metal speciation in soil extracts: measurement and mechanistic multicomponent modeling. *Environ Sci Technol* 42:1123–1130.

- Koopmans, G.F., Römkens, P.F.A.M., Fokkema, M.J., Song, J., Luo, Y.M., Japenga, J., Zhao, F.J., 2008. Feasibility of phytoextraction to remediate Cd and Zn contaminated soils. *Environmental Pollution* 156, 905-914.
- Krauss, M., Wolfgang, W., Kobza, J., Zech, W., 2002. Predicting heavy metal transfer from soil to plant: Potential use of Freundlich-type functions. *Journal of Plant Nutrition and Soil Science* 165, 3-8.
- Krishnamurti, G.S.R., McArthur, D.F.E., Wang, M.K., Kozak, L.M., Huang, P.M., 2005. Biogeochemistry of soil Cd and the impact on terrestrial food chain contamination. In: Huang, P.M., Gobran, G.R. (Eds), *Biogeochemistry of Trace Elements in the Rhizosphere*, Elsevier, pp. 197-257.
- Kukier, U., Chaney, R.L., 2002. Growing rice grain with controlled Cd concentrations. *Journal of Plant Nutrition* 25, 1793-1820.
- Lamothe P.J., Fries T.L., Consul J.J., 1986. Evaluation of a microwave oven system for the dissolution of geologic samples. *Anal Chem* 58:1881–1886
- Li P, Wang X.X., Zhang T.L., Zhou D.M., He Y.Q., 2009. Distribution and accumulation of Cu and Cd in soil-rice system as affected by soil amendments. *Water Air Soil Pollut*, 196:29–40.
- Li, Z., Li, L., Pan, G., Chen, J., 2005. Bioavailability of Cd in a soil–rice system in China: Soil type versus genotype effects. *Plant and Soil* 271, 165-173.
- Lin, C.W., 2002. Mapping soil lead and remediation needs in contaminated soils. *Environmental Geochemistry and Health* 24, 23-33.
- Lin, Y.P., Teng, T.P., Chang, T.K., 2002. Multivariate analysis of soil heavy metal pollution and landscape pattern in Changhua county in Taiwan. *Landscape and Urban Planning* 62, 19-35.
- Liu, J., Qian, M., Cai, G., Yang, J., Zhu, Q., 2007. Uptake and translocation of Cd in different rice cultivars and the relation with Cd accumulation in rice grain. *Journal of Hazardous Materials* 143, 443-447.
- Liu, J., Zhu, Q., Zhang, Z., Xu, J., Yang, J., Wong, M.H., 2005. Variations in Cd accumulation among rice cultivars and types and the selection of cultivars for reducing Cd in the diet. *Journal of the Science of Food and Agriculture* 85, 47-153.
- Lofts S., Spurgeon D.J., Svendsen C., Tipping E., 2004. Deriving soil critical limits for Cu, Zn, Cd, and Pb: a method based on free ion concentrations. *Environ Sci Technol* 38:3623-3631.
- Lu LT, Chang I.C., Hsiao T.Y., Yu Y.H., Ma H.W., 2007. Identification of pollution source of Cd in soil. Application of material flow analysis and a case study in Taiwan. *Environ Sci Pollut Res* 14:49-59.
- Ma, Y.B., Uren, N.C., 2006. Effect of aging on the availability of Zn added to a calcareous clay soil. *Nutrient Cycling in Agroecosystems* 76, 11-18.
- McBride M.B., 1994. *Environmental chemistry of soils*. Oxford University Press.
- McBride M.B., Barrett K.A., Kim B., Hale B., 2006. Cd sorption in soils 25 years after amendment with sewage sludge. *Soil Sci* 171:21-28.
- McGrath, S.P., Zhao, F.J., Dunham, S.L., Crosland, A.R., Coleman, K., 2000. Long-term changes in the extractability and bioavailability of Zn and Cd after sludge application. *Journal of Environmental Quality* 29, 875-883.
- McLaughlin, M.J. M. Whatmuff, M. Warne, D. Heemsbergen, G Barry, M. Bell, D. Nash, D. Pritchard. 2006. *Field Investigation of Solubility and Food Chain*

- Accumulation of Biosolid-Cadmium Across Diverse Soil Types. *Environ. Chem.* 3, 428–432.
- Meers E., Du Laing G., Unamuno V., Ruttens A., Vangronsveld J., Tack F.M.G., Verloo M.G., 2007. Comparison of cadmium extractability from soils by commonly used single extraction protocols. *Geoderma* 141:247–259.
- Mohr, E.C.J., Van Baren, F.A., Van Schuylenborgh, J., 1972. *Tropical Soils: A Comprehensive Study of Their Genesis*. Mouton-Ichtiar Baru-Van Hoeve Press, Den Haag, the Netherlands, 481 p.
- Muchuweti, M., J.W. Birkett, E. Chinyanga, R. Zvauya, M.D. Scrimshaw and J.N. Lester. 2006. Heavy metal content of vegetables irrigated with mixture of wastewater and sewage sludge in Zimbabwe: implications for human health. *Agriculture, Ecosystem and Environment* 112 (2006), pp. 41–48.
- Naidu, R., Kookana, R.S., Rogers, S., Bolan, N.S., Adriano, D., 2003. Bioavailability of metals in the soil–plant environment and its potential role in risk assessment. In: Naidu, R., Gupta, V.V.S.R., Rogers, S., Kookana, R.S., Bolan, N.S., Adriano, D. (Eds.), *Bioavailability, Toxicity, and Risk Relationships in Ecosystems*. Science Publishers, Inc., Enfield, New Hampshire, pp. 46-81.
- Nelson, J.L., Boawn L.C., Viets F.G. Jr., 1959. A method for assessing Zn status of soils using acid extractable Zn and ‘titratable alkalinity’ values. *Soil Sci* 88:275-283
- NEPAC, 1995. Environmental quality standard for soils. GB 15618. National Environmental Protection Agency of China.
- NPF, 2004. Exploration of preservation and utilization of hillside. Research report no. 093-004, National Policy Foundation, Taipei, Taiwan.
- Peijnenburg W.J.G.M., Zablotskaja M., Vijver M.G., 2007. Monitoring metals in terrestrial environments within a bioavailability framework and a focus on soil extraction. *Ecotoxicol Environ Saf* 67:163–179.
- Peijnenburg, W.J.G.M., Posthuma L., Zweers P.G.P.C., de Baerselman R., de Groot A.C., Van Veen R.P.M., Jager T., 1999. Prediction of metal bioavailability in Dutch field soils for the oligochaete *Enchytraeus crypticus*. *Ecotoxicol. Environ Saf* 43:170–186.
- Pueyo M., Lopez-Sanchez J.F., Rauret G., 2004. Assessment of CaCl<sub>2</sub>, NaNO<sub>3</sub>, and NH<sub>4</sub>NO<sub>3</sub> extraction procedures for the study of Cd, Cu, Pb and Zn extractability in contaminated soils. *Anal Chim Acta* 504:217–226.
- Qin F., Wen B., Shan X-Q., Xie Y-N., Liu T., Zhang S-Z., Khan S.U., 2006. Mechanisms of competitive adsorption of Pb, Cu, and Cd on peat. *Environ Pollut* 144:669-680.
- Rattan R.K., Datta S.P., Chhonkar P.K., Suribabu K., Singh A.K., 2005. Long-term impact of irrigation with sewage effluents on heavy metal content in soils, crops and groundwater - a case study. *Agric Ecosyst Environ* 109:310-322.
- Rietra, R.P.J.J. and P.F.A.M. Römkens, 2007. Cadmium and zinc in soils and arable crops in the Kempen Area (NL) in 2006: results for schorzonera, carrots and leek (in Dutch). Alterra report no. 1422. Alterra, Wageningen UR, the Netherlands, 39 p.
- Römkens P.F.A.M., Groenenberg J.E., Bril J., Vries W., 2004. Derivation of partition equations to calculate heavy metal speciation and solubility in soils. Report no. 305, Alterra, Wageningen, the Netherlands.

- Römken, P.F.A.M., H.Y. Guo, C.L. Chu, T.S.Liu, C.F. Chiang, and G.F. Koopmans. 2009a. Prediction of Cadmium uptake by brown rice and derivation of soil-plant transfer models to improve soil protection guidelines. *Environmental Pollution* 157 (2009) 2435–2444.
- Römken, P.F.A.M., Guo, H.Y., Chu, C.L., Liu, T.S., Chiang, C.F., Koopmans, G.F. 2009b. Characterization of soil metal pools in paddy fields in Taiwan: chemical extraction and solid-solution partitioning. *Journal of Soils and Sediments* 9:216–228.
- Sauvé S, N Cook, W.H. Hendershot M.B. McBride, 1996. Linking plant tissue concentrations and soil copper pools in urban contaminated soils. *Environmental Pollution* 94 (2) 153-157.
- Saxe J.K., Impellitteri C.A., Peijnenburg W.J.G.M., Allen H.E., 2001. Novel model describing trace metal concentrations in the earthworm, *Eisenia Andrei*. *Environ Sci Technol* 35: 4522-4529.
- Schröder T.J., Hiemstra T., Vink J.P.M., Van Der Zee S.E.A.T.M., 2005. Modeling of the solid-solution partitioning of heavy metals and arsenic in embanked flood plain soils of the rivers Rhine and Meuse. *Environ Sci Technol* 39:7176-7184.
- Shentu, J., Z. He, X. Yang, and T. Li, 2008. Accumulation Properties of Cadmium in a Selected Vegetable-Rotation System of Southeastern China. *J. Agric. Food Chem.*, 2008, 56 (15), 6382-6388
- Shimada, M., Piscator, T., Iwata-Nishino, H., 1977. Urine analysis for detection of Cd-induced renal changes, with special reference to  $\beta_2$ -microglobulin. *Environmental Research* 13, 407-424.
- Simmons R.W., Noble A.D., Pongsakul P., Sukreeyapongse O., Chinabut N., 2008. Analysis of field-moist Cd contaminated paddy soils during rice grain fill allows reliable prediction of grain Cd levels. *Plant Soil* 302:125–137.
- Simmons R.W., Pongsakul P., Saiyasitpanich D., Klinphoklap S., 2005. Elevated levels of Cd and Zn in paddy soils and elevated levels of Cd in rice grain downstream of a Zn mineralized area in Thailand: implications for public health. *Environ Geochem Health* 27:501-511.
- Singhal J.P., Gupta G.K., 1978. Reactions of zinc with acid and base saturated dickites. *Clay Clay Miner* 26:365-371.
- Sun, H., Zhang, T.L., Sun, B., 2001. Assessment of pollution of heavy metals on paddy field with sewage irrigation in Guixi, Jiangxi province. *Agro-environmental Protection* 20, 405-407. (In Chinese, with English abstract).
- Thakali, S., H. Allen, D. M Di Toro, A.A. Ponizovsky, C. R. Rooney, F.J. Zhao and S. P. McGrath, 2006. A terrestrial Biotic Ligand Model. 1 Development and Application to Cu and Ni Toxicities to Barley Root Elongation in Soils. *Environmental Science and Technology* 40, 7085-7093.
- Thornton, I., Farago, M.E., Thums, C.R., Parrish, R.R., McGill, R.A.R., Breward, N., Fortey, N.J., Simpson, P., Young, S.D., Tye, A.M., Crout, N.M.J., Hough, R.L., Watt, J., 2008. Urban geochemistry: Research strategies to assist risk assessment and remediation of brownfield sites in urban areas. *Environmental Geochemistry and Health* 30, 565–576.
- Tipping E., Lawlor A.J., Lofts S., Shotbolt L., 2006. Simulating the long-term chemistry of an upland UK catchment: heavy metals. *Environ Pollut* 141:139-150.

- Tipping E., Rieuwerts J., Pan G., Ashmore MR, Lofts S, Hill MTR, Farago ME, Thornton I., 2003. The solid–solution partitioning of heavy metals (Cu, Zn, Cd, Pb) in upland soils of England and Wales. *Environ Pollut* 125:213–225.
- Tiwari, R.C., Kumar, B.M., 1982. A suitable extractant for assessing plant-available copper in different soils (peaty, red and alluvial). *Plant and Soil* 68, 131-134.
- Tsukahara T., Ezaki T., Moriguchi J., Furuki K., Shimbo S., Matsuda-Inoguchi N., Ikeda M., 2002. Rice as the most influential source of Cd intake among general Japanese population. *Sci Total Environ* 305:41-51.
- USDA, 1996. Soil Survey Laboratory Methods Manual. Soil Survey Investigations Report no. 42, USDA, Washington, D.C., US.
- US-EPA, 2002. Methods for the determination of total organic carbon (TOC) in soils and sediments. NCEA-C-1282 EMASC-001, US-EPA, Las Vegas, US.
- Wang, G.Q., Koopmans, G.F., Song, J., Temminghoff, E.J.M., Luo, Y.M., Zhao, Q.G., Japenga, J., 2007. Mobilization of heavy metals from contaminated paddy soil by EDDS, EDTA, and elemental sulfur. *Environmental Geochemistry and Health* 29, 221-235.
- Watanabe T., Zhang Z.W., Moon C.S., Shimbo S., Nakatsuka H., Matsuda-Inoguchi N., Higashikawa K., Ikeda M., 2000. Cadmium exposure of women in general populations in Japan during 1991-1997 compared with 1977-1981. *Int. Arch. Occup. Environ. Health* 73:26-34.
- Weng L., Temminghoff E.J.M., Van Riemsdijk W.H., 2001. Contribution of individual sorbents to the control of heavy metal activity in sandy soil. *Environ Sci Technol* 35:4436 -4443.
- Xu R., 2000. Particle characterization: light scattering methods. Particle Technology Series Vol 13, Kluwer Academic Publishers, Dordrecht, the Netherlands.
- Yang, S.Y., Chang, W.L., 2005. Use of finite mixture distribution theory to determine the criteria of Cd concentrations in Taiwan farmland soils. *Soil Science* 170, 55-62.
- Yu, H., Wang, J., Fang, W., Yuan, J., Yang, Z., 2006. Cd accumulation in different rice cultivars and screening for pollution-safe cultivars of rice. *Science of the Total Environment* 370, 302-309.
- Zheng, N., Q. Wang, X. Zhang, D. Zheng, Z. Zhang, S. Zhang, 2007. Population health risk due to dietary intake of heavy metals in the industrial area of Huludao city, China *Science of the Total Environment* 387 (2007) 96–104.
- Zhou, Q., Wang, X., Liang, R., Wu, Y., 2003. Effects of Cd and mixed heavy metals on rice growth in Liaoning, China. *Soil and Sediment Contamination* 12, 851-864.
- Zhu, H.P., Zhang, L.J., Wu, W.G., 1992. A study on heavy metal content of soil in Taiwan. In: *Proceedings of 3<sup>rd</sup> Workshop of Soil Pollution and Prevention, Taiwan*, pp 1-14. (in Chinese).

## Appendix 1 Overview of model coefficients used to calculate Cd in brown rice from soil data (soil model)

Table A1.1 Model coefficients of the soil-plant model to predict Cd levels in the grain of brown rice using Aqua Regia, 0.43 N HNO<sub>3</sub> or 0.1 M HCl.

AR	number	Jap/Ind.	Constant	AR	pH	CEC
Tainung No.70	1	J	1.37	0.76	-0.25	-0.53
Taiken No.8	2	J	0.63	0.75	-0.13	-0.42
Tainung No.72	3	J	0.92	0.82	-0.20	-0.43
Kaohsiung No.143	4	J	0.89	0.84	-0.23	-0.26
Taitung No.30	5	J	1.32	0.77	-0.24	-0.57
Tainung Sen No.20	6	I	1.33	0.76	-0.21	-0.38
Tainung No.71	7	J	0.54	0.84	-0.14	-0.32
Tainung No.67	8	J	0.94	0.78	-0.20	-0.46
Kaohsiung Sen Yu No.1151	9	I	0.95	0.68	-0.18	-0.22
Taichung Sen Waxy No.1	10	I	1.34	0.86	-0.23	-0.23
Taichung Sen No.10	11	I	0.97	0.96	-0.17	-0.32
Kaohsiung No.144	12	J	0.76	0.81	-0.15	-0.54
Japonica – all cultivars		J	0.93	0.80	-0.19	-0.44
Indica – all cultivars		I	1.17	0.83	-0.20	-0.30

Table A1.2 Model coefficients of the soil-plant model to predict Cd levels in the grain of brown rice using 0.43 N HNO<sub>3</sub>

0.43 N HNO <sub>3</sub>			Constant	HNO <sub>3</sub>	pH	CEC
Tainung No.70	1	J	1.44	0.71	-0.24	-0.52
Taiken No.8	2	J	0.73	0.70	-0.13	-0.42
Tainung No.72	3	J	1.06	0.75	-0.22	-0.38
Kaohsiung No.143	4	J	0.94	0.79	-0.22	-0.26
Taitung No.30	5	J	1.36	0.74	-0.22	-0.60
Tainung Sen No.20	6	I	1.32	0.72	-0.18	-0.43
Tainung No.71	7	J	0.63	0.79	-0.13	-0.37
Tainung No.67	8	J	0.85	0.72	-0.17	-0.41
Kaohsiung Sen Yu No.1151	9	I	0.95	0.62	-0.15	-0.24
Taichung Sen Waxy No.1	10	I	1.33	0.78	-0.19	-0.25
Taichung Sen No.10	11	I	1.11	0.86	-0.16	-0.31
Kaohsiung No.144	12	J	0.76	0.74	-0.13	-0.53
Japonica – all cultivars		J	0.98	0.74	-0.18	-0.43
Indica – all cultivars		I	1.20	0.76	-0.17	-0.32

Table A1.3 Model coefficients of the soil-plant model to predict Cd levels in the grain of brown rice using 0.1 M HCl.

0.1 M HCl			Constant	HCl	pH	CEC
Tainung No.70	1	J	1.38	0.67	-0.23	-0.50
Taiken No.8	2	J	0.75	0.70	-0.12	-0.46
Tainung No.72	3	J	1.04	0.74	-0.20	-0.39
Kaohsiung No.143	4	J	0.94	0.79	-0.21	-0.30
Taitung No.30	5	J	1.44	0.74	-0.23	-0.63
Tainung Sen No.20	6	I	1.44	0.70	-0.20	-0.42
Tainung No.71	7	J	0.73	0.79	-0.13	-0.41
Tainung No.67	8	J	0.92	0.72	-0.17	-0.46
Kaohsiung Sen Yu No.1151	9	I	0.98	0.61	-0.15	-0.27
Taichung Sen Waxy No.1	10	I	1.36	0.79	-0.20	-0.26
Taichung Sen No.10	11	I	1.11	0.86	-0.16	-0.33
Kaohsiung No.144	12	J	0.86	0.71	-0.13	-0.58
Japonica – all cultivars		J	1.01	0.73	-0.18	-0.46
Indica – all cultivars		I	1.25	0.75	-0.18	-0.33



## Appendix 2 Look-up tables for individual cultivars




Figure A2.1 Examples of look-up tables for individual cultivars

Note: All tables are constructed using a CEC of  $20 \text{ cmol}^+\text{kg}^{-1}$ , a range in soil pH from 5 to 7, Cd-soil (Aqua Regia) ranging from 1 to  $5 \text{ mg kg}^{-1}$ .



## Appendix 3 Overview of soil data per site

Table A3.1 General Soil data

Site	pH H <sub>2</sub> O				pH 0.01 M CaCl <sub>2</sub>				CEC (cmol <sup>+</sup> kg <sup>-1</sup> )				OS (%)			
	mean	min	max	media n	mean	min	max	media n	mean	min	max	media n	mean	min	max	media n
CH6	5.0	4.2	6.9	5.0	4.7	3.8	6.8	4.7	16.9	5.0	19.8	17.0	6.7	2.5	8.3	6.7
CH7	6.3	5.0	7.5	6.3	6.1	4.7	7.2	6.1	11.1	5.7	19.7	11.0	4.6	2.2	6.4	4.6
CH8	6.5	5.2	7.1	6.5	6.3	4.6	6.9	6.4	19.0	4.0	21.6	19.3	6.3	1.5	7.2	6.3
HC3	5.7	4.8	6.2	5.7	5.4	4.4	5.8	5.4	5.5	2.8	9.3	5.6	2.7	2.0	4.9	2.7
HC4	5.9	5.2	6.5	5.9	5.6	4.7	5.9	5.6	8.7	5.0	13.2	8.9	4.5	3.0	5.7	4.5
HC5	5.5	4.7	6.0	5.5	5.1	4.0	5.7	5.1	6.9	3.0	10.9	7.1	3.5	2.3	4.4	3.5
HM3	6.3	5.0	7.0	6.4	6.2	4.9	7.0	6.3	19.1	5.6	22.5	19.3	7.5	2.9	8.6	7.5
HM4	6.1	4.7	6.8	6.2	5.9	4.5	6.6	6.0	16.2	2.6	25.1	16.3	6.2	4.8	7.9	6.2
HM5	5.6	4.5	6.8	5.7	5.3	4.2	6.4	5.4	15.1	6.2	18.5	15.1	5.9	4.4	7.6	5.8
LK1	5.7	4.7	6.8	5.7	5.3	4.4	6.6	5.3	5.9	2.6	19.4	5.7	2.7	1.3	7.3	2.7
LK2	6.1	5.2	6.6	6.1	5.8	4.8	6.3	5.8	10.6	4.9	18.9	10.7	4.4	2.3	6.6	4.3
PDA	5.7	4.8	6.1	5.7	5.3	4.4	5.9	5.4	13.1	8.5	19.7	13.2	7.1	6.1	8.3	7.1
PDB	5.5	4.5	6.2	5.6	5.2	4.1	5.8	5.2	14.4	6.9	24.2	14.5	7.5	6.4	9.1	7.5
PDC	5.6	4.8	6.1	5.7	5.2	4.1	5.5	5.2	13.3	6.7	16.1	13.7	6.8	5.8	8.0	6.8
PDD	5.4	4.6	5.7	5.4	5.0	3.9	5.3	5.0	14.6	10.7	18.6	14.7	7.8	6.8	8.9	7.8
PDE	5.5	4.8	5.8	5.6	5.1	4.3	5.4	5.2	14.1	8.3	18.8	14.3	7.5	6.3	9.0	7.5
PDF	5.6	4.7	5.9	5.6	5.2	4.3	5.5	5.3	15.2	7.2	20.4	15.3	7.8	6.5	8.9	7.7
PDG	5.4	4.6	5.8	5.5	5.1	4.1	5.4	5.1	14.9	9.6	20.9	15.3	7.7	6.8	8.5	7.7
PDH	5.3	4.7	5.7	5.3	5.0	4.3	5.4	5.0	16.4	8.8	20.9	16.8	8.7	7.4	10.2	8.7

Table A3.2 Cadmium

Site	Extract (all in mg kg <sup>-1</sup> dm)																			
	Cd Aqua Regia				Cd 0.43 M HNO <sub>3</sub>				Cd 0.1 M HCl				Cd 0.05 M EDTA				Cd 0.01 M CaCl <sub>2</sub>			
	mean	min	max	med	mean	min	max	med	mean	min	max	med	mean	min	max	med	mean	min	max	med
CH6	0.6	0.3	1.2	0.5	0.4	0.2	1.1	0.3	0.4	0.1	1.0	0.3	0.3	0.2	1.0	0.3	0.069	0.003	0.186	0.065
CH7	0.4	0.2	1.4	0.4	0.2	0.1	0.5	0.2	0.2	<0.1	0.4	0.2	0.2	0.1	0.5	0.2	0.011	0.001	0.093	0.008
CH8	0.4	0.2	2.1	0.4	0.2	0.1	0.6	0.2	0.2	0.1	0.6	0.2	0.2	0.1	0.5	0.2	0.006	0.001	0.098	0.005
HC3	2.7	0.2	6.8	2.6	2.3	0.1	5.0	2.2	2.2	0.1	5.3	2.1	1.8	0.1	3.3	1.7	0.446	0.033	0.989	0.428
HC4	0.4	0.2	2.0	0.4	0.3	0.1	1.9	0.3	0.3	0.2	1.9	0.3	0.3	0.2	1.3	0.3	0.046	0.007	0.414	0.041
HC5	0.2	0.1	0.5	0.2	0.1	0.1	0.3	0.1	0.1	0.1	0.3	0.1	0.1	0.1	0.3	0.1	0.035	0.012	0.087	0.033
HM3	0.7	0.3	2.1	0.6	0.5	0.3	1.6	0.4	0.5	0.2	1.5	0.4	0.5	0.2	1.6	0.4	0.012	0.001	0.061	0.008
HM4	1.1	0.3	2.9	0.9	0.8	0.2	2.0	0.7	0.8	0.1	1.9	0.6	0.7	0.2	1.9	0.6	0.024	0.004	0.088	0.023
HM5	0.5	0.2	2.5	0.5	0.3	0.1	1.7	0.3	0.3	<0.1	1.7	0.3	0.3	0.1	2.0	0.2	0.033	0.005	0.083	0.030
LK1	0.3	0.1	0.8	0.3	0.2	0.1	0.6	0.2	0.2	<0.1	0.5	0.2	0.2	0.1	0.6	0.2	0.031	0.002	0.100	0.031
LK2	0.3	0.1	1.7	0.3	0.2	0.1	1.4	0.2	0.2	<0.1	1.3	0.1	0.2	0.1	1.3	0.2	0.010	0.001	0.073	0.008
PDA	14.1	5.3	20.9	14.0	11.7	4.6	17.5	11.5	11.2	4.4	16.8	11.1	8.5	3.9	12.5	8.5	0.940	0.206	1.996	1.018
PDB	14.8	6.2	29.4	13.8	11.9	5.8	19.9	11.3	11.7	5.7	25.7	11.1	8.3	3.7	16.4	8.1	0.896	0.211	2.181	0.876
PDC	5.9	4.5	14.0	5.9	5.2	3.5	11.9	5.2	4.8	3.7	13.6	4.7	4.2	2.3	8.4	4.2	0.586	0.192	0.923	0.606
PDD	5.9	4.0	21.6	5.6	5.0	3.1	15.2	4.8	4.7	3.2	14.5	4.4	3.8	2.7	12.9	3.7	0.549	0.161	1.482	0.560
PDE	11.6	6.6	23.2	10.3	10.0	4.5	20.6	8.8	9.4	4.1	19.0	8.5	7.7	3.7	15.6	7.0	1.325	0.392	3.232	1.254
PDF	4.6	3.7	5.7	4.6	3.9	2.7	5.1	3.7	3.6	2.7	4.6	3.6	3.1	1.8	4.0	3.0	0.327	0.071	1.253	0.303
PDG	4.2	3.4	5.5	4.2	3.5	2.5	4.5	3.5	3.4	2.5	4.7	3.4	2.9	1.0	3.9	2.9	0.430	0.136	1.263	0.420
PDH	6.3	3.3	12.6	5.7	5.2	2.7	11.0	4.7	5.0	2.3	12.2	4.5	4.1	1.5	7.9	3.7	0.488	0.114	1.378	0.472

Table A3.3 Zinc

Site	Extract (all in mg kg <sup>-1</sup> dm)																			
	Zn Aqua Regia				Zn 0.43 M HNO <sub>3</sub>				Zn 0.1 M HCl				Zn 0.05 M EDTA				Zn 0.01 M CaCl <sub>2</sub>			
	mean	min	max	med	mean	min	max	med	mean	min	max	med	mean	min	max	med	mean	min	max	med
CH6	431.9	274.6	964.5	420.5	115.8	48.2	344.6	112.2	104.5	31.2	247.0	102.3	99.6	24.6	328.0	95.7	24.6	0.05	54.7	22.6
CH7	461.2	195.3	995.5	364.8	115.4	40.7	335.2	81.0	99.5	31.3	300.3	70.7	86.9	20.7	255.1	63.8	4.8	0.01	31.9	2.3
CH8	287.5	190.6	795.0	269.6	52.0	36.0	254.3	49.5	41.3	30.7	227.2	39.2	30.0	20.2	180.0	27.8	0.2	0.01	6.9	0.2
HC3	95.5	63.1	185.5	92.1	25.8	18.6	46.8	24.5	22.2	11.9	43.4	21.9	19.1	10.7	39.0	18.5	4.5	2.42	10.6	4.3
HC4	198.7	88.7	296.5	194.3	48.0	24.3	81.0	46.6	41.6	22.2	76.8	40.1	37.8	15.9	67.0	36.9	5.6	1.49	11.5	5.2
HC5	116.0	72.2	229.6	105.2	29.7	17.3	76.2	27.0	25.9	15.8	53.4	23.6	23.9	13.8	51.6	21.6	8.0	1.81	32.5	7.2
HM3	358.9	220.0	970.3	317.8	77.8	43.3	447.5	68.3	64.5	34.8	344.8	56.8	54.6	29.9	285.9	47.0	1.0	0.02	38.3	0.3
HM4	438.6	217.9	1164.1	372.1	94.8	38.6	278.1	79.2	81.4	33.1	225.7	68.0	69.5	26.5	189.7	58.8	1.6	0.14	38.9	1.2
HM5	328.5	230.3	1052.4	294.1	82.3	41.7	249.1	77.0	72.6	40.1	200.7	68.3	66.0	30.9	186.2	61.7	7.1	0.20	16.3	6.6
LK1	511.3	179.8	1225.2	367.8	150.6	34.3	574.2	86.2	134.6	31.1	537.3	76.9	112.5	27.7	383.6	69.2	16.8	0.12	96.9	14.0
LK2	302.9	156.9	1189.1	253.8	77.2	35.6	396.6	61.2	68.8	28.0	348.3	53.1	63.1	27.5	304.6	50.0	3.2	0.21	30.5	2.4
PDA	419.2	204.6	555.3	418.6	193.9	79.8	327.2	191.8	162.9	70.4	308.0	161.1	147.3	61.5	223.7	146.4	17.8	4.96	36.7	18.4
PDB	306.4	179.3	484.6	292.3	119.3	54.7	314.3	114.3	102.4	61.5	211.3	98.5	87.0	45.3	169.9	83.8	10.2	3.17	22.8	9.9
PDC	221.8	149.3	297.1	223.0	87.2	61.0	118.5	87.0	71.9	46.6	101.1	72.0	68.6	39.8	86.4	68.4	7.9	1.22	13.2	8.0
PDD	172.0	132.5	390.1	168.7	62.6	43.2	147.3	61.0	52.9	22.3	134.5	50.7	47.7	26.6	127.5	46.1	5.9	1.63	13.4	6.2
PDE	263.2	181.0	420.6	240.3	97.9	48.7	175.6	90.2	81.3	43.4	137.9	76.2	73.3	40.1	134.9	68.3	10.1	2.35	21.6	9.8
PDF	151.4	135.5	178.5	150.9	49.8	33.9	64.2	49.6	41.1	31.9	50.9	40.7	36.2	22.7	49.8	35.5	2.5	0.58	9.4	2.4
PDG	141.5	120.1	167.9	141.0	42.6	32.3	52.5	42.4	36.8	29.9	47.5	36.2	33.1	12.6	41.6	33.2	3.9	0.55	12.0	3.8
PDH	172.4	128.9	334.1	161.6	61.5	39.9	112.3	55.8	52.3	31.7	92.1	48.1	47.7	23.9	87.3	44.0	5.6	1.95	16.5	5.3

Table A3.4 Nickel

Site	Extract (all in mg kg <sup>-1</sup> dm)																			
	Ni Aqua Regia				Ni 0.43 M HNO <sub>3</sub>				Ni 0.1 M HCl				Ni 0.05 M EDTA				Ni 0.01 M CaCl <sub>2</sub>			
	mean	min	max	med	mean	min	max	med	mean	min	max	med	mean	min	max	med	mean	min	max	med
CH6	176.6	45.6	461.4	164.8	58.4	6.7	96.4	57.9	54.6	5.0	101.7	54.7	57.7	5.2	89.1	57.7	12.3	0.12	29.4	11.8
CH7	138.1	53.6	247.5	134.5	32.8	7.8	69.5	33.8	28.5	6.8	64.7	29.5	26.8	8.1	66.1	27.5	2.2	<0.01	11.0	1.5
CH8	51.7	32.4	207.1	48.0	8.1	4.6	46.2	7.4	7.2	4.0	63.5	6.3	7.3	3.5	33.3	6.7	0.1	<0.01	1.7	0.1
HC3	66.4	42.4	110.6	64.0	31.8	23.2	43.9	32.1	26.9	18.3	37.9	26.5	27.5	17.7	35.1	27.5	6.1	3.26	10.5	6.1
HC4	91.3	50.0	129.9	88.7	34.4	11.4	46.5	34.8	30.3	20.9	42.7	30.7	31.0	22.6	38.8	31.1	4.7	1.38	8.7	4.5
HC5	86.3	51.3	139.5	82.6	40.0	24.4	57.2	39.9	34.9	24.0	49.5	34.1	37.0	22.7	53.1	37.1	10.9	2.40	28.3	10.4
HM3	158.0	33.3	373.7	140.9	41.1	23.9	94.7	38.8	36.4	21.0	73.5	36.0	35.7	20.2	60.7	34.1	0.8	0.03	9.9	0.5
HM4	194.4	47.9	445.6	171.5	46.0	25.9	96.7	44.7	41.1	24.2	80.8	40.6	39.2	15.8	68.0	39.4	1.4	0.21	19.6	1.2
HM5	176.0	107.1	417.1	157.2	51.6	17.1	78.1	50.8	47.8	14.7	69.3	47.6	49.6	13.5	73.9	50.0	5.8	0.38	11.8	5.6
LK1	60.1	25.5	221.0	51.6	15.7	3.2	60.8	11.9	13.3	3.3	59.2	9.8	12.4	3.3	57.1	10.1	1.9	0.10	8.1	1.7
LK2	209.1	51.7	391.0	191.0	59.4	10.9	102.2	57.5	53.1	10.1	97.0	52.5	51.4	9.4	84.7	52.0	3.6	0.04	15.3	3.4
PDA	31.0	19.0	56.0	30.9	5.5	3.5	13.8	5.4	4.5	3.1	9.1	4.4	4.6	3.0	7.6	4.5	0.5	0.08	1.1	0.5
PDB	29.9	18.4	49.7	29.7	3.9	2.0	8.9	3.8	3.2	2.1	5.1	3.1	3.2	1.3	5.0	3.2	0.3	0.04	0.7	0.3
PDC	28.2	18.6	37.2	28.3	3.8	2.3	5.1	3.7	3.0	2.1	4.1	3.0	3.2	2.0	4.3	3.2	0.3	0.04	0.4	0.3
PDD	29.6	17.9	86.8	29.3	3.3	2.3	4.4	3.3	2.6	0.8	3.7	2.5	2.7	1.7	3.8	2.7	0.2	0.01	0.4	0.2
PDE	33.0	19.5	46.8	32.4	5.2	3.0	9.5	5.0	4.3	2.3	7.7	3.8	4.2	2.2	8.7	4.0	0.4	0.15	0.9	0.4
PDF	30.1	21.2	39.6	29.9	2.6	1.7	3.4	2.6	1.9	0.9	2.8	1.9	1.9	0.8	2.9	1.9	0.1	0.01	0.5	0.1
PDG	29.3	19.9	61.9	28.2	2.7	1.9	3.6	2.7	2.2	1.6	3.0	2.1	2.2	1.1	3.4	2.2	0.2	<0.01	0.8	0.2
PDH	31.1	16.8	42.6	30.7	3.5	2.0	8.2	3.0	2.8	1.4	5.6	2.5	2.9	1.2	6.1	2.5	0.2	0.02	1.7	0.2

Table A4.5 Lead

Site	Extract (all in mg kg <sup>-1</sup> dm)																			
	Pb Aqua Regia				Pb 0.43 M HNO <sub>3</sub>				Pb 0.1 M HCl				Pb 0.05 M EDTA				Pb 0.01 M CaCl <sub>2</sub>			
	mean	min	max	med	mean	min	max	med	mean	min	max	med	mean	min	max	med	mean	min	max	med
CH6	62.1	30.4	91.1	61.9	35.8	14.5	57.9	35.8	20.2	4.1	34.2	19.5	26.4	5.0	43.8	25.8	0.08	0.001	0.29	0.07
CH7	35.2	23.5	84.0	33.0	22.1	15.3	37.2	20.8	10.9	3.2	22.4	10.5	17.4	7.0	34.0	16.6	0.09	0.001	0.34	0.05
CH8	55.4	24.5	159.9	53.8	26.7	15.2	40.9	26.6	11.5	3.1	20.2	11.6	19.7	9.9	27.6	19.9	0.08	0.001	0.34	0.06
HC3	23.4	16.4	36.0	23.5	13.5	8.9	20.7	13.3	7.7	3.2	11.6	7.8	9.3	2.0	16.3	9.4	0.09	0.001	0.50	0.08
HC4	34.7	23.2	54.4	33.6	20.2	8.8	33.8	20.2	11.7	4.2	20.7	11.2	15.5	8.2	24.3	15.1	0.11	0.001	0.23	0.11
HC5	22.5	10.8	46.2	20.4	13.3	6.7	30.8	12.0	6.9	0.8	17.8	6.3	9.6	3.0	19.9	8.8	0.10	0.005	0.30	0.10
HM3	32.7	21.2	97.3	31.5	18.3	10.8	55.9	17.7	6.5	0.3	42.1	6.2	11.3	4.1	45.5	10.8	0.08	0.001	0.30	0.05
HM4	33.2	23.0	64.0	32.6	18.0	11.2	54.6	17.3	6.8	0.7	22.8	6.6	10.8	4.5	30.7	10.5	0.08	< 0.001	0.29	0.06
HM5	36.2	22.2	97.4	35.7	19.0	11.7	29.8	18.6	7.7	1.9	15.3	7.6	12.1	4.5	23.2	12.2	0.08	0.001	0.34	0.05
LK1	51.6	18.3	217.4	42.3	30.5	8.0	66.3	26.1	21.9	5.4	49.9	19.3	25.1	5.0	57.1	21.7	0.08	< 0.001	0.28	0.06
LK2	33.0	17.5	122.2	28.4	19.4	8.9	49.1	17.1	10.8	1.2	43.5	8.7	16.5	6.2	54.4	14.0	0.08	< 0.001	0.32	0.05
PDA	686.9	280.7	933.8	678.6	479.4	183.8	715.4	480.3	425.8	135.4	821.7	383.6	419.7	151.6	624.9	415.2	0.43	0.060	0.93	0.43
PDB	551.0	290.7	848.4	540.6	388.7	172.0	723.3	373.8	320.6	139.1	762.3	291.0	329.7	149.7	569.2	326.5	0.46	0.076	1.05	0.45
PDC	344.6	177.7	595.7	333.9	233.8	132.4	397.4	225.1	171.3	79.9	389.1	167.3	197.4	100.0	316.4	195.7	0.24	0.061	0.45	0.23
PDD	225.0	152.6	567.4	210.9	153.2	105.5	413.8	146.0	103.4	68.6	317.4	96.9	128.3	63.8	372.9	120.7	0.23	0.065	0.47	0.21
PDE	453.5	263.1	1171.4	395.3	318.2	155.5	755.6	283.8	237.8	100.2	789.0	204.6	251.8	145.4	549.5	224.0	0.40	0.060	0.94	0.39
PDF	156.1	121.7	234.4	152.9	102.5	64.5	133.2	102.1	64.0	48.7	79.9	64.4	85.4	57.2	109.0	84.9	0.12	0.001	0.32	0.12
PDG	162.1	126.9	208.5	160.8	105.6	80.5	143.1	102.8	67.6	49.4	99.2	66.4	88.2	39.8	138.9	86.7	0.17	0.024	0.43	0.19
PDH	208.5	86.3	544.7	173.8	136.6	57.5	373.3	117.3	94.6	32.2	370.7	73.6	119.1	40.3	308.1	97.1	0.21	0.004	0.54	0.20

Table A3.6 Chromium

Site	Extract (all in mg kg <sup>-1</sup> dm)																			
	Cr Aqua Regia				Cr 0.43 M HNO <sub>3</sub>				Cr 0.1 M HCl				Cr 0.05 M EDTA				Cr 0.01 M CaCl <sub>2</sub>			
	mean	min	max	med	mean	min	max	med	mean	min	max	med	mean	min	max	med	mean	min	max	med
CH6	115.6	56.7	483.0	106.0	8.5	2.1	82.4	7.0	2.7	0.31	31.58	2.17	1.02	0.10	6.96	0.88	0.018	< 0.001	0.153	0.016
CH7	101.1	50.2	200.3	90.4	10.0	2.9	28.8	6.9	4.0	0.83	12.77	2.83	2.00	0.06	6.96	1.23	0.013	< 0.001	0.036	0.012
CH8	66.5	41.2	146.7	61.9	2.4	0.9	22.9	1.9	0.7	0.04	11.65	0.57	0.22	0.01	5.32	0.13	0.013	< 0.001	0.043	0.010
HC3	87.9	56.3	149.8	83.8	8.6	4.4	16.7	8.0	2.6	0.53	4.81	2.45	1.85	0.70	3.29	1.78	0.036	0.015	0.080	0.034
HC4	159.7	90.3	259.1	153.6	13.7	6.3	34.1	11.7	5.4	1.54	18.98	4.20	2.47	1.24	5.20	2.26	0.034	0.006	0.061	0.034
HC5	99.9	50.1	267.4	85.5	7.3	2.9	37.5	5.3	1.8	0.26	17.29	1.01	1.43	0.29	5.53	1.07	0.041	0.012	0.185	0.036
HM3	120.8	62.3	345.7	100.8	9.0	3.0	42.1	6.0	3.3	0.96	13.96	2.26	0.92	0.03	5.08	0.67	0.016	< 0.001	0.043	0.016
HM4	153.2	64.0	468.4	126.3	13.4	3.4	60.6	8.2	4.6	1.05	20.80	3.22	1.20	0.02	5.27	0.89	0.014	< 0.001	0.045	0.012
HM5	153.7	64.9	450.9	125.1	11.4	3.7	67.6	7.4	3.8	0.12	17.52	2.71	1.32	0.00	5.63	0.89	0.018	< 0.001	0.043	0.017
LK1	93.6	44.3	189.0	86.4	9.0	2.0	33.5	6.4	3.7	0.62	15.93	2.47	1.59	0.11	5.89	1.20	0.015	< 0.001	0.044	0.014
LK2	130.1	57.1	343.5	103.1	13.1	4.4	68.3	8.3	4.9	1.49	19.30	3.38	2.23	0.56	9.15	1.55	0.014	< 0.001	0.044	0.011
PDA	54.0	43.1	93.3	53.6	4.2	2.5	8.1	4.1	0.6	0.03	1.86	0.60	0.14	0.02	0.50	0.10	0.013	< 0.001	0.030	0.013
PDB	53.3	43.0	76.2	53.3	2.6	1.0	4.7	2.5	0.2	0.01	0.80	0.10	na	na	na	na	0.013	< 0.001	0.037	0.013
PDC	47.3	38.9	58.0	46.9	2.4	1.8	3.3	2.4	0.1	0.01	0.52	0.08	na	na	na	na	0.013	< 0.001	0.030	0.013
PDD	51.3	41.4	149.1	50.4	2.1	1.3	4.1	2.1	0.2	0.01	0.58	0.07	na	na	na	na	0.013	< 0.001	0.031	0.013
PDE	52.1	38.9	70.2	52.0	2.5	1.6	4.4	2.4	0.1	0.01	0.55	0.08	na	na	na	na	0.012	< 0.001	0.038	0.012
PDF	53.1	41.1	95.3	52.9	2.2	1.4	3.0	2.1	0.1	0.01	0.37	0.09	na	na	na	na	0.014	< 0.001	0.030	0.013
PDG	52.6	44.1	104.4	51.9	2.1	1.6	3.2	2.1	0.1	0.03	0.07	0.06	0.07	0.07	0.07	0.07	0.014	< 0.001	0.053	0.014
PDH	52.9	43.3	72.0	52.6	2.0	1.4	3.3	1.8	0.1	0.01	0.22	0.15	na	na	na	na	0.012	< 0.001	0.032	0.012



Table A3.7 Copper

Site	Extract (all in mg kg <sup>-1</sup> dm)																			
	Cu Aqua Regia				Cu 0.43 M HNO <sub>3</sub>				Cu 0.1 M HCl				Cu 0.05 M EDTA				Cu 0.01 M CaCl <sub>2</sub>			
	mean	min	max	med	mean	min	max	med	mean	min	max	med	mean	min	max	med	mean	min	max	med
CH6	130.0	72.2	765.9	109.8	76.5	42.9	573.8	64.6	65.3	44.5	367.6	55.9	76.4	39.5	347.3	66.5	0.277	0.009	4.466	0.170
CH7	315.3	64.1	903.5	215.8	202.8	45.3	635.8	129.0	187.8	39.3	640.3	113.3	195.4	35.1	539.3	132.3	0.518	0.012	9.242	0.210
CH8	162.3	72.8	695.5	158.4	87.5	47.6	438.9	84.6	71.4	40.9	401.4	69.5	88.4	46.5	429.1	86.3	0.052	<0.001	0.833	0.037
HC3	32.1	22.9	196.0	29.6	18.6	12.1	128.6	17.3	15.0	10.0	117.5	13.5	15.7	9.3	118.0	14.5	0.065	0.013	0.717	0.057
HC4	205.7	29.8	310.7	202.3	134.9	17.3	211.1	132.2	121.2	14.6	206.7	119.9	117.0	13.1	198.7	118.9	0.328	0.037	0.753	0.287
HC5	105.8	32.9	299.8	90.7	67.7	17.3	259.2	53.1	57.0	14.9	188.2	44.9	60.0	14.6	204.7	47.2	0.370	0.015	2.180	0.252
HM3	134.9	71.8	420.9	102.9	83.7	45.0	289.8	63.6	71.9	36.4	261.6	54.1	82.9	41.3	263.7	63.8	0.052	<0.001	0.971	0.029
HM4	136.1	55.3	556.2	105.7	80.6	34.6	292.1	61.2	70.4	27.5	262.0	53.8	78.9	29.5	285.4	62.1	0.043	<0.001	0.401	0.023
HM5	112.5	57.9	610.5	87.0	67.6	34.0	417.6	54.2	58.7	30.7	386.7	46.5	67.0	24.1	418.1	54.5	0.094	0.004	0.844	0.075
LK1	173.3	51.5	509.1	142.2	113.2	29.7	336.4	88.9	101.3	27.6	299.2	81.4	111.0	31.7	315.0	93.7	0.478	0.006	2.533	0.364
LK2	179.5	79.4	489.5	137.8	113.6	54.8	338.8	87.9	102.7	48.2	313.2	77.3	116.4	58.0	307.5	91.1	0.151	0.003	1.009	0.095
PDA	124.2	49.8	245.7	123.5	63.6	27.0	128.1	62.5	52.7	23.2	102.3	52.3	47.6	19.6	82.9	50.4	0.082	0.001	0.192	0.096
PDB	58.8	34.1	163.9	55.5	31.4	14.5	281.9	26.7	24.4	12.4	180.9	22.2	23.5	9.9	54.9	21.9	0.039	<0.001	0.161	0.039
PDC	55.1	35.1	67.8	56.2	30.6	19.6	41.8	30.6	24.3	14.9	31.2	24.2	27.9	14.8	36.9	28.2	0.047	0.001	0.092	0.052
PDD	35.8	26.0	62.3	34.8	17.6	11.4	56.8	16.6	13.9	10.5	33.0	13.3	15.7	11.0	36.3	15.1	0.030	0.005	0.083	0.030
PDE	48.4	36.1	78.4	45.5	26.0	15.9	45.8	24.8	20.2	13.1	34.1	18.7	23.0	14.4	40.3	21.5	0.044	0.003	0.100	0.050
PDF	33.6	26.6	40.6	33.8	16.6	11.6	24.1	16.6	12.8	9.5	15.3	12.9	14.8	8.9	18.4	14.9	0.035	0.004	0.133	0.034
PDG	31.3	25.5	42.2	30.9	14.9	11.2	21.4	14.7	11.7	9.1	17.3	11.3	14.0	5.0	21.9	13.8	0.032	0.001	0.157	0.034
PDH	33.5	23.7	56.5	32.0	14.8	9.8	29.3	13.6	11.4	6.7	23.3	10.3	13.6	6.5	24.9	12.9	0.027	0.001	0.164	0.024



## Appendix 4 Overview of plant data

Table A4.1 Cadmium

JAPONICA

Site	Roots						Stem						Leaf						Husk						Rice					
	Min	5%	Median	Mean	95%	Max	Min	5%	Median	Mean	95%	Max	Min	5%	Median	Mean	95%	Max	Min	5%	Median	Mean	95%	Max	Min	5%	Median	Mean	95%	Max
CH6	5.13	42.48	101.92	107.81	225.65	272.20	0.72	2.38	7.99	9.87	21.35	30.79	0.45	3.49	10.83	11.50	22.16	30.24	1.59	2.33	4.60	7.56	17.12	20.06	1.52	2.41	4.01	4.36	7.48	9.69
CH7	1.15	9.60	33.22	42.51	119.07	178.44	0.26	0.79	2.56	3.01	6.90	10.30	0.31	0.76	2.00	2.26	4.21	5.98	0.59	0.88	2.39	2.91	5.94	16.99	0.41	0.67	1.80	1.86	3.30	4.45
CH8	0.16	3.02	9.03	14.35	31.63	348.54	0.01	0.20	1.32	1.71	3.95	11.42	0.08	0.38	1.36	1.86	4.77	16.49	0.47	0.64	1.34	1.54	2.41	17.66	0.51	0.61	1.12	1.35	3.23	4.42
HC3	28.40	31.68	43.85	45.95	65.88	81.86	2.32	4.09	8.52	8.98	14.71	17.21	1.71	2.23	4.41	4.69	7.80	11.29	2.15	2.61	4.59	4.67	7.45	8.54	2.78	2.98	4.52	4.59	6.51	7.54
HC4	21.99	26.38	33.90	35.49	48.94	58.47	2.77	3.71	7.87	8.13	12.67	16.14	0.19	0.61	2.65	2.91	5.28	6.30	1.68	2.00	3.60	3.74	5.14	9.93	2.01	2.24	3.04	3.40	5.09	11.27
HC5	18.24	22.06	49.65	49.76	86.73	117.88	3.26	5.38	12.47	13.59	24.96	32.06	0.62	1.59	4.77	5.53	11.48	13.95	1.63	2.64	5.14	5.22	8.33	12.97	1.90	2.29	4.38	4.81	8.15	9.56
HM3	2.77	16.39	45.42	52.51	101.66	193.58	0.25	0.52	2.39	2.95	7.54	11.31	0.27	0.87	2.64	2.86	5.73	10.53	0.71	0.88	2.59	2.85	5.63	7.96	0.53	0.91	1.97	1.97	3.07	4.07
HM4	17.29	25.33	54.96	62.38	119.39	183.20	0.05	0.57	2.91	3.94	11.75	14.65	0.09	0.78	2.61	3.42	8.40	12.49	0.70	1.39	3.06	3.66	7.99	11.62	0.28	1.50	2.54	2.77	4.85	5.65
HM5	10.65	35.18	78.52	83.32	135.51	311.93	0.65	1.69	5.08	5.83	12.34	15.22	0.49	1.10	4.07	4.37	9.41	12.87	1.10	1.86	4.94	5.27	10.17	12.47	0.96	1.91	3.34	3.44	5.31	7.37
LK1	13.45	18.70	45.97	53.38	107.50	141.32	0.55	1.41	4.47	5.40	11.83	17.13	1.16	1.65	4.61	5.87	13.72	18.27	1.34	2.12	3.92	5.28	11.34	21.08	0.81	1.45	2.81	2.96	4.86	5.97
LK2	34.06	50.07	96.09	103.28	194.32	319.07	0.09	1.00	4.20	5.24	12.40	17.53	0.37	2.85	7.42	8.61	18.01	25.52	1.37	1.96	3.98	4.87	9.69	13.67	0.69	1.71	3.00	3.27	5.76	7.53
PDA	0.18	1.37	6.21	6.70	13.47	18.35	0.01	0.25	1.42	1.56	3.15	4.61	0.05	0.31	1.29	1.39	2.80	3.77	0.62	0.70	1.37	1.70	3.12	6.08	0.23	0.31	0.80	0.83	1.34	2.39
PDB	0.10	1.26	5.96	7.16	16.75	25.16	0.06	0.16	1.26	1.50	3.67	6.14	0.04	0.13	1.03	1.15	2.67	3.42	0.41	0.50	0.90	1.03	1.88	2.79	0.00	0.12	0.52	0.51	0.91	1.01
PDC	0.56	1.19	6.55	6.51	12.08	15.14	0.14	0.20	1.46	1.49	2.88	4.85	0.23	0.50	1.42	1.51	2.90	3.04	0.68	0.74	1.21	1.73	3.54	5.35	0.03	0.12	0.84	0.84	1.31	2.06
PDD	0.13	0.55	6.09	7.03	17.89	18.98	0.05	0.11	1.02	1.27	2.97	4.99	0.10	0.23	1.31	1.31	2.56	2.67	0.48	0.51	0.90	1.13	2.19	2.40	0.07	0.23	0.62	0.64	1.12	1.44
PDE	0.18	0.99	6.55	6.94	14.94	17.00	0.11	0.19	1.23	1.30	3.19	3.86	0.25	0.38	1.49	1.49	2.76	2.99	0.68	0.80	1.43	1.53	2.74	3.27	0.39	0.56	0.95	1.02	1.62	1.89
PDF	0.08	0.56	4.47	5.05	11.25	16.68	0.11	0.23	0.97	1.11	2.46	2.94	0.05	0.26	0.88	1.03	2.16	3.53	0.44	0.46	0.68	0.98	1.99	2.06	0.00	0.03	0.41	0.38	0.82	0.95
PDG	0.18	0.38	5.19	5.73	12.29	14.35	0.03	0.08	1.34	1.37	3.02	4.24	0.07	0.09	1.47	1.39	3.07	3.46	0.49	0.55	0.89	1.05	1.86	1.92	0.06	0.20	0.56	0.56	1.04	1.07
PDH	0.05	0.88	6.12	6.62	14.09	20.52	0.09	0.23	1.14	1.30	2.67	3.46	0.02	0.07	1.21	1.16	2.48	3.05	0.36	0.45	0.97	1.10	2.14	3.54	0.01	0.06	0.58	0.58	1.06	1.88

INDICA

Site	Root						Stem						Leaf						Husk						Rice					
	Min	5%	Median	Mean	95%	Max	Min	5%	Median	Mean	95%	Max	Min	5%	Median	Mean	95%	Max	Min	5%	Median	Mean	95%	Max	Min	5%	Median	Mean	95%	Max
CH6	53.94	80.54	117.78	139.83	294.76	368.08	1.72	4.27	12.46	13.73	28.69	36.47	1.61	4.36	9.95	11.07	19.02	26.02	1.44	2.28	3.86	5.19	10.77	17.40	0.83	2.16	4.60	4.91	8.15	8.89
CH7	5.65	10.88	41.96	56.82	140.34	275.00	0.46	0.73	3.74	3.77	7.56	11.23	0.10	0.51	2.40	2.60	5.29	16.91	0.45	0.61	1.73	1.79	3.13	4.31	0.60	0.79	2.43	2.47	4.05	5.33
CH8	2.17	3.94	11.87	15.27	42.58	98.95	0.05	0.42	2.08	2.46	5.63	7.41	0.01	0.19	1.46	1.76	4.14	5.14	0.19	0.58	1.25	1.61	3.83	10.64	0.79	0.95	1.80	2.08	3.35	7.35
HC3	30.76	33.29	40.31	42.67	54.85	57.54	6.34	6.68	9.82	9.94	14.54	16.56	3.37	3.59	6.09	6.28	9.69	10.22	3.84	3.90	5.12	5.35	7.68	9.27	3.99	4.43	6.53	6.55	9.60	9.90
HC4	21.46	23.12	33.53	33.00	46.93	47.55	3.57	4.19	8.63	8.88	14.05	16.80	1.15	1.66	4.21	4.15	6.23	10.01	2.24	2.57	3.58	3.64	5.10	5.22	2.78	2.95	4.39	4.31	5.96	6.15
HC5	3.13	22.34	53.48	55.90	93.16	113.70	3.81	4.83	14.79	16.50	35.53	47.41	0.67	1.59	7.08	7.84	16.91	21.85	1.96	2.28	4.91	5.26	9.11	13.68	2.46	2.78	6.37	6.59	11.76	13.65
HM3	15.20	24.06	47.55	50.76	85.50	132.97	0.15	0.53	2.47	3.35	8.06	10.31	0.02	0.15	2.71	2.82	5.66	7.56	0.57	0.80	1.74	1.90	3.85	4.32	0.61	0.79	2.18	2.45	4.41	5.56
HM4	25.46	33.93	57.45	65.52	137.79	175.11	0.35	0.80	4.34	5.89	14.73	15.97	0.18	0.31	3.65	3.78	8.04	9.80	0.78	1.35	2.39	2.75	5.09	7.14	0.89	1.55	3.09	3.26	5.50	6.05
HM5	36.04	41.46	87.12	91.89	187.51	236.37	0.49	3.52	8.40	10.09	21.37	23.29	1.96	2.27	4.99	5.67	11.56	17.22	1.88	2.33	3.69	4.13	6.61	8.49	0.79	2.80	4.53	4.60	7.17	8.05
LK1	23.98	30.64	50.04	58.57	116.19	149.72	1.27	2.94	7.31	8.15	16.05	19.90	0.89	1.94	5.91	6.66	13.90	19.13	1.58	2.04	3.51	4.28	9.08	10.06	0.58	2.11	3.77	3.90	6.21	7.38
LK2	19.54	49.33	105.72	111.37	189.12	366.76	1.11	1.91	5.80	6.70	14.04	16.44	1.50	3.30	6.77	7.81	15.75	22.89	1.05	1.47	2.69	2.95	5.39	7.83	1.06	2.14	3.82	3.84	5.95	6.58
PDA	0.02	1.29	5.65	6.42	13.60	46.60	0.31	0.66	1.80	1.97	3.45	4.52	0.08	0.26	1.60	1.55	2.70	3.25	0.53	0.65	1.17	2.07	4.06	4.88	0.04	0.04	0.92	0.92	1.88	3.42
PDB	0.09	0.62	3.61	3.89	8.13	10.19	0.11	0.24	1.17	1.22	2.57	3.29	0.03	0.10	1.14	1.04	1.95	3.24	0.38	0.48	0.84	1.43	3.02	3.37	0.17	0.20	0.60	0.94	1.70	9.35
PDC	1.01	1.02	4.96	4.51	7.36	7.47	0.29	0.35	1.09	1.20	2.29	2.85	0.30	0.56	1.12	1.57	3.21	4.49	0.65	0.73	1.45	1.82	3.21	3.38	0.03	0.25	1.05	1.19	2.36	2.58
PDD	0.41	0.77	3.41	3.50	7.67	7.82	0.21	0.21	0.89	1.18	2.89	3.46	0.55	0.66	1.11	1.38	3.02	3.41	0.53	0.56	0.87	1.40	2.81	2.95	0.03	0.03	0.64	0.68	1.45	1.48
PDE	1.70	2.24	5.12	5.63	10.68	11.17	0.03	0.13	1.36	1.32	2.70	3.29	0.15	0.52	1.16	1.39	2.61	3.43	0.60	0.83	1.63	1.64	2.68	2.74	0.95	1.21	1.46	1.55	2.25	2.50
PDF	0.18	0.63	2.86	3.33	7.07	8.08	0.14	0.15	0.63	0.74	1.47	1.59	0.10	0.11	0.84	1.00	2.13	2.50	0.36	0.39	0.60	0.94	1.77	1.90	0.00	0.02	0.34	0.51	1.21	1.26
PDG	0.02	0.74	3.84	3.98	8.44	8.81	0.14	0.18	1.32	1.19	2.10	2.34	0.02	0.06	0.79	0.91	2.09	2.33	0.44	0.50	1.14	1.14	1.91	1.99	0.25	0.32	0.63	0.79	1.27	1.47
PDH	0.24	0.47	4.77	4.62	8.52	9.51	0.13	0.33	1.30	1.27	2.35	2.43	0.09	0.25	1.08	1.26	2.63	3.41	0.36	0.45	1.07	1.30	2.65	3.86	0.11	0.12	1.00	0.96	1.87	2.34

Table A4.2 Zinc

JAPONICA

Site	Roots						Stem						Leaf						Husk						Rice					
	Min	5%	Median	Mean	95%	Max	Min	5%	Median	Mean	95%	Max	Min	5%	Median	Mean	95%	Max	Min	5%	Median	Mean	95%	Max	Min	5%	Median	Mean	95%	Max
CH6	35.88	56.54	119.99	126.09	245.00	297.50	106.20	172.30	452.90	502.20	884.80	4465.90	27.07	50.50	137.60	133.10	231.80	366.00	26.54	29.67	53.83	57.05	91.03	117.60	17.56	23.84	29.67	30.16	38.01	45.34
CH7	18.22	24.19	68.59	74.39	168.00	248.80	36.40	47.50	192.30	229.90	487.00	2200.00	19.55	34.36	96.00	116.80	248.70	437.70	20.40	25.18	57.22	62.92	117.95	182.10	17.60	20.33	27.46	27.81	35.40	42.75
CH8	20.39	25.10	46.81	57.08	110.70	294.70	48.10	57.50	132.90	189.70	422.80	1750.60	15.83	24.96	42.00	51.70	107.60	259.00	13.00	16.49	24.15	25.37	37.09	103.60	9.93	13.45	19.98	20.13	27.45	34.39
HCS	28.68	32.53	52.63	56.06	90.70	102.30	56.50	72.00	151.50	170.00	315.90	345.20	32.54	34.19	83.40	89.40	198.50	268.40	26.42	29.97	40.26	40.88	56.07	60.90	20.22	21.60	27.38	26.60	31.27	32.51
HC4	32.17	38.12	56.31	59.64	94.70	119.60	138.90	171.10	240.60	256.60	400.90	447.30	39.14	54.31	99.40	111.90	186.50	208.50	24.29	25.45	42.30	46.18	77.83	91.40	22.50	24.71	28.94	28.96	34.15	44.72
HC5	19.25	25.47	55.64	58.90	101.60	164.30	60.60	105.90	248.60	257.50	459.80	587.80	35.86	48.15	121.90	128.60	241.90	397.80	25.04	27.06	42.40	45.06	68.33	213.40	20.38	21.24	28.06	27.51	34.20	37.24
HM3	20.54	24.53	47.04	51.69	79.70	210.90	29.00	47.00	183.50	208.00	430.60	1014.80	32.78	38.31	59.20	63.20	95.40	301.60	21.05	25.62	36.86	38.38	54.35	155.20	16.69	18.74	25.31	25.01	31.44	38.49
HM4	32.48	36.57	71.08	77.40	140.70	434.00	41.90	119.00	199.80	250.50	508.30	752.50	34.89	49.90	75.30	86.10	132.30	330.00	14.08	33.77	45.30	49.55	74.46	323.30	17.39	21.07	25.96	26.61	34.38	43.86
HM5	27.16	38.10	63.02	70.54	134.30	185.50	90.90	113.60	213.00	241.70	491.00	1034.20	28.88	39.14	65.40	72.00	119.60	230.90	23.98	27.04	37.38	38.93	57.37	81.90	16.32	20.62	26.79	26.86	35.27	37.10
LK1	27.06	77.21	196.42	209.87	384.20	600.10	132.50	310.50	693.00	733.20	1322.70	1937.20	28.24	86.82	189.20	213.00	433.70	925.50	31.35	44.51	64.13	75.61	153.62	223.30	22.13	26.13	33.06	33.68	42.98	50.41
LK2	30.66	41.42	66.08	75.10	131.40	262.30	74.00	90.20	163.80	198.50	476.50	787.30	28.15	33.73	63.80	73.10	140.90	225.10	20.86	23.97	38.30	72.43	71.40	4107.00	14.24	20.15	27.89	28.08	38.28	63.32
PDA	48.92	70.17	109.49	109.48	158.80	193.30	80.30	225.20	459.70	506.40	917.30	1096.00	45.84	65.78	153.40	160.90	308.10	412.00	22.62	29.62	50.24	55.45	95.00	129.20	24.39	28.22	34.80	35.32	43.77	52.90
PDB	12.57	24.77	64.75	63.18	96.90	113.30	30.80	87.40	230.40	231.20	387.00	506.30	19.52	28.96	79.00	82.50	149.60	188.30	15.61	20.92	37.66	40.54	67.71	90.90	20.29	22.49	30.02	30.10	38.62	43.91
PDC	40.17	43.71	74.74	77.38	116.60	121.60	190.20	199.50	318.90	339.90	525.70	799.40	40.61	46.95	93.20	97.80	173.10	260.30	24.48	28.32	45.59	53.20	100.65	165.40	27.88	29.65	33.74	35.46	45.17	48.52
PDD	22.56	26.57	44.55	47.36	76.70	85.60	76.30	89.70	194.50	192.90	303.10	363.70	38.84	45.99	78.90	78.30	115.40	134.30	23.73	25.95	40.61	41.92	60.70	79.00	21.10	23.51	29.12	29.76	38.65	39.70
PDE	36.82	40.32	58.96	62.13	84.60	165.20	111.00	115.40	184.40	187.20	270.90	368.30	32.94	39.66	68.20	71.50	100.50	265.50	26.37	28.98	39.11	41.57	58.15	71.90	19.45	22.72	31.06	31.51	39.94	44.89
PDF	17.08	20.77	35.46	37.21	60.90	69.70	80.40	93.10	156.90	163.70	277.00	322.40	34.16	40.33	68.90	67.70	93.20	118.60	25.14	27.50	39.71	41.82	60.54	68.50	21.89	23.06	29.16	29.49	37.93	39.93
PDG	17.03	23.77	34.30	35.57	48.70	69.80	76.50	91.10	124.10	136.60	244.40	271.40	32.82	35.51	52.80	54.60	79.60	110.50	22.87	26.93	39.69	40.48	57.60	64.80	20.67	23.17	30.85	30.81	39.38	42.93
PDH	15.92	22.28	41.39	43.08	66.60	125.20	70.80	84.40	146.90	157.60	261.70	378.90	33.57	42.86	62.60	65.80	101.20	153.20	20.56	26.19	40.51	44.22	65.46	318.30	20.81	23.78	29.72	30.27	38.17	44.62

INDICA

Site	Root						Stem						Leaf						Husk						Rice					
	Min	5%	Median	Mean	95%	Max	Min	5%	Median	Mean	95%	Max	Min	5%	Median	Mean	95%	Max	Min	5%	Median	Mean	95%	Max	Min	5%	Median	Mean	95%	Max
CH6	75.55	89.91	142.25	147.50	230.40	344.30	222.50	276.70	611.40	748.30	1833.30	2486.60	50.42	62.48	117.30	142.30	303.80	335.40	19.44	25.18	40.66	52.03	93.21	354.60	15.32	22.67	27.38	28.30	36.62	50.74
CH7	23.42	28.79	96.77	101.05	188.60	311.40	44.40	61.40	262.50	313.50	1008.70	1265.30	11.18	33.59	113.50	132.60	356.50	494.90	19.05	21.26	41.79	56.26	166.68	192.70	15.42	20.05	27.31	27.70	35.77	47.97
CH8	24.36	31.99	68.47	74.78	165.30	178.30	68.10	76.90	228.20	280.20	633.10	869.80	20.39	25.00	47.60	57.90	131.80	173.70	15.55	16.40	24.65	28.39	46.78	120.20	14.22	17.24	22.47	22.52	28.64	40.50
HCS	33.83	38.56	72.63	73.09	116.50	129.10	76.90	103.00	243.90	229.30	372.40	407.10	35.05	38.46	111.70	105.60	234.30	261.00	23.15	26.12	38.19	40.74	62.94	70.10	21.13	24.10	28.61	29.02	34.32	40.91
HC4	27.07	38.40	79.30	77.55	121.00	148.80	157.20	176.10	333.00	369.80	682.60	1000.90	63.03	65.30	103.00	122.10	254.30	275.10	22.02	23.98	31.80	36.72	60.33	66.60	23.20	24.52	27.47	27.60	31.19	32.96
HC5	20.12	22.18	72.13	67.19	112.60	140.80	88.60	105.90	362.70	395.90	748.20	925.30	36.97	51.94	107.10	122.70	267.00	288.70	23.22	25.72	34.10	37.29	59.74	64.00	20.45	21.14	26.06	26.29	32.03	35.52
HM3	24.17	30.84	59.41	63.33	105.20	135.90	33.40	73.50	233.50	275.80	624.80	836.10	34.88	45.41	61.30	66.10	106.30	139.60	19.19	22.43	28.98	35.49	49.53	289.70	15.66	19.20	23.95	24.80	32.35	39.21
HM4	44.19	56.35	99.84	103.45	174.30	210.70	72.00	147.00	375.70	490.40	1151.60	2474.40	32.44	46.79	82.30	92.60	193.00	246.00	22.16	24.12	33.32	34.64	47.06	63.90	13.30	20.87	23.74	25.20	34.93	39.06
HM5	46.39	56.93	97.34	103.74	176.60	421.60	191.40	225.30	399.50	531.40	1349.20	2657.80	40.79	46.78	73.80	86.00	147.00	450.30	21.28	26.10	33.78	34.52	48.91	51.60	15.61	20.17	25.71	25.99	33.56	38.42
LK1	68.89	96.31	230.16	253.99	546.80	594.70	177.10	280.60	906.70	1084.70	2590.70	3945.70	70.88	84.46	246.90	265.50	514.00	616.40	30.34	37.80	51.86	60.11	121.03	213.10	20.30	26.20	33.60	34.45	43.51	57.84
LK2	44.24	47.85	86.71	90.00	188.60	309.60	98.50	127.20	250.60	310.60	602.20	1668.80	35.30	38.50	63.00	72.10	120.70	278.20	20.21	22.89	29.44	31.96	48.29	59.90	10.82	21.64	25.71	26.77	35.86	41.38
PDA	47.50	50.80	119.58	122.00	187.80	206.80	144.70	267.20	676.30	764.00	1433.10	1668.80	62.91	88.31	157.80	175.50	316.30	406.30	22.39	24.51	44.15	64.94	136.27	137.20	25.40	26.51	37.21	37.02	52.72	65.39
PDB	29.08	46.05	71.94	75.08	116.30	177.40	131.90	154.50	313.10	322.70	618.80	654.80	44.66	55.62	101.00	102.30	169.10	189.30	16.09	19.62	36.47	52.04	107.40	124.80	22.51	23.69	32.39	35.09	55.24	117.50
PDC	53.01	58.66	100.85	95.73	125.70	134.20	193.80	225.00	455.80	479.70	1125.20	1234.50	70.92	75.72	125.10	129.40	207.90	249.50	23.29	27.05	44.40	53.88	89.31	112.40	28.79	28.88	37.67	38.44	53.66	54.06
PDD	27.17	28.18	49.01	50.62	72.50	93.30	102.80	105.40	241.50	260.40	525.60	530.30	42.90	47.65	100.50	104.10	171.50	175.10	20.09	20.82	36.86	45.27	82.06	83.00	22.80	22.93	30.26	31.72	44.95	45.03
PDE	54.67	56.67	75.35	76.65	112.70	114.20	118.20	128.50	265.40	269.10	450.20	481.50	40.85	44.44	105.60	104.80	194.40	217.30	21.36	25.30	43.86	44.74	70.42	76.80	25.21	27.41	35.77	34.56	41.89	45.46
PDF	19.46	25.70	34.76	36.20	51.00	53.20	108.50	119.40	170.40	202.90	394.50	431.30	51.73	54.47	70.60	84.90	147.90	182.00	17.91	19.28	30.25	40.61	68.72	69.90	22.15	24				

Table A4.3 Nickel

JAPONICA

Site	Roots						Stem						Leaf						Husk						Rice					
	Min	5%	Median	Mean	95%	Max	Min	5%	Median	Mean	95%	Max	Min	5%	Median	Mean	95%	Max	Min	5%	Median	Mean	95%	Max	Min	5%	Median	Mean	95%	Max
CH6	5.13	42.48	101.92	107.81	225.65	272.20	0.72	2.38	7.99	9.87	21.35	30.79	0.45	3.49	10.83	11.50	22.16	30.24	1.59	2.33	4.60	7.56	17.12	20.06	1.52	2.41	4.01	4.36	7.48	9.69
CH7	1.15	9.60	33.22	42.51	119.07	178.44	0.26	0.79	2.56	3.01	6.90	10.30	0.31	0.76	2.00	2.26	4.21	5.98	0.59	0.88	2.39	2.91	5.94	16.99	0.41	0.67	1.80	1.86	3.30	4.45
CH8	0.16	3.02	9.03	14.35	31.63	348.54	0.01	0.20	1.32	1.71	3.95	11.42	0.08	0.38	1.36	1.86	4.77	16.49	0.47	0.64	1.34	1.54	2.41	17.66	0.51	0.61	1.12	1.35	3.23	4.42
HC3	28.40	31.68	43.85	45.95	65.88	81.86	2.32	4.09	8.52	8.98	14.71	17.21	1.71	2.23	4.41	4.69	7.80	11.29	2.15	2.61	4.59	4.67	7.45	8.54	2.78	2.98	4.52	4.59	6.51	7.54
HC4	21.99	26.38	33.90	35.49	48.94	58.47	2.77	3.71	7.87	8.13	12.67	16.14	0.19	0.61	2.65	2.91	5.28	6.30	1.68	2.00	3.60	3.74	5.14	9.93	2.01	2.24	3.04	3.40	5.09	11.27
HC5	18.24	22.06	49.65	49.76	86.73	117.88	3.26	5.38	12.47	13.59	24.96	32.06	0.62	1.59	4.77	5.53	11.48	13.95	1.63	2.64	5.14	5.22	8.33	12.97	1.90	2.29	4.38	4.81	8.15	9.56
HM3	2.77	16.39	45.42	52.51	101.66	193.58	0.25	0.52	2.39	2.95	7.54	11.31	0.27	0.87	2.64	2.86	5.73	10.53	0.71	0.88	2.59	2.85	5.63	7.96	0.53	0.91	1.97	1.97	3.07	4.07
HM4	17.29	25.33	54.96	62.38	119.39	183.20	0.05	0.57	2.91	3.94	11.75	14.65	0.09	0.78	2.61	3.42	8.40	12.49	0.70	1.39	3.06	3.66	7.99	11.62	0.28	1.50	2.54	2.77	4.85	5.65
HM5	10.65	35.18	78.52	83.32	135.51	311.93	0.65	1.69	5.08	5.83	12.34	15.22	0.49	1.10	4.07	4.37	9.41	12.87	1.10	1.86	4.94	5.27	10.17	12.47	0.96	1.91	3.34	3.44	5.31	7.37
LK1	13.45	18.70	45.97	53.38	107.50	141.32	0.55	1.41	4.47	5.40	11.83	17.13	1.16	1.65	4.61	5.87	13.72	18.27	1.34	2.12	3.92	5.28	11.34	21.08	0.81	1.45	2.81	2.96	4.86	5.97
LK2	34.06	50.07	96.09	103.28	194.32	319.07	0.09	1.00	4.20	5.24	12.40	17.53	0.37	2.85	7.42	8.61	18.01	25.52	1.37	1.96	3.98	4.87	9.69	13.67	0.69	1.71	3.00	3.27	5.76	7.53
PDA	0.18	1.37	6.21	6.70	13.47	18.35	0.01	0.25	1.42	1.56	3.15	4.61	0.05	0.31	1.29	1.39	2.80	3.77	0.62	0.70	1.37	1.70	3.12	6.08	0.23	0.31	0.80	0.83	1.34	2.39
PDB	0.10	1.26	5.96	7.16	16.75	25.16	0.06	0.16	1.26	1.50	3.67	6.14	0.04	0.13	1.03	1.15	2.67	3.42	0.41	0.50	0.90	1.03	1.88	2.79	0.00	0.12	0.52	0.51	0.91	1.01
PDC	0.56	1.19	6.55	6.51	12.08	15.14	0.14	0.20	1.46	1.49	2.88	4.85	0.23	0.50	1.42	1.51	2.90	3.04	0.68	0.74	1.21	1.73	3.54	5.35	0.03	0.12	0.84	0.84	1.31	2.06
PDD	0.13	0.55	6.09	7.03	17.89	18.98	0.05	0.11	1.02	1.27	2.97	4.99	0.10	0.23	1.31	1.31	2.56	2.67	0.48	0.51	0.90	1.13	2.19	2.40	0.07	0.23	0.62	0.64	1.12	1.44
PDE	0.18	0.99	6.55	6.94	14.94	17.00	0.11	0.19	1.23	1.30	3.19	3.86	0.25	0.38	1.49	1.49	2.76	2.99	0.68	0.80	1.43	1.53	2.74	3.27	0.39	0.56	0.95	1.02	1.62	1.89
PDF	0.08	0.56	4.47	5.05	11.25	16.68	0.11	0.23	0.97	1.11	2.46	2.94	0.05	0.26	0.88	1.03	2.16	3.53	0.44	0.46	0.68	0.98	1.99	2.06	0.00	0.03	0.41	0.38	0.82	0.95
PDG	0.18	0.38	5.19	5.73	12.29	14.35	0.03	0.08	1.34	1.37	3.02	4.24	0.07	0.09	1.47	1.39	3.07	3.46	0.49	0.55	0.89	1.05	1.86	1.92	0.06	0.20	0.56	0.56	1.04	1.07
PDH	0.05	0.88	6.12	6.62	14.09	20.52	0.09	0.23	1.14	1.30	2.67	3.46	0.02	0.07	1.21	1.16	2.48	3.05	0.36	0.45	0.97	1.10	2.14	3.54	0.01	0.06	0.58	0.58	1.06	1.88

INDICA

Site	Root						Stem						Leaf						Husk						Rice					
	Min	5%	Median	Mean	95%	Max	Min	5%	Median	Mean	95%	Max	Min	5%	Median	Mean	95%	Max	Min	5%	Median	Mean	95%	Max	Min	5%	Median	Mean	95%	Max
CH6	53.94	80.54	117.78	139.83	294.76	368.08	1.72	4.27	12.46	13.73	28.69	36.47	1.61	4.36	9.95	11.07	19.02	26.02	1.44	2.28	3.86	5.19	10.77	17.40	0.83	2.16	4.60	4.91	8.15	8.89
CH7	5.65	10.88	41.96	56.82	140.34	275.00	0.46	0.73	3.74	3.77	7.56	11.23	0.10	0.51	2.40	2.60	5.29	16.91	0.45	0.61	1.73	1.79	3.13	4.31	0.60	0.79	2.43	2.47	4.05	5.33
CH8	2.17	3.94	11.87	15.27	42.58	98.95	0.05	0.42	2.08	2.46	5.63	7.41	0.01	0.19	1.46	1.76	4.14	5.14	0.19	0.58	1.25	1.61	3.83	10.64	0.79	0.95	1.80	2.08	3.35	7.35
HC3	30.76	33.29	40.31	42.67	54.85	57.54	6.34	6.68	9.82	9.94	14.54	16.56	3.37	3.59	6.09	6.28	9.69	10.22	3.84	3.90	5.12	5.35	7.68	9.27	3.99	4.43	6.53	6.55	9.60	9.90
HC4	21.46	23.12	33.53	33.00	46.93	47.55	3.57	4.19	8.63	8.88	14.05	16.80	1.15	1.66	4.21	4.15	6.23	10.01	2.24	2.57	3.58	3.64	5.10	5.22	2.78	2.95	4.39	4.31	5.96	6.15
HC5	3.13	22.34	53.48	55.90	93.16	113.70	3.81	4.83	14.79	16.50	35.53	47.41	0.67	1.59	7.08	7.84	16.91	21.85	1.96	2.28	4.91	5.26	9.11	13.68	2.46	2.78	6.37	6.59	11.76	13.65
HM3	15.20	24.06	47.55	50.76	85.50	132.97	0.15	0.53	2.47	3.35	8.06	10.31	0.02	0.15	2.71	2.82	5.66	7.56	0.57	0.80	1.74	1.90	3.85	4.32	0.61	0.79	2.18	2.45	4.41	5.56
HM4	25.46	33.93	57.45	65.52	137.79	175.11	0.35	0.80	4.34	5.89	14.73	15.97	0.18	0.31	3.65	3.78	8.04	9.80	0.78	1.35	2.39	2.75	5.09	7.14	0.89	1.55	3.09	3.26	5.50	6.05
HM5	36.04	41.46	87.12	91.89	187.51	236.37	0.49	3.52	8.40	10.09	21.37	23.29	1.96	2.27	4.99	5.67	11.56	17.22	1.88	2.33	3.69	4.13	6.61	8.49	0.79	2.80	4.53	4.60	7.17	8.05
LK1	23.98	30.64	50.04	58.57	116.19	149.72	1.27	2.94	7.31	8.15	16.05	19.90	0.89	1.94	5.91	6.66	13.90	19.13	1.58	2.04	3.51	4.28	9.08	10.06	0.58	2.11	3.77	3.90	6.21	7.38
LK2	19.54	49.33	105.72	111.37	189.12	366.76	1.11	1.91	5.80	6.70	14.04	16.44	1.50	3.30	6.77	7.81	15.75	22.89	1.05	1.47	2.69	2.95	5.39	7.83	1.06	2.14	3.82	3.84	5.95	6.58
PDA	0.02	1.29	5.65	6.42	13.60	46.60	0.31	0.66	1.80	1.97	3.45	4.52	0.08	0.26	1.60	1.55	2.70	3.25	0.53	0.65	1.17	2.07	4.06	4.88	0.04	0.04	0.92	0.92	1.88	3.42
PDB	0.09	0.62	3.61	3.89	8.13	10.19	0.11	0.24	1.17	1.22	2.57	3.29	0.03	0.10	1.14	1.04	1.95	3.24	0.38	0.48	0.84	1.43	3.02	3.37	0.17	0.20	0.60	0.94	1.70	9.35
PDC	1.01	1.02	4.96	4.51	7.36	7.47	0.29	0.35	1.09	1.20	2.29	2.85	0.30	0.56	1.52	1.57	3.21	4.49	0.65	0.73	1.45	1.82	3.21	3.38	0.03	0.25	1.05	1.19	2.36	2.58
PDD	0.41	0.77	3.41	3.50	7.67	7.82	0.21	0.21	0.89	1.18	2.89	3.46	0.55	0.66	1.11	1.38	3.02	3.41	0.53	0.56	0.87	1.40	2.81	2.95	0.03	0.03	0.64	0.68	1.45	1.48
PDE	1.70	2.24	5.12	5.63	10.68	11.17	0.03	0.13	1.36	1.32	2.70	3.29	0.15	0.52	1.36	1.39	2.61	3.43	0.60	0.83	1.63	1.64	2.68	2.74	0.95	1.21	1.46	1.55	2.25	2.50
PDF	0.18	0.63	2.86	3.33	7.07	8.08	0.14	0.15	0.63	0.74	1.47	1.59	0.10	0.11	0.84	1.00	2.13	2.50	0.36	0.39	0.60	0.94	1.77	1.90	0.00	0.02	0.34	0.51	1.21	1.26
PDG	0.02	0.74	3.84	3.98	8.44	8.81	0.14	0.18	1.32	1.19	2.10	2.34	0.02	0.06	0.79	0.91	2.09	2.33	0.44	0.50	1.14	1.14	1.91	1.99	0.25	0.32	0.63	0.79	1.27	1.47
PDH	0.24	0.47	4.77	4.62	8.52	9.51	0.13	0.33	1.30	1.27	2.35	2.43	0.09	0.25	1.08	1.26	2.63	3.41	0.36	0.45	1.07	1.30	2.65	3.86	0.11	0.12	1.00	0.96	1.87	2.34

Table A4.4 Lead

JAPONICA

Site	Roots						Stem						Leaf						Husk						Rice					
	Min	5%	Median	Mean	95%	Max	Min	5%	Median	Mean	95%	Max	Min	5%	Median	Mean	95%	Max	Min	5%	Median	Mean	95%	Max	Min	5%	Median	Mean	95%	Max
CH6	4.89	9.98	20.22	24.51	55.38	136.40	0.02	0.39	1.81	2.11	5.19	8.01	0.05	0.56	2.86	3.20	7.19	10.07	0.06	0.21	1.26	2.12	7.54	12.10	0.01	0.05	0.17	0.20	0.43	1.03
CH7	3.14	3.68	8.10	10.37	27.80	36.30	0.05	0.33	1.31	1.82	4.66	8.81	0.19	0.46	2.03	2.89	5.93	9.14	0.27	0.51	2.28	3.31	9.48	25.72	0.00	0.03	0.19	0.23	0.44	1.81
CH8	2.19	3.44	7.23	8.87	19.01	44.70	0.11	0.24	1.47	1.95	5.11	6.19	0.11	0.56	3.90	4.54	11.17	20.53	0.17	0.24	0.89	4.06	14.28	24.87	0.02	0.06	0.18	0.20	0.43	1.27
HC3	3.17	3.93	7.00	7.50	12.45	15.90	0.04	0.25	1.80	1.76	3.28	4.28	0.42	0.85	2.40	2.48	4.20	5.00	0.32	0.42	0.76	0.90	1.55	7.36	0.02	0.04	0.10	0.11	0.20	0.42
HC4	3.12	4.70	9.98	10.87	22.00	27.00	0.08	0.53	2.27	2.28	4.60	5.05	0.30	0.70	2.56	2.44	4.04	4.40	0.35	0.39	0.66	0.74	1.22	2.08	0.00	0.02	0.10	0.10	0.20	0.68
HCS	2.82	4.02	6.60	7.50	13.49	20.30	0.09	0.87	2.56	2.51	4.12	4.66	0.08	0.49	2.32	2.22	4.15	5.15	0.32	0.39	0.67	0.76	1.54	2.65	0.00	0.02	0.09	0.10	0.18	1.14
HM3	2.45	4.42	9.09	9.93	17.55	33.50	0.04	0.20	1.21	1.94	5.40	12.72	0.04	0.32	2.43	2.68	6.06	11.07	0.22	0.32	1.70	2.38	6.44	13.57	0.01	0.04	0.17	0.20	0.40	1.84
HM4	2.72	3.73	8.12	9.42	20.06	29.20	0.04	0.28	1.02	1.38	4.01	5.41	0.03	0.32	2.29	2.56	5.31	9.61	0.19	0.36	1.47	2.19	6.10	10.64	0.00	0.02	0.15	0.19	0.45	1.99
HMS	3.18	4.36	8.57	9.94	17.83	37.90	0.04	0.11	1.10	1.62	5.04	7.78	0.10	0.30	2.99	3.35	6.62	15.60	0.25	0.37	1.61	2.31	6.51	9.02	0.00	0.02	0.14	0.17	0.42	0.83
LK1	3.16	11.48	31.64	36.53	88.52	154.70	0.01	0.30	1.84	2.48	6.03	27.71	0.18	1.16	4.26	7.04	21.56	99.53	0.21	0.65	3.44	7.13	30.88	68.24	0.01	0.08	0.22	0.25	0.48	1.89
LK2	3.32	4.93	11.34	13.29	30.27	65.30	0.08	0.20	1.01	1.43	4.08	6.65	0.04	0.62	3.29	3.38	6.93	11.09	0.08	0.19	2.61	3.39	10.40	19.03	0.03	0.06	0.16	0.23	0.58	1.96
PDA	51.95	69.40	131.19	136.60	218.05	269.60	9.65	13.35	20.53	21.38	33.67	57.95	2.65	3.84	6.75	6.80	10.50	12.45	0.48	0.74	1.33	1.54	2.85	10.13	0.04	0.08	0.17	0.20	0.43	1.19
PDB	19.48	41.15	105.68	106.77	204.96	257.40	0.92	10.07	21.88	22.80	42.97	67.75	2.15	3.28	6.72	7.07	12.63	19.02	0.57	0.81	1.26	1.83	5.66	11.74	0.00	0.06	0.16	0.20	0.47	1.65
PDC	33.35	35.69	55.44	59.38	84.91	224.80	0.71	4.40	11.19	12.13	21.22	37.37	1.89	2.48	4.37	4.48	7.34	10.10	0.52	0.82	1.05	1.38	2.88	11.12	0.00	0.01	0.13	0.18	0.30	2.15
PDD	15.22	25.20	43.75	45.64	70.57	73.30	0.48	5.34	10.21	11.09	18.88	28.29	1.43	2.11	3.86	4.10	7.20	8.75	0.57	0.65	1.00	1.11	1.66	3.73	0.02	0.04	0.13	0.19	0.48	1.38
PDE	40.70	44.78	67.32	80.58	160.65	247.00	5.00	6.04	10.83	11.80	20.76	36.93	2.45	2.79	4.47	4.83	7.48	11.60	0.59	0.63	1.23	1.24	2.10	2.26	0.01	0.04	0.14	0.17	0.41	0.57
PDF	19.28	20.74	30.10	33.46	56.83	77.60	3.89	5.23	9.18	9.69	16.52	19.85	0.92	1.73	3.88	3.91	6.33	7.05	0.46	0.61	1.07	1.09	1.68	2.28	0.00	0.03	0.13	0.13	0.27	0.30
PDG	12.87	15.80	26.92	29.01	53.34	60.70	2.32	3.53	6.35	7.40	14.49	18.26	1.00	1.51	3.21	3.26	4.87	5.58	0.45	0.60	0.99	1.09	1.82	2.14	0.01	0.03	0.13	0.16	0.35	0.63
PDH	12.10	13.41	38.10	46.42	106.16	302.60	2.39	3.28	8.71	9.54	19.90	33.04	0.03	1.04	3.87	4.12	9.72	11.76	0.45	0.67	1.02	1.09	1.67	2.21	0.01	0.04	0.13	0.16	0.31	0.79

INDICA

Site	Root						Stem						Leaf						Husk						Rice					
	Min	5%	Median	Mean	95%	Max	Min	5%	Median	Mean	95%	Max	Min	5%	Median	Mean	95%	Max	Min	5%	Median	Mean	95%	Max	Min	5%	Median	Mean	95%	Max
CH6	8.89	14.00	25.51	31.65	73.83	120.20	0.02	0.78	1.87	2.13	4.75	5.97	0.27	0.87	2.96	3.47	8.33	9.22	0.08	0.25	0.62	2.14	6.92	12.77	0.01	0.03	0.19	0.32	1.42	2.37
CH7	3.11	4.66	9.41	12.03	28.60	49.70	0.02	0.18	1.22	1.94	6.19	9.12	0.10	0.30	2.50	3.44	11.23	12.31	0.34	0.44	2.03	2.57	6.09	11.24	0.03	0.10	0.22	0.27	0.58	1.37
CH8	3.30	4.31	9.32	10.05	18.40	41.00	0.53	0.55	1.98	2.51	5.74	8.28	0.32	0.58	4.16	4.48	10.30	11.00	0.07	0.15	0.88	4.00	14.30	18.01	0.04	0.11	0.24	0.35	1.56	2.20
HC3	3.42	3.84	7.81	8.02	12.41	15.70	0.49	0.58	1.61	1.70	3.09	4.53	0.59	0.78	2.01	2.28	4.38	6.61	0.49	0.56	0.80	0.99	2.27	3.16	0.02	0.04	0.13	0.14	0.26	0.27
HC4	3.77	5.09	9.29	10.30	18.92	19.50	0.47	0.67	1.82	2.01	3.52	4.29	0.08	0.15	1.52	1.57	2.79	3.10	0.36	0.39	0.67	0.82	1.37	2.61	0.02	0.03	0.11	0.12	0.21	0.23
HCS	3.32	4.99	7.42	11.97	15.84	206.90	0.17	0.37	2.20	2.28	4.05	5.99	0.17	0.34	2.07	2.15	3.97	4.35	0.39	0.43	0.73	0.84	1.45	3.96	0.03	0.04	0.12	0.13	0.27	0.36
HM3	3.43	4.30	8.03	9.62	16.47	42.90	0.01	0.22	1.02	1.34	3.55	3.78	0.07	0.30	2.35	2.77	5.98	8.40	0.26	0.34	1.17	1.81	5.14	7.24	0.02	0.05	0.20	0.24	0.55	1.62
HM4	3.36	4.10	8.57	10.86	25.08	72.50	0.05	0.33	1.18	1.52	5.52	6.91	0.59	1.12	2.82	3.22	7.26	7.86	0.22	0.32	1.08	2.01	7.18	11.85	0.03	0.06	0.16	0.27	1.04	1.61
HMS	3.66	5.11	9.97	11.68	23.52	47.50	0.02	0.15	1.36	1.75	5.26	7.70	0.00	0.20	2.80	3.14	6.79	13.46	0.21	0.27	0.92	2.35	7.78	14.45	0.01	0.05	0.19	0.36	1.50	3.05
LK1	5.51	12.78	29.92	37.32	77.61	122.20	0.24	0.46	2.10	2.59	5.06	25.06	0.05	0.34	4.33	5.64	15.48	25.58	0.25	0.35	3.53	5.25	18.33	26.60	0.06	0.09	0.21	0.24	0.54	0.70
LK2	4.65	6.05	12.50	16.61	39.05	82.20	0.08	0.42	1.35	2.03	6.32	17.75	0.10	0.71	2.59	3.08	6.40	7.63	0.12	0.19	0.86	2.27	7.77	9.87	0.02	0.05	0.17	0.22	0.72	0.87
PDA	10.92	84.39	129.31	128.38	183.49	200.90	9.65	14.68	24.90	26.11	47.61	55.35	2.15	3.65	5.69	6.26	9.93	10.92	0.55	0.58	1.22	1.52	3.40	6.19	0.00	0.01	0.15	0.16	0.32	0.45
PDB	39.39	70.34	108.38	106.49	147.70	185.30	8.92	11.88	20.70	21.29	35.17	43.54	1.43	2.57	5.54	5.99	9.99	10.64	0.49	0.54	0.98	1.26	2.62	3.56	0.03	0.04	0.13	0.17	0.42	0.85
PDC	40.71	42.91	62.55	63.13	85.28	121.00	7.99	8.07	13.49	14.12	20.23	24.39	1.38	1.61	4.33	4.37	6.72	6.90	0.54	0.55	0.78	1.07	2.35	2.42	0.01	0.04	0.15	0.19	0.64	0.78
PDD	21.51	26.27	42.50	44.27	69.54	73.90	5.17	5.80	10.86	10.97	17.03	17.10	1.82	1.92	3.47	3.66	5.95	6.27	0.46	0.52	0.79	0.86	1.38	1.51	0.00	0.01	0.09	0.10	0.27	0.30
PDE	53.95	57.74	79.37	84.37	128.87	150.10	6.44	8.53	16.51	16.00	26.63	30.44	1.52	2.35	4.67	5.14	8.54	8.75	0.49	0.56	0.87	1.12	2.48	2.81	0.05	0.07	0.17	0.17	0.29	0.32
PDF	19.73	21.08	33.74	36.08	64.80	76.70	4.51	4.59	8.50	10.38	20.53	24.57	1.48	1.59	3.86	3.97	7.10	7.34	0.46	0.50	0.81	0.95	1.71	1.84	0.00	0.03	0.17	0.17	0.41	0.46
PDG	21.59	22.44	32.04	32.24	41.18	46.20	2.42	4.53	7.79	8.44	16.68	17.20	0.90	1.20	3.02	3.14	5.45	6.13	0.51	0.53	0.93	0.93	1.58	1.70	0.00	0.02	0.14	0.17	0.44	0.67
PDH	14.66	17.71	40.73	47.85	100.25	130.60	2.31	4.17	8.99	11.48	22.27	45.87	0.37	0.82	2.95	3.57	7.33	15.12	0.50	0.52	0.85	0.95	1.68	2.30	0.04	0.04	0.12	0.16	0.40	0.53

Table A4.5 Chromium

JAPONICA

Site	Roots						Stem						Leaf						Husk						Rice					
	Min	5%	Median	Mean	95%	Max	Min	5%	Median	Mean	95%	Max	Min	5%	Median	Mean	95%	Max	Min	5%	Median	Mean	95%	Max	Min	5%	Median	Mean	95%	Max
CH6	1.89	4.22	16.59	19.67	48.00	131.49	0.01	0.06	0.43	0.52	1.36	2.27	0.01	0.12	0.66	0.94	2.76	9.10	0.09	0.14	0.30	0.44	1.20	4.43	0.10	0.13	0.21	0.22	0.34	0.60
CH7	0.72	4.52	11.56	15.08	33.92	83.49	0.02	0.05	0.42	0.56	1.85	3.86	0.01	0.18	0.70	0.76	1.47	2.27	0.10	0.12	0.31	0.51	1.68	3.82	0.10	0.17	0.26	0.27	0.41	0.71
CH8	0.06	1.23	6.54	7.63	17.20	26.52	0.02	0.08	0.45	0.54	1.18	4.03	0.00	0.08	0.52	0.69	1.56	6.60	0.10	0.14	0.33	0.46	1.35	4.16	0.10	0.14	0.21	0.23	0.34	0.65
HC3	5.11	6.41	15.40	16.97	31.43	72.58	0.13	0.77	2.73	3.19	6.71	10.31	0.01	0.06	0.51	0.57	1.19	1.63	0.24	0.27	0.43	0.44	0.66	0.83	0.06	0.08	0.15	0.15	0.25	0.48
HC4	7.47	10.16	19.40	19.94	33.80	49.42	0.35	0.58	2.41	2.64	5.60	7.40	0.01	0.04	0.43	0.55	1.47	2.03	0.24	0.29	0.41	0.47	0.88	1.93	0.07	0.09	0.14	0.15	0.23	0.30
HC5	3.61	5.52	12.76	14.94	26.86	93.61	0.16	0.66	1.86	2.45	6.13	8.32	0.01	0.06	0.40	0.45	1.00	1.85	0.24	0.28	0.41	0.48	0.83	4.26	0.05	0.08	0.14	0.14	0.20	0.29
HM3	1.85	5.08	14.94	19.41	49.40	104.95	0.01	0.05	0.39	0.47	1.08	1.92	0.03	0.17	0.94	1.00	2.13	4.37	0.07	0.15	0.40	0.55	1.62	4.29	0.10	0.16	0.25	0.26	0.39	0.72
HM4	1.34	5.00	17.73	23.15	65.53	139.49	0.01	0.04	0.34	0.43	1.22	2.59	0.01	0.07	0.54	0.65	1.32	5.85	0.08	0.14	0.34	0.41	0.80	3.10	0.02	0.13	0.25	0.26	0.43	0.76
HM5	4.83	9.02	26.39	31.16	61.21	203.59	0.01	0.04	0.41	0.49	1.45	1.95	0.01	0.04	0.49	0.57	1.25	3.30	0.08	0.13	0.33	0.42	1.21	2.28	0.11	0.16	0.23	0.25	0.46	0.83
LK1	2.62	6.65	25.74	27.71	54.53	80.63	0.02	0.07	0.39	0.43	0.93	1.36	0.10	0.22	0.91	0.92	1.71	2.23	0.07	0.13	0.45	0.54	1.35	2.86	0.10	0.12	0.21	0.23	0.39	0.92
LK2	4.66	9.26	20.66	26.66	60.88	155.55	0.01	0.10	0.42	0.54	1.33	3.12	0.04	0.12	0.76	0.91	1.98	5.55	0.10	0.12	0.41	0.62	2.29	4.95	0.09	0.14	0.21	0.22	0.32	0.77
PDA	0.01	0.78	4.12	4.36	8.12	11.58	0.07	0.15	0.56	0.67	1.42	2.08	0.04	0.16	0.53	0.59	1.22	2.13	0.28	0.34	0.48	0.50	0.76	1.31	0.08	0.08	0.15	0.16	0.31	0.46
PDB	0.01	0.47	2.88	3.22	7.45	9.92	0.04	0.20	0.73	0.87	2.09	7.39	0.01	0.08	0.57	0.61	1.25	1.73	0.27	0.32	0.44	0.50	0.85	1.73	0.06	0.09	0.15	0.16	0.24	0.40
PDC	0.38	0.75	4.76	4.96	11.23	12.96	0.09	0.14	0.55	0.69	1.81	2.23	0.04	0.08	0.54	0.58	1.18	1.69	0.21	0.31	0.41	0.45	0.81	1.00	0.00	0.02	0.16	0.15	0.24	0.32
PDD	0.05	0.12	2.06	2.53	5.36	10.10	0.02	0.07	0.55	0.81	2.33	3.36	0.05	0.08	0.55	0.62	1.33	1.71	0.21	0.28	0.42	0.43	0.68	1.05	0.06	0.08	0.15	0.17	0.28	0.42
PDE	0.12	0.65	3.83	4.33	8.93	10.91	0.02	0.07	0.49	0.65	1.81	3.22	0.01	0.07	0.47	0.52	1.18	1.30	0.26	0.27	0.41	0.42	0.55	0.78	0.07	0.08	0.15	0.16	0.25	0.51
PDF	0.13	0.36	3.52	3.88	7.66	21.28	0.09	0.11	0.53	0.62	1.48	1.60	0.01	0.10	0.41	0.45	1.04	1.24	0.26	0.31	0.44	0.45	0.63	0.83	0.06	0.08	0.15	0.15	0.25	0.41
PDG	0.07	0.19	3.07	3.21	6.22	7.35	0.04	0.05	0.49	0.65	1.89	2.65	0.02	0.07	0.50	0.52	1.18	1.23	0.28	0.32	0.43	0.45	0.69	1.04	0.07	0.08	0.15	0.17	0.32	0.47
PDH	0.05	0.52	2.88	3.56	8.27	11.09	0.03	0.07	0.46	0.60	1.81	2.22	0.01	0.10	0.48	0.54	1.24	2.02	0.21	0.29	0.42	0.42	0.58	0.71	0.06	0.08	0.14	0.15	0.26	0.32

INDICA

Site	Root						Stem						Leaf						Husk						Rice					
	Min	5%	Median	Mean	95%	Max	Min	5%	Median	Mean	95%	Max	Min	5%	Median	Mean	95%	Max	Min	5%	Median	Mean	95%	Max	Min	5%	Median	Mean	95%	Max
CH6	6.99	10.77	23.77	26.88	51.88	119.76	0.01	0.10	0.41	0.50	1.24	1.58	0.07	0.18	0.82	1.02	1.84	6.55	0.08	0.13	0.33	0.49	1.60	4.90	0.11	0.15	0.24	0.26	0.44	0.71
CH7	4.12	5.76	16.47	19.57	47.21	76.78	0.01	0.06	0.35	0.40	0.97	1.65	0.04	0.12	0.57	0.77	1.51	6.87	0.11	0.13	0.26	0.36	0.74	2.42	0.15	0.17	0.27	0.28	0.41	0.44
CH8	0.39	2.20	9.37	10.68	23.75	29.97	0.08	0.11	0.48	0.63	1.45	5.93	0.02	0.09	0.56	0.85	3.63	4.93	0.08	0.13	0.35	0.39	0.81	1.07	0.13	0.15	0.23	0.26	0.51	0.80
HC3	3.36	4.51	10.77	12.12	21.22	27.83	0.10	0.20	1.58	1.73	3.60	6.01	0.04	0.06	0.44	0.47	1.01	1.36	0.31	0.32	0.43	0.46	0.64	1.17	0.09	0.10	0.16	0.16	0.27	0.32
HC4	6.48	10.49	18.84	18.83	28.85	30.21	0.23	0.34	1.65	1.97	4.58	5.05	0.02	0.07	0.50	0.53	0.98	2.02	0.30	0.30	0.47	0.59	1.73	2.80	0.07	0.08	0.15	0.15	0.25	0.27
HC5	1.58	2.97	13.17	13.90	26.13	38.79	0.09	0.31	1.32	1.75	5.13	6.46	0.02	0.18	0.47	0.52	1.12	1.43	0.26	0.31	0.53	0.55	0.91	1.22	0.07	0.08	0.15	0.15	0.22	0.32
HM3	0.88	6.91	16.31	19.48	49.79	96.56	0.02	0.14	0.42	0.46	0.99	1.46	0.10	0.26	1.11	1.21	2.57	3.28	0.12	0.15	0.32	0.47	1.02	4.35	0.09	0.13	0.24	0.25	0.39	0.81
HM4	7.91	11.27	22.65	28.73	73.19	113.60	0.01	0.03	0.49	0.53	1.34	1.99	0.06	0.13	0.60	0.72	1.45	3.64	0.09	0.14	0.31	0.40	0.73	3.29	0.13	0.15	0.25	0.27	0.45	0.70
HM5	6.95	9.90	26.83	31.82	69.75	171.41	0.01	0.01	0.35	0.34	0.64	0.91	0.03	0.12	0.65	1.06	3.55	9.62	0.07	0.12	0.30	0.35	0.79	1.59	0.11	0.15	0.23	0.25	0.43	0.87
LK1	8.73	11.61	29.26	36.87	90.04	114.38	0.04	0.10	0.46	0.62	1.58	6.41	0.01	0.10	0.79	0.94	2.34	4.49	0.09	0.14	0.40	0.54	1.16	3.69	0.09	0.14	0.23	0.25	0.37	0.95
LK2	5.93	10.60	25.21	31.78	71.60	104.91	0.06	0.12	0.50	0.61	1.32	2.62	0.01	0.07	0.81	0.93	2.37	5.00	0.13	0.14	0.33	0.40	0.69	2.03	0.13	0.15	0.24	0.24	0.32	0.39
PDA	0.04	0.53	3.41	3.99	7.34	25.80	0.02	0.10	0.57	0.60	1.19	1.35	0.01	0.18	0.70	0.69	1.29	1.60	0.34	0.38	0.48	0.51	0.72	0.78	0.00	0.00	0.17	0.16	0.25	0.40
PDB	0.04	0.24	2.39	2.75	6.34	7.28	0.04	0.13	0.57	0.61	1.31	1.57	0.03	0.07	0.45	0.52	1.04	1.34	0.30	0.34	0.46	0.48	0.66	0.90	0.08	0.10	0.18	0.19	0.28	0.42
PDC	0.06	0.10	2.78	3.40	8.10	11.00	0.04	0.05	0.29	0.39	1.10	1.13	0.08	0.08	0.49	0.48	1.17	1.17	0.32	0.33	0.47	0.49	0.74	0.99	0.00	0.06	0.17	0.16	0.23	0.27
PDD	0.30	0.32	3.23	3.30	6.71	7.35	0.03	0.10	0.57	0.60	1.22	1.34	0.07	0.09	0.38	0.59	2.28	5.21	0.32	0.33	0.45	0.46	0.63	0.68	0.00	0.00	0.16	0.14	0.23	0.24
PDE	0.12	0.23	2.92	2.97	6.16	7.57	0.03	0.03	0.59	0.63	1.52	2.00	0.02	0.06	0.43	0.45	0.92	1.09	0.34	0.37	0.45	0.48	0.71	0.76	0.09	0.09	0.17	0.17	0.25	0.27
PDF	0.12	0.24	2.73	2.98	7.00	7.06	0.07	0.08	0.35	0.41	1.13	1.36	0.05	0.07	0.26	0.41	1.38	2.06	0.38	0.38	0.48	0.49	0.66	0.68	0.00	0.06	0.18	0.16	0.22	0.24
PDG	0.01	0.16	2.31	2.71	6.06	8.18	0.06	0.07	0.43	0.45	0.93	0.95	0.01	0.05	0.51	0.48	0.86	0.87	0.33	0.33	0.47	0.47	0.58	0.63	0.08	0.09	0.17	0.18	0.28	0.53
PDH	0.25	0.35	2.34	2.45	5.27	6.76	0.03	0.06	0.33	0.41	0.97	1.02	0.03	0.05	0.38	0.43	0.84	1.36	0.26	0.35	0.47	0.46	0.61	0.65	0.06	0.09	0.17	0.16	0.22	0.26

Table A4.6 Copper

JAPONICA

Site	Roots						Stem						Leaf						Husk						Rice					
	Min	5%	Median	Mean	95%	Max	Min	5%	Median	Mean	95%	Max	Min	5%	Median	Mean	95%	Max	Min	5%	Median	Mean	95%	Max	Min	5%	Median	Mean	95%	Max
CH6	11.60	18.44	29.21	37.09	86.25	199.36	3.05	4.82	12.56	12.38	21.77	24.70	3.14	3.81	7.87	8.15	14.08	26.37	2.76	3.29	5.46	5.52	7.65	12.82	3.62	3.83	4.63	4.84	6.34	7.77
CH7	14.39	21.18	49.54	66.89	165.49	352.44	5.07	7.01	14.13	14.34	23.89	28.44	4.55	6.73	9.14	9.39	13.42	15.68	4.22	4.51	6.13	6.21	8.58	10.86	2.83	3.88	5.74	5.83	8.32	9.68
CH8	17.31	21.52	36.99	42.68	97.69	134.44	4.57	9.44	14.37	14.96	21.01	42.85	3.97	6.78	13.31	13.67	23.43	32.05	3.79	4.47	6.27	6.63	9.60	10.40	3.81	4.56	6.19	6.20	8.90	10.54
HC3	9.63	10.38	17.80	18.29	27.42	39.74	6.49	7.95	15.81	16.13	26.03	36.59	2.48	3.14	4.21	4.38	6.14	7.78	1.70	2.04	3.45	3.51	4.77	5.94	3.31	3.54	4.23	4.29	5.14	5.37
HC4	15.04	16.57	32.00	32.19	50.79	53.12	14.08	15.84	27.95	29.52	48.55	53.79	3.56	3.74	6.37	6.76	10.90	13.83	2.86	3.33	4.45	4.60	6.15	8.86	3.42	3.63	4.90	4.98	6.53	10.81
HCS	6.28	11.14	32.75	32.36	59.25	72.40	5.40	10.26	31.45	32.71	57.79	68.78	1.93	2.79	6.97	7.53	15.06	18.37	2.00	2.62	4.04	4.39	6.90	8.90	2.51	2.68	4.95	5.06	8.02	9.17
HM3	7.80	10.42	20.63	27.06	67.52	167.76	2.35	3.34	7.88	8.33	14.57	20.44	5.72	6.46	9.99	12.40	26.03	32.02	4.19	4.76	6.22	6.97	11.48	15.08	3.24	3.42	4.39	4.59	6.30	10.83
HM4	10.45	12.51	22.67	28.39	63.90	127.38	1.24	2.54	6.69	7.45	15.90	19.76	2.39	3.49	6.70	7.68	14.90	19.33	2.97	3.86	4.86	5.16	7.73	9.79	0.63	3.28	4.16	4.44	7.43	10.83
HMS	11.57	14.43	25.22	29.25	56.18	140.61	2.10	3.56	8.09	8.70	16.83	27.84	3.60	5.70	15.34	16.14	32.92	45.32	3.57	4.34	7.79	7.73	11.93	13.87	2.74	3.73	4.56	4.72	6.40	7.73
LK1	17.69	25.53	81.89	94.35	209.86	281.27	5.74	9.45	17.71	19.85	34.77	167.65	5.30	10.91	22.44	29.61	69.29	306.08	5.50	7.60	10.61	11.73	20.68	31.42	3.87	4.59	6.46	6.70	9.56	10.90
LK2	13.96	20.68	34.31	45.52	98.55	205.40	2.59	3.16	6.64	8.12	16.39	26.72	4.10	5.38	10.91	11.85	21.91	29.63	3.03	3.99	5.63	6.16	9.59	11.92	3.07	3.57	4.61	4.87	7.04	9.15
PDA	6.09	9.38	18.20	17.83	26.62	29.14	5.22	11.50	17.52	17.99	25.61	31.15	2.42	4.45	6.17	6.38	8.76	10.24	2.30	2.74	4.47	4.72	7.78	10.38	3.46	3.86	5.07	5.19	6.79	7.47
PDB	0.06	2.45	9.80	10.17	17.82	21.27	1.58	3.19	10.45	10.81	18.23	23.45	1.14	2.48	3.94	3.99	5.57	7.02	1.21	1.92	3.33	3.37	4.89	6.59	0.77	2.02	4.09	3.97	5.29	5.94
PDC	8.83	9.00	12.79	13.74	20.73	24.90	4.60	5.27	11.82	12.38	19.52	29.54	3.26	3.85	4.70	4.78	6.11	6.54	2.44	2.52	4.06	4.15	5.71	9.49	0.08	1.15	4.63	4.70	6.74	7.18
PDD	2.59	3.76	9.44	9.27	14.43	16.43	5.43	6.62	10.43	10.70	15.83	21.57	2.42	2.66	3.82	3.87	5.55	6.11	1.74	2.00	3.14	3.31	4.98	6.17	2.77	3.35	4.10	4.20	5.12	5.59
PDE	4.52	6.33	9.46	10.20	15.54	21.75	3.94	4.44	7.79	8.54	16.00	17.97	2.25	2.73	3.68	3.85	5.05	6.78	2.00	2.28	3.38	3.55	5.09	5.58	3.16	3.60	4.33	4.49	5.76	6.36
PDF	2.47	3.78	8.45	8.70	13.73	15.52	4.90	6.43	9.91	10.50	16.18	27.16	2.02	2.62	3.63	3.79	5.32	6.24	2.20	2.23	3.53	3.50	5.40	6.09	3.10	3.46	3.99	4.16	5.51	6.17
PDG	3.03	3.97	7.36	8.73	12.20	14.54	3.00	4.87	7.74	8.73	16.40	25.11	2.61	2.81	3.44	3.47	4.21	4.95	2.06	2.31	3.35	3.33	4.61	4.91	3.03	3.56	4.07	4.19	5.52	6.02
PDH	1.79	2.98	8.17	8.32	17.16	20.89	3.00	4.00	7.93	8.22	13.63	21.00	1.86	2.21	3.24	3.32	4.56	6.38	1.64	2.03	3.19	3.33	4.70	13.00	2.80	3.08	3.91	4.08	5.71	6.03

INDICA

Site	Root						Stem						Leaf						Husk						Rice					
	Min	5%	Median	Mean	95%	Max	Min	5%	Median	Mean	95%	Max	Min	5%	Median	Mean	95%	Max	Min	5%	Median	Mean	95%	Max	Min	5%	Median	Mean	95%	Max
CH6	18.63	19.84	32.77	43.53	92.90	260.65	9.43	10.81	22.62	22.83	33.52	44.15	4.77	5.51	10.09	10.55	16.73	20.76	3.32	3.51	5.49	5.75	8.34	13.93	2.60	4.45	6.07	6.10	8.06	10.62
CH7	20.24	21.46	60.36	77.16	159.97	383.44	7.76	8.91	20.13	21.59	42.40	52.14	5.74	7.69	11.07	12.79	21.39	36.36	4.22	4.49	5.96	6.09	8.22	10.36	2.67	4.60	6.75	6.98	10.37	11.21
CH8	16.92	24.26	44.01	51.29	104.04	123.78	12.35	16.20	26.98	27.10	37.40	42.23	5.51	7.23	13.58	16.20	30.27	36.55	3.94	4.80	6.55	6.84	9.72	11.99	3.40	4.82	7.75	7.60	10.11	10.72
HC3	14.04	16.41	26.41	25.67	33.84	34.73	11.82	13.09	23.48	23.34	34.78	38.01	3.78	4.48	5.76	5.86	7.68	8.50	3.01	3.27	4.38	4.40	5.87	6.25	4.19	4.27	5.01	5.28	7.04	7.42
HC4	19.41	19.88	44.50	41.36	62.30	66.17	13.81	17.86	44.75	46.84	79.98	81.44	4.16	5.47	9.06	9.81	16.41	16.70	4.26	4.42	5.64	5.75	7.81	8.00	4.39	4.72	6.43	6.82	9.22	9.49
HCS	9.35	13.57	42.22	39.16	60.08	70.21	12.84	16.31	53.80	49.35	87.22	93.41	3.39	3.84	10.60	10.51	17.77	28.42	2.65	3.16	5.23	5.29	8.36	8.96	3.05	3.56	6.52	6.35	9.56	10.34
HM3	8.18	11.03	19.20	21.80	42.01	62.88	2.85	6.44	11.52	12.08	20.43	26.59	6.02	7.86	13.26	14.00	23.10	30.32	3.69	4.42	6.44	6.68	10.10	11.31	2.59	4.06	5.09	5.42	7.80	8.44
HM4	13.77	14.48	27.20	32.06	57.26	174.90	2.40	5.99	13.86	15.60	29.53	33.56	2.90	4.35	10.61	10.68	18.59	27.31	3.09	3.61	5.41	5.44	7.64	10.29	3.35	3.81	5.01	5.78	9.85	11.69
HMS	18.31	18.92	29.57	33.00	69.58	105.54	6.24	9.44	17.61	18.23	29.89	32.02	5.33	8.30	15.39	17.23	30.07	80.24	4.49	4.75	6.96	7.11	9.80	10.04	3.44	4.35	5.43	5.75	8.49	8.98
LK1	23.01	41.75	88.56	106.06	226.68	432.40	9.66	17.73	29.82	30.69	48.27	55.33	9.01	15.30	29.41	35.20	82.82	110.17	8.21	8.48	12.53	14.51	23.64	74.93	5.23	6.74	9.78	9.90	12.50	19.97
LK2	17.27	19.35	43.00	52.57	130.16	277.09	5.64	6.67	13.43	14.30	25.75	36.56	5.30	6.30	11.62	13.63	25.28	40.24	3.86	4.05	5.20	5.81	8.86	11.24	3.40	4.08	5.21	5.88	9.44	9.83
PDA	7.15	9.13	19.82	19.92	29.77	47.00	6.08	17.34	32.04	31.55	46.96	67.67	4.10	4.96	8.82	8.64	12.57	13.49	2.75	2.94	4.93	6.51	11.42	22.57	0.12	0.17	6.03	5.40	8.01	8.48
PDB	3.11	5.21	13.30	13.22	21.18	21.80	6.12	8.11	18.78	18.84	31.13	35.91	2.80	3.31	5.30	5.45	7.81	8.69	2.12	2.34	3.66	4.60	7.82	9.31	2.77	3.33	5.03	5.02	6.68	11.32
PDC	11.51	11.71	16.95	17.86	27.57	28.25	8.94	12.91	26.66	26.78	45.05	46.31	4.51	4.61	6.78	6.80	8.38	8.52	3.26	3.31	4.88	4.89	6.69	7.35	0.18	3.15	5.22	5.54	8.03	8.40
PDD	3.04	3.24	12.16	11.62	18.84	22.48	6.79	7.22	18.02	19.20	36.43	39.50	3.30	3.35	5.03	5.45	8.37	8.93	2.32	2.41	3.96	4.28	6.28	7.21	0.12	0.12	4.45	3.98	6.22	6.39
PDE	10.68	10.84	15.63	16.42	25.89	26.94	8.24	8.47	25.74	24.11	42.49	43.74	3.12	3.35	4.95	5.28	7.57	7.97	2.00	2.65	4.21	4.18	5.31	5.89	3.93	4.24	4.94	5.04	6.13	6.30
PDF	2.83	3.48	10.03	9.93	17.83	20.08	8.35	8.85	17.53	17.45	26.56	39.71	2.70	2.89	5.15	5.04	6.85	7.82	2.02	2.12	3.20	3.76	6.10	6.52	0.11	2.83	4.72	4.55	6.01	6.53
PDG	7.38	7.45	11.09	12.13	19.37	21.92	9.48	10.19	19.32	20.05	30.81	31.45	3.92	4.13	4.94	5.11	6.62	7.17	2.67	2.67	3.95	4.12	6.05	6.85	3.82	4.03	5.11	5.08	6.21	6.51
PDH	3.85	5.86	11.57	11.80	21.00	26.82	8.71	9.72	17.09	18.97	36.33	41.21	2.81	3.09	4.34	4.62	6.71	7.64	2.15	2.22	4.04	4.02	6.18	7.61	2.87	3.18	4.82	4.90	6.74	6.87



## Appendix 5 Theory behind the derivation of P50, P90 and P95 confidence levels of soil quality standards

In figure A5.1 the concept of the derivation of the P50, P90 and P95 Soil Quality Standard is shown.




Figure A5.1 Conceptual representation of the meaning of the P50, P90 and P95 SQS

The relation between Cd in soil or soil solution ( $\text{CaCl}_2$ ) is represented by a solid line showing an increase in the Cd level in rice with increasing soil or soil solution Cd levels. As such this line represents the predicted Cd content in rice at a given pH and CEC in case of the soil model or at a given Zn content in case of the  $\text{CaCl}_2$  model.

Based on the food quality standard (FQS) used (either  $0.2$  or  $0.4 \text{ mg kg}^{-1}$ ) the crossing of the model predictions and this FQS line represents the soil quality standard (P50 SQS). This means that at the pH and CEC given, the soil Cd content at the P50 SQS level will, on average result in a Cd level in rice equal to the FQS. Due to model uncertainty however, the actual Cd content in rice at the P50 SQS Cd level in soil will be higher (or lower) than the average model predicted lines. The upper limit of the predicted levels is represented by the 90 and 95% lines. This means that at a given soil Cd content, 95 out of 100 samples of rice will be below the level represented by the 95 percentile line. The quality of the model obviously determines how large the difference between the average model predictions and the 90 or 95% limits are. In case of a perfect model, the 95% line is equal to the model predictions (no error of prediction). Obviously this is never the case for models as discussed here.

The P50 SQS therefore implies that there is a chance of 50% that the real Cd level in rice still exceeds the FQS. On the other hand there is also a 50% chance that the real Cd level in rice will be below the FQS. To increase the security that the measured Cd level in rice is below the FQS with a confidence of 90 or 95% one has to decrease the SQS to the level where the 90% or 95% predictions lines cross the FQS line. Obviously, the P90 SQS and P95 SQS will be lower than the P50 SQS. Again, the

degree to which the P90 or P95 are below the P50 SQS depends on the model quality. In chapter 6 it was shown that for the data derived in this study P90 levels derived by the soil model or the CaCl<sub>2</sub> model are roughly half the value of the P50 SQS.