

# The development, evaluation and application of O<sub>3</sub> flux and flux-response models for additional agricultural crops.

Emberson, L.D.,<sup>1</sup> Massman, W.J.,<sup>2</sup> Büker, P.,<sup>1</sup> Soja, G.<sup>3</sup> and van de Sand, I.<sup>1</sup> Mills, G.,<sup>4</sup> Jacobs, C.<sup>5</sup>

<sup>1</sup> Stockholm Environment Institute, University of York, York YO10 5DD, UK

<sup>2</sup> USDA Forest Service, Rocky Mountain Research Station, Fort Collins, CO, USA

<sup>3</sup> Research Centre Seibersdorf, Dept. of Environmental Research, Seibersdorf, A-2444, Austria

<sup>4</sup> Centre for Ecology and Hydrology, Bangor, Gwynedd, LL57 2UP, UK

<sup>5</sup> Alterra, Droevendaalsesteeg 3, 6708 PB Wageningen, The Netherlands

\* correspondence to: [l.emberson@york.ac.uk](mailto:l.emberson@york.ac.uk)

## 1. Introduction

Currently, stomatal O<sub>3</sub> flux and flux-response models only exist for wheat and potato (LRTAP Convention, 2004), as such there is a need to extend these models to include additional crop types. The possibility of establishing robust stomatal flux models for five agricultural crops (tomato, grapevine, sugar beet, maize and sunflower) was investigated. These crops were selected on consideration of their distribution across Europe, sensitivity to ozone and economic value. The stomatal flux models would be based on the DO<sub>3</sub>SE<sup>1</sup> stomatal conductance ( $g_s$ ) multiplicative algorithm (MM  $g$ ) as described in the revised UNECE Mapping Manual, (LRTAP Convention, 2004) and hence require a number of different  $g_s$  parameters and  $g_s$  relationships with environmental variables to be identified. To establish the availability of parameterisation data, a comprehensive literature search was conducted for each species. In addition, authors of scientific papers that presented  $g_s$  data in their publications were contacted in an attempt to obtain the original datasets for inclusion in parameter setting boundary line analysis. On the basis of this work, it was deemed possible to develop MM  $g$  models for three of the five crops selected for investigation, namely grapevine, sunflower and tomato. For the other species the current data availability was considered too limited for the definition of robust models.

The DO<sub>3</sub>SE model has been developed for application within the EMEP photo-oxidant model (Simpson et al, 2003) and is able to estimate O<sub>3</sub> dry deposition to both stomatal and non-stomatal components of vegetated surfaces. The stomatal component of this DO<sub>3</sub>SE model is that upon which the MM  $g_s$  models for wheat and potato are based. Canopy stomatal deposition is an important driver of total deposition, especially during the period of greatest physiological activity of the surface vegetation. As such, comparisons of modelled with observed total O<sub>3</sub> flux/deposition to homogenous vegetation-covered surfaces provides a useful tool to both evaluate the DO<sub>3</sub>SE model's predictive capacity, but also to infer the role of the stomatal component of deposition and assess the importance of parameterisation of this model component. Environmental, stomatal and O<sub>3</sub> flux data collected during a campaign conducted in California (referred to here as the CODE91 campaign) are used to evaluate the DO<sub>3</sub>SE model for grapevine, and specifically the revised multiplicative  $g_s$  model parameterisation presented here. This field campaign recorded observations of total O<sub>3</sub> deposition (e.g. Massman & Grantz, 1995), enabling us to compare estimates of O<sub>3</sub> deposition with observed values.

However, the ability to faithfully simulate total deposition and O<sub>3</sub> stomatal flux is not an end in itself; to understand the impacts of absorbed O<sub>3</sub> dose requires some means of translating O<sub>3</sub> dose into effects. For wheat and potato, flux-response relationships have been established by relating absorbed O<sub>3</sub> doses estimated using the MM  $g_s$  model to observed effects (yield losses in grain and tuber for wheat and potato respectively). In a similar manner, the  $g$  models established for the three crops presented here could, in theory, also be used to derive flux-response relationships for these crops were suitable datasets identified (i.e. that provide hourly records of key environmental data and O<sub>3</sub> fumigation regimes). The possible establishment of such a relationship is discussed here in relation to an open top chamber O<sub>3</sub> fumigation study conducted by Soja et al. (2004) which provides three years worth of environmental and O<sub>3</sub> concentration

data. Re-analysis of this dataset using the updated DO<sub>3</sub>SE g<sub>s</sub> multiplicative model parameterisation could provide the opportunity to establish a flux-response model for grapevine.

## 2. Development of stomatal flux models.

The stomatal conductance model described in the UNECE Mapping Manual (LRTAP Convention, 2004) (MM g<sub>s</sub>) is shown in Eq. 1.

$$g_s = g_{\max} * [\min(f_{\text{phen}}, f_{\text{O}_3})] * f_{\text{light}} * \max\{f_{\text{min}}, (f_{\text{temp}} * f_{\text{VPD}} * f_{\text{SWP}})\} \quad [1]$$

where g<sub>s</sub> is the actual stomatal conductance (mmol O<sub>3</sub> m<sup>-2</sup> sunlit projected leaf area (PLA) s<sup>-1</sup>) and g<sub>max</sub> is the species-specific maximum stomatal conductance (mmol O<sub>3</sub> m<sup>-2</sup> PLA s<sup>-1</sup>). The parameters f<sub>phen</sub>, f<sub>O<sub>3</sub></sub>, f<sub>light</sub>, f<sub>temp</sub>, f<sub>VPD</sub> and f<sub>SWP</sub> are all expressed in relative terms (i.e. they take values between 0 and 1) as a proportion of g<sub>max</sub>. These parameters allow for the modifying influence of phenology and ozone, and four environmental variables (irradiance, temperature, water vapour pressure deficit and soil water potential) on g to be estimated. The f<sub>O<sub>3</sub></sub> function is not considered further here since its parameterisation would require an extensive g<sub>s</sub> dataset collated under O<sub>3</sub> fumigation conditions. Results of the literature search to identify data necessary for parameterisation of the MM g<sub>s</sub> model for the five agricultural crop species (tomato, grapevine, sugar beet, maize and sunflower) are shown in Table 1.

**Table 1.** Parameterisation of the multiplicative g<sub>s</sub> model for five agricultural crop species based on a comprehensive literature search. The numbers in brackets refer to the published papers upon which the parameterisations are based. The grey shading indicates parameterisation founded on data with large variability.

Parameter	Units	Grapevine ( <i>Vitis vinifera</i> )	Tomato ( <i>Lycopersicum esculentum</i> )	Sunflower ( <i>Helianthus annuus</i> )	Maize ( <i>Zea mays</i> )	Sugar beet ( <i>Beta vulgaris</i> )
g <sub>max</sub>	mmol O <sub>3</sub> m <sup>-2</sup> PLA s <sup>-1</sup>	215 (16)	285 (8)	370 (15)	320 (3)	270 (10)
f <sub>min</sub>	(fraction)	0.01 (1)	0.01 (1)	0.05 (1)	0.06 (2)	0.05
f <sub>phen_a</sub>	(fraction)	0.2 (2)	-	0.62 (4)	-	-
f <sub>phen_b</sub>	(fraction)	0.5 (2)	-	0.41 (4)	-	-
f <sub>phen_c</sub>	days	60 (2)	-	34 (4)	-	-
f <sub>phen_d</sub>	days	45 (2)	-	34 (4)	-	-
f <sub>phen_e</sub>	°C days	-	-	-	-	-
f <sub>phen_f</sub>	°C days	-	-	-	-	-
light <sub>a</sub>	(constant)	0.0076 (5)	0.0175 (4)	0.002 (2)	0.0035	0.0025 (3)
T <sub>min</sub>	°C	9 (6)	0 (2)	-	2 (2)	-
T <sub>opt</sub>	°C	30 (6)	21 (2)	-	25 (2)	-
T <sub>max</sub>	°C	43 (6)	35 (2)	-	48 (2)	-
VPD <sub>max</sub>	kPa	1.6 (4)	1 (3)	1.2 (5)	-	-
VPD <sub>min</sub>	kPa	6.2 (4)	2.7 (3)	4.0 (5)	-	-
SVPD <sub>crit</sub>	kPa	-	-	-	-	-
SWP <sub>max</sub>	MPa	-1.2 (5)	-0.3 (3)	-0.25 (8)	-0.12	-
SWP <sub>min</sub>	MPa	-0.35 (5)	-1.0 (3)	-1.65 (8)	-0.8	-

On the basis of this work it has been possible to develop reasonably robust flux models for grapevine, sunflower and tomato, however, it should be noted that it has not been possible to find an f<sub>temp</sub> relationship for sunflower (a suitable surrogate would need to be identified to apply the sunflower g<sub>s</sub> model) and it has not been possible to establish an f<sub>phen</sub> relationship for either tomato or sunflower. Maize and sugar beet are missing key parameters due to a lack of suitable data describing g<sub>s</sub> relationships with phenology and important environmental variables. Table 1 also shows that it is not possible to parameterise the MM g<sub>s</sub> model for any of the crop species for thermal time determined phenology (indicated by f<sub>phen\_e</sub> and f<sub>phen\_f</sub>)

since the data needed to parameterise such relationships are unavailable in the literature; only analysis of datasets that have recorded  $g_s$  over entire growing seasons in conjunction with associated mean daily temperatures would offer the possibility of deriving these coefficients. Similarly, it has not been possible from the literature to identify the SVPD since derivation of this parameter requires sufficient diurnal  $g_s$  measurements and associated VPD values. As such, these parameters are only presented in grey font in Table 1. For tomato, it should also be noted that there was some inconsistency in the data that have been used to derive the  $f_{\text{light}}$  and  $f_{\text{VPD}}$  relationships.


Previous evaluations of the multiplicative  $g_s$  models have found the identification of an appropriate value for  $g_{\text{max}}$  to be crucial in determining the predictive abilities of the model. In view of this we present the data describing  $g_{\text{max}}$  for grapevine, sunflower and tomato in Figure 1 to indicate how robust these values are.  $g_{\text{max}}$  is assumed the median  $g_{\text{max}}$  value of all observations. In general, the lack of information describing  $g_{\text{max}}$  for tomato (based only on 8 studies) is not related to problems in identifying maximum  $g_s$  in papers but in the lack of clarification of two essential pieces of information: i) the gas for which the conductance data were recorded and ii) the leaf surface area (projected or total) on which the measurements were based. Without this information it is not possible to use the  $g_s$  data to identify  $g_{\text{max}}$  since values could be mis-represented by as much as 100%. For sunflower, it is evident that there is rather large uncertainty in the derivation of  $g_{\text{max}}$  with values taken from the literature ranging from 150 to over 1000  $\text{mmol O}_3 \text{ m}^{-2} \text{ PLA s}^{-1}$ . The use of the median value in effect assumes the extremely high values to be outliers.

For grapevine, further details of the  $g_s$  model parameterisation are provided since a complete  $g_s$  dataset was made available (described in Jacobs et al. 1996) which allowed more robust boundary line relationships for this species to be defined. As such, the parameterisation of the flux model for grapevine is considered the most robust of all species investigated since the use of both published data and  $g_s$  measurement datasets provides more certainty in the fitting of the boundary lines. Figure 2 shows the data and the DO<sub>3</sub>SE model parameterisation for the  $f$  function relationships with irradiance (PPFD,  $\mu\text{mol m}^{-2} \text{ s}^{-1}$ ), air temperature ( $^{\circ}\text{C}$ ) and air vapour pressure deficit (VPD, kPa) and soil water potential (inferred from measurements of pre-dawn leaf water). The main change to the grapevine parameterisation shown here compared to that described in Emberson et al. (2000) is in the  $f_{\text{VPD}}$  relationship which is now far less sensitive. This likely reflects the more southerly distribution of grapevines and their acclimation to high atmospheric water deficits. These data have also been used in a comparison of multiplicative and photosynthesis based  $g_s$  models by B ker et al. (this volume) to identify the most appropriate  $g_s$  algorithm for use in DO<sub>3</sub>SE.


### 3. Evaluation of grapevine DO<sub>3</sub>SE model

The DO<sub>3</sub>SE model has been evaluated against observations of total O<sub>3</sub> flux and  $g_s$  for a number of different ecosystem types representative of central and northern European locations (e.g. Tuovinen et al., 2001). These have generally shown that the model performs well, although improvements in the model predications could be achieved through “tuning” the model parameterisation for local conditions. However, the module has not been extensively tested under Mediterranean conditions with only one comparative study (Tuovinen et al., 2004) having been conducted to date for wheat growing in Italy. It is imperative that further evaluations should be performed, firstly since the model should be tested under all European climate types and secondly, since these regions are prone to co-occurring elevated ozone concentrations and high soil and atmospheric water deficits. As such, these conditions may lead to large differences in the accumulation of exposure versus flux-based indices that may be especially relevant to European emission abatement formulation.

**Figure 1.** Data used to establish  $g_{max}$  for three agricultural crop species. The mean and standard deviation by species are grapevine (mean=229, S.D.=50.98); tomato (mean=284, S.D.=73.83); sunflower (mean=436, S.D.=229.07).




**Figure 2.** The grapevine  $DO_3SE$   $f$  function relationships for irradiance (PPFD,  $\mu mol m^{-2} s^{-1}$ ), air temperature ( $^{\circ}C$ ) and air vapour pressure deficit (VPD, kPa) shown in relation to the data, collated from the literature and  $g_s$  datasets, used in their establishment.



A Californian vineyard dataset (Massman & Grantz, 1995) collected as part of the California Ozone Deposition Experiment (CODE) during July and August of 1991, provides an opportunity to i) infer an evaluation of the DO<sub>3</sub>SE grapevine  $g_s$  model parameterisation through a comparison of observed and modelled total O<sub>3</sub> deposition and ii) extend the number of DO<sub>3</sub>SE model evaluation studies conducted under “Mediterranean style” conditions. Measurements were taken at a grape vineyard site (*Vitis vinifera* L. cv. Thompson seedless) located in the San Joaquin Valley in California (36°51'36"N, 120°6'7"W). There was no precipitation during the study period, but the plants had been irrigated before the start of the experiment. The sky remained virtually cloud free for the duration of the investigation. There was almost no growth in the vineyard plants, since they had reached their maximum vegetative state (LAI = 3.4, vegetation height = 1.7m). Flux data for O<sub>3</sub>, heat, H<sub>2</sub>O, CO<sub>2</sub> and momentum were measured half hourly using eddy covariance; corresponding measurements of meteorological variables were also made. Further details of the site measurement and data descriptions for the CODE 91 experiment are given in Massman et al. (1994).

The DO<sub>3</sub>SE model as described in Emberson et al. (2000) was applied using observed reference height O<sub>3</sub> concentration and meteorological data, with the exception that the grapevine parameterisation described in Table 1 was used in place of that described in Emberson et al. (2000). Figure 3 shows a scatter plot and seasonal course of total ozone flux values modelled (using the “new” grapevine DO<sub>3</sub>SE model parameterisation) in comparison with the corresponding measured O<sub>3</sub> flux data available throughout the study period. It is clear from the R<sup>2</sup> values and seasonal profile that the DO<sub>3</sub>SE model is able to reproduce the seasonal diurnal profile but that the model consistently overestimates total deposition (the slope of the linear regression is approximately 0.6). The use of local parameterisation for  $g_{\max}$  (i.e. alteration of the value from 215 mmol O<sub>3</sub> m<sup>-2</sup> s<sup>-1</sup> to 165 mmol O<sub>3</sub> m<sup>-2</sup> s<sup>-1</sup> (i.e. within the range of  $g_{\max}$  extracted from the published literature) improves the prediction of the seasonal amplitude in total O<sub>3</sub> flux, although the highest modelled O<sub>3</sub> fluxes are still overestimated by approximately 20% (data not shown). The overestimation could be due to soil moisture deficit limiting actual  $g_s$ ; this could not be introduced into the modelling since the necessary integrated root depth SWP data were not available. It may also be that the canopy  $g_s$  is overestimated as all canopy leaves are assumed to be of the same age and hence have the same  $f_{\text{phen}}$  relationship. In reality older leaves with lower  $g_s$  may occur within the canopy, a situation that could be modelled by dividing the canopy into different leaf populations with specific  $f_{\text{phen}}$  functions (as in Tuovinen et al. 2004 for wheat).

**Figure 3.** Scatter plot and seasonal course showing observed versus modelled total ozone flux/deposition (nmol O<sub>3</sub> m<sup>-2</sup> PLA s<sup>-1</sup>) for grapevine data collected as part of the California Ozone Deposition Experiment (CODE) during July and August of 1991.



#### 4. Development of flux-response models

The establishment of robust flux models for grapevine, tomato and sunflower identify these species for targeted future development of flux-response models. However, the establishment of such models ordinarily requires the identification of suitable O<sub>3</sub> fumigation datasets. An appropriate dataset has been obtained for

grapevines, full details of this dataset are provided in Soja et al. (2004). In summary, the data describe experiments conducted on grapevines (*Vitis vinifera* L., cv. Welschriesling) that had been pre-cultivated for two years under field conditions in Eastern Austria, 30 km south of Vienna. The plants were transplanted into containers and moved to open top chambers (OTCs), ozone fumigation was started in 1994 and was continued during the growth periods until 1996. Four fumigation regimes were compared: charcoal filtered air, non-filtered air, non-filtered air + 25 nmol mol<sup>-1</sup>, non-filtered air + 50 nmol mol<sup>-1</sup>. Response parameters investigated were grape yield and sugar yield, the latter being defined on chemical analysis of grape juice for soluble carbohydrates. As such this dataset provides an excellent opportunity to develop flux based response relationships and re-analysis of this dataset with the revised grapevine DO<sub>3</sub>SE g<sub>s</sub> model described in this paper will be conducted. To date, no datasets that may be appropriate for the derivation of flux-response relationships for either tomato or sunflower have been identified.

## 5. Conclusions

This paper has described the development of stomatal O<sub>3</sub> flux models for additional crop species (grapevine, tomato and sunflower) to those for which flux, and flux-response models already exist (namely, wheat and potato as described in the UNECE Mapping Manual (LRTAP Convention, 2004). The paper has highlighted the importance of the stomatal component of deposition on application of the revised grapevine DO<sub>3</sub>SE g<sub>s</sub> model parameterisation in a study to compare modelled and measured total O<sub>3</sub> deposition values. This comparison suggests that the g<sub>max</sub> is an important driver of deposition, particularly when deposition is high during the middle of the day; as such, its parameterisation is crucial both to total deposition for O<sub>3</sub> mass balance modelling but also for stomatal deposition for effects modelling. In terms of effects modelling, the development of additional flux-response models should be a future priority. To this end, an O<sub>3</sub> fumigation dataset has been identified for grapevine that may be used with the revised grapevine DO<sub>3</sub>SE g<sub>s</sub> model parameterisation to establish flux-effect relationships for this species. The issue of identifying an appropriate g<sub>max</sub> for use in this re-analysis can be dealt with by using the g<sub>s</sub> dataset collected during the fumigation study.

## References

- Emberson L.D., Simpson D., Tuovinen J.-P., Ashmore M.R. and Cambridge H.M. 2000. Towards a model of ozone deposition and stomatal uptake over Europe. Norwegian Meteorological Institute, Oslo, EMEP MSC-W Note X/00, 57p.
- Jacobs C.M.J., van den Hurk B.J.J.M., de Bruin H.A.R. Stomatal behaviour and photosynthetic rate of unstressed grapevines in semi-arid conditions. *Agricultural and Forest Meteorology* 80: 111-134
- LRTAP Convention UNECE Revised manual on methodologies and criteria for mapping critical levels/loads and geographical areas where they are exceeded, Chapter 3: Mapping Critical Levels for Vegetation. 2004. Umweltbundesamt, Berlin, Germany. <http://www.oekodata.com/icpmapping/html/manual.html>, 53p.
- Massman, W.J., Pederson, J., Delany, A., Grantz, D., den Hartog, G., Oncley, S.P., Pearson, Jr. R., Shaw, R. 1994. An evaluation of the RADM surface module for ozone uptake at three sites in the San Joaquin Valley of California. *Journal of Geophysical Research* 99: 8281-8294
- Simpson D., Fagerli H., Jonson J.E., Tsyro S. and Wind P. 2003. Transboundary Acidification, Eutrophication and Ground Level Ozone in Europe. Part I Unified EMEP Model Description. Norwegian Meteorological Institute, Oslo, EMEP MSC-W Note 1/2003, [http://www.emep.int/publ/common\\_publications.html](http://www.emep.int/publ/common_publications.html), 104p.
- Soja G., Reichenauer T.G., Eid M., Soja A.-M., Schaber R. and Gangl H. Long-term ozone exposure and ozone uptake of grapevines in open-top chambers. *Atmospheric Environment* 38: 2313-2321
- Tuovinen J.-P., Simpson D., Mikkelsen T. N., Emberson L. D., Ashmore M. R., Aurela M., Cambridge H. M., Hovmand M. F., Jensen N. O., Laurila T., Pilegaard K. and Ro-Poulsen H. 2001. Comparisons of measured and modelled ozone deposition to forests in Northern Europe. *Water, Air and Soil Pollution: Focus* 1: 263-274
- Tuovinen J.P., Ashmore M.R., Emberson L.D., Simpson D. 2004. Testing and improving the EMEP ozone deposition module. *Atmospheric Environment* 38: 2373-2386.