

ORIGINAL PAPER

T. H. Park · V. G. A. A. Vleeshouwers · D. J. Huigen
E. A. G. van der Vossen · H. J. van Eck
R. G. F. Visser

Characterization and high-resolution mapping of a late blight resistance locus similar to *R2* in potato

Received: 7 January 2005 / Accepted: 15 April 2005 / Published online: 21 May 2005
© Springer-Verlag 2005

Abstract Identification of resistance (*R*) genes to *Phytophthora infestans* is an essential step in molecular breeding of potato. We identified three specific *R* genes segregating in a diploid mapping population. One of the *R* genes is located on chromosome 4 and proved phenotypically indistinguishable from the *Solanum demissum*-derived *R2*, although *S. demissum* is not directly involved in the pedigree of the population. By bulked segregant analysis combined with a resistance assay, a genetic linkage map of the *R2*-like locus was constructed with 30 coupling and 23 repulsion phase AFLP markers. Two markers flanking the *R2*-like locus were applied to screen an extended population of 1,586 offspring. About 103 recombinants were selected, and an accurate high-resolution map was constructed. The *R2*-like resistance was localized in a 0.4 cM interval and was found co-segregating with four AFLP markers, which can be used to isolate the *R2*-like gene by map-based gene cloning. By analyzing race-specificity and *R* gene-specific molecular markers, we also found that an *R1*-like gene and an additional unknown *R* gene are segregating in the population.

Introduction

Phytophthora infestans is the causal agent of late blight and one of the most important pathogens in cultivated potato (*Solanum tuberosum* L.). The economic value of the loss of global annual production and the cost of crop protection are estimated to be US \$5 billion annually (Duncan 1999). In early breeding programmes for late blight resistance, 11 resistance (*R*) genes originating from the hexaploid wild species *S. demissum* were introduced into *S. tuberosum* (Black et al. 1953; Malcolmson and Black 1966). These *R* genes confer race-specific resistance, and they are controlled by major single dominant factors (Mastenbroek 1953; Malcolmson and Black 1966). The potato-*P. infestans* interaction is following the gene-for-gene model as proposed by Flor (1971). Although the *R* genes do not give durable resistance, the isolation of these *R* genes is of importance to understand the mechanism of the specific plant defence. Several *R* genes have been localized, e.g. *R1* on chromosome 5 (Leonards-Schippers et al. 1992) and *R3*, *R6* and *R7* clustered on chromosome 11 (El-Kharbotly et al. 1994, 1996). *R1* and *R3* have been isolated (Ballvora et al. 2002; Huang 2005). The *R2* locus has been localized on chromosome 4 using a tetraploid mapping population EJ96-4061 comprising 86 offspring (Li et al. 1998). Eleven AFLP markers linked to the *R2* locus have been identified using bulked segregant analysis [(BSA) Michelmore et al. 1991]), and the *R2* linkage group has been assigned using the reference map of AM3778-16. Three AFLP markers that are tightly linked to *R2* are present in the diploid mapping population RHAM026, which is used in this study.

Recently many different haplotypes of *R3*-like genes were discovered at the *R3* locus in *S. demissum* and ample variation for functional *R* genes to late blight appears to exist in *Solanum* species (Huang et al. 2004, Huang 2005). In this research we identified resistances by assaying race specificities and focused on a locus that is indistinguishable from *R2* in the diploid mapping

Communicated by G. Wenzel

T. H. Park · V. G. A. A. Vleeshouwers · D. J. Huigen
H. J. van Eck · R. G. F. Visser (✉)
Department of Plant Sciences, Laboratory of Plant Breeding,
Graduate School Experimental Plant Sciences,
Wageningen University, Binnenhaven 5, 6709 PD,
Wageningen, The Netherlands
E-mail: Richard.Visser@wur.nl
Tel.: +31-317-482857
Fax: +31-317-483457

E. A. G. van der Vossen
Plant Research International BV, P.O. Box 16, 6700 AA,
Wageningen, The Netherlands

population RHAM026. Because the resistance was introgressed from different wild species, we designate the gene as *R2-like*. From the *R2-like* locus, a high-resolution map was constructed.

Materials and methods

Plant materials

The diploid mapping population RHAM026 was obtained from a cross between AM3778-16 (AM) and RH89-039-16 (RH). The dihaploid female parent AM is derived from the tetraploid breeding clone AM78-3778, which has various wild *Solanum* species in its pedigree, including *S. tuberosum* spp. *andigena*, *S. vernei*, *S. vernei* spp. *ballsii*, *S. oplocense* and *S. edinense*. The detailed pedigree of the mapping population is shown in Fig. 1.

The mapping population was generated from seeds that were sown in vitro after they were sterilized. When the seeds germinated and the seedlings reached approximately 3–5 cm, they were duplicated. One was used for maintenance and another for DNA isolation.

Inoculum preparation and resistance assay

Four different isolates were used to identify race-specific resistance (Table 1). For experiments, a plug of mycelium was transferred to a fresh plate with rye sucrose agar medium (Caten and Jinks 1968). One to 2 weeks later, ice-cold water was added to the mycelium that covered the agar plate. The sporangiospore suspension was pipetted into a tube and incubated at 4°C for 3 h.




Fig. 1 Pedigree of the diploid mapping population RHAM026 (RH) used in this study

After the release of zoospores, the concentration of inoculum was adjusted to 5×10^4 spores/ml.

A resistance assay was performed by a detached leaf assay. Fully expanded and healthy leaves were collected from greenhouse plants and inoculated on wet paper in humid trays. Droplets of 10 μ l inoculum were applied to the leaves, which were subsequently incubated at 15°C (Vleeshouwers et al. 1999). The symptoms on the leaves were evaluated 3 days and 5 days after inoculation.

DNA isolation

A high-throughput DNA isolation procedure was followed using a Retsch machine (Retsch, Haan, Germany) and 96-deep-well COSTER microtiter plates (Corning, Corning, N.Y., USA). Fresh leaf tissue was obtained by harvesting in vitro plants and grinding with two steel balls in the presence of nuclear lysis buffer (0.2 M Tris-HCl, 0.05 M EDTA, 2 M NaCl, 2% CTAB, with an end pH of 7.5), DNA extraction buffer (100 mM Tris-HCl, 5 mM EDTA, 0.35 M Sorbitol, 20 mM sodium bisulfite, with an end pH of 7.5) and 5% sarcosyl, followed by incubation at 65°C in a water bath for 1 h. Ice-cold chloroform isoamyl alcohol (24:1) was added, and the samples were mixed and centrifuged. The supernatant was transferred to new tubes and an equal volume of isopropanol was added. Another centrifugation step allowed the precipitation of DNA. The DNA pellet was dried and dissolved in $T_{0.1}E$ -buffer (+0.5 μ g RNase).

AFLP analysis

The AFLP analysis was essentially carried out as described by Vos et al. (1995). Primary template DNA was prepared using *EcoRI* and *MseI* and adaptors fitting to the restriction enzyme sites. Template DNA was diluted ten times prior to the selective pre-amplification with single-nucleotide extended primers, which decreases marker density. For the selective amplification, AFLP reactions with three nucleotides extended primers (*Eco*+3/*Mse*+3 primers) were performed to find markers linked to the resistance locus. The AFLP bands were separated on a 6% polyacrylamide gel in a Li-cor sequencer (Li-cor, Lincoln, Neb., USA).

The AFLP markers were named with the enzyme, the primer combination and the mobility of the fragment as described in reference autoradiograms created by Keygene NV, Wageningen, The Netherlands.

Table 1 Isolates of *Phytophthora infestans* and virulence factors

Isolate	Race specificity	Source
IPO-0	0	W. Flier, Plant Research International, The Netherlands
99018	1, 4	F. Govers, Wageningen University, The Netherlands
90128	1, 3, 4, 7, 8, 11	F. Govers, Wageningen University, The Netherlands
USA618	1, 2, 3, 6, 7, 11	W.E. Fry, Cornell University, USA





Fig. 3 Genetic linkage map of the *R2-like* locus. The resistance locus is underlined. Two flanking markers used for screening the extended population are dot-underlined. All 30 coupling phase markers, but only four repulsion phase markers and the resistance locus, are included in the map

tance locus. One is EATC/MCGA_204, which was mapped 1.3 cM telomeric from the resistance locus. Another is EATG/MCAG_215, which was mapped 2.5 cM centromeric from the resistance locus in the 78 offspring (Fig. 4). By screening the expanded population of 1,582 genotypes, 103 recombinants were selected resulting in a larger interval of a 6.4-cM genetic distance between these two AFLP markers. We screened the 103 recombinants with 14 AFLP markers located between EATC/MCGA_204 and EATG/MCAG_215 (Fig. 3) and tested them for their resistance to *P. infestans* isolate 90128. Subsequently a high-resolution genetic map of

Fig. 4 Schematic representation for recombinant screening towards high-resolution mapping. The AFLP gel images of the two flanking AFLP markers EATC/MCGA_204 and EATG/MCAG_215, which were used to select recombinants in 1,582 genotypes. One example of a recombinant is indicated by X



the 6.4-cM interval containing the resistance locus was constructed (Fig. 5), in which the AFLP markers are precisely separated and mapped. Closest markers on both sides of the resistance locus are EACA/MCAC_105 (three recombinants in 0.2 cM) and a group containing five markers (three recombinants in 0.2 cM). Four AFLP markers are still co-segregating with the resistance locus.

Comparative genetics

The *R2* and *R2-like* loci derived from two different mapping populations were compared. The bridged markers EATC/MCGA_186 and EACT/MCAC_189 are 0.2 cM apart in the *R2-like* diploid population (three recombinants in 1,582 genotypes) and 1.2 cM in the *R2* tetraploid population (one recombinant in 86 genotypes) (Li et al. 1998). One AFLP marker, EATC/MCGA_186, co-segregates with the *R2-like* locus, but is 1.2 cM separated from the *R2* locus. Another AFLP marker EACT/MCAC_189 is 0.2 cM separated from the *R2-like* locus, but co-segregates with the *R2* locus. The position of flanking markers that connect the two genetic linkage maps confirmed that the *R2* and *R2-like* loci are located at the same region of chromosome 4.

Discussion

In this research, we identified three resistance loci in the diploid segregating population RHAM026. One locus was localized on chromosome 4, where the *R2* locus from *S. demissum* was previously mapped using a tetraploid mapping population EJ96-4601 (Li et al. 1998). Accurate specificity studies with a panel of four different *P. infestans* isolates revealed that *R2-like* resistance of




Fig. 5 High-resolution map of the *R2-like* locus. The 1,582 genotypes were screened with two AFLP markers, EATC/MCGA_204 and EATG/MCAG_215, which are *dot-underlined*, and 103 recombinants were selected. Four markers are co-segregating with the *R2-like* locus

this population is phenotypically indistinguishable from *R2*.

The origin of the AFLP markers EACT/MCAC_189 and EATC/MCGA_186 was previously thought to be *S. demissum*-specific, because the AFLP markers were identified in the fingerprints of three accessions of *S. demissum* (Li et al. 1998). However, we observed that EACT/MCAC_189 is also present in *S. acaule* accessions and EATC/MCGA_186 in *S. bulbocastanum* accessions (data not shown) and therefore conclude that these markers are present in a broader range of *Solanum* germplasm. Although *S. demissum* is not directly present in the ancestors of the *R2-like* diploid population, the introgression fragment could be of wild species origin, e.g. *S. edinense*, which is a natural hybrid between *S. tuberosum* and *S. demissum* (Hawkes 1990). It remains unknown what the origin of the *R2-like* locus is. Even if it is derived from *S. demissum*, definite identity with *R2* (Li et al. 1998) can only be determined after cloning both genes.

To construct a marker saturation and high-resolution map at the *R2-like* locus, we used a diploid population. Compared to the tetraploid level, mapping studies at the diploid level can avoid the complexities of tetrasomic

inheritance and makes the study of potato genetics more feasible (Jacobs et al. 1995; Li et al. 1998). The number of markers is at least fourfold higher in a diploid population than in a tetraploid population, because marker alleles with linkage in repulsion can be used at the diploid level (Wu et al. 1992). We identified 69 AFLP markers with linkage to *R2-like*, using 256 primer combinations. A similar high efficiency was achieved in another study in which 29 AFLP markers with linkage to *R1*, using 108 primer combinations were identified at the diploid level (Meksem et al. 1995). At the tetraploid level, much lower efficiency was achieved. Li et al. (1998) detected only 11 AFLP loci with linkage to *R2*, using 205 primer combinations.

Many R genes and resistance gene homologues are clustered in the genome (Meyers et al. 1998; Michelmore and Meyers 1998). In the potato genome, for instance, at least five R genes against diverse pathogens are located at the gene cluster on chromosome 5 (Leister et al. 1996; Marano et al. 2002), i.e. *Gpa* conferring resistance to the potato cyst nematode *Globodera pallida* (Kreike et al. 1994), *Grp1* conferring resistance to the potato cyst nematode *Globodera rostochiensis* (Roupe van der Voort et al. 1998), *Nb* conferring resistance to potato virus X (de Jong et al. 1997), *Rx2* (Ritter et al. 1991) conferring resistance to potato virus X and *R1* conferring resistance to *P. infestans* (Leonards-Schippers et al. 1992). Also near the *R2-like* locus on chromosome 4, various known R genes are located. The *Hero* gene, which confers broad-spectrum resistance against *G. rostochiensis* in tomato, is located between CT229 and TG370 (Ganal et al. 1995), *Gpa4* conferring resistance to *G. pallida* (Bradshaw et al. 1998) and *R2* (Li et al. 1998), *Rpi-abpt* (Park et al. 2004) and *Rpi-blb3* (Park et al. 2004) conferring resistance to *P. infestans* are mapped at the same region on chromosome 4.

One of the most challenging methods for gene cloning in a plant species with a large genome is map-based cloning, which can be considered routine in model species such as *Arabidopsis* with a small genome (Lukowitz et al. 2000). Accurate mapping of the target gene and identification of recombinants in the population are essential steps for map-based cloning, and here we report a reliable high-resolution map of the *R2-like* locus within a 0.4-cM interval on chromosome 4 in the potato breeding line RHAM026. The initial research for tightly linked AFLP markers was facilitated using AFLP technology (Vos et al. 1995) in combination with BSA (Michelmore et al. 1991). The *R2-like* locus is flanked by one AFLP marker on the telomeric side, a group of five AFLP markers on the centromeric side and co-segregating with four AFLP markers (Fig. 5). The flanking and co-segregating markers will facilitate isolation of the *R2-like* gene by BAC walking. A 0.2-cM (4/2,109 recombinants) interval was covered by one BAC clone of 100 kb in the cloning of the potato late blight resistance gene *Rpi-blb1* (van der Vossen et al. 2003) and in the cloning of *R1*, a 0.2-cM (2/1049 recombinants) interval was estimated at 250–300 kb (Ballvora et al. 2002). We estimate that the 0.4-cM

interval spanning the *R2-like* gene could be covered by a single BAC clone or a few BAC-walking steps.

Currently a BAC library of diploid genotype AM is being constructed to establish a physical contig with the co-segregating markers and the flanking markers and isolation of *R2-like* gene in near future is quite feasible. After that, physical analysis of the gene cluster containing the *R2-like* gene might accelerate isolation of the homologous genes conferring resistance to *P. infestans* in different genetic backgrounds.

References

- Ballvora A, Ercolano MR, Weiß J, Meksem K, Bormann CA, Oberhagemann P, Salamini F, Gebhardt C (2002) The *R1* gene for potato resistance to late blight (*Phytophthora infestans*) belongs to the leucine zipper/NBS/LRR class of plant resistance genes. *Plant J* 30:361–371
- Black W, Mastenbroek C, Mills WR, Peterson LC (1953) A proposal for an international nomenclature of races of *Phytophthora infestans* and of genes controlling immunity in *Solanum demissum* derivatives. *Euphytica* 2:173–240
- Bradshaw JE, Hackett CA, Meyer RC, Milbourne D, McNicol JW, Phillips MS, Waugh R (1998) Identification of AFLP and SSR markers associated with quantitative resistance to *Globodera pallida* (Stone) in tetraploid potato (*Solanum tuberosum* subsp. *tuberosum*) with a view to marker-assisted selection. *Theor Appl Genet* 97:202–210
- Caten CE, Jinks JL (1968) Spontaneous variability of single isolates of *Phytophthora infestans* I. Cultural variation. *Can J Bot* 46:329–347
- Duncan JM (1999) *Phytophthora*—an abiding threat to our crops. *Microbiol Today* 26:114–116
- El-Kharbotly A, Leonards-Schippers C, Huigen DJ, Jacobsen E, Pereira A, Stiekema WJ, Salamini F, Gebhardt C (1994) Segregation analysis and RFLP mapping of the *R1* and *R3* alleles conferring race-specific resistance to *Phytophthora infestans* in progeny of dihaploid potato parents. *Mol Gen Genet* 242:749–754
- El-Kharbotly A, Palomino-Sanchez C, Salamini F, Jacobsen E, Gebhardt C (1996) *R6* and *R7* alleles of potato conferring race-specific resistance to *Phytophthora infestans* (Mont.) de Bary identified genetic loci clustering with the *R3* locus on chromosome XI. *Theor Appl Genet* 92:880–884
- Flor HH (1971) Current status of the gene-for-gene concept. *Annu Rev Phytopathol* 9:275–296
- Ganal MW, Simon R, Brommonschenkel S, Arndt M, Phillips MS, Tanksley SD, Kumar A (1995) Genetic mapping of a wide spectrum nematode resistance gene (*Hero*) against *Globodera rostochiensis* in tomato. *Mol Plant Microbiol Interact* 8:886–891
- Gebhardt C, Ritter E, Barone A, Debener T, Walkemeier B, Schachtschabel U, Kaufmann H, Thompson RD, Bonierbale MW, Ganal MW, Tanksley SD, Salamini F (1991) RFLP maps of potato and their alignment with the homologous tomato genome. *Theor Appl Genet* 83:49–57
- Hawkes JG (1990) The potato: evolution, biodiversity and genetic resources. Smithsonian Institution Press, Washington
- Huang S (2005) The discovery and characterization of the major late blight resistance complex in potato: genomic structure, functional diversity and implications. PhD thesis, Wageningen University, Wageningen
- Huang S, Vleeshouwers VGAA, Werij JS, Hutten RCB, van Eck HJ, Visser RGF, Jacobsen E (2004) The *R3* resistance to *Phytophthora infestans* in potato is conferred by two closely linked R genes with distinct specificities. *Mol Plant Microbiol Interact* 17:428–435
- Jacobs JMF, van Eck HJ, Arens P, Verkerk-Bakker B, te Lintel Hekkert B, Bastiaanssen HJM, El-Kharbotly A, Pereira A, Jacobsen E, Stiekema WJ (1995) A genetic map of potato (*Solanum tuberosum*) integrating molecular markers, including transposons and classical markers. *Theor Appl Genet* 91:289–300
- de Jong W, Forsyth A, Leister D, Gebhardt C, Baulcombe DC (1997) A potato hypersensitive resistance gene against potato virus X maps to a resistance gene cluster on chromosome 5. *Theor Appl Genet* 95:246–252
- Kreike CM, De Koning JRA, Vinke JH, van Ooijen JW, Stiekema WJ (1994) Quantitatively-inherited resistance to *Globodera pallida* is dominated by one major locus in *Solanum spagazzinii*. *Theor Appl Genet* 88:764–769
- Leister D, Ballvora A, Salamini F, Gebhardt C (1996) A PCR-based approach for isolating pathogen resistance genes from potato with potential for wide application in plants. *Nat Genet* 14:421–429
- Leonards-Schippers C, Gieffers W, Salamini F, Gebhardt C (1992) The *R1* gene conferring race-specific resistance to *Phytophthora infestans* in potato is located on potato chromosome V. *Mol Gen Genet* 233:278–283
- Li X, van Eck HJ, Rouppe van der Voort JNAM, Huigen DJ, Stam P, Jacobsen E (1998) Autotetraploids and genetic mapping using common AFLP markers: the *R2* allele conferring resistance to *Phytophthora infestans* mapped on potato chromosome 4. *Theor Appl Genet* 96:1121–1128
- Lukowitz W, Gillmor CS, Scheible WR (2000) Positional cloning in *Arabidopsis*. Why it feels good to have a genome initiative working for you. *Plant Physiol* 123:795–805
- Malcolmson JF, Black W (1966) New R genes in *Solanum demissum* Lindl. and their complementary races of *Phytophthora infestans* (Mont.) de Bary. *Euphytica* 15:199–203
- Marano MR, Malcuit I, de Jong W, Baulcombe DC (2002) High-resolution genetic map of *Nb*, a gene that confers hypersensitive resistance to potato virus X in *Solanum tuberosum*. *Theor Appl Genet* 105:192–200
- Mastenbroek C (1953) Experiments on the inheritance of blight immunity in potatoes derived from *Solanum demissum* Lindl. *Euphytica* 2:197–206
- Meksem K, Leister D, Peleman J, Zabeau M, Salamini F, Gebhardt C (1995) A high-resolution map of the vicinity of the *R1* locus on chromosome V of potato based on RFLP and AFLP markers. *Mol Gen Genet* 249:74–81
- Meyers BC, Chin DB, Shen KA, Sivaramakrishnam S, Lavelle DO, Zhang Z, Michelmore RW (1998) The major resistance gene cluster in lettuce is highly duplicated and spans several megabases. *Plant Cell* 10:1817–1832
- Michelmore RW, Meyers BC (1998) Clusters of resistance genes in plants evolve by divergent selection and a birth-and-death process. *Genome Res* 8:1113–1130
- Michelmore RW, Paran I, Kesseli RV (1991) Identification of markers linked to disease-resistance genes by bulked segregant analysis: a rapid method to detect markers in specific genomic regions by using segregating populations. *Proc Natl Acad Sci USA* 88:9828–9832
- Os H van, Buntjer J, Stam P, van Eck HJ (2000) Evaluation of two algorithms for the construction of genetic linkage maps. International Conference of Plant Animal Genome Research, PAG VIII, 9–12 January 2000, San Diego (<http://www.intl-pag.org/pag/8/abstracts/pag8621.html>)
- Park TH, van der Vossen E, Vleeshouwers VGAA, Tan A, Visser RGF, van Eck HJ (2004) Major resistance genes for tuber and leaf resistance to *Phytophthora infestans* in potato: an outline of a PhD project. *Crop Functional Genomics* 2004, April 2004, Jeju, Korea, p 93
- Ritter E, Debener T, Barone A, Salamini F, Gebhardt C (1991) RFLP mapping on potato chromosomes of two genes controlling extreme resistance to potato virus X (PVX). *Mol Gen Genet* 227:81–85
- Rouppe van der Voort J, Lindeman W, Folkertsma R, Hutten R, Overmars H, van der Vossen E, Jacobsen E, Bakker J (1998) A

- QTL for broad-spectrum resistance to cyst nematode species (*Globodera* spp.) maps to a resistance gene cluster in potato. *Theor Appl Genet* 96:654–661
- Stam P (1993) Construction of intergrated genetic linkage maps by means of a new computer package: JoinMap. *Plant J* 3:739–744
- Vleeshouwers VGAA, van Dooijweert W, Paul Keizer LC, Sijpkens L, Govers F, Colon LT (1999) A laboratory assay for *Phytophthora infestans* resistance in various *Solanum* species reflects the field situation. *Eur J Plant Pathol* 105:241–250
- Voorrips RE (2002) MapChart: software for the graphical presentation of linkage maps and QTLs. *J Hered* 93:77–78
- Vos P, Hogers R, Bleeker M, Reijans M, van de Lee T, Hornes M, Frijters A, Pot J, Peleman J, Kuiper M, Zabeau M (1995) AFLP: a new technique for DNA fingerprinting. *Nucleic Acids Res* 23:4407–4414
- van der Vossen E, Sikkema A, Hekkert BL, Gros J, Stevens P, Muskens M, Wouters D, Pereira A, Stiekema W, Allefs S (2003) An ancient R gene from the wild potato species *Solanum bulbocastanum* confers broad-spectrum resistance to *Phytophthora infestans* in cultivated potato and tomato. *Plant J* 36:867–882
- Wu KK, Burnquist W, Sorrells ME, Tew TL, Moore PH, Tanksley SD (1992) The detection and estimation of linkage in polyploids using single-dose restriction fragments. *Theor Appl Genet* 83:294–300