

# Transformation of Folate-Consuming *Lactobacillus gasseri* into a Folate Producer

Arno Wegkamp, Marjo Starrenburg, Willem M. de Vos, Jeroen Hugenholtz,\*  
and Wilbert Sybesma

Wageningen Centre for Food Sciences and NIZO food research, Ede, The Netherlands

Received 14 August 2003/Accepted 20 February 2004

**Five genes essential for folate biosynthesis in *Lactococcus lactis* were cloned on a broad-host-range lactococcal vector and were transferred to the folate auxotroph *Lactobacillus gasseri*. As a result *L. gasseri* changed from a folate consumer to a folate producer. This principle can be used to increase folate levels in many fermented food products.**

Folate is an essential component in the human diet, and adequate intake of folate may prevent the occurrence of diseases and syndromes like neural tube defect, coronary heart disease, anemia, and certain types of cancer (9). Food products like green vegetables, meat, and fermented dairy products contain significant folate levels. Despite this, folate deficiency occurs throughout the world, including several well-developed countries. Recently, it has been shown that metabolic engineering can be used to increase folate levels in fermented foods (12).

Lactic acid bacteria such as *Lactococcus lactis* and *Lactobacillus plantarum* have the ability to synthesize folate, which is a biological cofactor involved in their amino acid and nucleotide metabolism (7, 11). The genes for folate biosynthesis have been identified (6, 12). The biosynthetic pathway includes seven consecutive steps, in which the precursor guanosine triphosphate is converted into tetrahydrofolate (10). However, some lactic acid bacteria, such as *Lactobacillus gasseri* strain ATCC 33323, cannot synthesize folate, because the genes involved in folate biosynthesis are lacking in the genome except for the two genes, *folA* and *folC*, involved in regeneration and retention of reduced folates taken up from the medium ([http://genome.jgi-psf.org/draft\\_microbes/lacga/lacga.draft.html](http://genome.jgi-psf.org/draft_microbes/lacga/lacga.draft.html)).

The folate biosynthetic genes of *L. lactis* MG1363 are organized in a folate gene cluster, consisting of six genes (*folA*, *folB*, *folKE*, *folP*, *ylgG*, and *folC*) (Fig. 1) (12). In the present work we describe the transformation of the folate-consuming *L. gasseri* into a folate producer by the transfer of a broad-host-range plasmid containing the folate gene cluster from *L. lactis*.

**Cloning and transformation of the folate gene cluster of *L. lactis* into *L. gasseri*.** The plasmid pNZ7017 (12) was digested by using XbaI and SphI (both Invitrogen, Paisley, United Kingdom) as restriction enzymes. The 3.1-kb DNA fragment that was obtained from the digestion consisted of the constitutive *pepN* promoter (14), a part of the multiple cloning site, chloramphenicol resistance marker, and replication genes that originated from pNZ12 (1). The folate gene cluster (*folB*, *folKE*, *folP*, *ylgG*, and *folC*) of *L. lactis* was amplified by PCR

by using high-fidelity *Pwo* polymerase (Invitrogen). The forward primer SphfolB-F (5'-AGGAAGCATGCCTTACAAAATAAACTTAATAATATG-3') was extended at the 5' end, creating an SphI restriction site overlapping the start codon of *folB*. The reverse primer folCXba-R (5'-TCTCTAGACTACTTTTCTTTTTTCAAAAATTCACG-3') was extended at the 5' end, creating an XbaI restriction site that overlapped the stop codon of *folC* (Fig. 1). The amplified PCR fragment was restricted with XbaI and SphI. Subsequently, the two fragments were ligated by using T4 ligase (Invitrogen), generating a translational fusion between the constitutive promoter of the *pepN* gene (14) and the folate gene cluster (Fig. 1). The resulting plasmid was designated pNZ7019. After transformation to *L. lactis* NZ9000 and subsequent cultivation of the strain, the plasmid was harvested as described previously (2).

*L. gasseri* (ATCC 33323) was transformed with purified pNZ7019 by using an established procedure (8) and was plated on MRS medium (Merck, Darmstadt, Germany) containing 10 µg of chloramphenicol/ml. After incubation for 40 h at 37°C, chloramphenicol-resistant colonies were examined for the presence of pNZ7019 by using restriction analyses. An *L. gasseri* colony harboring pNZ7019 was used for renewed cultivation by using the same growth conditions as previously described. Random amplified polymorphic DNA fingerprint analysis was used to confirm the identity of the transformant harboring pNZ7019 as *L. gasseri* ATCC 33323 (results not shown).

**Conversion of folate consumer into folate producer.** A modified Folic Acid Casei Medium (FACM) (Difco, Becton Dickinson and Co., Sparks, Md.) was developed for growth and subsequent folate analysis of the *L. gasseri* wild-type strain and the *L. gasseri* strain harboring pNZ7019. The FACM was enriched with 1 mg of vitamin B<sub>12</sub> (Sigma-Aldrich Chemie, GmbH, Steinheim, Germany)/liter and 1 ml of Tween 80 (Merck, Darmstadt, Germany)/liter. The wild-type strain could not grow at 37°C unless folate was added (1.0 mg/liter), whereas the strain harboring pNZ7019 showed folate-independent growth.

Folate levels were quantified by using the *Lactobacillus casei* microbiological assay, including enzymatic deconjugation of the polyglutamate tail (5, 11, 12). The *L. gasseri* strain harboring pNZ7019 produced significant intracellular and extracellu-

\* Corresponding author. Mailing address: NIZO food research, P.O. Box 20, 6710 BA Ede, The Netherlands. Phone: 31-318-659540. Fax: 31-318-650400. E-mail: jeroen.hugenholtz@nizo.nl.




FIG. 1. Construction of pNZ7019 by restriction of pNZ7017 and the amplification of the folate gene cluster from chromosomal DNA of *L. lactis* MG1363.

lar folate levels (Fig. 2). As expected, the wild-type strain consumed folate from the medium and intracellular folate concentrations remained below the detection limit (Fig. 2).

**Stability of the folate production in the *L. gasseri* strain.** *L. gasseri* strain ATCC 33323 harboring pNZ7019 was cultivated for approximately 30 generations on MRS medium supple-

mented with 10 µg of chloramphenicol/ml at 37°C. The culture was plated on MRS agar plates supplemented with 10 µg of chloramphenicol/ml. Subsequently, 100 colonies were transferred to folate-free FACM plates containing 10 µg of chloramphenicol/ml. Since all colonies grew on these plates, it appears that the folate biosynthesis is stably maintained in the




FIG. 2. Folate production of the *L. gasseri* wild-type strain (side marked 1) and the *L. gasseri* strain harboring pNZ7019 (side marked 2), grown on modified FACM. White bar, extracellular folate levels; gray bar, intracellular folate levels; and black bar, total folate levels. The *L. gasseri* wild-type strain could not grow without supplementation of folate. Therefore, folate levels depicted in the figure are corrected for folate added to the medium. Error bars indicate the standard deviation of the folate microbiological assay over two independent measurements.

pNZ7019 vector for more than 30 generations of growth in the presence of folate. Sequential cultivation in folate-rich medium resulted in decreased folate production by the transformant (data not shown). This is presumably a result of instability of the folate gene cluster.

**Conclusion.** The five genes, i.e., *folB*, *folKE*, *folP*, *ylgG*, and *folC*, directing folate biosynthesis in *L. lactis* were transferred to *L. gasseri* by using a derivative of the broad-host-range vector pNZ12 (1). These genes are sufficient to introduce a folate biosynthesis pathway in this folate auxotroph lactic acid bacterium, thereby transforming a folate consumer into a folate producer. *L. gasseri* is currently marketed as a probiotic (4), and when the described strategy is used, this lactic acid bacterium can be used to enrich (fermented) foods with the essential B vitamin, folate, in addition to conferring its health-promoting effect on the consumer (3, 13).

#### REFERENCES

1. De Vos, W. M. 1987 Gene cloning and expression in lactic streptococci. *FEMS Microbiol. Rev.* **46**:281–295.
2. De Vos, W. M., P. Vos, H. de Haard, and I. Boerrigter. 1989. Cloning and expression of the *Lactococcus lactis* SK11 gene encoding an extracellular serine proteinase. *Gene* **85**:169–176.
3. Fernandez, M. F., S. Boris, and C. Barbes. 2003. Probiotic properties of human lactobacilli strains, to be used in the gastrointestinal tract. *J. Appl. Microbiol.* **94**:449–455.
4. Holzapfel, W. H., P. Haberer, J. Snel, U. Schillinger, and J. H. J. Huis in't Veld. 1998. Overview of gut flora and probiotics. *Int. J. Food Microbiol.* **41**:85–101.
5. Horne, D. W., and D. Patterson. 1988. Lactobacillus casei microbiological assay of folic acid derivatives in 96-well microtiter plates. *Clin. Chem.* **34**:2357–2359.
6. Kleerebezem, M., J. Boekhorst, R. van Kranenburg, D. Molenaar, O. P. Kuipers, R. Leer, R. Turchini, S. A. Peters, H. M. Sandbrink, M. W. Fiers, W. Stiekema, R. M. Lankhorst, P. A. Bron, S. M. Hoffer, M. N. Groot, R. Kerkhoven, M. de Vries, B. Ursing, W. M. de Vos, and R. J. Siezen. 2003. Complete genome sequence of *Lactobacillus plantarum* WCFS1. *Proc. Natl. Acad. Sci. USA.* **100**:1990–1995.
7. Lin, M. Y., and C. M. Young. 2000. Folate levels in cultures of lactic acid bacteria. *Int. Dairy J.* **10**:409–414.
8. Luchansky, J. B., P. M. Muriana, and T. R. Klaenhammer. 1988. Application of electroporation for transfer of plasmid DNA to *Lactobacillus*, *Lactococcus*, *Leuconostoc*, *Listeria*, *Pediococcus*, *Bacillus*, *Staphylococcus*, *Enterococcus*, and *Propionibacterium*. *Mol. Microbiol.* **2**:637–646.
9. Lucock, M. 2000. Folic acid: nutritional biochemistry, molecular biology, and role in disease processes. *Mol. Genet. Metab.* **71**:121–138.
10. Suzuki, Y., and G. M. Brown. 1974. The biosynthesis of folic acid. XII. Purification and properties of dihydroneopterin triphosphate pyrophosphohydrolase. *J. Biol. Chem.* **249**:2405–2410.
11. Sybesma, W., M. Starrenburg, L. Tijsseling, M. H. N. Hoefnagel, and J. Hugenholtz. 2003. Effects of growth conditions on folate production by lactic acid bacteria. *Appl. Environ. Microbiol.* **69**:4542–4548.
12. Sybesma, W., M. Starrenburg, M. Kleerebezem, I. Mierau, W. M. de Vos, and J. Hugenholtz. 2003. Increased production of folate by metabolic engineering of *Lactococcus lactis*. *Appl. Environ. Microbiol.* **69**:3069–3076.
13. Usman and A. Hosono. 2001. Hypocholesterolemic effect of *Lactobacillus gasseri* SBT0270 in rats fed a cholesterol-enriched diet. *J. Dairy Res.* **68**:617–624.
14. Van Alen-Boerrigter, I. J., R. Baankreis, and W. M. de Vos. 1991. Characterization and overexpression of the *Lactococcus lactis pepN* gene and localization of its product, aminopeptidase N. *Appl. Environ. Microbiol.* **57**:2555–2561.