

# 1 **Model Choice and Crucial Tests. On the Empirical Epistemology** 2 **of the Higgs Discovery**

3

4 Peter Mättig<sup>1</sup> & Michael Stöltzner<sup>2</sup>

5 -

6

7 *Abstract:* Our paper discusses the epistemic attitudes of particle physicists on the  
8 discovery of the Higgs boson at the Large Hadron Collider (LHC). It is based on  
9 questionnaires and interviews made shortly before and shortly after the  
10 discovery in 2012. We show, to begin with, that the discovery of a Standard  
11 Model (SM) Higgs boson was less expected than is sometimes assumed. Once the  
12 new particle was shown to have properties consistent with SM expectations –  
13 albeit with significant experimental uncertainties –, there was a broad  
14 agreement that ‘a’ Higgs boson had been found. Physicists adopted a two-  
15 pronged strategy. On the one hand, they treated the particle as a SM Higgs boson  
16 and tried to establish its properties with higher precision; on the other hand,  
17 they searched for any hints of physics beyond the SM. This motivates our first  
18 philosophical thesis: the Higgs discovery, being of fundamental importance and  
19 establishing a new kind of particle, represented a crucial experiment if one  
20 interprets this notion in an appropriate sense. By embedding the LHC into the  
21 tradition of previous precision experiments and the experimental strategies thus  
22 established, Duhemian underdetermination is kept at bay. Second, our case study  
23 suggests that criteria of theory (or model) preference should be understood as  
24 epistemic and pragmatic values that have to be weighed in factual research  
25 practice. The Higgs discovery led to a shift from pragmatic to epistemic values as  
26 regards the mechanisms of electroweak symmetry breaking. Complex criteria,  
27 such as naturalness, combine epistemic and pragmatic values, but are coherently  
28 applied by the community.

29

30

## 31 **1. Introduction**<sup>3</sup>

32

33 The discovery of a<sup>4</sup> Higgs boson at the Large Hadron Collider (LHC) of the European  
34 Laboratory CERN, announced in July 2012, is arguably one of the most important

---

<sup>1</sup> Bergische University of Wuppertal, Department of Mathematics and Natural Science, Gaußstrasse 20, 42119 Wuppertal; now at Department of Physics and Astronomy, University of Bonn, Nussallee 12, 53115 Bonn, Germany; Email: peter.mattig@cern.ch.

<sup>2</sup> Department of Philosophy, University of South Carolina, Columbia, SC 29208, USA; Email: stoeltzn@sc.edu.

<sup>3</sup> The study was performed as part of the project ‘Model Dynamics’ supported by the DFG (project no. MA 2793/2-1). It is based on questionnaires that were developed by the authors, Arianna Borrelli, Robert Harlander, and Friedrich Steinle, and interviews conducted by Arianna Borrelli. Karsten Egger assisted in the evaluation of the questionnaire. We acknowledge the help of Annette Holtkamp in obtaining the SPIRES email list. We acknowledge discussions within the recently established DFG Research Unit ‘Epistemology of the LHC’ and detailed comments by Robert Harlander, Martin King, and Gregor Schiemann. Some other results from this study can be found in (Borrelli 2016) and her presentation at CERN available under <https://indico.cern.ch/event/232108/>. We thank the anonymous referees for their most helpful criticism and manifold suggestions. We thank Sophie Ritson for having made us aware of Baetu (2017).

35 scientific achievements of the past few decades. The discovery received world-wide  
36 attention; two of the inventors of the Higgs mechanism, François Englert and Peter  
37 Higgs, were awarded the 2013 Nobel Prize in physics. The Higgs boson had been the  
38 final piece of the so-called Standard Model of particle physics (SM) not observed by  
39 previous experiments. The Higgs mechanism in the SM was required to generate masses  
40 of the elementary particles in a consistent way. Even though with the discovery of the  
41 Higgs boson, the SM – terminology notwithstanding – has now become one of the most  
42 successful scientific theories of contemporary physics, all particle physicists agree that it  
43 will not be the final word. There are both compelling internal and external reasons to  
44 postulate physics beyond the SM (BSM).

45  
46 In retrospect, the discovery of the Higgs boson might seem to be just the final step in a  
47 long series of discoveries and precision tests in which stronger and stronger accelerator  
48 experiments confirmed all particles of the SM and scrutinized their interactions. The  
49 present paper argues that as regards the community of elementary particle physics this  
50 picture needs qualification. In actual fact, even shortly before the Higgs discovery a  
51 significant percentage of physicists raised concerns whether it would at all be found at  
52 the LHC and expressed preferences for other explanations of the particle masses.

53  
54 In this paper, results of questionnaires and interviews with LHC physicists shortly  
55 before (autumn 2011) and shortly after (autumn 2012) the discovery are presented and  
56 analysed. From these empirical sources, we reconstruct the physicists' beliefs in the  
57 adequacy of certain models, in the outcome of the LHC experiments, and concerning the  
58 possible impacts of the LHC experiments on those models. This will help us to  
59 understand the epistemic attitudes of particle physicists, and the principles and  
60 strategies guiding their research. Our empirically informed epistemological  
61 investigation also promises new insights for a philosophical analysis of how actual and  
62 expected experimental findings, on the one side, and pragmatic quality criteria of  
63 models, on the other, influence the research agendas of particle physicists.

64  
65 We have limited the scope of the present paper to a specific part of the empirical  
66 material available in the questionnaires and interviews: to the Higgs mechanism and  
67 competing accounts of mass generation, and to the presently most discussed pragmatic  
68 quality criterion, naturalness. It must be said, however, that the LHC was, from the very  
69 beginning, designed not only to search for the Higgs boson but also to probe the deep  
70 TeV energy range and find signs of BSM physics. Whereas the first objective has now  
71 been achieved, no 'new physics' BSM has been observed to date.<sup>5</sup>

72  
73 The specific descriptive questions addressed in this paper are as such:

- 74
- 75 1. Did physicists in 2011 expect the Higgs boson to be discovered at the LHC and how  
76 did they evaluate the Higgs candidate in 2012, that is, before its properties were  
77 known to a sufficient extent? What was their assessment of alternative models for  
78 mass generation in 2011 and in 2012?
  - 79 2. How important was the naturalness problem, a major guiding principle to develop  
80 models of physics beyond the SM (BSM), in shaping physicists' attitudes and  
81 preferences?

82  
83 We will show that, in 2011, physicists were rather undecided whether the SM Higgs

---

<sup>4</sup> We adopt the usual terminology and address the SM Higgs as 'the' Higgs, whereas those models with a potentially more complicated Higgs sector as containing 'a' Higgs boson.

<sup>5</sup> The attempts at finding physics BSM and their effects on the thinking of LHC physicists will be discussed in a separate paper.

84 boson would eventually be found, that is, even a few months before the first evidence  
85 was reported. However, once a candidate had been observed in 2012, they quickly  
86 embraced the notion that ‘a’ Higgs boson had been found. Its discovery immediately  
87 affected the research directions in particle physics. The experimental results pulled in  
88 different directions as regards the naturalness problem. There was, on the one hand,  
89 less motivation to search for alternatives to the Higgs mechanism. On the other hand,  
90 after finding the Higgs boson, the naturalness problem posed by the scalar Higgs particle  
91 changed from a virtual into a real problem, that is, there existed empirical results  
92 directly relevant for it. But since 2012 no BSM effect to cure this problem has been  
93 found. This has led some physicists to develop a more critical attitude as to naturalness’  
94 significance for elementary particle physics.

95  
96 The physical developments prompt the following philosophical questions.

- 97  
98 3. What do the epistemic attitudes of particle physicists shown in the questionnaire  
99 and the interviews mean for the significance and application of criteria of theory (or  
100 model)<sup>6</sup> choice and the principles and epistemic values guiding model development?  
101 4. What does the comparison of the situations before and after the discovery of the  
102 Higgs boson signify for the relationship between theory (or models) and  
103 experiment? In particular, was the Higgs discovery a crucial experiment for the SM?  
104

105 The paper is organised as follows. After a brief introduction into the theoretical  
106 motivation for the Higgs mechanism and the experimental attempts to find evidence for  
107 a Higgs boson (Section 2), we provide the background of the philosophical problems  
108 raised (Section 3) and discuss the methodology of our study (Section 4). The  
109 presentation of the results will be subdivided into the outcomes of the questionnaire  
110 and the interviews in 2011 (Section 5) and in 2012 (Section 6) respectively. Finally  
111 (Section 7), we outline our answers to the above-mentioned four questions.  
112  
113

## 114 ***2. The physics of electroweak symmetry breaking***

115  
116 Several articles of both physicists and philosophers discuss the emergence of what is by  
117 now called the ‘Higgs’ mechanism (Cf. Ellis, Gaillard, Nanopoulos, 2015; Nobel laudatio  
118 2013; Karaca 2013b). Here, only a brief account of the motivation and the concepts  
119 behind the Higgs boson can be given. In the early 1960s, various models were developed  
120 to unify two interactions governing the subnuclear world, the electromagnetic and the  
121 weak ones. These unifications adopted the concept of local gauge symmetry that had  
122 previously been applied successfully to quantum electrodynamics (QED). In brief, this  
123 symmetry means that the theory is invariant under a specific space-time dependent  
124 transformation of the quantum fields. Assuming this symmetry in the SM leads to a  
125 consistent interacting field theory, which for particle physicists means free of any  
126 infinities after renormalization.  
127

128 However, a major problem that physicists were facing in applying local gauge symmetry  
129 to weak interactions was that observations implied that the corresponding gauge  
130 bosons have a non-vanishing mass. As such, gauge boson masses break the symmetry  
131 explicitly, thus leading to theoretical inconsistencies, such as the violation of unitarity.  
132 To remedy this, in the 1960s, physicists used the concept of spontaneous symmetry  
133 breaking (SSB) to generate gauge boson masses in a gauge invariant way at the cost of

---

<sup>6</sup> In line with the current philosophical literature, we consider models as autonomous entities in scientific theorizing, not as the logical models of a theory. For how one can apply this conception to elementary particle physics, cf. Borrelli & Stöltzner (2013) and Stöltzner (2014).

134 introducing an additional scalar, i.e. spin-less, particle, which became known as the  
135 Higgs boson. This particle was discovered at the LHC some 50 years after its invention.  
136 The Higgs sector of the SM is a novel element in physics, in that it describes the mass of  
137 elementary particles in terms of their interaction with an elementary scalar field.

138  
139 Whereas the weak and the electromagnetic components of the ‘electroweak’ theory have  
140 almost the same strength at very high energies, they are substantially different at low  
141 energies, since only the weak interaction invokes a massive interacting particle.  
142 Therefore, the mechanism of mass generation is also referred to as ‘electroweak  
143 symmetry breaking’ (EWSB). The Higgs mechanism was originally only devised to give  
144 mass to the weak gauge bosons  $W^{+/-}$  and  $Z^0$ . (The latter represents the electrically  
145 neutral component of the weak interaction, which, however, has an admixture of an  
146 electromagnetic component.) It turned out that the Higgs mechanism could also be  
147 applied to give masses to fermions, through a Yukawa interaction, albeit without  
148 predicting their numerical values.

## 149 150 **2.1 The experimental search for the Higgs boson**

151  
152 The general conception of the Higgs mechanism just outlined was developed into  
153 phenomenological predictions<sup>7</sup> opening the way for experimental searches of the Higgs  
154 boson. Given the masses of the  $W$  and  $Z$  bosons, the Higgs mechanism introduced just  
155 one additional parameter to the SM that had to be determined by experiment, notably by  
156 measuring the Higgs mass<sup>8</sup>. Whereas the theory did not provide a prediction for this  
157 mass, it did lead to an upper bound of 800 GeV to maintain theoretical consistency.  
158 Depending on its mass, it could be unambiguously predicted how the Higgs boson is  
159 produced and the way it can be seen by experiments. Since a Higgs boson would only  
160 exist for small fractions of a second, it would decay, depending on its mass, mainly into  
161 massive fermions and  $W$  and  $Z$  bosons.

162  
163 As a result, a clear strategy for finding the Higgs boson was devised.<sup>9</sup> However, this did  
164 not make Higgs searches easy. Essentially no experiment before the start of CERN’s  
165 Large Electron Positron Collider (LEP) in 1989 was sensitive to the Higgs boson. At the  
166 end of the LEP data taking no significant signal was observed. However, the sensitivity  
167 of LEP was such that a Higgs of 114.4 GeV or less should have been found, allowing  
168 physicists to place a lower limit on the SM Higgs mass. Between the end of LEP and the  
169 start of LHC, an additional small mass interval around 160 GeV could be excluded at the  
170 Tevatron. The outstanding precision of the LEP data and theoretical calculations based  
171 on the SM provided an indirect sensitivity to the Higgs mass by quantum fluctuations,  
172 e.g. loop corrections to the  $W$  and  $Z$  bosons, bounding it to be lighter than 157 GeV.

173  
174 In 2010, data taking at the LHC started for the final assault. It was clear that the LHC had  
175 the sensitivity to observe the Higgs boson in the remaining allowed mass range, using  
176 the decay modes that were unambiguously predicted for a SM Higgs. Relatively soon one  
177 could exclude a high mass Higgs of 200-600 GeV – in full agreement with LEP’s indirect  
178 limits. With the rapid increase in data rate, both the ATLAS and CMS experiments

---

<sup>7</sup> Cf. Ellis, Gaillard, Nanopoulos (1976).

<sup>8</sup> More precisely, the Higgs potential has two parameters, one of which is related to the masses of the  $W$  and  $Z$  – bosons and connected to the vacuum expectation value  $v$  leaving one additional parameter that had not been determined before the Higgs discovery. It should be noted that in the SM the Yukawa couplings  $g$ , i.e. the Higgs couplings to the fermions are given by  $g = \sqrt{2} * m_f / v$ ,  $m_f$  being the mass of the fermion. Therefore, the Yukawa couplings re-express the fermion masses but are not counted as extra free parameters.

<sup>9</sup> One publication was even titled “The Higgs Hunter’s Guide” (Gunion et al. 1990).

179 reported, at a CERN colloquium on December 13, 2011, an excess of events that could be  
180 taken as initial evidence for a new particle around 126 GeV. On the other hand, the  
181 probability that this would be just a background fluctuation was still too high to claim an  
182 observation. However, half a year later much more data had been accumulated, such  
183 that both detectors presented, at a special CERN seminar, a signal of 5 standard  
184 deviations each. The data correspond to a background fluctuation probability of about  
185  $10^{-9}$ , where the background is considered as SM without Higgs. By convention in particle  
186 physics, this was sufficiently small to claim a discovery, an observation.<sup>10</sup> A few weeks  
187 later, the two experiments published their data. (Aad et al. 2012, Chatrchan et al. 2012).

188  
189 Still, the data were not sufficient to definitely claim this to be the long-awaited Higgs  
190 boson. Some important properties had not yet been confirmed, and the precision of the  
191 measurements on production and decay properties was still marginal. On the other  
192 hand, those properties that were observed corresponded to what is expected for a SM  
193 Higgs boson. For instance, the particle had been found in two decay modes with rates  
194 consistent with the expectation, and it had a mass in agreement with the direct and  
195 indirect limits known from previous experiments. As of today (2017), more properties  
196 of the discovered particle have been studied, the decay modes and the mass have been  
197 measured to higher precision, in accordance with the SM. Even though there is still need  
198 for further measurements, the majority among physicists now considers the new  
199 particle is indeed the Higgs boson.

200  
201

## 202 ***2.2 Alternatives to the Higgs boson***

203

204 Already shortly after the invention of the Higgs mechanism, several authors expressed  
205 discontent because this solution of the SSB problem appeared largely ad-hoc. For  
206 example, it has limited predictive power in that it cannot determine the quark and  
207 lepton masses. Concern was also raised that the Higgs mechanism introduces a new  
208 concept into the theory for the one and only purpose of mass generation. Over the years,  
209 the list of issues cited by physicists in this respect has expanded. (Cf. Friederich,  
210 Harlander, & Karaca 2014, sect. 3).

211

212 Several alternative mechanisms of EWSB have emerged over the past decades. They  
213 used a scalar particle and a Higgs-like potential to generate mass. However, in many  
214 cases, the conceptual framework of the alternative models was very different from, and  
215 implied physics beyond the SM. These BSM models will be considered in this article only  
216 in relation to mass generation.

217

218 A fairly straightforward modification of the original Higgs mechanism was to extend the  
219 Higgs sector. Originally, one complex Higgs doublet was assumed, leading to four fields  
220 one of which would be the observable Higgs boson, whereas the others would not be  
221 directly observable. However, one can also introduce, e.g., a second doublet leading to  
222 five physical elementary Higgs bosons with no change in the principal mechanism of  
223 mass generation. Such models allow the different Higgs bosons to assume different  
224 roles.

225

226 The two Higgs doublet model is of special interest in BSM considerations since it is the  
227 minimally required Higgs sector in the framework of Supersymmetry, the most often  
228 discussed extension of the SM. Supersymmetry assumes a new fundamental symmetry  
229 of particles with integer and half-integer spins. In the context of LHC physics,

---

<sup>10</sup> On the criteria when particle physicists claim ,evidence', versus ,observation' or ,exclusion' see the 'Prologue' to Franklin (2013).

230 Supersymmetry typically is studied in several variants of the MSSM ('minimal  
231 supersymmetric SM') that contain a rather broad range of features that allow one solve  
232 some basic problems of the SM. Among those is providing a Dark Matter candidate and  
233 solving the naturalness problem (see below). Moreover, the MSSM is the only BSM  
234 model that makes a firm prediction on the upper limit of the Higgs mass: it has to be  
235 lighter than about 130 GeV, a limit that is much tighter than the range allowed by the  
236 SM. (This limit of 130 GeV is valid for all SUSY models considered at the LHC and for  
237 SUSY scales of some 1 TeV.)<sup>11</sup>

238

239 Another class of models assumes the Higgs boson to be a composite, i.e. made up of sub-  
240 constituents. The first model of this kind was devised at the end of the 1970s by  
241 essentially copying concepts known from the strong interactions that explain hadron  
242 masses. This mechanism was dubbed 'Technicolour'; since it involved strong  
243 interactions, it was considered as a type of 'strong' or 'dynamical' EWSB. The realisation  
244 of these models led to inconsistencies with measurements, such that this approach by  
245 now has become disfavoured. However, the concept of composite Higgs particles has  
246 been implemented within multiple frameworks invoking additional symmetries, new  
247 interactions, or additional spatial dimensions (e.g. Csaki and Tanedo 2016).

248

249 All these alternative models assume scalar particles like the SM Higgs boson to generate  
250 the masses of gauge bosons and fermions. However, the properties of these scalars are  
251 different, albeit sometimes by a rather small amount given by tuneable free parameters.  
252 All of them also lead to new phenomena, e.g. more scalars and more fermions.

253

254

### 255 **2.3 The Naturalness problem**

256

257 From a theoretical perspective, the existence of an elementary scalar Higgs boson  
258 introduces an 'unnaturalness' into the SM. The concept 'naturalness' was introduced in  
259 slightly different forms by 't Hooft (1979) and Susskind (1979). The problem itself has a  
260 longer history (cf. Giudice 2008) and reaches beyond the context of the Higgs problem  
261 (cf. Giudice 2013). During the past decades, naturalness has developed from a merely  
262 technical problem into an influential guiding principle for BSM physics; that is,  
263 extensions of the SM were developed with the explicit aim to remedy the naturalness  
264 problem.

265

266 In a nutshell, the naturalness problem is this: since the fundamental equations of the SM  
267 can only be solved in a perturbative expansion, at each order a theoretically well-  
268 defined correction has to be applied to compensate for quantum fluctuations that would  
269 modify a physical quantity like mass or charge. Such 'renormalisation' is a standard  
270 technical procedure in theoretical particle physics. For the SM particles of spin  $\frac{1}{2}$  or 1  
271 these corrections are of a few percent. In the case of the Higgs boson, which is a scalar,  
272 however, the correction to the square of the Higgs mass grows quadratically with  
273 energy.

274

275 Introducing a cut-off mass where the theory would break down, leads to finite  
276 corrections. In the case of the SM, this could be at the rather high Planck scale, where  
277 gravity becomes important and the SM is known to be insufficient. Assuming such a  
278 scale within the SM, in case of the Higgs mass, makes these corrections appear 'dramatic  
279 and even bizarre' (Peskin and Schroeder 1995, p. 788); for instance, in order to keep the  
280 square of the Higgs mass at its measured value of 125 GeV, corrections have to be

---

<sup>11</sup> There exists a small (logarithmic) dependence of this bound on the masses of the SUSY particles. Even if more Higgs multiplets exist, the bound would only rise to 150 GeV.

281 invoked that are more than  $10^{30}$  times higher than the Higgs mass itself. Furthermore,  
282 these corrections have to be fine-tuned over many decimal places. Although  
283 theoretically viable and consistent, the magnitude of these corrections is considered  
284 'unnatural'. Once this correction is defined the theory is completely consistent and any  
285 dependence on the scale is eliminated.

286  
287 During the past two decades, naturalness has arguably become the most influential  
288 guiding principle for constructing and motivating BSM models. Or more specifically,  
289 many physicists believed that if a SM Higgs boson existed, it would come with new  
290 phenomena to keep the theory 'natural'. For instance, new symmetries, extra spatial  
291 dimensions, or a composite Higgs boson built from smaller objects would avoid  
292 unnaturalness. Allowing for corrections of just a few percent – as for the other sectors of  
293 the SM – these new phenomena should be in the mass range of 1 TeV that is well  
294 covered by the LHC. One has to be aware that there is no clear definition of when a  
295 theory would become unnatural and there is a large freedom how much fine tuning is  
296 considered acceptable. Yet once a bound on the acceptable fine tuning is set, it  
297 determines the mass range at which new phenomena are expected. At any rate, thus far  
298 there has neither been a direct observation nor any clear indirect indication from  
299 precision studies that such a new effect exists.

300  
301

### 302 ***3. Philosophical Background: Theory Choice and Crucial Experiments***

303

304 Our empirical study allows us to address two longstanding problems in philosophy of  
305 science from the perspective of the actual practice of scientists. First (in 3.2), we discuss  
306 the relationship between epistemic and pragmatic (including aesthetic) criteria of  
307 theory choice in the contexts of models of electroweak symmetry breaking. Presently  
308 most discussed among these criteria is naturalness. Second (in 3.3.), we discuss under  
309 which conditions complex experiments, such as the Higgs discovery, are considered  
310 decisive or even crucial. We begin this section, however, by showing that the present  
311 debates about naturalness represent a case in point about the influence of criteria of  
312 theory choice. The general aim of the present section is to give a short survey of the  
313 current philosophical discussion that provides the basis for Section 7.

314

315

#### 316 ***3.1. The Philosophical Challenge of Naturalness***

317

318 Several facets of the naturalness problem have attracted philosophers' attention; among  
319 them are its precise content and to what extent it influences current research in particle  
320 physics. Porter Williams (2015) has distinguished four (closely related) ways to  
321 formulate the naturalness problem: (i) quadratic divergences in renormalisation; (ii) 't  
322 Hooft's (1979) suggestion that setting a small parameter to zero must increase the  
323 symmetry of the system; (iii) a specific version of the problem of fine-tuning of  
324 fundamental constants; (iv) an aesthetic criterion, whose force is derived from various  
325 factors prevailing within the scientific community. Williams argues that none of his four  
326 reformulations captures the whole naturalness problem and believes that it is rather an  
327 expression of the central dogma of effective field theories according to which widely  
328 separated scales should eventually decouple.

329

330 The physicist James Wells (2015) considers (i) as the root of the problem, but  
331 subsequently emphasizes the significant difference between the technical naturalness  
332 (ii) and the absolute naturalness involved in fine-tuning that eventually goes back to  
333 Dirac's classical worries about large dimensionless numbers. He elaborates an example  
334 of an exotically augmented quantum electrodynamics (QED) that consistently

335 instantiates absolute naturalness at the expense of “more parameters, more fields, and  
336 more complexity in the theory.” (2015, 107) He admits that this principle is  
337 controversial, but believes, more generally, “that in the era of the Standard Model’s  
338 ascendancy, the influence of simplicity and Ockham’s razor to theory construction has  
339 paled in comparison to Naturalness.” (2015, 104)<sup>12</sup>

340  
341 Grinbaum (2012) instead has argued that – in virtue of its complex nature – naturalness  
342 is exclusively an aesthetic criterion. Williams (2015) rejects Grinbaum’s interpretation  
343 because aesthetic criteria are notoriously ambiguous. Supersymmetry, for instance, is  
344 considered most promising by many physicists, even though it is aesthetically attractive  
345 in the unbroken state but aesthetically unattractive after its breaking produces a large  
346 number of new constants. Borrelli (2015) argues, that it is precisely the vagueness of the  
347 concept of naturalness that allows it to function as a useful common narrative of the  
348 different subcultures of particle physics, the experimentalists and theoreticians

349  
350 The goal of the present paper is not to analyse all facets of naturalness. Instead we take  
351 it as the currently most important example of a guiding principle for a ‘good’ model  
352 within contemporary particle physics and provide empirical results about its  
353 relationship with other guiding principles. More specifically, we will compare the  
354 relatively new and quantitative concept of naturalness with the more familiar  
355 pragmatic, aesthetic, and qualitative criteria of elegance and simplicity – Ockham’s razor  
356 being one of its manifestations.

### 357 358 ***3.2 Epistemic and pragmatic criteria of theory choice***

359  
360 Philosophers have traditionally distinguished epistemic and pragmatic criteria of theory  
361 choice (or preference). The former, among them empirical adequacy and theoretical  
362 consistency, are held to be rationally compelling. Pragmatic criteria have instead been  
363 seen as a way to decide among epistemically equivalent alternatives by appealing to a  
364 theory’s simplicity or other aesthetic features, or to its fruitfulness for further research.  
365 Among the classical examples are the choice between a geocentric and a heliocentric  
366 world view at the time of Copernicus and the early philosophical debates about the  
367 nature and alleged conventionality of space and time. The philosophical significance of  
368 these criteria of theory choice arises from the problem of underdetermination of theory  
369 by empirical evidence that Pierre Duhem illustrated at the parallelism between  
370 Newton’s corpuscular theory and Huygens’s wave theory of light. Duhem argued that  
371 experimental data never uniquely determine a particular hypothesis because setting up  
372 and confirming a hypothesis presupposes the correctness of many other hypotheses  
373 including the theories governing the measurement devices. If one accepts some version  
374 of the underdetermination argument, pragmatic and aesthetic criteria become more  
375 relevant or even inevitable.

376  
377 Underdetermination is also discussed under the rubrics of theory-ladenness of data or –  
378 following Neurath and Quine – confirmational holism. This means that any experimental  
379 result confirms or refutes both the theory or model under investigation and a large set  
380 of other assumptions that are assumed to be true. Especially in Quine’s hands,  
381 underdetermination and holism took a logical and semantic tack that not only ruled out  
382 that empirical evidence could deductively entail scientific theories, but that additionally  
383 seemed to imply that any theory could be rationally retained in the face of recalcitrant  
384 evidence. Laudan has pointed out that, while the latter may be logically possible,

---

<sup>12</sup> Ockham’s razor instructs us not to add basic entities without any need to do so. It has originally been a metaphysical principle, but the term is nowadays used more broadly. Cf. the following section; see also Wells (2017).



385 scientists do not act “in an evaluative vacuum.” (1990, 276) To his mind, the non-  
386 uniqueness of theory resulting from Duhemian underdetermination can be accepted  
387 without adopting an egalitarian approach towards rival theories. Laudan and Leplin,  
388 more generally, held “that the epistemic bearing of evidence on theory is ... subject to  
389 reinterpretation as science grows and may be indeterminate at a particular point in the  
390 process of growth.” (1991, 455) Norton, moreover, has argued that ampliative  
391 inferences remain valid if underdetermination focuses locally, “on the confirmation of  
392 hypotheses by scientists in actual scientific practice,” (2008, 23) rather than being taken  
393 as a global challenge to its rationality. But overall the topic remains controversial (cf.  
394 Stanford 2017). Since the present paper is concerned with the analysis of experimental  
395 and theoretical practice, we are following Laudan in focusing on the scientific-practical  
396 aspects of underdetermination This focus on scientific practice is also a better basis for  
397 assessing the role of pragmatic and epistemic criteria of theory choice than debates  
398 about the rationality of science globally.  
399

400 Speaking of theory choice, philosophers of science have traditionally set pragmatic  
401 criteria firmly apart from the epistemic criterion of empirical adequacy and all other  
402 scientific questions that can be resolved within an explicitly formulated theoretical  
403 framework (cf. Carnap 1950). Thomas S. Kuhn (1977) rejected this separation and  
404 advocated a broader list of characteristics of a good scientific theory. It includes:  
405 empirical “accuracy, consistency [internally and with respect to other theories], scope,  
406 simplicity, and fruitfulness.” (1977, 322) These five criteria of theory choice are not  
407 mutually independent; they are often context-dependent and may point in opposite  
408 directions. For instance, an increase in accuracy can trivially be obtained by adding  
409 additional parameters; yet scientists may prefer to make do with a smaller number of  
410 fundamental quantities – or with a simpler law – even at the expense of some accuracy.  
411 Thus, scientists have to assess the relative weight of these criteria when deployed  
412 together. Both their form and the relative weight, to Kuhn’s mind, contain contextual  
413 and idiosyncratic (psychological) factors. Kuhn was however at pains to argue that such  
414 subjectivity does not render theory choice irrational or a mere matter of taste. Theory  
415 choice, we might add, was not a major battle in the conflict between historical rationality  
416 and historical contingency waged during the 1970s. Kuhn’s point was the historical and  
417 factual nature of theory choice, not its contingency or arbitrariness. Historians often find  
418 an increasing unanimity of individual choices in a certain field. Such factual unanimity  
419 does not establish rationally binding criteria for theory choice. Instead of being rules of  
420 an algorithm, the criteria of theory choice function “as values, which influence it. ...; they  
421 do specify a great deal: what each scientist must consider in reaching a decision.” (1977,  
422 331)<sup>13</sup>  
423

424 Heather Douglas has proposed a finer-grained account in order to restore the separation  
425 between epistemic and pragmatic cognitive values and reduce conflicts between them.  
426 She distinguishes (i) minimal criteria applied to the theory per se, among them internal

---

<sup>13</sup> Laudan reads Kuhn’s analysis of theory choice against the backdrop of scientific revolutions that represent breaks in rational justification. This rehearses, to Laudan’s (1990) mind, the holistic and egalitarian reading of underdetermination and provides a justification for the sociologizing of epistemology. Without entering into a broader Kuhn debate, it seems to us that once we limit ourselves to an epistemic or local understanding of underdetermination, the Kuhnian analysis of the values of a good scientific theory can still provide important insights into scientific practice. As Kuhn himself has emphasized, these values are only one element of theory choice, alongside sociological factors and inductive reasoning. Moreover, our goal here is not to find all determinants of theory choice, but to focus on the role of the epistemic and pragmatic criteria or values in the preference of models in elementary particle physics.

427 consistency; (ii) minimal criteria applied to the relation of theory and evidence, among  
428 them empirical adequacy; (iii) desiderata applied to theories per se, among them scope,  
429 simplicity, and potential explanatory power of a theory that largely “fall under the rubric  
430 of the fruitfulness of the theory”. (2013, p. 800); (iv) desiderata applied to the relation of  
431 theory and evidence, among them being supported by a broad range of empirical  
432 evidence and not being contrived to match a small domain of facts in an ad hoc fashion.  
433 While the values in categories (i) and (ii) are epistemic, category (iii) contains “strategic  
434 or pragmatic values” (2013, p. 800) that help in “deciding which theory to pursue next”  
435 (2013, 804)). Instead, group (iv) “provides assurance that our scientific claims are more  
436 likely to be reliable.” (2013, p. 800) Moreover: “While simplicity, scope, and explanatory  
437 power are often thought to pull against each other when considering theories alone  
438 (group iii), they pull together when considering a theory in relation to evidence (group  
439 iv).” (2013, 803)

440

441 Perhaps, the most important pragmatic criterion in the history of particle physics is  
442 simplicity. Most influential has been the quest for a simple unified theory of all  
443 fundamental forces.<sup>14</sup> Simplicity also stands behind particle physicists’ long-time  
444 worries about the many parameters that are needed to make the SM empirically  
445 adequate. As Baker (2013) rightly observes, it is quite challenging to pin down the  
446 notion precisely. Many authors distinguish elegance (typically attributed to a theory)  
447 and parsimony (Ockham’s razor that directs us not to introduce unnecessary entities).  
448 Both aspects of simplicity may come into conflict. For instance, the introduction of  
449 supersymmetric partners to all fundamental particles reduces the basic components  
450 into chiral super multiplets, thus reducing the complexity of the theory. The elegance of  
451 an exact symmetry between fermions and bosons in the unbroken theory disappears  
452 once a breaking mechanism is introduced, which leads to a large number of additional  
453 parameters.

454

455 From the interviews and questionnaires, we will analyse in Sect 7.3 and 7.4 how particle  
456 physicists understand and weigh epistemic and pragmatic values and how they assess  
457 the criterion of naturalness in BSM models. Applying the philosophical debate about  
458 those values to model preferences within a variegated model landscape has certain  
459 consequences on how to interpret such preferences further. We are following Kuhn and  
460 Douglas in speaking about values rather than criteria, and will also speak about  
461 preference instead of choice even in cases, such as supersymmetry or not, where the  
462 latter terminology could be appropriate.

463

464

### 465 ***3.3 Making experiments crucial***

466

467 The second classical philosophical problem relevant for the present paper concerns the  
468 interaction between theory (or models) and experiment. LHC’s first task consisted in a  
469 definitive and crucial test of the SM, i.e. to find the Higgs boson or exclude its existence.  
470 Since the Higgs boson is an essential part of the SM and since LHC would cover the  
471 whole energy scale relevant for direct searches, not finding it should have eventually  
472 implied that the SM was refuted. Thus, a large majority of elementary particle physicists  
473 interviewed expressed the conviction that a Higgs discovery or non-discovery at LHC  
474 represented a crucial and decisive test for the SM.

475

---

<sup>14</sup> Note that some philosophers – and some physicists, perhaps – would argue that there are metaphysical reasons or some a priori principle of rationality that imply that a simpler theory is more likely to be true. Such questions are, however, outside the scope of the present paper.

476 This widely shared conviction among physicists prompts the question whether the  
477 Higgs discovery represented a crucial experiment in a philosophical perspective? Let us  
478 take a closer look. The term ‘crucial experiment’ originated with Francis Bacon and  
479 became influential through Newton and his demonstration that sunlight consisted of  
480 rays exhibiting different behaviours. A crucial experiment, in this traditional  
481 understanding, unambiguously and definitively confirms a hypothesis or decides  
482 between rivalling hypotheses. Pierre Duhem objected on the basis of the  
483 underdetermination argument.

484  
485 This philosophical context has made scholars wary about crucial experiments, especially  
486 if they understood underdetermination as a primarily logical and global problem and  
487 followed Duhem in allowing only deductive inferences between theories and data.<sup>15</sup>  
488 While some emphasized that falsifications of a theory were more likely to be crucial  
489 experiments than corroborations, Lakatos famously objected to this asymmetry and  
490 bluntly stated: “No experiment is crucial at the time it is performed (except perhaps  
491 psychologically).” (1974, 320) His main argument was that the assessment of each  
492 experiment can only be performed against the backdrop of the entire research program  
493 it is embedded into and against its competitors. Thus, designating an experiment as  
494 crucial is partly a historical assessment.

495  
496 The idea that a crucial experiment is embedded into a broader program is also the core  
497 of a recent debate about crucial experiments in biology. Weber (2009) defends the  
498 characterization of an experiment as crucial, not within the traditional contexts of  
499 deductive reasoning and the refutation of alternative hypotheses, but by developing “an  
500 experimentalist version of inference to the best explanation.” (2009, 21) Hypotheses are  
501 not refuted, but positively selected as those best supported by the evidence. Weber’s  
502 strategy to defend the Meselson-Stahl experiment as crucial is now to show that both  
503 parts of Duhem’s problem, the problem of untested auxiliaries and the problem of an  
504 exhaustive partition of theoretical alternatives (including the unconceived ones)<sup>16</sup>, can  
505 be kept at bay. To this end he develops a holistic account of experimental mechanism  
506 that includes both a model of the mechanisms producing the phenomena and parts of  
507 the experimental system, among them “the characteristic manipulations and  
508 measurement devices used.” (2009, 34) Baetu (2017) has criticised Weber’s  
509 reconstruction and argued that the Meselson-Stahl experiment was inconclusive for the  
510 hypotheses considered. Instead, “it was part of a broader research project aiming to  
511 elucidate the mechanisms of DNA replication” (2017, 4.) – which ultimately led to the  
512 development of new experimental techniques. “Thus understood, the experiment  
513 extended over a decade or more. However, the crucial experiment account attributes all  
514 or most of the impact of the whole series of experiments to a single set of experimental  
515 results.” (2017) In the same vein as Lakatos put it, an experiment becomes crucial only  
516 in historical reconstruction and within the context of a broader research program.

517  
518 We believe that the Weber-Baetu debate rightly follows the trend diagnosed in Sect. 3.2.  
519 to view underdetermination and crucial experiments as an epistemic and factual  
520 problem rather than a logical and semantic one. In this way, the first aspect of Duhem’s  
521 problem, the auxiliary hypotheses, becomes embedded into an experimental research

---

<sup>15</sup> Note that Duhem actually believed that experiments could be crucial. But this could not be inductively inferred from the data, but required the *bon sens* of the physicists. While *bon sens* might have been a useful notion in Duhem’s days, it seems to us too vague for large-scale experiments in particle physics. At best one might take *bon sense* as an umbrella term for the detailed set of experimental strategies given by Franklin (2013).

<sup>16</sup> Cf. Stanford (2006) who shows that it is difficult to find cases where the underdetermination was not eventually resolved.

522 program. The reference to turn to in the present context is of course Franklin's (2013)  
523 philosophical reconstruction of the history of modern particle physics. There Franklin  
524 distils a list of reliable strategies that in effect allow one to keep Duhem's problem at bay  
525 and address the related problem of theory-ladenness of large-scale particle experiments.  
526 Beauchemin's (2017) autopsy of measurements with the ATLAS detector<sup>17</sup> can be read  
527 as a continuation, into the days of LHC, of Franklin's (2013) history of the reliable  
528 experimental strategies and rules of data analysis that characterize contemporary  
529 elementary particle physics. We will take up some of these strategies in Section 7.2. and  
530 discuss how they permit us to consider the Higgs discovery as a crucial experiment. Let  
531 us however first assess Franklin's assessment of crucial experiments.

532  
533 Franklin and Perovic (2015) compare two ground-breaking particle physics  
534 experiments. While they classify the discovery of parity violation as a crucial  
535 experiment, the discovery of CP-violation represented only a 'persuasive experiment'.  
536 "The difference lies in the length and complexity of the derivation linking the hypothesis  
537 to the experimental result, or to the number of auxiliary hypotheses required for the  
538 derivation." (2015, 85) Indeed, physicists had speculated about parity violation before,  
539 and the observed effect was maximal. CP-violation was completely unexpected, but most  
540 theoreticians quickly settled for it. Franklin and Perovic consider this acceptance as a  
541 "pragmatic solution of the Duhem-Quine problem." (2015, 84). In the case of the Stern-  
542 Gerlach experiment, as reconstructed by Franklin and Perovic, the diagnosis of cruciality  
543 underwent several changes. By discovering the space quantization  
544 [*Richtungsquantelung*] predicted by the Bohr-Sommerfeld quantum theory, it became a  
545 crucial watershed between classical and quantum physics, but not by confirming the  
546 latter theory. For what Stern and Gerlach actually measured was a new quantum  
547 phenomenon, electron spin, that was only postulated after the experiment. Thus, the  
548 experiment "was regarded as crucial at the time it was performed, but, in fact, wasn't. ...  
549 A new theory [quantum mechanics] was proposed and although the Stern-Gerlach result  
550 initially also posed problems for the new theory, after a modification of that new theory  
551 [the integration of spin], the result confirmed it. In a sense, it was crucial after all. It just  
552 took some time." (2015, 40-41)<sup>18</sup>

553  
554 These examples also indicate that establishing experimental evidence and deciding  
555 whether an experiment is conclusive or even crucial, is largely a factual question and  
556 involves different time scales. Acquiring precision data sometimes represents a long-  
557 term process that involves previous experiments and is continued in the experiment  
558 itself. The actual discovery of a particle instead represents a precisely dated event;  
559 scientists decide after a detailed statistical analysis that the evidence is sufficient.

560  
561 Using the Higgs discovery, in 7.2 we will argue that the diagnosis of Franklin and Perovic  
562 seems to us counterintuitive because it makes the characterization of an experiment as  
563 crucial or not depend on short-term development of scientific theorizing. In Section 7.2.,  
564 we will provide a different characterization according to which all three examples  
565 mentioned qualify as crucial experiments.

566  
567 **4. The methods of this project**

---

<sup>17</sup> Note that Beauchemin's concept of theory-ladenness is wider than the one typically used in the philosophical literature, where theory-ladenness represents a problem for empirical science, not a feature that can be exploited by clever experimenters.

<sup>18</sup> In philosophical discussions about quantum mechanics, spin is considered as the quantum mechanical quantity par excellence and the Stern-Gerlach apparatus as its paradigmatic experiment. Notice that while Stern wanted to test quantum theory, Gerlach himself considered the experiment as part of a broader experimental research program.

568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621

Against the backdrop of the different experimental situations in 2011 and 2012, and the various solutions of the EWSB (including the Higgs mechanism) proposed by theoretical model builders, our project investigated the general attitudes and preferences of the LHC physicists by quantitative and qualitative empirical methods. In questionnaires and interviews LHC physicists were asked about their views of the status of particle physics, their anticipation of what the LHC will ultimately find, and the ways experimentalists and theorist interact.

Questionnaires were sent via e-mail to some 15000 physicists related to particle physics in August 2011 and September 2012. Each contained eight groups of questions, which were to be answered by either assigning a subjective probability for the correctness of a certain statement or by choosing an answer among various options. These were (i) the probability to find the SM Higgs particle (respectively confirm a minimal SM Higgs), (ii) the possible explanations of new physics found at LHC, (iii) the preference for certain BSM models independently of the LHC results, (iv) the criteria guiding the researcher's answers to this question, (v) the most critical flaws to the SM, (vi) the signatures in which LHC would most likely find new physics, (vii) general features of particle physics for whose understanding LHC will be most important, (viii) the interaction between experimentalists and theoreticians.

A large fraction of the questions within the above-mentioned groups were identical for the two periods, however, from experience with the first one, modifications were made for the second questionnaire. In the first questionnaire some answers could be ranked up to four times. This was considered less meaningful for the second questionnaire and modified. In 2012, a question was also added to address the Higgs boson candidate. The precise list of the questions can be found in appendix 2.

The lists of physicists to which the questionnaires were sent both in 2011 and 2012 were obtained from the INSPIRE data base (Dallmeier-Tiessen, S., Hecker, B., Holtkamp, A; 2016) maintained centrally at CERN. This data base is established by surveying journals, conferences, books, theses etc. in the pertinent fields and listing all authors. For the purpose of the questionnaire, authors in the categories 'hep-ph' (phenomenology), 'hep-th' (theory), 'hep-ex' (experiment) were contacted. In total this amounted to some 15000 authors. About half of the authors are theorists belonging to about the same amount to either the 'th' or 'ph' category, the other half experimentalists. Taking into account that some 8000 experimental physicists are directly involved in the LHC experiments, with an additional number of several thousand theorists, the list of physicists included probably almost all those who are actively working on LHC physics. Certainly, some physicists on the list were somewhat remote from LHC experiments or theory, e.g. mathematical theorists or accelerator physicists, but also some retired physicists or those who had left the field. It is difficult to assess, how large a fraction this was.

The anonymous replies were collected and statistically evaluated at Wuppertal. There were 1435, respectively 903 replies to the two questionnaires, which corresponds to a return rate of 10%, respectively 6% which is acceptable for empirical studies that are combined with interviews. Our goal had not been to obtain a truly representative sample in the sense of quantitative sociology. Still, there seems to be no strong bias in our replies: the regional distribution of respondents is consistent with the regional distribution of physicists working in LHC experiments, and also the fraction of theorists and experimentalists agrees with the fraction in the list. Yet, there are discrepancies as regards seniority: only few PhD students (<5% of the replies) have answered the questionnaires, whereas they amount to about a third in the LHC experiments. In the

622 following, the replies were considered separately for experimentalists and theorists  
623 because this promised some interesting insights. In addition, the comparison of the  
624 replies before and after the discovery should indicate certain trends in the thinking of  
625 the LHC physicists.

626  
627 In addition to the questionnaires, 9 (6) LHC physicists were interviewed around April  
628 2011 (September 2012). Both groups included experimentalists from different LHC  
629 experiments and theorists. Furthermore, it was attempted to cover a wide range of  
630 interests and responsibilities within the LHC project. There is only a small overlap  
631 between the physicists in the two rounds; this was done deliberately in order to obtain a  
632 broader picture. The physicists interviewed and their respective roles at the time of the  
633 interviews are listed in Appendix 2. In the following discussion, no names will be  
634 assigned to the respective citations.

635  
636 Each interview took about an hour. A few topics were addressed in every interview, for  
637 instance: in 2011, the prospects of a Higgs discovery, the perceived status of super-  
638 symmetry, and the chances to find new physics; in 2012, the impact of the Higgs  
639 discovery on the interviewee's research. On the other hand, the interviews were kept  
640 flexible to better understand the reasoning and preferences of each interviewee. This  
641 included, depending on the answers of the counterpart, also questions about the work  
642 environment, the methods of research, which outcome is expected at the LHC and why,  
643 and which outcome would be preferred on theoretical or pragmatic-aesthetic grounds.

644  
645

## 646 ***5. The physicists' expectations in autumn 2011***

647

648 At the beginning of our empirical study, the physical situation was characterized by an  
649 excellent performance of the LHC and its experiments. The year 2011 brought an  
650 unexpectedly large amount of data at the energy of 7 TeV. Based on this understanding,  
651 the LHC physicists performed simulation studies predicting that the whole range  
652 pertinent to the mass of a Higgs boson could be covered at the LHC within two years. On  
653 the other hand, although a broad range of searches for new effects had been performed  
654 by fall 2011, no sign for any of the many postulated extensions of the SM had been  
655 found. In particular, no indication for Supersymmetry was observed. Supersymmetry  
656 had been highly favoured by theorists, and it was predicted that its particles could be  
657 detected shortly after the LHC launch. Supersymmetry is the only BSM model that  
658 provides a strict constraint on the highest allowed mass of the Higgs boson (of about  
659 130 GeV). The sensitivity of many of these searches for new physics reached the energy  
660 scale of about 1 TeV, at which the naturalness problem should have been resolved  
661 before the corrections become too high.

662

### 663 ***5.1 Outcome of questionnaires***

664

665 In total 1435 physicists answered the questions, with the number of theorists (769) and  
666 experimentalists (696) being about the same. The number of replies to each of the  
667 questions differed only by a small amount. Assuming multinomial distributions and an  
668 outcome for an answer of 50%, these numbers imply a typical error margin on the  
669 answers of 1.5% for the total sample and 2% for each subgroup. The precise uncertainty  
670 depends on the number of answers given; the fewer there are, the larger is the relative  
671 uncertainty. Where relevant, the exact uncertainties will be provided.

672

673


#### 674 ***5.1.1 The importance of the origin of mass***

675

676 The high expectations that physicists had in the LHC to understand the mechanism of  
 677 mass generation become most apparent in the replies to a question about the  
 678 importance of LHC results for several key problems of current physics. Participants  
 679 were asked whether they fully agreed, somewhat agreed, were undecided, somewhat  
 680 disagreed, or fully disagreed with the statement: '*LHC results will be very important to*  
 681 *understand ...*'. Close to 50% (48%/49% of the theorists/experimentalists) chose to  
 682 'fully agree', and close to 80% (77%/80%) at least 'somewhat agreed' on the importance  
 683 of the LHC for the 'origin of mass'. Comparable results were obtained for two other  
 684 topics from the SM, 'strong interactions' and 'flavour physics', while the outcomes for  
 685 BSM physics were much lower. The as of then only undiscovered element of the SM was  
 686 accordingly given the highest priority among all the potential features that could be  
 687 found at the LHC.

688  
 689  
 690  
 691

### 5.1.2 Expectation on finding the Higgs Boson at LHC




692  
 693  
 694  
 695  
 696  
 697

Fig.1 Percentage of answers of theorists (blue) and experimentalists (red) assigning probabilities in intervals of 20% on the chance that the LHC will find a Standard Model Higgs Boson (Questionnaire of 2011)

698  
 699  
 700  
 701  
 702  
 703  
 704  
 705  
 706  
 707  
 708  
 709  
 710  
 711  
 712  
 713  
 714

Given the importance of the origin of mass and the fact that the LHC was expected to provide the ultimate sensitivity for finding the SM Higgs boson the questionnaire asked physicists: '*What is your personal estimate of the probability [that] the LHC will find the Standard Model Higgs boson?*'. This (subjective) probability was to be given in terms of percentage intervals of 20%, which represented the respondent's current degree of belief. The replies did not reveal any strong tendency towards either discovery or non-discovery, but instead were rather uniformly distributed over all probability values (see Fig. 1). Some 35% (34% of theorists/35% of experimentalists) assigned a chance of at most 40% that the SM Higgs boson will be discovered, whereas only a few more (41%/36%) expected it to be found with 60% probability or more; the values for more than 80% probability were even lower (22%/15%). Thus, although simulation results showed that the LHC, in virtue of its foreseeable performance, had the potential to find the SM Higgs boson if it at all existed, a large fraction of LHC physicists assumed that it would not be found. These assessments were largely identical for experimentalists and theorists.

715 A second question addressed the ‘*personal estimate of the probability ... that the LHC will*  
 716 *rule out the Standard Model Higgs boson*’. In this case 59%/46% of the  
 717 theorists/experimentalists considered the probability low (i.e. smaller than 40%). On  
 718 the other hand, only 19%/26% (uncertainty about 2.5%) estimated that the SM Higgs  
 719 boson could be ruled out with high probability (i.e. larger than 60%). Low probability  
 720 here means either that the SM Higgs boson will eventually be found or that a candidate  
 721 is found whose properties cannot be measured precisely enough to rule out other  
 722 interpretations. High probability instead means that the LHC will be able to definitively  
 723 rule out the SM Higgs particle because there is no such particle or it will find one or  
 724 more candidates that accomplish mass generation with properties different from the SM  
 725 expectations. The responses showed that, in 2011, theorists were more sceptical about  
 726 the LHC to rule out the SM Higgs boson than experimentalists.  
 727  
 728  
 729  
 730



731  
 732  
 733 *Fig.2 Percentage of answers of theorists (blue) and experimentalists (red) assigning probabilities in*  
 734 *intervals of 20% on the chance that the LHC will rule out a Standard Model Higgs Boson*  
 735 *(Questionnaire of 2011)*  
 736

737  
 738 Although the two last questions are closely connected, there are subtle differences that  
 739 lead to somewhat different replies. Firstly, the fraction of physicists that assigned an at  
 740 least 60% probability to *find* the SM Higgs boson is smaller than the fraction of  
 741 physicists who assigned an at least 60% chance that it will *not be ruled out*. Secondly,  
 742 whereas the answers of theorists and experimentalists were rather consistent with the  
 743 first question, a significantly larger portion of theorists than experimentalists  
 744 considered it unlikely that the SM boson will be ruled out.  
 745

746 The first difference is probably related to the much stricter requirement to confirm not  
 747 only the existence of a new particle, but to determine all of its properties to a precision  
 748 that allows one, e.g., to distinguish it from alternative models of EWSB. Especially in the  
 749 case of a more complicated Higgs group structure (as favoured by many physicists – see  
 750 below) it will be more difficult to unambiguously identify the particle to be a SM Higgs  
 751 boson than to rule it out. How-to interpret the differences between experimentalists and  
 752 theorists is more difficult. In general, the replies – and the interviews below – indicate a  
 753 greater reluctance of experimentalists to commit themselves to what their data will  
 754 finally reveal.




755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781

This exemplifies the first lesson from the 2011 questionnaire. In contrast to public perception, interpreting the newly found particle as being the SM Higgs boson was not a simple yes/no alternative to be decided promptly. Physicists were largely prepared for a more complicated outcome that achieved all that the SM Higgs mechanism was designed for. Thus, finding a particle consistent with a SM Higgs would only be the first step in further investigating the properties of the new particle. The second conclusion from these two questions is that there existed a substantial scepticism among physicists as to the existence of a SM Higgs at this stage. This means that, although the LHC was expected to cover the whole allowed mass range for the SM Higgs particle, the LHC community was rather undecided if it exists. Taking both lessons together shows that there was no significant asymmetry in physicists' expectations between refuting and confirming the SM.

### 5.1.3 Expectations on various EWSB models

The questionnaire also addressed potential scenarios for 'new physics', i.e. a process or particle that is not part of the SM. Physicists were asked 'Assuming that the LHC finds new physics, which (if any) of the following models do you think has the best chance of explaining it'. The physicists had two ranked choices; here we will typically just provide the first choice, the second gives fairly similar results. Several models, including those rather remote from EWSB, like string theory, extra spatial dimensions, or 4<sup>th</sup> generation models, were also considered. (cf. Appendix 2) In the following, we focus on the three most popular groups that were also those most closely related to EWSB. (Fig.3)



782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794

Fig.3 Percentage of answers of theorists (blue) and experimentalists (red) on the most probable model that the LHC might find. Only answers with relation to the electroweak symmetry breaking are given, the remaining 51/58% refer to different models (Questionnaire of 2011)

Fractions of 10/11% of theorists/experimentalists opted for an extended Higgs sector, i.e., more than one Higgs boson. About twice as many (23%/24%) voted for the favourite theory of Supersymmetry, which also requires an extended Higgs sector. Therefore, about one third of physicists were expecting new physics in extended Higgs sectors either without or within the context of an explicit model. Both of these answers assume Higgs bosons and expect them to be elementary as the SM Higgs. As mentioned

795 above, one of the Higgs bosons in the extended sector might have properties very  
796 similar to the SM Higgs.


797

798 In addition, a sizeable fraction of theorists ( $16\pm 1.3\%$ ) expected a dynamically generated  
799 electroweak symmetry breaking, leading to a composite scalar particle with several  
800 properties that are distinctively different from the SM Higgs boson. As discussed in Sect.  
801 2.3., at least the historically first such model, Technicolour – a model that also contained  
802 the least additional assumptions –, had been strongly disfavoured by data. This might be  
803 the reason why only  $7(\pm 1)\%$  of the experimentalists chose this option.

804

805 The follow-up question (Fig. 4) was ‘*which preference*’ [physicists] have ‘*independently of*  
806 *the expectations regarding LHC results*’, i.e., irrespective of the sensitivity of LHC itself.  
807 Whereas the replies alluding to an extended sector remained rather the same as to the  
808 previous question, dynamical EWSB was now even more favoured by theorists (19%),  
809 while the fraction of experimentalists preferring this alternative was unchanged at 7%.<sup>19</sup>

810



811

812

813 Fig.4 Percentage of answers of theorists (blue) and experimentalists (red) on the preferred model  
814 for physics beyond the Standard Model. Only answers with relation to the electroweak symmetry  
815 breaking are given, the remaining 50/58% refer to different models (Questionnaire of 2011).

816

817 The difference between the answers as to the importance of dynamical EWSB as a vision  
818 for the LHC data, on the one hand, and in a general perspective, on the other, shows that  
819 physicists’ preferences are also guided by nonfactual and non-epistemic aspects. The  
820 perceived ‘beauty’ of a theoretical framework, or other pragmatic values of theory  
821 preference, weigh significantly relative to the chances of confirmation or  
822 disconfirmation by soon-to-be-available experimental data.

823

824

825


#### 826 5.1.4 The importance of the naturalness criterion

827

828 The question as to the preferred model was followed by the question ‘*which (if any) of*  
829 *the following criteria have guided you in answering the previous question?*’. Four ranked

<sup>19</sup> Although the present paper focuses on EWSB, it is worth noting that the largest percentages for new physics are obtained for the option ‘None of those, but something totally unexpected’ (28% for both theoreticians and experimentalists). The perspectives for new physics will be discussed in a separate paper that will also analyse some of the models neglected here.

830 choices were allowed (Fig. 5). Considering only the first choice, the criterion that a  
 831 model ‘solves the naturalness problem’ was preferred by 21%/17%. It is thus  
 832 considered as important as the classical pragmatic, or rather aesthetic, criteria of  
 833 ‘elegance’ (22%/18%) and ‘simplicity’ (16%/20%)<sup>20</sup>. More ‘factual’ criteria like the  
 834 model ‘will provide a better fit to the data’, or ‘makes specific predictions’ or even ‘has a  
 835 candidate for dark matter’ are much less considered (each one below 10% for both  
 836 experimentalists and theorists).  
 837  
 838  
 839




840  
 841  
 842 *Fig.5 Percentage of answers of theorists (blue) and experimentalists (red) on the criteria to choose*  
 843 *the preferred model. (Questionnaire of 2011).*  
 844

845 Physicists were further asked ‘*what (if any) are the most critical flaws of the Standard*  
 846 *Model*’, and could make up to three unranked choices (Fig. 6). Indeed the problem of  
 847 ‘*quadratic divergences in corrections to the Higgs mass*’, causing the naturalness problem,  
 848 was mentioned often by theorists (14%), while only by 10% of the experimentalists  
 849 (statistical uncertainty of difference 2.2%). However, quadratic divergences are  
 850 considered less of a flaw of the SM than its many parameters (18%/19%), the absence of  
 851 gravity within the Standard Model framework (18%/21%), or that it does not include a  
 852 Dark Matter candidate (17%/17%).  
 853

---

<sup>20</sup> It should be noted that physicists were not given any specific definition of these concepts; hence the replies were based on the intuition of the individual physicist. We do not consider this as too problematic for our purpose, not least because many philosophical authors who provide a definition - cf. Baker 2013 discussed in Section 3.2 - simultaneously emphasize that the terminology often is all over the place.



854  
855  
856  
857  
858

*Fig.6 Percentage of answers of theorists (blue) and experimentalists (red) on most critical flaw of the Standard Model (Questionnaire of 2011). The three answers were summed up and normalized.*

859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873

Both of these questions refer to SM properties that point beyond its limits. But they do so from a somewhat different perspective. The first (Fig. 5) had asked for the motivations of model preferences in Fig. 4, that is irrespective of the chances to soon find solutions at LHC. The second (Fig. 6) asked for flaws of the SM, irrespective of the existence of a credible model or strategy to resolve them. In both questions naturalness, respectively quadratic divergences, scored within the top group of the list and matched their counterparts, the pragmatic values of preference simplicity and elegance, and respectively (among the flaws), the many parameters of the SM.<sup>21</sup> However, the differences in the relative weights for other elements pointing BSM, e.g. for Dark Matter, are significant. To our mind, this has to do with the different perspectives of the questions. There are several proposals of physics BSM, however no universally agreed upon Dark Matter candidate. Nevertheless, it is a significant flaw of the SM, even though it may be only solved in the longer term. Naturalness, instead, is of immediate relevance to problems of model builders and guides expectations for BSM at the LHC. Experimentalists and theorists largely agree in this attitude.

874  
875

## 5.2 Responses in interviews

876  
877  
878  
879  
880  
881  
882  
883  
884

The questionnaires were complemented by interviews with nine theorists and experimentalists<sup>22</sup>. Overall, their statements were consistent with the outcomes of the questionnaire. Yet they provide a deeper insight into the reasoning of elementary particle physicists at the time. In particular, they illustrate the rather diverse set of attitudes and the broad variety of expectations among the physicists. The following selected quotes are related to the mechanism of EWSB.

<sup>21</sup> The “many parameters” of the SM (Fig. 6) are traditionally seen as a principal lack of simplicity and motivate physicists to devise other models. In the same vein, the “quadratic corrections to the Higgs mass” (Fig. 6) amount to a peculiar technical feature in the renormalization scheme for the scalar Higgs boson that motivates models “solving the naturalness problem” (Fig. 5). In Section 5.2.3. we will provide some evidence from interviews that notwithstanding the philosophical distinctions discussed in Section 3.1., naturalness is largely treated in the same fashion throughout the community of particle physicists.

<sup>22</sup> For the list of names, see the table in Appendix 1.

885 **5.2.1 Crucial, Long-Awaited, but Uncertain: Does the Higgs boson exist?**

886

887 The interviews were conducted at a time, when the allowed mass range for the SM Higgs  
888 boson was rapidly shrinking and the experiments were close to completely covering the  
889 remaining parameter space. No wonder that in all interviews the suspense whether the  
890 Higgs boson would be discovered or some alternative mechanism of EWSB would  
891 become visible, played a pivotal role. Here are two typical examples. One physicist  
892 stated that a discovery of the Higgs boson would amount *'to a revolution ... We*  
893 *understand the mass, we understand a lot of things'*. Another one assessed the *'Higgs*  
894 *problem'* as a *'key question'*. The measurement of its mass should *,be a very important*  
895 *clue to what sort of theory maybe goes beyond it'*.

896

897 Although accordingly an experimental verdict, a crucial and long-awaited test for the  
898 SM, was in sight, opinions diverged on what its outcome would ultimately be. In this  
899 respect, the answers span a broad range. At one end of the spectrum, an experimentalist  
900 argued that in this situation one should *'press theorists'* to answer the question: *'if there*  
901 *is no such thing [the Higgs boson], then what?'*. Being a few femtobarns away from the  
902 final call about the Higgs boson, this represented the mood of some physicists that one  
903 had to move to a *'provocative question'*. The interviewee even identified a *,change of*  
904 *mind-set'* because the - to date unsuccessful - experimental searches led to a general  
905 doubt whether the Higgs was a *,done deal''*.

906

907 Other interviewees emphasized the personal and even emotional aspects of this  
908 increasingly pressing insecurity. E.g. *'I don't know, I don't know': 'we have been waiting so*  
909 *long for this, .... there is ... no concrete criterion to really judge whether [it] is more or less*  
910 *likely and emotionally, needless to say, I would like to see that as soon as possible, so I hope*  
911 *it's more likely that it comes out, but it is purely emotional because I do not want to wait*  
912 *another five years, but I have no idea.'* Another interviewee diagnosed a change of  
913 attitude. Previously colleagues might have argued that it *'is much more exciting to see*  
914 *nothing. But it was before LHC started. Now that things work so well, people are sure that*  
915 *the Higgs will be found in 2012, 2013, public opinion you know changed dramatically'*. It is  
916 *'psychologically very interesting'*. This strong desire to 'find something' also reflected the  
917 increasing gap between the enormous success of the SM predictions during the past 40  
918 years and the fact that quite a few still unsolved questions remained.

919

920 Other experimental interviewees were rather optimistic to find the Higgs boson. One  
921 stated *'I would be more surprised if they don't find the Standard Model Higgs because I*  
922 *think that the Standard Model with the Higgs mechanism at the moment is one of the best*  
923 *ways of explaining the masses of the gauge bosons and particle masses, so I would really*  
924 *like the Standard Model like Higgs'*, or more pronounced *'I will be surprised if it is not*  
925 *found. I think it will be found at 120 GeV.'* (This was in accordance with the indirect and  
926 direct limits at that time.)

927

928 The attitude, according to which the Higgs is 'the best way' to explain masses had  
929 already been strengthened by indirect measurements disfavouring otherwise preferred  
930 alternative models. It also becomes apparent in replies from a theorist, who held that  
931 *'there is a lot of circumstantial evidence in favour of that [i.e. the Higgs boson], the case is*  
932 *not proved but that might well happen'*. This factual statement, however, is immediately  
933 put into perspective, when the same interviewee points to his preference *'I would find a*  
934 *lot of intellectual attraction in the dynamical symmetry breaking models'*.

935

936 **5.2.2 Is the Higgs mechanism attractive?**

937

938 While the expectation as to the possible discovery of the Higgs boson was an issue of

939 considerable suspense at that time, the motivations to expect or reject the Higgs  
940 mechanism, differ among the physicists. Already in the above statements, it became  
941 apparent that the mechanism, even if expected, was regarded with some reservations.  
942

943 There were several values of theory preference in play. One theorist held that the Higgs  
944 mechanism is also disfavoured *,because of its minimal predictive power'*. Instead the  
945 alternative scenario of strong EWSB is intellectually favoured. *'Whereas the dream would  
946 be, in a dynamical theory of electroweak-symmetry breaking ..., there is at least a  
947 conceivable possibility of making definite predictions, however they should turn out.'* This  
948 requires a *'somewhat bigger theory'*. The scepticism is also shared by experimentalists.  
949 One stated drastically, *'the Higgs is a totally ad hoc thing. .... If the Higgs is not there, it  
950 will not surprise me.'* He argued that *'people had faith in it the way people have faith in  
951 God'*. Another experimentalist held, that even if the *'Higgs will be found, ... the Higgs  
952 mechanism seems not elegant'* and there are *'very attractive theories without the Higgs'*.  
953

954 In fact, only a few physicists interviewed emphasized the broader virtues of the Higgs  
955 boson beyond merely giving a solution for one specific problem. One theorist focussed  
956 on its role in the more encompassing theory of Supersymmetry, which as mentioned  
957 gives an upper bound of the Higgs mass. It would be *'very disappointing to find Higgs at a  
958 mass compatible with SM at high energies [above 130 GeV]'*. *'If the Higgs boson is light as  
959 suggested by the data [i.e. indirect measurements and left-over phase space masses], then  
960 presumably ..... super-symmetry is a prototype of such a weakly interacting extension of the  
961 standard Higgs mode'*, i.e. the virtue of the Higgs mechanism is its accordance with a  
962 larger and generally favoured theory. On the other hand, the interviewee points out, that  
963 the favoured variant of supersymmetry, has been so tightly tested without finding  
964 anything. Therefore, one finds oneself in *'a weird situation '*.  
965

966 Moreover, one theorist rejected the statement that the Higgs mechanism is complicated  
967 and ad hoc, but emphasised the virtue of introducing spontaneous symmetry breaking  
968 into particle physics in general. The Higgs discovery would then be seen as something  
969 new *'in the sense of new particles, .... but it is a break-through since you have the  
970 experimental test of spontaneous symmetry breaking'*. Moreover, this general idea would  
971 be *'not immediately thrown away'*, if the Higgs boson would not be found.  
972

### 973 **5.2.3 The problem of naturalness**

974

975 Several interviews stressed the value of naturalness as a pragmatic guideline. Yet as  
976 regards its aesthetic aspects, opinions differed. One theorist became a proponent of  
977 Supersymmetry once it was shown to solve the naturalness problem; *'so for me the big,  
978 sort of change in my world view came when people pointed out that super-symmetrical  
979 particles could potentially control the quantum corrections and make the theory more  
980 manageable.'* Another theorist, who was asked whether the naturalness or the hierarchy  
981 problem were serious, answered more cautiously: *'now to assess this, one goes back to  
982 these convictions somehow that the progress of science is always driven by an aesthetic  
983 judgement .... that goes beyond mechanical relations between formulas, equations and the  
984 need to see beyond. In other words, when you see some recurrence, when you see some  
985 "accident", it is natural for a scientist to consider the possibility that it is not an accident  
986 but there is something beyond and then this accident becomes natural. Now, this is not  
987 always correct, there are many accidents that we witness around that are not driven by the  
988 first principles but just accidents. So in that respect one can be wrong, but for the issue of  
989 naturalness, all of it, so called problems of the standard model, the picture is quite  
990 compelling.'* Only one experimentalist was explicitly asked about naturalness. Again,  
991 there was no strong commitment, but instead it was *'take[n] easy, it is a matter of taste'*.  
992 Note that the interviewees did not distinguish between naturalness, fine tuning of the



993 quantum corrections, and quadratic divergences.

994

995

## 996 **6. The physicists' response to the discovery of a Higgs candidate**

997

998 A year after the first questionnaire was sent out and the interviews had been performed,

999 a sufficient amount of data was collected by the LHC experiments to provide the desired

1000 sensitivity for a SM Higgs boson in almost the whole remaining mass range. Indeed, in

1001 July 2012, the observation of a new boson was announced. Since the signatures were

1002 consistent with the expectations, there was a very broad consensus that this particle

1003 was a very strong candidate for a SM Higgs. However, the properties known by then

1004 were few, and the precision of the measurements was still marginal<sup>23</sup>. The simultaneous

1005 searches for new effects BSM remained inconclusive, although the mass reach and

1006 sensitivity was extended.

1007

1008 Shortly after the announcement, a new questionnaire was sent to the same mailing list

1009 and a new round of interviews was performed. One of the main aims was to understand

1010 if, respectively how, the views and expectations of physicists had changed.

1011

### 1012 **6.1 Outcome of the Questionnaire in 2012**

1013

1014 The second questionnaire was sent out in September 2012. To a large part, the

1015 questions were identical to those of the first questionnaire. In this survey 903 physicists

1016 replied, among them 464 theorists and 439 experimentalists. The typical statistical

1017 uncertainty in the replies is therefore 1.7% for the whole sample and 2.4% for each of

1018 the subsamples. The relative uncertainty of the answers between the two

1019 questionnaires depends on whether the same or different physicists replied. In the

1020 former case the relative uncertainty would be very small, in the latter case some 2.3%

1021 (for the whole sample). Since the answers were given anonymously, there was no way to

1022 tell. Compared to the first round the possibility of ranked answers was restricted to a

1023 single choice only or to unranked options. The subjective degrees of belief in a statement

1024 (cf. 4.1.1.) were rephrased as 'fully agree', 'somewhat agree', etc.

1025

1026

#### 1027 **6.1.1 What is the new particle?**

1028

1029 Reacting to the discovery of the new particle, a set of questions was directed at its likely

1030 significance for the SM and beyond. The first statement to be evaluated was *After the*

1031 *discovery of the new particle at 125 GeV, the LHC will confirm the minimal Higgs sector*.

1032 The majority of both experimentalists and theorists (63%/63%) fully or somewhat

1033 agreed with this statement (see Fig. 7). Compared to the first questionnaire<sup>24</sup>, this is, not

1034 surprisingly, a significant increase from the 41%/36%. Still, only 19%/23% 'fully


1035 agreed' that LHC will confirm the minimal Higgs sector of the SM.

1036

---

<sup>23</sup> It was only half a year later, after more data became available and more studies had been made that the particle lost its status of being a candidate and was indeed considered a Higgs boson. This became apparent in a CERN press release (CERN press office, 2013) in which recent results were summarized.


<sup>24</sup> As a reminder, the exact wording of the first questionnaire was *'What is your personal estimate ... that the LHC will find the Standard Model Higgs boson'*.



1037  
 1038  
 1039  
 1040  
 1041  
 1042  
 1043  
 1044  
 1045  
 1046  
 1047  
 1048  
 1049  
 1050  
 1051  
 1052  
 1053  
 1054  
 1055  
 1056  
 1057

Fig.7 Percentage of answers of theorists (blue) and experimentalists (red) assigning probabilities in intervals of 20% on the chance that the LHC will confirm a minimal Model Higgs Boson (Questionnaire of 2012)

This question combined two aspects: whether the observed particle was indeed a Higgs boson and whether it would remain the only Higgs boson, i.e. the SM Higgs boson. This ambiguity could be somewhat resolved by asking a second question, to wit, whether the LHC will 'find a more complicated Higgs sector'; for the new particle could be one of many Higgs bosons (Fig. 8). A sizeable fraction of 30%/31% (theorist/experimentalists) expected this to be the case. This is almost the remainder of those who fully or somewhat agreed with the first statement. However, almost half (46%/47%) of the responses were 'undecided', consistent for theorists and experimentalists. As in 2011 a more complicated Higgs sector appeared to be a very attractive option for many physicists. One may speculate about the reason for this rather neutral opinion. Certainly, the data were too scarce at the time of the questionnaire; moreover, the physicists may have been considering the probably limited precision of the LHC measurements.




1058  
 1059  
 1060  
 1061  
 1062  
 1063

Fig.8 Percentage of answers of theorists (blue) and experimentalists (red) assigning probabilities in intervals of 20% on the chance that the LHC will find a more complicated Higgs sector (Questionnaire of 2012)



1064 The third question asked if the LHC will *'find an alternative mechanism of EWSB'* (Fig. 9).  
 1065 It was only a minority of roughly 10% that agreed at least 'somewhat' with this  
 1066 statement. Full agreement was at the 1% level. Given the discovery of a Higgs candidate  
 1067 shortly before the questionnaire, such a result is not surprising. Even though the LHC  
 1068 data available at that stage were still marginal, the consistency with what is expected for  
 1069 a SM Higgs boson disfavoured a radically different mechanism already then. Although a  
 1070 third of the replies were undecided, the vast majority of physicists no longer expected  
 1071 any radically new physics to emerge in the sector of mass generation.  
 1072




1073  
 1074  
 1075 *Fig.9 Percentage of answers of theorists (blue) and experimentalists (red) assigning probabilities in*  
 1076 *intervals of 20% on the chance that the LHC will find an alternative mechanism of EWSB*  
 1077 *(Questionnaire of 2012)*

### 1080 **6.1.2 Which alternatives to the SM Higgs are still considered?**

1081  
 1082 Whereas the new particle was largely considered to be a Higgs boson, its discovery  
 1083 initially did not preclude it to be, or involve, an element of new physics. Hence physicists  
 1084 were again asked: *'Assuming that the LHC finds new physics, which (if any) of the*  
 1085 *following models do you think has the best chance of explaining it?'*. However, while in  
 1086 2011 two ranked choices were possible, only one was allowed in 2012 (Fig. 10). To  
 1087 compare the two surveys, only the first choice of 2011 is considered here.  
 1088


1089 Some 40% favoured extended Higgs sectors either with or without Supersymmetry, an  
 1090 increase from the about 34% in 2011. Going into more detail, the fraction of those who  
 1091 assumed Supersymmetry to explain new physics did not change with the discovery  
 1092 (25%/24% from 23%/24% as the first choice in 2011). Given that Supersymmetry was  
 1093 the only theory to predict such a light Higgs boson (cf. Sect. 2), its discovery could be  
 1094 seen to have strengthened the case for supersymmetry. On the other hand, none of the  
 1095 expected direct signals of Supersymmetry had been found, seemingly balancing the  
 1096 indirect support from the Higgs mass. General extended Higgs sectors have gained some  
 1097 ground in 2012 (increase from 10%/11%, as first choice in 2011, to 15%/14% in 2012).  
 1098



1099  
 1100  
 1101  
 1102  
 1103  
 1104  
 1105  
 1106  
 1107  
 1108  
 1109  
 1110  
 1111  
 1112  
 1113  
 1114  
 1115  
 1116  
 1117  
 1118  
 1119  
 1120

Fig.10 Percentage of answers of theorists (blue) and experimentalists (red) on the most likely model for physics beyond the Standard Model. Only answers with relation to the electroweak symmetry breaking are given, the remaining 50/58% refer to different models (Questionnaire of 2012).

In contrast, the fraction of physicists considering dynamical EWSB as the best chance was reduced to almost half between 2011 and 2012. Just 10%/4% of theorists/experimentalists advocated for it after the observation of a Higgs candidate (previously 16%/7%). Even in spite of the limited parameter space of composite Higgs models, the number of proponents was still remarkable among theorists. The replies to the question *Independently of your expectations regarding LHC results, which (if any) of the following models do you prefer?* (Fig. 11), hardly changed. There was only a small decrease in the responses for dynamical electroweak symmetry breaking (14%/5% after 19%/7% in 2011). Thus, a significant minority among theorists prefers a solution of EWSB that is different from the Higgs mechanism, even though many do not believe this to be realised at LHC energies. This testifies the tenacity of theories that are attractive on internal grounds, that have strong pragmatic virtues, notwithstanding negative empirical results as long as there remain at least some options to adapt them to the data. (cf. 5.1.3.)



1121  
 1122  
 1123  
 1124  
 1125  
 1126  
 1127

Fig.11 Percentage of answers of theorists (blue) and experimentalists (red) on the preferred model for physics beyond the Standard Model. Only answers with relation to the electroweak symmetry breaking are given, the remaining 50/58% refer to different models (Questionnaire of 2012).

1128 **6.1.3 How was the naturalness problem seen after the Higgs candidate?**


1129

1130 With the discovery of the Higgs candidate, i.e. the likely existence of an elementary  
 1131 scalar, it appeared that the naturalness problem had changed from a potential problem  
 1132 to an actual one. It could no longer resolve itself by the absence of a scalar from the set  
 1133 of fundamental particles. Furthermore, no BSM signal had been found to alleviate these  
 1134 concerns. We therefore tried to understand whether the physicists' assessment had  
 1135 changed after the Higgs discovery.

1136

1137 As in 2011, physicists were asked for the criteria, 'which have guided' their selection of  
 1138 the preferred model, irrespective of the chances of confirming it at LHC. (Fig. 12) The  
 1139 attitude towards naturalness, however had hardly changed after the observation of a  
 1140 Higgs candidate. Naturalness was mentioned in 17%/18% of all answers, only mildly  
 1141 behind the criterion of 'elegance' (20%/17%), but clearly ahead of simplicity  
 1142 (12%/10%).

1143



1144


1145

1146 *Fig.12 Percentage of answers of theorists (blue) and experimentalists (red) on the criteria to choose*  
 1147 *the preferred model (Questionnaire of 2012). The three choices were added up.*

1148

1149 The other question pertinent to assessing the physicists' attitude towards naturalness  
 1150 was again the 'most critical flaw of the Standard Model'. (Fig 13) As in the 2011 survey  
 1151 up to three choices could be given. As before 'quadratic divergences in corrections to  
 1152 Higgs mass' remained at 15%/11% (compared to 14%/10% in 2011) as one of the three  
 1153 major flaws. Also most of the other assessments were fairly similar. A notable exception  
 1154 was that experimentalists now tended to consider the absence of a dark matter  
 1155 candidate' (21% after 17%) to become more critical. In both cases the replies from  
 1156 theorists did not change significantly. After the discovery of the Higgs boson, it seems,  
 1157 experimentalists shifted their interest to the next problem, which they thought to be in  
 1158 reach of the LHC, dark matter, even though there was no consensus about a suitable  
 1159 theoretical model, or whether such a dark matter candidate would be at all in the energy  
 1160 range of the LHC.

1161



1162  
1163

1164 *Fig.13 Percentage of answers of theorists (blue) and experimentalists (red) on the criteria to choose*  
1165 *the preferred model (Questionnaire of 2012). The three answers were added up and normalized.*  
1166 *Note that the classifications are abbreviated the exact questions are listed in Appendix 2*  
1167

1168 Both questions show that the perceived importance of the naturalness problem has not  
1169 been affected by the change in the problem's specific status. A possible reason for this  
1170 stability is that epistemic and pragmatic criteria, once adopted by the community,  
1171 usually operate on a longer time scale. It takes a certain record of scientific successes to  
1172 support them – as has been the case with the naturalness principle, at least in the eyes of  
1173 the particle physics community.<sup>25</sup> That said, one may expect that after years of  
1174 unsuccessful searches for BSM physics, naturalness would become much less attractive.  
1175

1176

## 1177 **6.2 Outcome from interviews in the light of the discovery**

1178

1179 As in 2011 the questionnaire was complemented by interviews, this time with five  
1180 theorists and two experimentalists. Only one of them had already been interviewed in  
1181 2011; this time more emphasis was given to less senior physicists. The interviews took  
1182 place in autumn 2012.  
1183

1184

1185 A large part of the interviewees characterized the Higgs discovery as an '*exciting*' event  
1186 that would have decisive implications on future research. Overwhelmingly physicists  
1187 cautioned to jump to immediate conclusions about the details of the SM Higgs boson.  
1188 One interview partner expressed this very clearly '*well, we still do not know what we*  
1189 *observed*'. But: '*It would have surprised me more if it would not be it [the Higgs boson].*  
1190 *This is in some sense paradoxical, since it is just something one got used to*'. Despite all  
1191 caution there was agreement about the next steps.

1192

### 1193 **6.2.1 From the observation to scrutiny**

1194

1195 The focus, both experimentally and theoretically, was now to qualify this boson and look  
1196 whether it was the SM Higgs boson or whether it had new physics in its wake. An  
1197 experimentalist noted '*At this stage we observe a new particle ... with properties*

<sup>25</sup> Guidice (2008) shows that there the record of the naturalness principle becomes mixed if one assumes a broader perspective; and one might even consider the fortunes of fine-tuning arguments more general. However, we are restricting our considerations to the understanding and role of naturalness within the community of elementary particle physicists.

1197 *consistent with the Standard Model Higgs. This can change .... if we find something that*  
1198 *does not fit into the Standard Model'. But also theory is required to improve the precision*  
1199 *of the predictions of Higgs properties 'If you don't see new physics directly, then maybe we*  
1200 *see it through precision measurements, .... indirect tests.'*

1201  
1202 In this sense, the new particle is seen as a potential harbinger of new physics, not the  
1203 closure of a research program. 'Higgs physics' changed from searching for the Higgs  
1204 boson towards measuring the properties of the new particle, assuming it to be 'largely'  
1205 the Higgs boson, but also searching for deviations from its precisely predictable  
1206 properties.

1207  
1208

### 1209 **6.2.2 The implication on other models**

1210

1211 As discussed above, Supersymmetry predicted the Higgs boson to be lighter than 130  
1212 GeV. It would have been in deep troubles, had the Higgs been found at a larger mass.  
1213 After unsuccessful searches for direct signs of Supersymmetry, the mere consistency of  
1214 the measurements with its prediction, was taken as indirect support. One theorist says '*if*  
1215 *it had been 140 GeV or 150 – that would have changed a lot, because then my SUSY-models*  
1216 *would all be dead. And I would stop working on them.'*

1217

1218 Another impact of the Higgs discovery on BSM models is that, within the experimental  
1219 uncertainties, it provides additional constraints. Indeed, in this perspective '*the*  
1220 *discovery of the Higgs' is the main... result that influences our work', a theorist explained;*  
1221 *'when we [work on] models we have to take... into account ... this particle and this*  
1222 *changes... the situation definitely'. Its existence and even the marginal precision of*  
1223 *autumn 2012 constrains the allowed parameter space of BSM models, 'the determination*  
1224 *of parameters of models or the testing of models I continue to do, now including the*  
1225 *information from the Higgs. And that has changed something in the interpretation'. E.g. to*  
1226 *determine the allowed parameter space in supersymmetric models. 'What would be a*  
1227 *125 GeV Higgs [in super symmetry]? Of course, a SUSY Higgs, but what a parameter space*  
1228 *would be compatible with this? It would be a very small one.'* In this sense the discovery of  
1229 the Higgs candidate has severe implications for many models, at least in limiting  
1230 significantly the allowed parameter range. This is of course an effect that becomes  
1231 poignant on a larger timescale when it may eventually squeeze out certain models  
1232 entirely. Such an effect was not yet visible in the changes of model preference between  
1233 the 2011 and 2012 questionnaires.

1234

1235

### 1236 **6.2.3 Shedding doubt on the previous guiding principle of naturalness?**

1237

1238 As mentioned in 5.1.3, the naturalness problem in 2012 turned from fiction to reality.  
1239 This can also be gathered from the interviews. '*I would say that now that it is certain*  
1240 *that there is a Higgs state at this mass, [the naturalness problem] is more alive than ever'.*  
1241 And at least some continue to consider it an important question. Naturalness '*is still an*  
1242 *important argument. I cannot see any reason, why should happen something like that, such*  
1243 *[fine – tuning] just in a natural way. .... we put this fine-tuning by hand, ... it cannot happen*  
1244 *in nature.'*

1245

1246 Naturalness continues to be seen as an important guiding principle for the development  
1247 of BSM models. "*We need the guidelines. Because, it's not just the experiment, it's not just*  
1248 *mathematics. It's something, which is between induction and deduction. ... And you need ...*  
1249 *some guidelines. One guideline could be this naturalness, ... which is a theoretical guideline.*  
1250 *Or, 'minimality', one theorist argued, that is, for a model to have the minimal number of*

1251 free parameters.

1252

1253 However, this was not the only reaction. The unnaturalness of the SM becomes more  
1254 acute since none of the anticipated solutions in terms of New Physics has shown up in  
1255 the energy range where it was expected, a fact that was considered highly disturbing.  
1256 This dilemma leads to a growing discussion about the status of the naturalness problem:  
1257 *'is this problem a real problem or just a fantasy of theoretical physicists? ... I've been*  
1258 *trained to look at it as a serious problem'*. Yet another theorist raises doubts: *'what we*  
1259 *thought was a main motivation to expect Supersymmetry at the LHC, this hierarchy*  
1260 *problem or naturalness problem, ... I'm not so sure anymore whether this is actually*  
1261 *something that leads us into the right direction.'* *'People have just accepted the fact that*  
1262 *there is more and more fine-tuning now, because of the limits that become larger and*  
1263 *larger. And it's not so clear to me whether it's still a good idea to consider that'*. Similarly  
1264 another theorist argued, *'We can discuss whether... we have to accept fine-tuning. What*  
1265 *could be behind a fine-tuning ... or if one wants a natural theory without fine-tuning. ...*  
1266 *This is the main argument which ... I think, drove ... the theoretical community for the last*  
1267 *twenty or thirty years. But it's not a solid argument'*.

1268

1269 To sum up, the discovery of the Higgs boson, together with the absence of any sign of  
1270 new physics, has turned naturalness from a potential into a real theoretical problem.  
1271 While before the discovery of the Higgs boson physicists might have expected that  
1272 naturalness could be restored by a different mechanism of EWSB, LHC has now  
1273 confirmed the unnatural Higgs mechanism, yet without finding evidence for a potential  
1274 solution outside the EWSB sector. It is true, naturalness could still serve as a guideline  
1275 for devising new models. But the absence of a cure for the naturalness problem of the  
1276 SM has made some physicists wonder whether it is actually a deep problem or whether  
1277 one should simply accept fine-tuning as a fact about nature and accept models that  
1278 violate naturalness. Therefore, the naturalness problem was still considered to be  
1279 important in the questionnaires, but the interviews showed that its previous importance  
1280 was put into question. Throughout the interviews, there was no indication that  
1281 physicists considered the naturalness problem as multifaceted or vague. Instead they  
1282 interchangeably denoted it as hierarchy problem, fine-tuning, or quadratic divergences.  
1283 In Section 7.4., we will discuss how this coherence in scientific practice squares with the  
1284 differences in philosophical analysis.

1285

1286

## 1287 **7. Some philosophical lessons**

1288

1289 In this Section we interpret the outcome of interviews and questionnaires in light of the  
1290 questions mentioned in the Introduction.

1291

### 1292 **7.1 Scepticism before and after a crucial test**

1293

1294 In retrospect, it might appear that the Higgs discovery had been largely expected. Our  
1295 studies show that, in actual fact, the expectations of the LHC physicists were fairly  
1296 diverse. At least shortly before a Higgs candidate was discovered, the community was  
1297 basically split whether to expect the observation of a SM Higgs or not. (cf. 5.1.2) They all  
1298 were aware that LHC had sufficient luminosity to accomplish such a crucial test. The  
1299 reluctance to embrace the SM Higgs boson is in line with wide-spread criticism of the  
1300 conceptual structure of the Higgs mechanism. None of the interviewees emphasised its  
1301 theoretical elegance, some even considered it an ad-hoc argument. This reluctance is in  
1302 contrast to especially Supersymmetry, which is frequently considered as too beautiful a  
1303 theory that nature should not have chosen it – regardless whether it is realised in the  
1304 LHC energy range. The proponents of the Higgs mechanism simply regarded it as the

1305 'best' solution for mass generation that had come to the physicists' mind over the course  
1306 of more than four decades. It did not contradict any measurement, was compatible with  
1307 a wider theory, and had only relatively few parameters.

1308  
1309 On the other hand, after a candidate with the 'right' mass and with properties consistent  
1310 with the expectations had been observed, most LHC physicists-almost immediately  
1311 embraced the notion that 'a' Higgs boson had been found – although they left it open  
1312 whether it would be the only one. This overwhelming acceptance came, although the  
1313 precision of the measurements still left quite some room for alternative solutions of  
1314 EWSB. (cf. 6.1.1. & 6.2.1) In fact, one of the main research directions after the  
1315 observation of the Higgs candidate, both experimentally and theoretically, became to  
1316 scrutinize how large a parameter space for alternative solutions would be left. Whereas  
1317 the vast majority did not consider solutions radically different from the SM, such as a  
1318 composite scalar, as realistic, there was considerable hope to find deviations from the  
1319 SM expectations that would give physicists a hint how to further investigate the complex  
1320 landscape of BSM models. (cf. 6.1.2) When doing so, most data analyses after 2012  
1321 assumed the existence of a SM Higgs boson at 125 GeV, at least as the best  
1322 approximation of the observed boson. This represented a significant discontinuity in the  
1323 actual experimental strategy.

## 1324 1325 **7.2. Was the Higgs Discovery a Crucial Experiment?**

1326  
1327 In the interviews, the discovery of a Higgs boson was widely considered as extremely  
1328 important for particle physics. This accords with the many statements in the literature  
1329 during the last decades<sup>26</sup> and the significant material and intellectual resources that  
1330 went into large experimental facilities to solve the problem of EWSB. This widely shared  
1331 conviction among physicists that the problem of EWSB was at the crossroads of particle  
1332 physics at large prompts the question as to whether the Higgs discovery represented a  
1333 crucial experiment in a philosophical perspective? In Section 3.2 and 3.3, we have  
1334 discussed underdetermination and its impact on the feasibility of crucial experiments  
1335 and other ground-breaking experiments.

1336  
1337 In this section, we argue that the underdetermination argument can be contained to  
1338 such an extent that it does not play a role in actual scientific practice. On this basis we  
1339 also argue that the Higgs discovery can indeed be considered as a crucial experiment.  
1340 Even though the Higgs discovery – as shown in Section 7.1 and emphasised in many  
1341 physics papers – was not a simple yes/no experiment, its cruciality, to our mind, rests  
1342 upon the following characteristics:

- 1343
- 1344 a. The Higgs boson was an essential and indispensable element of the SM.  
1345 Moreover, the Higgs discovery was the final confirmation of a theoretical  
1346 framework that had been developed over decades. Of course, the SM would also  
1347 have broken down if, e.g. no  $Z^0$  boson would have been found. But the Higgs,  
1348 belongs to a sector of the SM that had not been seen before and is based on the  
1349 additional concept of spontaneous symmetry breaking, for which no evidence  
1350 had been observed before.
  - 1351
  - 1352 b. The Higgs boson is fundamentally different from all particles found up to date  
1353 because it is an elementary scalar. It is not the first scalar found, but e.g. the pion  
1354 has been shown to be composed of two quarks and gluons.
  - 1355

---

<sup>26</sup> See, e.g. Ellis, J., Gaillard, M.K., and Nanopoulos D.V. (2012), Quigg (2007).



1356 c. In view of the importance of the EWSB mechanism, a plethora of alternative  
1357 models had been constructed. The discovery of the Higgs boson basically  
1358 eliminated several families of these alternative models, and reshaped the  
1359 direction of future research in a fundamental way.

1360  
1361 Whereas a. and b. emphasise the crucial importance of a Higgs discovery for the SM and  
1362 the general concept of elementary particle, characteristic c. widens the traditional  
1363 philosophical understanding of crucial experiments which required that only one model  
1364 survives. However, even if some alternatives may remain viable, the crucial experiment  
1365 drastically reshapes the field. In the next paragraphs we argue that the crucial nature of  
1366 an experiment as complex as LHC has to be judged against the backdrop of the historical  
1367 development of the respective field.

1368  
1369 Let us assess the acceptance of the observed particle being a Higgs in more detail.  
1370 Immediately after the announcement at CERN in 2012 and the first measurements of its  
1371 mass and decay modes, there was a flood of theoretical analyses, significantly reducing  
1372 the possible parameter space of the many previously developed BSM models. Some  
1373 models, among them 'higgsless models' and 'higgs-gauge unification', were strongly  
1374 disfavoured and did not play a role in the subsequent discussions. Others, like 2HDM  
1375 were more difficult to reject at this stage. As becomes apparent from the questionnaire,  
1376 even only a few months after the discovery was announced, alternative models for  
1377 EWSB were only expected by less than 10% of the respondents. (cf. 6.1.1.).

1378  
1379 During the following year – with more data being analysed and additional properties  
1380 being searched for, especially the spin – the notion of 'having found a Higgs boson' was  
1381 adopted by the majority of the LHC physicists. The notion of 'a' Higgs boson leaves open  
1382 the option of having a more complicated Higgs sector than the SM Higgs. In principle this  
1383 can be resolved by higher experimental precision and additional searches, however, the  
1384 precision will never be perfect in the future, such that small deviations from the SM  
1385 Higgs cannot be ruled out with 100% certainty.

1386  
1387 In this situation physicists are moving forward in their research accepting the SM Higgs  
1388 boson to exist. This consensus was not based on logical inference, in the sense that  
1389 physicists waited until all alternative solutions were definitively excluded.<sup>27</sup> It contained  
1390 a certain dose of pragmatism in choosing promising research strategies. Such a  
1391 'pragmatic solution' to the Duhem-Neurath-Quine problem, as Franklin and Perovic  
1392 (2015, 84) have aptly put it, does not preclude intensive future scrutiny of the signal  
1393 both from an experimental and theoretical side. Just the opposite: this scrutiny leads to  
1394 the emergence of a very significant new research direction. This persistent search for  
1395 potential deviations is not, to our mind, in conflict with the Higgs discovery being  
1396 crucial. To the contrary, such explorative searches that do not test models already  
1397 proposed by theorists, can be seen as one way to address the problem of unconceived  
1398 alternatives.

1399  
1400 Let us discuss now more specifically how Duhemian underdetermination is openly  
1401 addressed within the statistical data analysis. First, the discrimination of two (or  
1402 several) hypotheses, framed in an identical theoretical environment, as the SM in  
1403 particle physics, with the same (kind) of experiments is (almost) completely free of  
1404 detailed theoretical considerations. For in such cases the well accepted and identical  
1405 procedures are applied to either of the hypotheses. Duhemian underdetermination and

---

<sup>27</sup> Wüthrich (2016) advocates a notion of diagnostic causal inference that partially dispenses with the explicit assumption of theories without sacrificing talk about causality in particle scattering.



1406 theory-ladenness are shielded off by referring in the same way to an entire experimental  
 1407 set-up. This vastly reduces their sway among scientists.<sup>28</sup> Second, theory ladenness is  
 1408 significantly alleviated by using precision data from LHC and other experiments and the  
 1409 familiarity with experimental strategies, among them the rules of data analysis and the  
 1410 knowledge about background processes. The Higgs discovery, accordingly, was part and  
 1411 parcel of a longer tradition of accelerator experiments and the associated theoretical  
 1412 research programs.

1413  
 1414 One can formalize the physicists' handling of Duhem's problem as such. We denote the  
 1415 observed number of events of a certain signature<sup>29</sup> of scattering process by  $O$ ,  $T_1$  and  $T_2$   
 1416 are two theoretical hypotheses and  $A_i$  a set of auxiliary hypotheses. The  $P_i$  denote the  
 1417 predictions to measure a number of events given hypothesis  $T_i$  and the auxiliary  
 1418 hypotheses. We then have<sup>30</sup> in a simplified notation omitting the uncertainties of  $T_i$ ,  $A_i$ ,

1419  
 1420  $(T_1 \& A_1 \& A_2 \& A_3 \& \dots A_n) \models P_1, O \models P_1$   
 1421

1422 Duhem argues correctly that  $O$  can also be inferred from (e.g.)

1423  
 1424  $(T_2 \& A_1 \& A_2 \& A_3^* \& \dots A_n) \models P_1, O \models P_1$   
 1425

1426  $A_3^*$  being an alternative auxiliary hypothesis. The two-pronged strategy by which  
 1427 particle physicists deal with the Duhem problem can be expressed as such. Using the  
 1428 identical auxiliary hypotheses for an experimental and theoretical environment allows  
 1429 one to test  $T_1$  and  $T_2$ . Most importantly, each of the auxiliary hypotheses has been  
 1430 experimentally tested under multiple conditions. Theoretical assumptions are kept  
 1431 minimal and, if needed, they also have been tested extensively. Therefore  $A_3^*$  can be  
 1432 excluded, leading to

1433  
 1434  $(T_2 \& A_1 \& A_2 \& A_3 \& \dots A_n) \models P_2, O \not\models P_2,$   
 1435

1436 which allows physicists eventually to discriminate between  $T_1$  and  $T_2$ .

1437  
 1438 Let us be more specific about the auxiliary hypotheses. They include the known physics,  
 1439 rules of data analysis, criteria of statistical significance (cf. Beauchemin 2017). Some of  
 1440 these auxiliary hypotheses have been extensively tested before LHC was even built,  
 1441 others can be tested in-situ using the redundancy of LHC experiments.

1442  
 1443 In the actual practice of particle physics, the  $T_i$  and  $A_i$  are only known to some statistical  
 1444 and systematic uncertainty. This implies that also the predictions  $P_i$  have uncertainties,  
 1445 i.e.  $(P_i \pm \delta_i)$ , where the  $\delta_i$  are convolutions of the uncertainties of the individual  $A_i$  and  $T_i$ .  
 1446 As a result, instead of a strict agreement or disagreement of the predictions with the  
 1447 observation  $O$ , only a finite likelihood  $p_i(O|P_i)$  can be assigned taking into account the  
 1448 uncertainty of  $P_i$ . Duhem's problem therefore reappears in probabilistic terms. To keep  
 1449 underdetermination at bay, to resolve Duhem's problem in scientific practice, two  
 1450 additional conditions must be met. The first one is to confirm not only the correctness of  
 1451 the  $A_i$ , but also all their individual uncertainties. This is done simultaneously with the

---

<sup>28</sup> Assuming the proper functioning of cables and switches is not to say that such errors do not occur. But it seems to us that those are not the kind of victories that advocates of a strong global notion of underdetermination (cf. 3.2.) would want to score. As Franklin (2013) shows, interesting failures of experiments are of a different kind.

<sup>29</sup> We will provide a more detailed analysis of the role of signatures in a subsequent paper that draws on the respective material from the questionnaire and the interviews.

<sup>30</sup> Note that  $\models$  does not amount to deductive entailment in the original Duhemian sense.

1452 extensive tests of the  $A_i$  (cf. Mättig 2019). Secondly, this probabilistic reasoning may in a  
1453 strict (and naïve) manner be interpreted as impossibility to decide between  $T_1$  and  $T_2$ ,  
1454 even if  $p_2$  is ridiculously small, say  $10^{-40}$ . To ‘exclude’ such meaningless hypotheses  $T_i$ ,  
1455 the actual scientific practice defines conventions on the minimum magnitude of the  $p_i$  to  
1456 accept a hypothesis.

1457  
1458 This shows that by embedding the actual experiment into a broader context and by  
1459 distinguishing the different layers of theorizing and experimentation, crucial  
1460 experiments are possible. Duhem’s underdetermination can be addressed in scientific  
1461 practice. This also agrees with Weber’s intuition discussed in Sect. 3.3. that experiments  
1462 are embedded into a broader experimental program and partially takes up Baetu’s point  
1463 that such a program may not only contain the purported crucial experiment. LHC has  
1464 always tested other models and done exploratory searches alongside testing the SM.  
1465

1466 Having dealt with Duhem’s problem, let us discuss more generally what it takes to make  
1467 an experiment in particle physics crucial. The Higgs discovery clearly was essential for  
1468 the SM, but why was it not simply a ‘persuasive experiment’ in the sense of Franklin and  
1469 Perovic? (cf. 3.3) After all, different mechanisms of EWSB are still being discussed even  
1470 though – as we have seen in Sect. 6.1. – they are starting to draw less interest. Franklin  
1471 and Perovic’s distinction is based on the complexity of the inference  $\models$  and the number  
1472 of auxiliary hypotheses required. But this seems to be problematic for large experiments  
1473 like LHC that use many auxiliary hypotheses that are tested by different research  
1474 groups. Taking into account also their analysis of the Stern-Gerlach experiment, it  
1475 appears to us that the requirement of an immediate acceptance or refutation of  
1476 competing models makes the notion of a crucial experiment subject to matters of short-  
1477 term historical developments in the sense that that the status of the P violation  
1478 experiment could have changed if some months later some alternative explanation had  
1479 emerged that required further experiments to be excluded. Would the discovery of CP  
1480 violation turn into a crucial experiment, if those alternatives had been devised before  
1481 the experiment? Does the ‘cruciality’ of an experiment depend on the number of models  
1482 developed before and afterwards?  
1483

1484 On the basis of these classical examples from particle physics, we are thus suggesting a  
1485 notion of crucial experiment that is closer to experimental practice and less dependent  
1486 on short-term developments in physical theory. This notion seems to us also in the spirit  
1487 of Weber’s (2009) discussion. But we are well aware that a substantive discussion of our  
1488 proposal would require a broader set of examples, both positive and negative ones.  
1489

1490 An experiment with a systematically and statistically significant outcome, is crucial in  
1491 some field of science, if it is

- 1492 a) seminal or decisive for the further development of this field,  
1493 and at least one of the following criteria are fulfilled.  
1494 b) It adds a new concept to the body of physics,  
1495 c) it implies a rejection of one or several theoretical solutions of a significant problem,  
1496 or refutes an established concept.

1497 The third criterion takes up characteristic b. in the above description of the Higgs  
1498 discovery. In the examples discussed, P and CP violation fulfil criteria a) and c), the  
1499 Stern-Gerlach experiment fulfilled criteria a), b), c) before the advent of quantum  
1500 mechanics, fulfilled a), c) until spin was fully integrated into the theory, after which it  
1501 fulfilled a) and b)<sup>31</sup>. The Higgs discovery fulfilled a) and b) by itself, but it also refuted  
1502 alternative models, such that it also fulfils c).

---

<sup>31</sup> In contrast to Franklin and Perovic, we believe that the experiment’s crucial character was present throughout. Initially, the experiment was proof of the phenomenon of space quantization

1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545

### 7.3. Principles and values of model choice

The questionnaires and the interviews have shown that in 2011 there was a large variety of epistemic and pragmatic values guiding the physicists' expectations about the Higgs searches, their concerns about the SM and their preferences for BSM physics. The discovery of the Higgs boson disfavoured certain models and strengthened the genuinely epistemic criteria. The example of dynamical symmetry breaking after the Higgs discovery showed that theoretical simplicity or other pragmatic criteria, e.g. fertility to calculate further particle masses, can motivate researchers even if the respective model is experimentally disfavoured. This corresponds to Kuhn's insight that the balancing of epistemic and pragmatic values is neither a perfect logical inference nor a matter of taste, but a fact in the history of scientific practice. Even though physicists are now largely accepting the SM as one of the most successful theories, they will keep looking for deviations that promise to be interesting.

Even taking into account tight experimental constraints, the 'expectation' to find dynamical EWSB was still significant among theorists before the observation of the Higgs candidate. It decreased after the observation, but remained remarkably high. Interestingly, the preference among theoreticians was, both in 2011 and 2012, always higher when asked 'independently of LHC'. (cf. 5.1.3. & 6.1.2.) Even though the chances for experimental evidence in the near future were low, dynamical EWSB remained a preferred solution for many. Notably, the preference of dynamical EWSB showed the largest difference in the surveys between experimentalists and theorists.

Let us evaluate possible consequences of applying the notion of values of preference to models instead of theories. Not all BSM models are as close to the status of a theory as the SM. Some of them are renormalizable and based on a sufficiently elaborated theoretical idea, such as Supersymmetry. But others are not; there is, for instance, considerable freedom in populating an extended Higgs sector. This does not prevent physicists from considering such models as worthy of pursuit, from choosing to investigate them, and indicate their motives for doing so. Finding evidence for such models would certainly have prompted theoretical investigations before physicists would commit themselves to the truth of that model in the same sense as one might commit oneself the existence of supersymmetry in nature. We do not see this as a weakness of our account, but as a consequence of the variegated model landscape.

Let us now look at our findings from the perspective of Douglas's classification of cognitive and pragmatic values. All models are physically consistent and are empirically adequate in the sense that they are not in conflict with the existing data. Thus, they fulfil the minimal criteria (i) and (ii), They also make specific predictions<sup>32</sup> and have a high predictive accuracy once basic parameter(s) are fixed. Furthermore, there is a consensus in the importance of the mechanism of EWSB. They are thus fulfilling the

---

predicted by the Bohr-Sommerfeld theory and a refutation of the classical views developed by Larmor. However, within quantum mechanics the concept of space quantization was replaced by a somewhat different notion of angular momentum, but this did not end the experiments crucial significance against pre-quantum theories (c). When quantum mechanics was fully established, including spin, there was so much spectroscopic evidence against the pre-quantum physics that the Stern-Gerlach experiment became less important in this respect (ending c)), but paradigmatic for the phenomenon of spin (fulfilling b with respect to this concept).

<sup>32</sup> The fractions of physicists choosing 'fit to data' and 'specific predictions' seems to be lower than one might expect but has to be seen on the background of these being not special to any model considered.

1546 desiderata (iv). The questionnaire has also revealed a high score for genuinely  
1547 pragmatic values that fall in Douglas's category (iii) and we have argued that they we  
1548 able to balance the bleak empirical prospects for certain models as long as they were not  
1549 ruled out in the sense of the minimal criterion (ii). Douglas admits that such a practice is  
1550 legitimate as long as it is done with "the full acknowledgement that the theory is  
1551 inadequate as it stands and that it must be corrected to meet the minimum  
1552 requirements as quickly as possible." (2013, 802) She also admits that there are  
1553 tensions in practice "between a well-supported theory (with group (iv) values  
1554 supporting it) and an underdeveloped theory (with lots of group (iii) values and thus  
1555 lots of potential)" (2013, 804). But these tensions are only pragmatic ones between  
1556 conservatives and risk takers that do not endanger the separation between groups (iii)  
1557 and (iv) because they "aim at different purposes" in the sense that "pragmatic criteria  
1558 have no bearing on what should be thought of as our best supported scientific  
1559 knowledge at the moment." (2013, 804)

1560  
1561 It seems to us that when applying the values of preference to a complex model  
1562 landscape, matters are a bit more complicated and Douglas' classification has to be  
1563 taken with a grain of salt. Let us start with the claim that "groups (i) and (ii) ... trump  
1564 groups (iii) and (iv)" (2013, 804) and are clearly distinct as necessary conditions and  
1565 desiderata. 'Internal consistency' (group (i)) in particle physics means that the theory is  
1566 free of any infinities and can be extrapolated to energies  $\Lambda \rightarrow \infty$ . Strictly speaking, given  
1567 the measurements of the Higgs and top mass this may even not be true for the SM itself,  
1568 with the Higgs potential breaking down at high energy, although energies significantly  
1569 beyond the Planck scale of  $10^{19}$  GeV. This does not play any role for the  $10^3$  GeV  
1570 reachable at the LHC and the empirical adequacy of the theory within the experimental  
1571 uncertainties. However, the same argument applies to the BSM models for EWSB. For all  
1572 models, some energy scale  $\Lambda$  is introduced at which some theory should exist - with  
1573 properties that are vaguely known - and where it is assumed to be fully renormalizable.  
1574 This  $\Lambda$  is, in general, far beyond the reach of the LHC and the models predict just some  
1575 'low energy' (i.e. in the LHC range) phenomena, where the details of the full high energy  
1576 theory do not play a role. As seen in the questionnaires and the interviews, this is not of  
1577 concern for physicists. What is more important to them are solutions of problems like  
1578 dark matter, unification, and naturalness (see next section), but also pragmatic criteria  
1579 like simplicity and elegance.

1580  
1581 In all these models, empirical adequacy is guaranteed by constructing them so as to  
1582 encompass the SM that had a very high degree of experimental confirmation, except for  
1583 the Higgs sector. While accordingly, empirical adequacy is accepted as a preeminent  
1584 value, in the practice of physicists, internal consistency plays only a role if it prevents  
1585 clear predictions. This means that group (iv), at least temporarily, can trump group 1. It  
1586 is true, physicists are fully aware of this fact, however to reach  $\Lambda_{\text{BSM}}$  may take decades, if  
1587 not centuries. To evade this, one either has to redefine the meaning of physical  
1588 consistency or retrigger the epistemic values. Wells (2012) considers mathematical  
1589 consistency (his term for internal consistency) to be as preeminent as empirical  
1590 consistency (i.e. adequacy), while not denying that this is not the general attitude among  
1591 physicists. To implement this preference, he advocates effective field theories (EFT) that  
1592 accomplish internal consistency by adding an infinite series of additional terms made up  
1593 of all fields of the model, implying also an infinite set of free parameters. Whereas this is  
1594 in principle correct it is hardly a practice followed widely in particle physics. When EFTs  
1595 are considered in the actual practice, only a limited set of terms is used destroying the  
1596 mathematical consistency but making the theory tractable.

1597  
1598  
1599

#### 1600 **7.4 The Guiding Principle ‘Naturalness’**

1601

1602 Our questionnaire has shown that naturalness is indeed on a par with the traditional  
1603 pragmatic values of ‘elegance’ and ‘simplicity’ – as for the guiding principles of model  
1604 preference – and on a par with ‘too many independent parameters’, the ‘missing dark  
1605 matter candidate’ and the ‘non-inclusion of gravity’ – as for the most critical flaws of the  
1606 SM. While the many independent parameters render the SM not simple – for decades  
1607 elementary particle physicists have been looking for a simple unifying theory – the two  
1608 other flaws concern empirical facts that cannot be accommodated by the SM.  
1609 From the interviews (cf. 6.2.3) we have concluded that naturalness is considered as  
1610 sufficiently well entrenched within the community to be considered as a coherent  
1611 guiding principle for scientific practice. But it operates both in an epistemic and a  
1612 pragmatic mode.

1613

1614 Renormalization is the way to guarantee the finiteness of the theory, that is, its  
1615 theoretical consistency. In principle the huge ‘unnatural’ renormalization corrections for  
1616 the elementary Higgs boson do not make the theory inconsistent in a strictly formal  
1617 sense. In practice, however, most physicists find this unacceptable - they do not accept  
1618 too much fine-tuning of SM parameters – and try to find a remedy by supplementing the  
1619 SM. This renders naturalness an epistemic value, in Kuhn’s and Douglas’s terminology.  
1620 However, in our understanding, naturalness also acts a pragmatic value. It is an  
1621 operationally relatively easy-to-apply quantitative criterion, at least once it is specified  
1622 how much fine-tuning is allowed, and it constrains models; e.g. it suggests new particles  
1623 with top flavour to compensate the main culprit for ‘unnaturalness’. This may also be the  
1624 reason why naturalness is maintained as an important criterion to devise BSM models.  
1625 Such a double-track value of preference complicates the grouping of values for model  
1626 preference. There is, to our mind, no clear separation of this complex criterion into more  
1627 elementary epistemic and pragmatic values. For despite its complexity, naturalness is  
1628 coherently applied as guiding principle by the physics community. It seems to us that  
1629 this diagnosis does not contradict the philosophical differentiations advocated by Wells  
1630 and Williams (cf. 3.3).

1631

1632 With the Higgs discovery ‘naturalness’ turned from a potential problem of the SM into a  
1633 real one. The positive empirical finding was not accompanied by an observation of new  
1634 particles in the TeV range that could resolve the problem in close temporal proximity to  
1635 the Higgs discovery. One might expect that after confirming the cause of the naturalness  
1636 problem, its solution should have been considered as more urgent. Such a trend was not  
1637 visible in the questionnaire; its high ranking as guiding principle or most critical flaw of  
1638 the SM did not change. The interviews revealed a more differentiated picture<sup>33</sup>. Some  
1639 physicists still regard the naturalness problem as a nuisance, but contemplate that it  
1640 might be an accidental feature of particle physics instead of a solid theoretical argument.  
1641 In a sense, physicists are becoming prepared to live with it.<sup>34</sup>

1642

1643 The resilience of the naturalness problem may result from the fact that there exists no  
1644 clear threshold when a theory becomes ‘unnatural’. At least before the first results of the  
1645 LHC folklore had it that fine-tuning requires new physics at an energy scale of 1 TeV. Yet  
1646 there is no prohibitive argument against changing this to 5 or 10 (or more) TeV, even if  
1647 this increases the amount of fine-tuning. Therefore, with some adjustments, the  
1648 naturalness problem can still be maintained for the forthcoming LHC data; moreover,  
1649 also the parameter space for new physics at the 1 TeV scale has not been completely

---

<sup>33</sup> This agrees with several articles by physicists reconsidering the Naturalness problem, e.g. Guidice (2013), Dine (2015).

<sup>34</sup> Cf. Friederich, Harlander & Karaca (2014, Sect. 7).

1650 covered by previous searches. Thus, many physicists defer the final word on the  
 1651 importance of naturalness to the higher energies and intensities that the LHC is about to  
 1652 enter. It remains to be seen whether BSM physics is found that indeed can cure the  
 1653 naturalness problem, If not, one may wonder how far the scale of new physics can be  
 1654 stretched or whether naturalness will eventually be abandoned. This again shows that it  
 1655 is much more specific than the usual pragmatic values of model preference.  
 1656  
 1657  
 1658  
 1659  
 1660

1661 **8. Conclusion**

1662  
 1663 Let us sum up the main results of our paper. First, the discovery of the Higgs boson and  
 1664 the confirmation of the SM were less expected than is often assumed. With the growing  
 1665 evidence that the newly discovered particle has properties consistent with the SM  
 1666 expectations, most physicists accepted it to be a Higgs, and at least tentatively, a SM  
 1667 Higgs boson. This does not contradict the fact that searches for possible deviations from  
 1668 the SM will be ongoing for a long time. Second, the Higgs discovery represented a crucial  
 1669 experiment for the SM if one interprets the notion in a sense that is appropriate for  
 1670 modern experiments. An experiment as complex as LHC cannot be properly understood  
 1671 without its embedding into a tradition of previous precision experiment and the  
 1672 tradition of reliable and established experimental strategies. These are crucial for  
 1673 keeping underdetermination at bay. Third, our case study suggests that criteria of  
 1674 theory choice be understood as epistemic and pragmatic values that have to be weighed  
 1675 in in factual practice. The Higgs discovery led to a certain shift from pragmatic to  
 1676 epistemic values as regards the mechanisms of EWSB. Complex criteria, such as  
 1677 naturalness, combine different values without becoming inconsistent or inapplicable by  
 1678 the scientific community.  
 1679  
 1680

1681 **Appendix 1 List of interview partners**

1682  
 1683 **March & April 2011**

1684

Prof. V.Sharma	UC San Diego, USA	experimentalist	Convenor of Higgs group in CMS expt.	Male
Prof. F.Gianotti	CERN	experimentalist	Spokeswoman of ATLAS expt.	Female
Prof. G.Tonelli	U of Pisa (Italy)	experimentalist	Spokesman of CMS expt.	Male
Prof. A.Golutvin	IC London (UK)	experimentalist	Spokesman of LHCb expt.	Male
Dr. J.Boyd	CERN	experimentalist	Coordinator data preparation in ATLAS expt.	Male
Prof. J.Ellis	CERN	theorist		Male
Prof. C.Quigg	Fermilab (USA)	theorist		Male
Prof. M.Mangano	CERN	theorist		Male
Dr. M.Mihalla	KIT (Germany)	theorist		Female

1685  
 1686  
 1687 **Fall 2012**  
 1688

Prof. M.Krämer	RWTH Aachen	theorist		Male
Prof. L.Feld	RWTH Aachen	experimentalist		Male
Dr. L. Di Luzio	KIT (Germany)	theorist		Male
Dr. F. Domingo	KIT (Germany)	theorist		Male
Prof. C.Issever	U of Oxford (UK)	experimentalist	Convenor exotics ATLAS expt.	Female
Dr. M.Mihalla	KIT (Germany)	theorist		Female

1689  
1690

1691 **Appendix 2: List of questions in questionnaires**

1692

1693 **In 2011**

1694

1695 **1. What is your personal estimate of the probability of the following**  
1696 **scenarios? The LHC will...**

1697 a. find the Standard Model Higgs boson

1698 b. rule out the Standard Model Higgs boson

1699 c. find indisputable evidence of new physics

1700 The probabilities to be assigned were in 20% intervals, i.e. 0-20, 20-40%, .....

1701

1702 **2. Assuming that the LHC finds new physics, which (if any) of the following**  
1703 **models do you think has the best chances of explaining it**

1704 a. extended Higgs sector

1705 b. supersymmetry

1706 c. extra-dimensions

1707 d. dynamical electroweak symmetry breaking

1708 e. 4<sup>th</sup> generation

1709 f. extended gauge symmetry (Z', Little Higgs)

1710 g. string theory

1711 h. other

1712 i. None of those, but something totally unexpected

1713 j. I don't know

1714 The questionnaire asked for two ranked choices.

1715

1716 **3. Independently of your expectations regarding LHC results, which (if any) of**  
1717 **the following models do you prefer?**

1718 a. extended Higgs sector

1719 b. supersymmetry

1720 c. extra-dimensions

1721 d. dynamical electroweak symmetry breaking

1722 e. 4<sup>th</sup> generation

1723 f. extended gauge symmetry (Z', Little Higgs)

1724 g. string theory

1725 h. other

1726 i. I don't know

1727 The questionnaire asked for two ranked choices.

1728

1729 **4. Which (if any) of the following criteria have guided you in answering the**  
1730 **previous question? (Four ranked answers were possible.)**

1731 a. The model solves naturalness/hierarchy problem

1732 b. The model is simple

1733 c. The model will provide a better fit to the data

1734 d. The model is elegant

1735 e. The model makes very specific predictions

1736 f. The model allows the unification of forces

1737 g. The model has a candidate for dark matter

1738 h. other

1739 i. none of the above

1740

1741 **5. Which (if any) are the most critical flaws of the Standard Model? (up to**  
1742 **three answers possible)**

1743 a. too many independent parameters

1744 b. small but nonzero neutrino masses



- 1745 c. replication of fermion families
- 1746 d. different magnitude of scales of fermion masses
- 1747 e. quadratic divergences in corrections to Higgs mass
- 1748 f. left-right asymmetry
- 1749 g. gravity is not included
- 1750 h. no unification of strong and electroweak forces
- 1751 i. CP violation
- 1752 j. No dark matter candidate

1753

1754 **6. In which of the following signatures (if any) do you think the LHC will most likely find new physics?**

- 1755
- 1756 a. signatures with bottom quarks
- 1757 b. signatures with top quarks
- 1758 c. signatures with tau leptons
- 1759 d. signatures with missing energy
- 1760 e. signatures with multi – jet topologies
- 1761 f. signatures with multi – lepton topologies
- 1762 g. soft events
- 1763 h. other
- 1764 i. I don't know

1765 Two ranked choices were asked for

1766

1767 **7. How much do you agree with the following statements? LHC results will be very important to understand...**

- 1768
- 1769 a. strong interactions
- 1770 b. flavour physics
- 1771 c. origin of mass
- 1772 d. quantum gravitational effects
- 1773 e. dark matter
- 1774 f. dark energy
- 1775 g. cosmology of the early universe

1776 The answers should be given for each field in terms of

1777 'fully agree', 'somewhat agree', 'undecided', 'somewhat disagree', 'fully disagree'

1778

1779 **8. How much do you agree with the following statements?**

- 1780 a. There is plenty of dialogue between theoretical and experimental physicists on LHC physics
- 1781 b. Theorists are fully prepared to tackle future new data from LHC
- 1782 c. Theorists are making helpful suggestions on how to collect and analyse LHC data
- 1783 d. Experimental physicists are sufficiently taking into account suggestions from theorists
- 1784 e. Experimental physicists are presenting their results in the most helpful way for theorists

1785 The answers should be given for each field in terms of 'fully agree', 'somewhat agree', 'undecided', 'somewhat disagree', 'fully disagree'

1786

1787 **In 2012**

1788

1789 **1. How much do you agree with the following statements? After the discovery of the new particle at 125 GeV, the LHC will...**

- 1790 a. confirm the minimal Higgs sector
- 1791 b. find a more complicated Higgs sector
- 1792 c. find an alternative mechanism for EWSB

1793

1799 d. find indisputable evidence of new physics  
1800 The answers should be given for each field in terms of 'fully agree', 'somewhat  
1801 agree', 'undecided', 'somewhat disagree', 'fully disagree'  
1802

1803 **2. Assuming that the LHC finds new physics, which (if any) of the following**  
1804 **models do you think has the best chances of explaining it**

- 1805 a. extended Higgs sector  
1806 b. supersymmetry  
1807 c. extra-dimensions  
1808 d. dynamical electroweak symmetry breaking  
1809 e. 4<sup>th</sup> generation  
1810 f. extended gauge symmetry (Z', Little Higgs)  
1811 g. string theory  
1812 h. other  
1813 i. None of those, but something totally unexpected  
1814 j. I don't know

1815 Only one choice was possible  
1816

1817 **3. Independently of your expectations regarding LHC results, which (if any) of**  
1818 **the following models do you prefer?**

- 1819 a. extended Higgs sector  
1820 b. supersymmetry  
1821 c. extra-dimensions  
1822 d. dynamical electroweak symmetry breaking  
1823 e. 4<sup>th</sup> generation  
1824 f. extended gauge symmetry (Z', Little Higgs)  
1825 g. string theory  
1826 h. other  
1827 i. I don't know

1828 Only one choice possible  
1829

1830 **4. Which (if any) of the following criteria have guided you in answering the**  
1831 **previous question?**

- 1832 a. The model solves naturalness/hierarchy problem  
1833 b. The model is simple  
1834 c. The model will provide a better fit to the data  
1835 d. The model is elegant  
1836 e. The model makes very specific predictions  
1837 f. The model allows the unification of forces  
1838 g. The model has a candidate for dark matter  
1839 h. other  
1840 i. none of the above

1841 Up to three answers were asked for  
1842

1843 **5. Which (if any) are the most critical flaws of the Standard Model? (up to**  
1844 **three answers possible)**

- 1845 a. too many independent parameters  
1846 b. small but nonzero neutrino masses  
1847 c. replication of fermion families  
1848 d. different magnitude of scales of fermion masses  
1849 e. quadratic divergencies in corrections to Higgs mass  
1850 f. left-right asymmetry  
1851 g. gravity is not included  
1852 h. no unification of strong and electroweak forces

- 1853 i. CP violation
- 1854 j. No dark matter candidate
- 1855
- 1856 **6. In which of the following signatures (if any) do you think the LHC will most**
- 1857 **likely find new physics?**
- 1858 a. signatures with bottom quarks
- 1859 b. signatures with top quarks
- 1860 c. signatures with tau leptons
- 1861 d. signatures with missing energy
- 1862 e. signatures with multi – jet topologies
- 1863 f. signatures with multi – lepton topologies
- 1864 g. soft events
- 1865 h. other
- 1866 i. I don't know
- 1867 Two ranked choices were asked for
- 1868
- 1869 **7. How much do you agree with the following statements? LHC results will be**
- 1870 **very important to understand...**
- 1871 a. strong interactions
- 1872 b. flavour physics
- 1873 c. origin of mass
- 1874 d. quantum gravitational effects
- 1875 e. dark matter
- 1876 f. dark energy
- 1877 g. cosmology of the early universe
- 1878 The answers should be given for each field in terms of 'fully agree', 'somewhat
- 1879 agree', 'undecided', 'somewhat disagree', 'fully disagree'
- 1880
- 1881 **8. How much do you agree with the following statements?**
- 1882 a. There is plenty of dialogue between theoretical and experimental physicists
- 1883 on LHC physics
- 1884 b. Theorists are fully prepared to tackle future new data from LHC
- 1885 c. Theorists are making helpful suggestions on how to collect and analyse LHC
- 1886 data
- 1887 d. Experimental physicists are sufficiently taking into account suggestions from
- 1888 theorists
- 1889 e. Experimental physicists are presenting their results in the most helpful way
- 1890 for theorists
- 1891 The answers should be given for each field in terms of 'fully agree', 'somewhat
- 1892 agree', 'undecided', 'somewhat disagree', 'fully disagree'
- 1893
- 1894
- 1895
- 1896

- 1897 **References:**  
1898  
1899 Aad, G. *et al.* (2012). Observation of a New Particle in the Search for the Standard Model  
1900 Higgs Boson with the ATLAS Detector at the LHC. *Physics Letters B*, 716, 1-29.  
1901 *arXiv:1207.7214v1 [hep-ex]*,  
1902 Baetu, T. (2017). On the Possibility of Crucial Experiments in Biology. *The British Journal*  
1903 *for the Philosophy of Science*, published online.  
1904 Chatrchyan, S. *et al.* (2012). Observation of a New Boson at a Mass of 125 GeV with the  
1905 CMS Experiment at the LHC. *Physics Letters B*, 716, 30-61.  
1906 Baker, A. (2013) "Simplicity", *The Stanford Encyclopedia of Philosophy* (Fall 2013  
1907 Edition), Edward N. Zalta (ed.), URL =  
1908 <<http://plato.stanford.edu/archives/fall2013/entries/simplicity/>>.  
1909 Beauchemin, P.-H. (2017). Autopsy of measurements with the ATLAS detector at the  
1910 LHC, *Synthese* 194, 275-312.  
1911 Borrelli, A. (2015). Between logos and mythos: narratives of "naturalness" in today's  
1912 particle physics community. In "Narrated Communities: Narrated Realities", H. Blume &  
1913 C. Leitgeb (Eds.)  
1914 Borrelli, A. (2016): Was Sie schon immer über das CERN wissen wollten, aber bisher  
1915 nicht zu fragen wagten – eine philosophische und soziologische Perspektive, in H. Satz,  
1916 P. Blanchard, C. Kommer (eds.), *Großforschung in neuen Dimensionen Denker unserer*  
1917 *Zeit über die aktuelle Elementarteilchenphysik am CERN*, Berlin-Heidelberg: Springer,  
1918 pp. 119-149.  
1919 Borrelli, A. & Stöltzner, M. (2013) Model Landscapes in the Higgs Sector. In *EPSA11*  
1920 *Perspectives and Foundational Problems in Philosophy of Science*, V. Karakostas & D.  
1921 Dieks (eds.), *The European Philosophy of Science Association Proceedings 2*, Dordrecht:  
1922 Springer, 241-252.  
1923 Carnap, R. (1950). Empiricism, Semantics, Ontology. *Revue Internationale de Philosophie*,  
1924 4, 20-40.  
1925 CERN press office (March 14, 2013) New results indicate that particle discovered at  
1926 CERN is a Higgs boson; [https://press.web.cern.ch/press-releases/2013/03/new-](https://press.web.cern.ch/press-releases/2013/03/new-results-indicate-particle-discovered-cern-higgs-boson)  
1927 [results-indicate-particle-discovered-cern-higgs-boson](https://press.web.cern.ch/press-releases/2013/03/new-results-indicate-particle-discovered-cern-higgs-boson)  
1928 Csaki, C. and Tanedo, P. (2016). Beyond the Standard Model Lectures at the 2013  
1929 European School of High Energy Physics, arXiv:1602.04228v1 [hep-ph] (2016)  
1930 Dallmeier-Tiessen, S. (2016), Hecker, B., Holtkamp, A. et al for the INSPIRE collaboration,  
1931 How partnership accelerates Open Science: High-Energy-Physics and INSPIRE, a case  
1932 study of a complex repository ecosystem, CERN-OPEN-2016-008.  
1933 Dine, M. (2015). Naturalness Under Stress, *Ann.Rev.Nucl.Part.Sci.* 65: 43-62,  
1934 arXiv:1501.01035  
1935 Douglas, H. (2013). The Value of Cognitive Values. *Philosophy of Science* 80: 796-806.  
1936 Ellis, J., Gaillard, M.K. and Nanopoulos D.V., A phenomenological profile of the Higgs  
1937 boson, *Nucl. Phys. B* 106 (1976) 292. (CERN preprint Nov. 1975)  
1938 Ellis, J., Gaillard, M.K., and Nanopoulos D.V. (2012). A Historical Profile of the Higgs  
1939 Boson. arXiv:1201.6045v1 [hep-ph].  
1940 Ellis, J., Gaillard, M.K., and Nanopoulos D.V. (2015). An Updated Historical Profile of the  
1941 Higgs Boson. <http://arxiv.org/abs/1504.07217>.  
1942 Franklin, A. (2013) *Shifting Standards. Experiments in Particle Physics in the Twentieth*  
1943 *Century*. Pittsburgh: University of Pittsburgh Press.

- 1944 Franklin, A. & Perovic, S. (2015). Experiment in Physics, *The Stanford Encyclopedia of*  
 1945 *Philosophy* (Fall 2013 Edition), Edward N. Zalta (ed.), URL =  
 1946 <http://plato.stanford.edu/archives/sum2015/entries/physics-experiment/>
- 1947 Friederich, S., Harlander R., and Karaca, K. (2014). Philosophical perspectives on ad hoc-  
 1948 hypotheses and the Higgs mechanism. *Synthese* 191:3897–3917.
- 1949 Giudice, G. (2008), Naturally Speaking: The Naturalness Criterion and Physics at  
 1950 the LHC, in G. Kane and A. Pierce (eds.), *Perspectives on LHC Physics*, Singapore:  
 1951 World Scientific, 2008, pp. 155-178; also , arXiv:0801.2562.
- 1952 Giudice, G. (2013), Naturalness after LHC8, PoS EPS-HEP2013 (2013) 163. 3,  
 1953 arXiv:1307.7879.
- 1954 Grinbaum, A. (2012). Which fine-tuning arguments are fine? *Foundations of Physics*,  
 1955 42(5), 615–631.
- 1956 Gunion, J.F, Haber, H.E., Kane, G., Dawson, S. (1990). *The Higgs Hunter's Guide*. Redford  
 1957 City: Addison-Wesley.
- 1958 INSPIRE <https://inspirehep.net/info/general/project/index>
- 1959 Karaca, K (2013a). The strong and weak senses of theory-ladenness of experimentation:  
 1960 Theory-driven versus exploratory experiments in the history of high-energy elementary  
 1961 particle physics. *Science in Context* 26: 93-136.
- 1962 Karaca, K. (2013b). The construction of the Higgs mechanism and the emergence of the  
 1963 electroweak theory. *Studies in History and Philosophy of Modern Physics*, 44 (1), 1-16.
- 1964 Kuhn, T.S. (1977). Objectivity, Value Judgment, and Theory Choice. In: *The Essential*  
 1965 *Tension*. Chicago: Chicago University Press, pp. 320-329.
- 1966 Lakatos, I. (1974) *The Role of Crucial Experiments in Science*. *Studies in History and*  
 1967 *Philosophy of Science* 4, 309-325.
- 1968 Laudan, L. (1990). Demystifying Underdetermination. In C. Wade Savage, ed., *Scientific*  
 1969 *Theories* (267-297). Minneapolis: University of Minnesota Press.
- 1970 Laudan, L., and J. Leplin. (1991). Empirical Evidence and Underdetermination. *Journal of*  
 1971 *Philosophy* 88: 449–472.
- 1972 Mättig P (2019) Validation of Simulation in Particle Physics in: Beisbart, C. & Saam, N. J.  
 1973 (eds.), *Computer Simulation Validation - Fundamental Concepts, Methodological*  
 1974 *Frameworks, and Philosophical Perspectives*, Cham: Springer, to appear 2019
- 1975 Norton, J. (2008). Must Evidence Underdetermine Theory? In M. Carrier, D. Howard, and  
 1976 J. Kourany, eds., *The Challenge of the Social and the Pressure of Practice: Science and*  
 1977 *Values Revisited* (pp. 17-44). Pittsburgh: University of Pittsburgh Press.
- 1978 Peskin, M and Schroeder, S, (1995) *An Introduction to Quantum Field Theory*, Cambridge  
 1979 (Massachusetts): Perseus Books
- 1980 Quigg, C. (2007), Spontaneous Symmetry Breaking as a Basis for Particle Masses,  
 1981 *Rept.Prog.Phys.* 70 (2007) 1019-1054, arXiv:0704.2232v2 [hep-ph]
- 1982 Stanford, K. (2006) *Exceeding Our Grasp: Science, History, and the Problem of*  
 1983 *Unconceived Alternatives*, New York: Oxford University Press.
- 1984 Stanford, Kyle, "Underdetermination of Scientific Theory", *The Stanford Encyclopedia of*  
 1985 *Philosophy* (Winter 2017 Edition), Edward N. Zalta (ed.), URL =  
 1986 <[https://plato.stanford.edu/archives/win2017/entries/scientific-](https://plato.stanford.edu/archives/win2017/entries/scientific-underdetermination/)  
 1987 [underdetermination/](https://plato.stanford.edu/archives/win2017/entries/scientific-underdetermination/)>.
- 1988 Stöltzner, M. (2014) Higgs Models and Other Stories about Mass Generation. *Journal for*  
 1989 *General Philosophy of Science* 45, 369-384

- 1990 Susskind, L. (1979). Dynamics of spontaneous symmetry breaking in the Weinberg-  
1991 Salam theory. *Physical ReviewD*, 20(November), 2619–2625.
- 1992 The Royal Swedish Academy of Science, The BEH-Mechanism, Interactions with Short  
1993 Range Forces and Scalar Particles, 8.10.2013,  
1994 [http://www.nobelprize.org/nobel\\_prizes/physics/laureates/2013/advanced.html](http://www.nobelprize.org/nobel_prizes/physics/laureates/2013/advanced.html)
- 1995 't Hooft, Gerard (1979). Naturalness, chiral symmetry, and spontaneous chiral symmetry  
1996 breaking. *NATO Advanced Science Institutes Series B:Physics*, 59 (PRINT-80-  
1997 0083(UTRECHT)),135.
- 1998 Weber, M. (2009). The Crux of Crucial Experiments: Duhem's Problems and Inference to  
1999 the Best explanation. *British Journal for the Philosophy of Science* 60: 10-49.
- 2000 Wells, J. (2012). Effective Field Theories and the Role of Consistency in Theory Choice,  
2001 arXiv 1211.0634
- 2002 Wells, J. (). *Effective Theories in Physics. From Planetary Orbits to Elementary Particle*  
2003 *Masses*. Heidelberg: Springer 2012.
- 2004 Wells, J. (2015). The utility of Naturalness, and how its application to Quantum  
2005 Electrodynamics envisages the Standard Model and the Higgs boson. *Studies in History*  
2006 *and Philosophy of Modern Physics* 49: 102-108.
- 2007 Wells, J. (2017). Higgs naturalness and the scalar boson proliferation instability  
2008 problem, *Synthese* 194, 477-490
- 2009 Williams, P. (2015). Naturalness, the autonomy of scales, and the 125 GeV Higgs. *Studies*  
2010 *in History and Philosophy of Modern Physics* 49: 102-108.
- 2011 Wüthrich, A. (2016). The Higgs discovery as a diagnostic causal inference, *Synthese* 194,  
2012 461-476.