

EXPERT REVIEW

Molecular psychiatry of zebrafish

AM Stewart^{1,2,8}, JFP Ullmann^{2,3,8}, WHJ Norton^{2,4,8}, CH Brennan⁵, MO Parker⁵, R Gerlai⁶ and AV Kalueff^{1,2,7}

Due to their well-characterized neural development and high genetic homology to mammals, zebrafish (*Danio rerio*) have emerged as a powerful model organism in the field of biological psychiatry. Here, we discuss the molecular psychiatry of zebrafish, and its implications for translational neuroscience research and modeling central nervous system (CNS) disorders. In particular, we outline recent genetic and technological developments allowing for *in vivo* examinations, high-throughput screening and whole-brain analyses in larval and adult zebrafish. We also summarize the application of these molecular techniques to the understanding of neuropsychiatric disease, outlining the potential of zebrafish for modeling complex brain disorders, including attention-deficit/hyperactivity disorder (ADHD), aggression, post-traumatic stress and substance abuse. Critically evaluating the advantages and limitations of larval and adult fish tests, we suggest that zebrafish models become a rapidly emerging new field in modern biological psychiatry research.

Molecular Psychiatry (2014) **00**, 000–000. doi:10.1038/mp.2014.128

INTRODUCTION

Psychiatric disorders affect many brain and behavioral processes, including thoughts, feeling, sociability and mood. These complex human diseases also cause a significant burden to society, especially as the available drug therapies have not sufficiently progressed.^{1,2} The etiology of psychiatric disorders is multifaceted, and involves the interaction of multiple genetic and environmental factors.

Genetic mutations trigger psychiatric disorders by altering gene expression and disrupting normal physiological functions of the brain. Mutations linked to psychiatric disorders are often found in genes expressed during embryogenesis, and affect early brain development, cell division, migration, differentiation or survival, as well as axon path finding, dendritic architecture and wiring of neural circuits. For example, Rett syndrome is caused by mutations in the X-linked *MECP2* gene, encoding methyl CpG binding protein-2 that binds to methylated DNA and controls gene expression.³ While *MECP2* is particularly critical during brain development, its mutations are also associated with other 'early-onset' developmental brain disorders, such as X-linked mental retardation and autism³ (also see the substantial contribution of mutations in genes with higher expression in early fetal life to schizophrenia⁴). Environmental factors also have an important role in psychiatric disorders, leading to well-established gene x environment (G x E) interactions.⁵

The notion that psychiatric disorders have fundamental underlying molecular mechanisms is critical for understanding complex brain illnesses, such as anxiety, depression, schizophrenia or addiction. The promise of such an approach is that once these mechanisms are identified and understood, the brain disorder can be treated, reversed or prevented.^{6,7}

Clearly 'mainstream' today, molecular psychiatry had a long journey before being accepted as a new, dynamic field of translational neuroscience.⁷ Utilizing various *in vivo* tests and applying a wide range of model organisms (Figure 1) is critical for uncovering novel mechanisms of brain disorders.⁸ Zebrafish (*Danio rerio*) have long been underrepresented in experimental biological psychiatry, and only recently emerged as a model organism in this field.^{9–11} It is therefore timely to discuss the molecular psychiatry of zebrafish and its implications for clinical as well as translational neuroscience research.

THE UTILITY OF ZEBRAFISH FOR TRANSLATIONAL NEUROSCIENCE

Understanding 'zebrafish psychiatry'

While zebrafish represent a powerful model for developmental neuroscience (Figure 1), a combination of well-characterized neural development and several cutting-edge genetic tools also makes them an ideal species to study complex psychiatric disorders. Although similar manipulations are often available in other vertebrates (for example, mice), the ease of generating large numbers of fish, their external fertilization and transparent embryonic/larval stages are particularly useful, also facilitating high-throughput screening and central nervous system (CNS) imaging.^{10,11} Zebrafish also have an excellent potential to link genes to CNS disorders, because the nucleotide sequence of zebrafish genes is homologous (usually exceeding 70%) to the corresponding human genes, and their function in these distantly related species is often similar.¹² As zebrafish genes can be mutated with high efficiency, the strains (lines) carrying interesting mutations may be detected using phenotype screens, and the mutated genes can be efficiently identified genetically (see

¹ZENEREI Institute, Slidell, LA, USA; ²International Zebrafish Neuroscience Research Consortium (ZNRC), Slidell, LA, USA; ³Centre for Advanced Imaging, University of Queensland, Brisbane, QLD, Australia; ⁴Department of Biology, College of Medicine, Biological Sciences and Psychiatry, University of Leicester, Leicester, UK; ⁵School of Biological and Chemical Sciences, Queen Mary University of London, London, UK; ⁶Department of Psychology, University of Toronto at Mississauga, Mississauga, Ontario, Canada and ⁷Research Institute for Marine Drugs and Nutrition, Guangdong Ocean University, Zhanjiang, Guangdong, China. Correspondence: Professor AV Kalueff, ZENEREI Institute and ZNRC, 309 Palmer Court, Slidell, LA 70458, USA.

E-mail: avkalueff@gmail.com

⁸These authors contributed equally to this work.

Received 6 July 2014; revised 27 August 2014; accepted 28 August 2014




Figure 1. A brief summary of zebrafish experimental models in neuroscience research. **(a)** The evolving nature of zebrafish models in the last 50 years, initially used mainly for basic genetic and neurodevelopmental studies, but more recently applied to developing *in vivo* models for complex brain disorders, such as autism, depression and psychoses, is shown. Inset: selected zebrafish strains useful in biological psychiatry research (top to bottom: adult wild-type zebrafish, *casper*, *spiegeldanio* and *nacre* mutants); photos courtesy of the Kalueff (ZENEREI Institute, USA), the Norton (University of Leicester, UK), the Parichy (University of Washington, USA) laboratories and Carolina Biological Supply Company (Burlington, USA). **(b)** Two research strategies which can both be applied to zebrafish models are illustrated. As a vertebrate species amenable to *in vivo* analyses and with high genetic/physiological homology to humans, zebrafish are ideal for 'intensive', mechanistically driven neuroscience research into conserved, core molecular pathways or neural circuits (photo). Due to their small size, ease of maintenance and short generation time, zebrafish also represent an excellent model for 'extensive' biomedical research, including low-cost high-throughput screening for small molecules or genetic mutations. Both strategies can lead to the development of novel therapies for major psychiatric disorders. Photo: visualizing zebrafish signaling as an example of intensive, pathway-oriented mechanistic research (the left image shows expression patterns of genetically encoded calcium indicator (GECI); zebrafish at 5 dpf show pan-neuronal expression of *HuC:Gal4; UAS:GCaMP5*, a zoomed-in image on the right shows a single-cell resolution: note that some cells are activated, as indicated by the increased fluorescence; scale bar 100 μ m). **(c)** The benefits of including inexpensive zebrafish models into preclinical screening batteries are illustrated. A substantial fiscal saving (shown as \$) can be achieved by narrowing screening to potentially active compounds using fish (rather than mice) as the first vertebrate model organism in the screening pipeline. In addition, the risks of missing promising 'hits' are lower because the subsequent screening in mammals will be 'confirmatory', and based on more valid data generated from zebrafish (rather than invertebrate or *in vitro*) tests. **(d)** Major neurotransmitter systems in the adult zebrafish brain are summarized: ACh—acetylcholine (dark blue), NA—noradrenergic (purple), DA—dopamine (green), HA—histamine (orange), 5-HT—serotonin (red); GABA— γ -aminobutyric acid (pink) and Glu—glutamate (light blue). The highlighted brain regions include olfactory bulb (OB), dorsal pallium (Dp), ventral pallium (Vp), habenula (Ha), optic nerve (ON), facial lobe (LVII) and vagal lobe (LX). Highlighted areas have been overlaid over a minimum deformation model of the adult zebrafish brain.




Figure 1. Continued.

further). Once genes involved in particular functions are identified using such 'forward genetic' strategy, their biological role(s) may be thoroughly investigated using 'reverse genetic' methods, specifically developed for zebrafish (for example, the TALEN ('transcription activator-like effector nuclease') system,¹³ allowing targeted mutation of pre-selected sequences). Thus, to the extent that a psychiatric disease can be modeled in any animal by the generation of a genetic mutant or by pharmacological manipulation, zebrafish present a useful, cost-effective and uniquely tractable system.

While aberrant neurotransmission is one of the main causes of psychiatric disorders,^{6,7} vertebrate neurochemistry is generally highly conserved.^{14–17} Zebrafish possess all primary neuromediator systems, including transmitters, receptors, transporters and the enzymes required for synthesis and metabolism of these

mediators.^{10,11} The spatial and temporal distribution of these systems has been well characterized in both larval and adult zebrafish for glutamate, γ -aminobutyric acid (GABA), acetylcholine and the aminergic neurotransmitters dopamine (DA), noradrenaline (NA), serotonin (5-hydroxytryptamine, 5-HT) and histamine (HA, Figure 2b)—key mediators implicated in multiple psychiatric diseases.

Major neurotransmitter systems in zebrafish

Glutamate is the primary excitatory CNS neurotransmitter involved in schizophrenia, post-traumatic stress disorder (PTSD) and epilepsy. Glutamate signaling is visualized using *in situ* hybridization for vesicular glutamate transporters (*VGLUT1* and *VGLUT2*),¹⁸ and glutamate metabotropic and ionotropic receptors, including




Figure 2. Neurobehavioral phenotypes in zebrafish relevant to modeling a diverse group of psychiatric disorders (also see Table 1). Axis represents five major phenotypic domains of brain disorders, and includes motor activity, cognition, affect/emotionality, reward and aggression. A dashed-line pentagon represents 'normal' healthy phenotype for each respective axis/domains. **(a)** Typical zebrafish responses relevant to ADHD (modified from Lange *et al.*^{125,126}) are shown. The bar diagram shows total distance swum in 5 min for control larvae and *Iphn3.1* morphants with and without methylphenidate (MPH) treatment. The right panel shows representative locomotor traces (recorded from the top view camera) in 6-dpf control and *Iphn3.1* morphant zebrafish (green color indicates swimming with normal speed, red color indicates bursts of 'impulsive' swimming with high acceleration). **(b)** Zebrafish mirror-induced aggression (MIA) responses in the mirror exposure test, and typical patterns of zebrafish aggressive confrontations (modified from Norton *et al.*¹³⁹) are illustrated. Note that MIA is particularly simple to record and quantify,¹⁶¹ making it also suitable for high-throughput screening (MIA images in the middle: courtesy of the Robison laboratory,¹⁶¹ University of Idaho, USA). **(c)** Zebrafish stress- and PTSD-related aversive behaviors, including neophobia (e.g., acute 5-min exposure to the unfamiliar blue marble in the novel object test) and predator avoidance (acute 5-min exposure to a predator fish) are illustrated. Note that zebrafish avoid a blue marble at the bottom of the tank (swim traces in the open-field test recorded by Noldus Ethovision XT8 from the top view), as well as display robust aversion of a live predator (e.g., Leaf fish), clearly spending more time in the opposite side of the tank. **(d)** Zebrafish reward-related addiction behaviors (in the conditioned place preference model, CPP) and molecular alterations in zebrafish brain following exposure to a substance of abuse are demonstrated. In the CPP, fish are initially tested for their preference for both sides of the apparatus, each of which contains a discriminative stimulus (i.e., spots vs no spots). They are then conditioned with drug to the least-preferred side over a number of sessions. Finally, their preference is assessed again in a probe trial, where the rewarding value of the drug can be ascertained by calculating the change in preference for the least-preferred side following conditioning (in addition to water immersion, also note the use of i.p. drug injections in some zebrafish CPP studies^{184,190}). The bottom right bar diagram also shows a classic dose-response curve for nicotine in adult fish (also note that mutant heterozygous for a non-sense mutation for the acetylcholinesterase gene (*ache*^{sb55/+}) do not show place preference for 6 μ M nicotine; Brock *et al.*, unpublished observations). Top right bar diagram shows fish that display persistent compulsive drug seeking (i.e., drug seeking despite adverse consequences/punishment) following chronic exposure to ethanol or nicotine, also demonstrating long-lasting drug-specific and conserved changes in gene expression (brain genes' microarray, inset).¹⁹⁷ **(e)** The reduction in zebrafish social behavior (e.g., shorter distance between a test fish and moving shoaling images delivered on a computer screen; inset) is shown. This reduction in shoaling response is abolished by acute administration of a high concentration of alcohol, which was inactive in fish chronically pre-treated with alcohol. Withdrawal from alcohol chronic exposure also abolishes the shoaling response. Note that alcohol-induced behavioral changes in zebrafish are closely paralleled by alterations in DA levels quantified from whole brain samples using HPLC (modified from^{203,230}).




Figure 2. Continued.

three subtypes *N*-methyl-D-aspartate, AMPA (α -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid receptor) and kainate. *VGLUT2* is found in zebrafish embryos beginning at 20–24 h post fertilization (hpf) in Rohon Beard neurons and in the mediodorsal domain of the zebrafish spinal cord.¹⁹ As the fish develops, there is an increase in existing areas of expression, as well as the formation of new domains throughout the spinal cord. By 4–5 days post fertilization (dpf), *VGLUT1* and *VGLUT2* are distributed throughout the majority of midbrain, hindbrain and spinal cord.¹⁹ Examination of the expression patterns of glutamate metabotropic receptors in zebrafish reveals a similar pattern to that seen in rodents, with glutamate metabotropic receptor expression in larval and adult zebrafish found in the olfactory bulb, optic tectum, hypothalamus, cerebellum and retina.²⁰ Zebrafish also possess high-affinity excitatory amino-acid transporters (EAATs), which regulate glutamate levels and prevent excitotoxicity. Four of the five EAAT-related genes are found in the zebrafish genome, including three EAAT1 paralogs, EAAT2, EAAT3 and two EAAT5 sequences.²¹

GABA is the primary inhibitory neurotransmitter in the adult CNS, modulating neuronal excitability and also exerting its action during development. Altered GABA and glutamic acid decarboxylase (the enzyme producing GABA from glutamate) are seen in many CNS disorders, including attention-deficit/hyperactivity disorder (ADHD), bipolar disorder and major depression.^{22–24} Expression patterns of GABA are usually investigated by direct immunostaining for GABA or the two glutamic acid decarboxylase isoforms (GAD65 and GAD67²⁵), as well as by visualizing the ionotropic and metabotropic GABA_{A/C} and GABA_B receptors.²⁶ In the embryonic zebrafish, GABA-immunoreactive (ir) cells can be detected very early during development. By 16 hpf, GABA-ir cells are found in the telencephalic nucleus, the nucleus of the tract of the postoptic commissure, the nucleus of the medial longitudinal fascicle and throughout the hindbrain and spinal cord.²⁷ At 24 hpf, existing ir-cell clusters increase in cell numbers, and new clusters are located in the dorsomedial nucleus of the posterior commissure and the primordium of the hypothalamus. As the zebrafish develops, three identifiable GABAergic axonal

projections include the supraoptic tract, the tract of the postoptic commissure and the medial longitudinal fascicle. By 3 dpf, the distribution of GABA-ergic neurons in the zebrafish brain is very similar to its mammalian counterparts at a corresponding developmental stage.²⁸ GABA-ir cells are widely distributed, locating in the olfactory bulb, subpallium, posterior preoptic area, the diencephalic basal plate, the central optic tectum, torus semicircularis, ventral mesencephalic tegmentum, valvula of the cerebellum and medulla oblongata.^{19,28} In adult zebrafish, GABA is broadly distributed. GABA-ergic expression in the telencephalon is very similar to embryonic zebrafish, with GABA-ir cells in the internal cellular layer of the olfactory bulb²⁹ and all regions of the subpallium, as well as the preoptic, pretectal, ventral, thalamic, hypothalamic, preglomerular and posterior tubercular nuclei, cerebellar corpus, valvus and vestibulolateral lobe.^{25,26}

DA and NA are the primary catecholamine neurotransmitters in zebrafish. DA has a key role in modulating learning, memory, motor control, food intake and motivation. Deficits of DA neurotransmission have been implicated in bipolar disorder, schizophrenia, ADHD and drug addiction. NA-ergic neurons serve an important function in the autonomic nervous system, but also modulate attention, arousal, reward, depression and schizophrenia. In zebrafish, the catecholaminergic signaling pathways have been characterized by visualizing the temporal and spatial expression of tyrosine hydroxylase (TH)^{30,31} or the *th* transcript³² in all catecholaminergic neurons, the DA transporter (*dat*) for DA neurons and DA beta hydroxylase (*dbh*) for NA neurons,^{33,34} and the three zebrafish DA receptors D1, D2 and D3.^{35,36} Zebrafish DA-ergic neurons are first formed in the telencephalon (olfactory bulb and subpallium) and diencephalon (preoptic area, pretectum, ventral thalamus, posterior tuberculum and hypothalamus) with projections that terminate locally.^{30,33} In adult zebrafish, *th1* and *th2* are differentially expressed, with *th1* expressed in the telencephalon, ventral thalamus and pretectum, *th2* more broadly distributed in the hypothalamus, and both genes strongly expressed in the posterior tuberculum and preoptic region.³² There are some notable differences between zebrafish and mammals: for example, zebrafish lack DA-ergic cell groups in the mesencephalon^{14,33} and ventral midbrain,³⁷ but possess an additional pretectal group.¹⁸ The zebrafish NA system is very similar to mammals, with NA cell groups located in the locus coeruleus, medulla oblongata and area postrema (AP).^{38,39} Zebrafish NA neurons can already be identified by 16 hpf, with few changes between larval and adult fish.⁴⁰ Projections from NA neurons are also conserved between zebrafish and mammals,³⁷ with neurons in the locus coeruleus projecting to telencephalon, diencephalon and mesencephalon, in the medulla projecting to hindbrain and spinal cord, and in the AP projecting to the same areas as the locus coeruleus.⁴⁰

HA also has a key role in regulating diverse brain functions (for example, higher cognitive functions, circadian rhythms and locomotor activity) and modulating other aminergic systems. The zebrafish HA-ergic system is very similar to other vertebrates, with L-histidine decarboxylase and three of the four HA receptors (H1, H2 and H3) found in the brain.⁴¹ The first HA-ir neurons detected in zebrafish embryos are located in the ventral hypothalamus at ~85 hpf, and by 90 hpf the first ir-fibers can be detected in the dorsal telencephalon.⁴² In adult zebrafish, HA-ir neurons are found in the ventrocaudal region of the hypothalamus surrounding the posterior recess¹⁶ and innervate every major region in the brain (except the cerebellum).¹⁴

5-HT mediates excitatory and inhibitory neurotransmission, serving a critical role in aggression, anxiety, cognition, learning, memory, mood and sleep. Perturbation of 5-HT levels during development have been implicated in abnormal psychiatric behaviors, such as anxiety spectrum disorders, schizophrenia and depression.⁴³ Zebrafish express four 5-HT receptors (*htr1aa*, *htr1ab*, *htr1bd* and *htr2c*),^{44,45} the spatial expression of which in

the brain is primarily visualized by immuno-histochemistry and *in situ* hybridization.⁴⁶ More recently, transgenic strains expressing enhanced green fluorescent protein under control of the zebrafish raphe-specific gene *pet1* have also been developed.⁴⁷ 5-HT neurons are expressed in the spinal cord of embryonic zebrafish by 1 dpf, with additional clusters distributed throughout the brain (including the telencephalon and the hindbrain) by 5 dpf.^{30,48} In adult zebrafish, 5-HT neurons are primarily located in the raphe nucleus with ascending projections terminating in telencephalon, diencephalon, tegmentum, hindbrain and spinal cord.⁴⁷ Additional groups of anatomically distinct clusters of 5-HT-ir cells are also found in the pineal, pretectal area, posterior tuberculum and hypothalamus, reticular formation, AP and spinal cord.^{14,47,49} 5-HT neurons have also been identified by the expression of other markers, such as three tryptophan hydroxylase genes (*tph1a*, *tph1b* and *tph2*),^{50,51} two 5-HT transporters genes (*serta* and *sertb*)^{44,52} and two vesicular monoamine transporter genes (*vmat1* and *vmat2*).⁵³ Examination of the transporter genes permits more specific analysis of temporal and spatial expression pattern of 5-HT, since it can be unique, with *serta* expressed in the pineal, pretectal area and raphe, and *sertb* in the retina, posterior tuberculum/hypothalamus and AP.⁵⁴

Acetylcholine has a fundamental role in CNS functions (for example, sleep, learning/memory and attention) and dysfunctions (for example, schizophrenia, ADHD and depression). The zebrafish cholinergic system is generally similar to other vertebrates, with both muscarinic⁵⁵ and the full set of nicotinic receptors.^{56,57} The distribution of cholinergic neurons in the zebrafish brain has been primarily described by mapping the enzymes choline acetyltransferase and acetylcholinesterase (AChE). Cholinergic-ir neurons first appear immediately after hatching in the zebrafish embryo optic tectum. By 60 hpf, additional cells are seen in the preoptic area and tegmentum, and by the completion of the embryonic stage, additional reactivity is seen in the isthmic region and medulla oblongata.⁵⁸ Adult zebrafish display widespread distribution of both choline acetyltransferase and AChE. AChE-ir neurons are found in the preoptic region, periventricular layer of the optic tectum, rostral tegmental, oculomotor, trochlear nuclei of the mesencephalic tegmentum, isthmic region, octaval nuclei and motor nuclei of the cranial nerves.⁵⁹ In contrast, choline acetyltransferase-ir neurons are located in the pretectum and the tectum, except the marginal layer, the mesencephalic tegmentum, isthmic region, in Purkinje and granule cells in the cerebellum, and throughout the medulla.

Finally, it is important to consider the fish-specific genome duplication that occurred around 450 million years ago.⁶⁰ In zebrafish, this duplication resulted in expressing more neurotransmitter receptors than other species. For example, zebrafish have an estimated 122 G-protein coupled receptors for biogenic amines (such as 5-HT, NA, DA, HA, adrenaline and trace amines), compared with 57 in mouse and 44 in humans.⁶¹ Importantly, not all zebrafish receptor paralogs have been identified and characterized. For example, to date only three 5-HT receptors have been cloned, despite the existence of seven distinct 5-HT receptor families in other animals.⁴⁴ Furthermore, the functional and pharmacological properties of zebrafish receptors have not been well studied,⁶² making it more difficult to translate information across species. One reasonably well-characterized group is the alpha-2 adrenoceptors (*adra2*). The zebrafish genome contains five *adra2* receptors, three of which (*adra2a*, *adra2b* and *adra2c*) are conserved with mammals.⁶³ However, the other two subtypes, *adra2da* and *adra2d*, are zebrafish specific, whereas mammalian *adra2a-c* paralogs are not found in fish. Strikingly, despite fairly low sequence homology (~55% overall, rising to ~75% in the trans-membrane domains), human and zebrafish receptors show similar ligand-binding profiles.⁶⁴ For example, the *adra2* agonist dexmedetomidine acts as a sedative in zebrafish and other species, an effect that can be inhibited by the *adra2*-specific

antagonist atipamezole.⁶⁵ Thus, while the function of these receptors appears to be conserved across species,⁶⁵ the increased number of neurotransmitter receptors in zebrafish (vs mammals) means that their pharmacological and functional properties must be carefully examined before translating them to other species. In particular, the above-mentioned sub-functionalization of receptor subtypes implies that manipulating genes linked to psychiatric disorders may not recapitulate all symptoms of a disease, and that some novel drugs characterized in fish may have different pharmacological properties in humans (see further).

Visualizing zebrafish CNS signaling pathways

Imaging neural patterns and circuits is a key strategy in biopsychiatry research, and zebrafish offer a highly amenable experimental model for investigating the spatiotemporal distribution of neurotransmitter systems. In addition to conventional immunohistochemistry for neurotransmitter proteins or *in situ* hybridizations for the transcripts, recent genetic and technological developments permit *in vivo* screening, high-throughput analyses and whole-brain assays in this model organism.

Larval zebrafish are particularly suited for translational models of psychiatric disease because they are small, undergo external development, and are transparent (until all major organs, including the brain, have fully developed).^{10,11} This permits neurons in larval fish to be easily observed *in vivo* and then ablated or manipulated. A wide range of genetic tools, developed or adapted to zebrafish to transiently alter gene expression and generate knockout disease models, includes antisense oligonucleotides (the morpholinos), zinc finger nucleases,⁶⁶ TALEN¹³ and TILLING ('targeting induced local lesions in genomes').⁶⁷ Similar to invertebrate models, novel mutants or small molecules can then be rapidly screened (in the thousands) using a combination of conventional fluorescent or confocal microscopy with automated fluidic screening platforms.^{68,69}

Complementing multiple genetic methods, a suite of sophisticated tools also help elucidate the anatomical alterations associated with novel phenotypic alterations. These techniques include *in vivo* monitoring of neural activity, functional analyses of specific neurons and tracing of neural circuits. Genetically encoded calcium indicators (GECIs, such as the GCaMP reporter constructs) fluoresce when calcium is present, thereby serving as an indicator for action potential firing.⁷⁰ By driving expression with the Gal4/UAS⁷¹ or transposon-based Tol2 systems,⁷² GECIs and optogenetic constructs, such as channelrhodopsin and halorhodopsin, can easily be inserted into the zebrafish genome to target specific cellular populations and perform functional analyses. With advanced microscope techniques (for example, light-sheet microscopy), functional imaging of the entire brain of a larval zebrafish can be performed in < 1.5 s,⁷³ a task unattainable in other vertebrate models. In contrast to rodents, this procedure is also almost entirely non-invasive, and does not require attaching fiber optic cables. Albeit currently used in zebrafish to address fundamental questions (for example, sensory processing⁷⁴ or motor control⁷⁵), this optogenetic toolbox can be similarly employed in this organism to model altered neural activity associated with psychiatric disease in humans.⁷⁶

Fluorescent markers (for example, green fluorescent protein) can also be utilized to resolve the neuronal projections and formation of neural circuits in larval zebrafish. Using combinatorial expression of several fluorescent proteins to individually label neurons and their projections,⁷⁷ Brainbow technology can be particularly useful for examining disease phenotypes related to miswired neural circuits (for example, autism⁷⁸). Briefly, Brainbow visualizes synaptic circuits by labeling neurons genetically, using Cre/lox recombination to create a stochastic choice of expression between several fluorescent proteins.^{79–81} Originally developed in mice, this transgenic technology has been successfully applied to

other model organisms, including zebrafish,⁷⁷ to map multiple individual cells within a population and provide a large-scale dissection of CNS circuitry.^{80,81} A recent modification of Brainbow, the ZebraBow, has been introduced for *in vivo* multicolor imaging in zebrafish.⁸² Applying several new transgenic strains for ubiquitous or tissue-specific multicolor labeling, this approach maximizes color diversity by optimizing Cre activity, and can be particularly useful for long-term color-based anatomical and lineage analyses in a wide variety of zebrafish tissues, including neurons and glia.⁸²

Furthermore, while calcium and voltage sensors are fundamental to functional analyses in the larval zebrafish brain, they do not specifically measure neurotransmitter release or concentration (typically measured by cyclic voltammetry, HPLC or ELISA). However, more recent specific markers enable *in vivo* measuring of neurotransmitters, based on placing a fluorescent label very close to the synapse, labeling the synaptic vesicle content or directly measuring extracellular neurotransmitters. Fluorescent false neurotransmitters are optical tracers that permit visualization of neurotransmitter release and uptake at presynaptic terminals.⁸³ Initially developed for the DA-ergic system, fluorescent false neurotransmitters are fluorescent substrates for targeting vesicular monoamine transporter 2, carrying monoamine neurotransmitters to synaptic vesicles. Fluorescent false neurotransmitter has also recently been extended to be a pH-responsive DAT substrate, which is able to identify DA cell bodies and dendrites and measures DAT activity and DA release at individual synapses.⁸⁴

Several optical sensors have also been developed to measure excitatory synaptic activity in the brain. The glutamate optical sensor is a fluorescent indicator made up of the glutamate-binding domain of the AMPA and a small fluorescent molecule bound near the glutamate-binding pocket.⁸⁵ When bound to glutamate the glutamate optical sensor changes its fluorescence intensity. Intensity-based glutamate-sensing fluorescent reports are a single-wavelength glutamate sensor that responds to glutamate *in situ* and therefore directly reports excitatory synaptic release.⁸⁶ This sensor can be genetically encoded and used in zebrafish to demonstrate the spatial organization of direction-sensitive synaptic activity in the optic tectum.⁸⁶ Finally, fluorescent indicators for acetylcholine volume transmission have also been created. Cell-based sensors of neurotransmitters are cultured cells that co-express M1 muscarinic receptors and the GECI TN-XXL, and (when implanted into the brain) provide a fluorescent readout of acetylcholine levels.⁸⁷

Imaging neurotransmitter systems in the adult zebrafish brain are significantly more difficult due to its increased size and lack of transparency. While transparent zebrafish strains (for example, *absolute*, *nacre* and *casper*, Figure 1a) exist,⁸⁸ the brains are still relatively opaque making it difficult to image the various photo-proteins described above. As a result, studies have typically been limited to sectioning the brain. However, numerous methods may render a fixed brain transparent and, when imaged with long working distance objectives, permit the examination of neurotransmitters *in situ*.^{89–96} These techniques use hydrophobic or hydrophilic chemical mixtures to remove lipids and reduce light scattering to visualize endogenous fluorescent markers in the brain.

Additional zebrafish neuroimaging tools

Several gene transcription factors, such as Pax3 (pair box 3), Fos and Otp (Orthopedia), represent a useful tool for visualizing CNS development and function in zebrafish. For example, Pax3 is a transcription factor that regulates cell proliferation, migration and neural development. Its gene is not only highly conserved (at the sequence level) between zebrafish and mammals, but also shows similar pattern of neural expression during embryogenesis.^{97–99} Thus, analyses of Pax3 expression in zebrafish can be particularly

Table 1. Examples of a diverse group of psychiatric disorders (Figure 2) which can be modeled in zebrafish (see Kalueff *et al.*¹³⁵ for a comprehensive catalog of zebrafish normal and pathological behaviors)

Selected disorders	Zebrafish phenotypes	References
Attention deficit hyperactivity disorder (ADHD)	Impulsive swimming with hyperactive locomotion and impulsivity-like bursts of accelerated swimming, which can be reduced by anti-ADHD drugs (Figure 2a)	125,126,217,218,231
Aggression	Social deficits (disrupted shoaling behavior), increased aggressive behaviors (follows, chases and biting), mirror-induced aggression (MIA) (Figure 2b)	135,139,161
Post-traumatic stress disorder (PTSD)	Reduced exploration following stress, increased avoidance (e.g., neophobia; Figure 2c), erratic behavior and freezing, elevated cortisol and brain <i>c-fos</i> (all highly sensitive to a wide range of anxiolytic and anxiogenic agents), social deficits (e.g., disrupted shoaling behavior)	144,146,147,232
Substance abuse, addiction (reward-related behavior)	Robust conditioned preference for commonly abused substances, compulsive drug seeking (i.e., despite adverse consequences; Figure 2d), withdrawal syndrome, stimulus-induced relapse following extinction, altered social behavior (Figure 2e)	197,198,201,202,205,233–235

useful for imaging early CNS development,^{97,98} which is relevant to modeling many neurodevelopmental disorders.

The homeodomain transcription factor *Otp* is an important regulator of neuroendocrine and dopaminergic cells in vertebrate species.^{100,101} *Otp* has two paralogs in zebrafish (*otpa* and *otpb*),¹⁰¹ and genetic manipulations suppressing *Otp* deplete certain dopaminergic neurons, enabling a functional dissection of dopaminergic contribution to CNS circuitry in early larval zebrafish.¹⁰²

Likewise, an early proto-oncogene *c-fos* serves as a well-established marker of neuronal activation in various species, including humans,¹⁰³ rodents^{104,105} and zebrafish.¹⁰⁶ For example, *c-fos* expression analyses of whole-brain zebrafish samples can be used to assess neuronal activation globally (for example, reflecting CNS excitation during epilepsy^{106,107} or following anxiogenic/anxiolytic or hallucinogenic treatments^{108,109}). Region-specific analyses of *c-fos* expression in zebrafish brain offer a more detailed functional mapping of neural circuits during various behavioral tests.¹¹⁰ For instance, in the light-dark test, *c-fos* neuronal activity mapping implicated several brain regions, including several telencephalic areas (the teleost homologs of the mammalian amygdala and striatum), during zebrafish 'anxiogenic' light avoidance behavior.¹¹⁰ Characterizing neuronal activity induced in zebrafish by acute amphetamine, drug-seeking or the light avoidance behavior, quantitative analyses of *c-fos* expression can also be combined with neuronal subtype-specific markers at single-cell resolution.¹¹¹ Showing the recruitment of the medial and the lateral pallium (structures homologous to the mammalian amygdala and hippocampus), these data suggest evolutionarily conserved function of amygdala-like structures in positive emotions and motivated behavior in zebrafish and mammals.¹¹¹ Collectively, this evidence strongly supports the utility of *c-fos* and other biomarkers in visualizing normal and aberrant CNS development and functioning.

Finally, with the increased use of zebrafish in disease genetics research, a quantitative voxel-based technique may help compare disease models with wild-type zebrafish. Magnetic resonance imaging is an inherent 3D tool that facilitates investigations in morphology, connectivity and function of the brain. Due to its 3D nature, multiple data sets can be mapped to a standard coordinate space and registered to produce a model representative of a population (thus, limiting individual morphological variance and instead demonstrating the average morphology of the entire population). Data collected from magnetic resonance imaging can include mean volumes of individual brain regions, the stereotaxic locations of individual structures and major fiber bundles, and functional activity. Importantly, magnetic resonance imaging models can also serve as a canonical space for registration of other imaging data sets. For example, recent whole-brain immunostaining and light microscope imaging (for example, SeeDB or Clarity), optical projection tomography and

microCT can all be registered to the same stereotaxic space, collectively enabling quantitative analysis of a wide range of markers in zebrafish brain.

ZEBRAFISH MODELS OF BRAIN DISORDERS

Bridging molecular events with behavior

Many psychiatric disorders are moderately or highly heritable, and have developmental trajectories.^{112,113} As behavior is controlled by activity within neural circuits, a major challenge is to map the genotype–phenotype relationship to understand how genetic variations affect neural circuits that control behavior. Various molecular tools have been effectively used for uncovering brain mechanisms in zebrafish. For example, in addition to traditional *N*-ethyl-*N*-nitrosourea-induced¹¹⁴ and viral-vector-mediated¹¹⁵ mutagenesis, new sophisticated 'gene-breaking transposon' (GBT)¹¹⁶ screens and 'clustered regularly interspaced short palindromic repeats' approaches for generating targeted zebrafish mutants have been developed.¹¹⁷ Combined with TILLING and TALEN tools, this makes the process of exploring genotype–phenotype relationships in zebrafish better targeted and less labor-intensive. With the establishment of translationally relevant zebrafish behavioral assays^{9–11} comes the ability to apply these molecular techniques to modeling psychiatric diseases. Here, we discuss how various zebrafish models can be utilized to target selected psychiatric disorders and their neurodevelopmental trajectories. ADHD, aggression, post-traumatic stress and drug abuse (Figure 2) were selected as examples, reflecting a wide and diverse spectrum of psychiatric disorders that can be modeled using the zebrafish.

Modeling ADHD

Zebrafish have recently been utilized to investigate some aspects of ADHD (Table 1; Figure 2a), a common psychiatric disorder affecting ~3–5% of children worldwide. ADHD symptoms (inattention, hyperactivity and impulsivity) persist into adulthood in ~50% of cases, reducing the patients' quality of life, academic performance and sociality.^{118,119} Altered DA, NA, 5-HT and glutamate neurotransmission in the prefrontal cortex, striatum, parietal cortex, vermis and the inferior lobes of the cerebellum are implicated in ADHD.^{120–123}

An interesting example of using larval zebrafish to model this disorder involves a human gene *LATROPHILIN3* (*LPHN3*), identified as candidate ADHD gene by linkage analysis.¹²⁴ In zebrafish, its homolog *latrophilin3.1* (*lphn3.1*) has a developmental role, and is expressed in differentiated neurons throughout the brain up to 6 dpf. The *lphn3.1* morphants display increased distance swum at 6 dpf (Figure 2a)^{125,126}—the hyperactivity phenotype also maintained during the night, demonstrating a permanent increase in

locomotion compared with control animals. Morphant larvae also show more bursts of accelerated swimming, indicative of motor impulsivity. Both hyperactivity and motor impulsivity can be rescued by the prototypical anti-ADHD drugs methylphenidate and atomoxetine (Table 1). While acute 1-h treatment with either drug had no effect on control-injected larvae at the doses used, it rescued morphant behavior bringing locomotion back to control levels. The *lphn3.1* morphants also display fewer cells in the posterior tuberculum, a prominent group of DA-ergic neurons in the ventral diencephalon that control larval locomotion and project both anteriorly and posteriorly, but mainly to the spinal cord.^{101,127} Thus, despite the difficulty of fully modeling ADHD in zebrafish, research in these animals shows an important role for *lphn3.1* in DA-ergic neuron development¹²⁸ (also see similar data in mice,¹²⁹ proving the feasibility of translating findings from zebrafish to other species).

Exposure of zebrafish embryos to chemicals during development may also provide insights into possible environmental causes of ADHD. For example, prenatal exposure to perfluorooctane sulfonate has been linked to clinical adolescent ADHD,¹³⁰ and treating zebrafish embryos with this agent during development elicits ADHD-like behaviors in 6-day-old larvae (i.e., hyperactivity, motor impulsivity and hyper-responsiveness to startle).¹³¹ This phenotype is reminiscent of the behavior of *lphn3.1* larvae and strikingly, these behavioral changes can be rescued by dexamfetamine, implicating DA signaling in this phenotype. Zebrafish are also used to study the consequences of long-term exposure to methylphenidate, displaying elevated DA, NE and 5-HT in the larval brain, and lasting behavioral changes (for example, impaired predatory escape and learning).¹³² Such ease of conducting embryonic manipulations and behavioral experiments in zebrafish strongly supports their potential for studying ADHD and related disorders.

Zebrafish models of aggression

Aggression, an adaptive animal behavior to monopolize resources, is essential for the organism's fitness and survival. However, pathological aggression is part of various brain disorders, and merits further studies *in vivo* models. Although aggression has long been studied in fishes, the genetic and neurobiological tools available in zebrafish make them an ideal species to study aggression. Zebrafish display aggression as a series of stereotypic body postures (for example, undulating body movements, short slaps of the caudal fin and bouts of swimming and biting directed against an opponent) which can easily be measured in the laboratory.¹³³ Aggressive incidents follow a highly structured pattern,¹³⁴ from extending the fins to circling, chasing and attempting to bite, further progressing to more frequent chases, bites and attempts to strike the opponent. The fish that loses such as encounter becomes submissive, exhibiting postures that include fleeing, freezing and retreating.^{134,135} In laboratory zebrafish, aggression can be measured by recording the interaction of two fish in a tank, or the interaction of a single fish with its own reflection in the mirror exposure test (mirror-induced aggression, MIA; Figure 2b).^{133,134}

Mounting evidence shows that zebrafish aggression has a genetic basis, and its moderate heritability suggests that environmental influences (for example, maternal care, rearing conditions and environmental complexity) also contribute to this behavior. For example, aggression/boldness profiles in several wild-type zebrafish strains (TM1, SH and Nadia) show strain differences in MIA, supporting the genetic basis of aggression in zebrafish.¹³⁶ A group of 40 genes linked to aggression have been identified in fish, with differences in expression level depending on both the area of the brain analyzed and the sex.¹³⁷ HPLC analyses of adult zebrafish brains following an aggressive interaction reveal altered DA, 5-HT and their metabolites in the telencephalon,

diencephalon and optic tectum. Other important brain areas include the preoptic area of the anterior hypothalamus¹³⁸ and the periventricular nucleus of the inferior hypothalamus.¹³⁹ Notably, similar brain areas have been linked to agonistic behavior in other vertebrates, highlighting the translatability of zebrafish research in this field. Like in other vertebrates, fish aggression is controlled by similar neurotransmitters, peptides and hormones, including DA, 5-HT, HA, 17 α -ethynylestradiol and arginine vasopressin/vasotocin. For example, addition of estrogen to tank water alters zebrafish aggression, implicating certain pathways and circuits,^{133,140} as reduced aggression in dominant male zebrafish correlates with altered 5-HT, DA, arginine vasopressin levels and the expression of the somatostatin pathway genes.¹⁴¹

A recently characterized novel aggressive *spiegeldanio* mutant zebrafish (*spd*, Figure 1a)¹⁴² harbors a mutation in the *fgf receptor 1a* gene, which disrupts the fibroblast growth factor signaling in the brain. Fibroblast growth factors are a large family of secreted signaling molecules with important roles during development.¹⁴³ Homozygous *spd* mutants show unaltered locomotion but increased aggression, boldness and exploration (Figure 2b), correlated with elevated breakdown of HA in the brain, and reduced neurotransmitter signaling in the hypothalamic periventricular nucleus. Importantly, pharmacological increase in HA rescues all three behavioral alterations in *spd* mutants.¹³⁹ Although *spd* zebrafish are always more aggressive than wild-type siblings, their behavior differs between the laboratories (for example, similar boldness but higher aggression in mutants raised in Germany vs France).¹³⁹ Taken together, these results demonstrate the importance of environmental influences on aggression, and that experimental modulation of selected target genes can strongly affect zebrafish aggressive phenotypes.

Post-traumatic stress disorder

Severe stress is the main cause of PTSD and other trauma/stress-related disorders. Zebrafish have recently been proposed as potential models of PTSD and trauma/stress-related disorders.^{144,145} The extensive, complex repertoire of behavioral responses of zebrafish to a wide variety of stress-evoking stimuli closely parallels that observed in mammalian models of PTSD (Figure 2c; Table 1).¹⁴⁵ For example, various PTSD-relevant phenotypes identified in zebrafish include robust anti-predatory responses^{146,147} and inhibitory or active avoidance conditioning.¹⁴⁸ Paralleling mammalian tests, an increasing number of zebrafish paradigms relevant to PTSD continue to be developed, based on fear conditioning^{148–152} and exposure to aggressive conspecifics,^{134,138} predators or other acute stressors (Figure 2).^{153–155} Moreover, the highly social nature of zebrafish^{156–158} offers further potential to examine pathological social phenotypes, representing the serious adverse clinical effects of PTSD/trauma/stress-related disorder. For instance, experimentally evoked disorganization in the social structure of zebrafish groups (shoals)^{158–161} may represent an interesting PTSD-related 'social withdrawal' phenotype.¹⁴⁵

The physiology of zebrafish also supports their suitability for modeling PTSD and other stress-related CNS disorders.^{120,121} Zebrafish rapid development enables time- and cost-efficient modeling of the developmental and 'temporal' pathogenetic aspects of PTSD. Moreover, given the strong involvement of the neuroendocrine stress axis in PTSD,¹⁶² the cortisol-driven neuroendocrine axis of zebrafish makes them well suited for studying glucocorticoid regulation in stress and PTSD.^{163,164} For example, zebrafish *gr*³⁵⁷ mutants, possessing non-functional glucocorticoid receptors, have recently offered a model of glucocorticoid resistance to study the developmental effects of glucocorticoid signaling deficits on stress physiology and related behaviors.^{165,166} Furthermore, an extensive array of homologs of clinical biomarkers of PTSD has also been identified in zebrafish, including corticotropin releasing factor,^{167,168} glucocorticoid receptor,^{169,170}

adrenocorticotrophic hormone,^{171,172} neuropeptide Y,^{173,174} catechol O-methyltransferase^{5,175,176} and monoamine oxidase A/B.^{171,177} Finally, the sensitivity of zebrafish sympathetic and parasympathetic responses (for example, altered heart rate) to pharmacological manipulation has recently been established,¹⁷⁸ offering the potential to study autonomic biomarkers of PTSD-relevant drug treatments. Overall, zebrafish emerge as a promising new model to complement traditional rodent models of PTSD.^{10,144,145}

Addiction, substance abuse

Zebrafish are increasingly utilized as a preclinical model to investigate molecular mechanisms of substance abuse. Drugs of abuse typically exert their effects by modulating DA pathways in the midbrain,¹⁷⁹ and the ascending DA systems (VTA-NAc reward pathway) in zebrafish share functional homology with those of mammals.^{180,181} Zebrafish express orthologs for many addiction-related human genes,¹⁸² including the majority of the DA-, 5-HT- and cholinergic receptor gene families. Zebrafish also display robust reward responses to translationally relevant doses of common drugs of abuse, including stimulants (cocaine,¹⁸³ amphetamine,¹⁸⁴ nicotine¹⁸⁵ and caffeine; Brock and Brennan, unpublished data), ethanol,¹⁸⁵ opiates (morphine,¹⁸⁶ fentanyl) and general anaesthetics (tetracaine and ketamine; Brock and Brennan, unpublished data). As drug-seeking behavior is evolutionarily conserved, zebrafish represent a powerful *in vivo* model to assess the contribution of genetic factors across specific endophenotypes or drugs of abuse.

The first systematic screens relevant to addiction, examining mutant alleles affecting catecholaminergic neurons,¹⁸⁷ showed that DA/hypothalamic neuronal development is regulated by forebrain embryonic zinc finger-like protein (*fezl*),¹⁸⁸ the mutation of which in zebrafish reduces opiate (but not food) reward.¹⁸⁹ Subsequent studies performed conditioned place preference (CPP, Figure 2d) screens of *N*-ethyl-*N*-nitrosourea-mutagenized F₂ adult fish for cocaine and amphetamine reward,¹⁸³ and found a number of families showing heritable reduced reward sensitivity.¹⁸⁴ For instance, the *nad*^{+/-} mutant zebrafish show insensitivity to the rewarding effects of amphetamine.¹⁹⁰ Microarray analysis, followed by qPCR confirmation, identified multiple genes that were upregulated/downregulated in the *nad* mutant, particularly relating to neurogenesis and brain development. Similar approaches have been applied to assess drug-induced reward and to identify novel genes and molecular pathways involved in zebrafish reward.^{184,189,191–194} For example, mutants heterozygous for a non-sense mutation in the AChE (*ache*+/-) gene¹⁸⁴ display reduced amphetamine-induced reward in a modified CPP paradigm, demonstrating the conservation of cholinergic regulation of reward pathways.

Although sensitivity to the rewarding effects of drugs is a fundamental aspect of vulnerability to addiction, addiction itself is characterized by the progression from occasional or transient to habitual drug taking, compulsive drug seeking and a tendency to relapse even after prolonged abstinence.¹⁹⁵ Both drug-seeking despite adverse consequences, and reinstatement of drug seeking following extinction are commonly used to model these behaviors in laboratory species.¹⁹⁶ Recent studies demonstrate persistent drug-seeking despite punishment¹⁹⁷ and relapse¹⁹⁸ in zebrafish chronically exposed to either nicotine or ethanol. Microarray analysis in this model reveals shared changes in gene expression, consistent with conservation of neuroadaptations and the establishment of dependence in zebrafish.¹⁹⁷ Collectively, these findings show that genetic and neural pathways are well conserved between mammals and fish, and that fish employ similar molecular adaptations following chronic drug exposure.¹⁹⁹

In addition to classical screening techniques in adult fish to identify strains with aberrant drug responsiveness, more targeted

approaches help identify specific genes involved in drug sensitivity. Fluorescently tagged GBT mutagenesis has been used to examine molecular influences on a measure of nicotine addiction, behavioral sensitization.¹¹⁶ This insertional method represents a leap forward from classic mutagenesis, enabling both visual identification of carriers and rapid identification of the disrupted gene. In a screen of GBT mutant strains for nicotine behavioral sensitization, 5-dpf larvae were exposed to nicotine acutely or 8 h following previous administration of this drug. Acute exposure caused an overt locomotor response, significantly accentuated in the larvae pre-treated with nicotine. GBT-mutagenized families with diminished acute and sensitized responses to nicotine, identified in this screen, included the *bdav* mutants with a truncation in chaperonin-containing protein 8 (*ccb8*, linked to nicotinic acetylcholine receptor), and the *hbog* zebrafish with a mutant ortholog of the GABA-B receptor (*gabbr1.2*).^{59,116} Furthermore, the behavioral phenotypes in these strains were reverted by cre-mediated recombination, revealing specific loci in zebrafish that may lead to translational targets for the treatment of tobacco addiction.¹¹⁶

Alcohol

Alcohol is the drug with the longest history of use and abuse in human society. Although legally accepted and culturally tolerated, it is a dangerous drug, and alcohol-related disorders present an enormous burden for the patient and society.²⁰⁰ Alcohol is a complex drug, affecting multiple mechanisms and behaviors in a dose- and use-dependent manner. This complexity is one of the reasons why effective therapies are still lacking, and alcohol-related disorders remain a serious medical concern. In general, findings from zebrafish alcohol research can be evaluated by three main criteria of a translationally acceptable model—face, construct and predictive validity. Addressing face validity, recent studies suggest that zebrafish respond to alcohol in a manner highly homologous to that of humans (Figure 2e).¹³³ For example, acute administration of alcohol makes fish agitated at intermediate doses, but lethargic at higher doses.²⁰¹ Zebrafish exhibit motor impairment in response to high doses of acute alcohol (for example, 1%), but show elevated aggression in response to intermediate (for example, 0.25–0.50%) doses.¹³³ Acute alcohol also impairs zebrafish response to social stimuli in a linear dose-dependent manner (Figure 2e), a reaction that is strikingly similar to that in humans.^{133,202} Chronic exposure to alcohol evokes tolerance in zebrafish, paralleling clinical findings.²⁰³ Withdrawal from alcohol also leads to responses (for example, elevated motor activity, anxiety, incoordination and reduced shoaling), phenotypically similar in zebrafish and humans (Figure 2e).^{203, 204} Likewise, embryonic alcohol exposure in zebrafish further parallels the clinical effects of embryonic alcohol exposure, including the social and learning impairment observed in fetal alcohol spectrum disorders.^{10,205}

Mounting evidence also suggests significant construct validity of zebrafish alcohol models. For example, following acute or chronic alcohol treatment, zebrafish show similar neurochemical alterations in the brain, compared with humans. Acute alcohol induces a rapid increase in brain DA, which is blunted after chronic alcohol exposure.²⁰³ Likewise, withdrawal from alcohol increases brain DA levels,²⁰³ whereas chronic alcohol exposure alters the expression of nearly 2000 genes in zebrafish, many of which have been implicated in human alcoholism.²⁰⁶

One of the greatest challenges for modeling the effects of alcohol in zebrafish is establishing predictive validity—the extent to which drugs (with particular known effects on human CNS) act in the same way in the zebrafish system. While the model system is expected to show efficacy of human drugs in the predicted manner, only a limited number of compounds expected to mediate the effects of alcohol has been assessed in zebrafish.

However, zebrafish have largely been shown to respond in a predictable manner to such compounds, paralleling clinical effects and suggesting a conservation of the main neurotransmitter signaling pathways between zebrafish and humans. Overall, while future studies are needed to further assess their face, construct and predictive validity as models of alcohol-related disorders, zebrafish already shows a significant potential in this field.

CURRENT CHALLENGES

The advantages and limitations of zebrafish in biopsychiatry research have recently been discussed in the literature.^{9–11,174} However, despite recent progress in zebrafish neurobiology, many important questions remain to be addressed. For example, the logical step in the development of zebrafish models with translational relevance to human addiction is to apply systematic screening of mutagenized families, to characterize genetic risk factors associated with drug abuse. Humans with addiction and other linked psychiatric disorders (for example, compulsive gambling, obsessive compulsive disorder, Tourette's syndrome or ADHD) often display deficits in general cognitive or executive function^{207–210} (which becomes relevant to studying the molecular basis of cognitive dysfunction linked to addiction). Emerging as a key endophenotype for cocaine addiction,^{211,212} trait impulsivity represents several behavioral constructs, including inability/unwillingness to wait, choice of smaller but more immediate (over larger, delayed) rewards, and motor impulsivity (failure to inhibit a pre-potent response).²¹³ In compulsive drug use in rodents, animals with high levels of motor impulsivity (assessed by premature responses on a test of sustained attention, the 5-CSRTT (five-choice serial reaction time task)²¹⁴) also show compulsive cocaine seeking²¹¹ and increased relapse after extinction training.²¹⁵ This task has recently been adapted for zebrafish,²¹⁶ resulting in a fully automated model,²¹⁷ validated pharmacologically by showing dose-dependent reductions in fish impulsivity following atomoxetine,²¹⁸ similar to rodents.²¹⁹ Collectively, this provides the exciting prospect of using three-generation screens (discussed above) or insertional transposon technologies to study impulsivity, thus fostering the search for the molecular basis of psychiatric disorders linked to impulsivity and other reward-related traits.

Likewise, the ability to develop novel zebrafish models of ADHD is also interesting and translationally relevant. However, the suitability of zebrafish models (for example, mutants lacking *lphn3.1* function) as an ADHD-like model must be further established by testing other behaviors, such as performance in the 5-CSRTT.²¹⁷ Although such a complex task will require a stable *lphn3.1* line to be established, this approach would be ideal to reinforce the utility of *lphn3.1* and other mutant fish as a tool to study the neurobiology of ADHD.

Because the formation, position and function of neurotransmitter signaling pathways sometimes differ between zebrafish and other vertebrates, comparative studies become important. For example, as already mentioned, similar to other teleosts, zebrafish have undergone a partial genome duplication, and some genes found in a single copy on the mammalian chromosome have two copies in the zebrafish chromosome.²²⁰ Such genetic redundancy can make it more difficult to analyze a disease gene's function (for example, if the effect of a mutated gene is masked by the unaltered 'sister' gene). While the function of some receptors and transporters is conserved across species,^{17,65} the increased spectrum of neurotransmitter signaling-related molecules in zebrafish (vs mammals) suggests that their functional properties merit further scrutiny in genetic and pharmacological studies. However, this redundancy can also present an opportunity (for example, permitting the analysis of the effects of genes whose null mutations lead to developmental lethality if in a single copy, a common problem in mouse knockout studies).

From an evolutionary perspective, there are also important physiological differences between mammals and zebrafish. For example, fish are poikilotherms and mammals are homeothermic animals, and substantial differences exist in metabolism and organ systems between species.²²¹ Likewise, there are also well-known differences in brain development between fish and mammals, which will be only briefly mentioned here. For instance, instead of the laminar cortex, telencephalic nuclei in teleost fishes have evolved as the pallium, containing similar but simpler neural circuits, compared with the mammalian cortex, which also has some additional circuits²²² (note, however, recent data showing that emotions and motivated behavior in zebrafish converge on an amygdala-like structure, similar to their mammalian counterparts¹¹¹). Another interesting example is the habenula—a key relay nucleus connecting the forebrain with the brainstem and modulating monoaminergic mechanisms of behavior and cognition.²²³ Although the development of its circuits appears to be conserved phylogenetically, some species differences in habenular asymmetry exist between fish and mammals²²³ (note, however, mounting evidence showing that the zebrafish habenula, like in humans,²²⁴ has an important role in the regulation of affective behavior and aversive memories^{152,225–227}). While not within the scope of this review, such physiological and neurobiological differences between species must be considered critically when translating zebrafish responses into human disorders.²²¹ Together with some genetic differences between species¹² and poorly characterized protein function differences (stemming from divergent evolution), this represents a serious current challenge for zebrafish molecular psychiatry.

Finally, perhaps the biggest challenge in multidisciplinary zebrafish studies is the behavioral models and tools.^{9,10} Despite a rapidly growing research in this field, the zebrafish is still a new model in behavioral neuroscience.¹¹ Will zebrafish analyses keep pace with the rapid advancement of molecular and neurobiological tools developed for this species? While the answer to this question is unclear, recent evidence offers grounds for optimism. For example, the rapidly increasing computer processing power, coupled with constantly improving 2D and 3D video-recording and imaging tools, generates behavioral data with ever increasing quality, precision and spatiotemporal resolution. Simultaneous video-tracking of a large number of fish (either swimming in separate compartments; Figure 2b or moving together in a group/shoal)^{10,160} now became a reality. As behavioral quantification becomes more precise, so does the delivery of stimuli that induce the behavioral responses. Because both stimulus delivery and behavioral response quantification are now increasingly computerized and standardized, the behavioral paradigms in zebrafish become more automated and scalable. Thus, running such multiple tasks in parallel makes even slow learning protocols high-throughput and efficient. Enhancing high-throughput screening of mutants and drugs that affect complex behavioral responses such as phenomics-based approaches in zebrafish foster further experimental modeling of human brain disorders.

CONCLUSION

Uncovering the molecular processes that underlie the formation of neural circuits and their activity to control the behavior is key to the understanding and treatment of psychiatric diseases.^{6,7} Recognizing the importance of zebrafish to molecular psychiatry is the critical step in our continued use of this model organism to study neurobiological mechanisms of human brain disorders. In addition to the growing availability of genetic and pharmacological assays and probes in zebrafish, there is also an urgent need for new behavioral assays and imaging neural pathways and circuits implicated in various pathological states in this species.^{228,229}

The availability of genetic, electrophysiological and optogenetic tools, as well as multiple mutant and transgenic strains, physiological biomarkers and robust protocols to measure behavior, collectively supports zebrafish potential for modeling psychiatric disorders. As both larval and adult zebrafish are now rapidly becoming key model organisms in neuroscience research, they represent a useful novel species for translational molecular psychiatry and CNS drug discovery.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ACKNOWLEDGMENTS

The authors' research has been supported by grants from the National Institutes of Health (NIAAA AA015325-01A2 to RG, NIDA DA030900-02 to AVK and NINDS NS077295-01 to JFPU), the EU FP7 Framework Program (Grant 602805 to WHJN), as well as the National Center for the Replacement, Reduction and Refinement of Animals in Research (NC3Rs, UK, G1000053 to CHB) and the Medical Research Council (MRC, UK to CHB). CHB is a Royal Society (UK) Industrial Research Fellow. The study was coordinated and facilitated by the International Zebrafish Neuroscience Research Consortium (ZNRC). AVK is the Director of ZENEREI Institute and Chair of ZNRC.

REFERENCES

- WHO. *The Global Burden of Disease: 2004 Update*. WHO (World Health Organization): Geneva, Switzerland, 2008.
- Griebel G, Holmes A. 50 years of hurdles and hope in anxiolytic drug discovery. *Nat Rev Drug Discov* 2013; **12**: 667–687.
- Gonzales ML, LaSalle JM. The role of MeCP2 in brain development and neurodevelopmental disorders. *Curr Psychiatry Rep* 2010; **12**: 127–134.
- Xu B, Ionita-Laza I, Roos JL, Boone B, Woodruff S, Sun Y *et al*. De novo gene mutations highlight patterns of genetic and neural complexity in schizophrenia. *Nat Genet* 2012; **44**: 1365–1369.
- Mitchell K. The miswired brain: making connections from neurodevelopment to psychopathology. *BMC Biol* 2011; **9**: 23.
- Duman RS, Heninger GR, Nestler EJ. Molecular psychiatry. Adaptations of receptor-coupled signal transduction pathways underlying stress- and drug-induced neural plasticity. *J Nerv Ment Dis* 1994; **182**: 692–700.
- Nestler EJ. The origins of molecular psychiatry. *J Mol Psychiatry* 2013; **1**: 3.
- Kalueff AV, Wheaton M, Murphy DL. What's wrong with my mouse model? Advances and strategies in animal modeling of anxiety and depression. *Behav Brain Res* 2007; **179**: 1–18.
- Kalueff AV, Echevarria DJ, Stewart AM. Gaining translational momentum: More zebrafish models for neuroscience research. *Progr Neuropsychopharmacol Biol Psychiatry* 2014 pii **S0278-5846**: 00037–2.
- Kalueff AV, Stewart AM, Gerlai R. Zebrafish as an emerging model for studying complex brain disorders. *Trends Pharmacol Sci* 2014; **35**: 63–75.
- Stewart AM, Braubach O, Spitsbergen J, Gerlai R, Kalueff AV. Zebrafish models for translational neuroscience research: from tank to bedside. *Trends Neurosci* 2014; **37**: 264–278.
- Howe K, Clark MD, Torroja CF, Torrance J, Berthelot C, Muffato M *et al*. The zebrafish reference genome sequence and its relationship to the human genome. *Nature* 2013; **496**: 498–503.
- Zu Y, Tong X, Wang Z, Liu D, Pan R, Li Z *et al*. TALEN-mediated precise genome modification by homologous recombination in zebrafish. *Nat Methods* 2013; **10**: 329–331.
- Kaslin J, Panula P. Comparative anatomy of the histaminergic and other aminergic systems in the zebrafish (*Danio rerio*). *J Comp Neurol* 2001; **440**: 342–377.
- Sallinen V, Sundvik M, Reenila I, Peitsaro N, Khrustal'ov D, Anichtchik O *et al*. Hyperserotonergic phenotype after monoamine oxidase inhibition in larval zebrafish. *J Neurochem* 2009; **109**: 403–415.
- Sundvik M, Panula P. Organization of the histaminergic system in adult zebrafish (*Danio rerio*) brain: neuron number, location, and cotransmitters. *J Comp Neurol* 2012; **520**: 3827–3845.
- Maximino C, Herculano AM. A review of monoaminergic neuropsychopharmacology in zebrafish. *Zebrafish* 2010; **7**: 359–378.
- Filippi A, Mueller T, Driever W. Vglut2 and gad expression reveal distinct patterns of dual GABAergic versus glutamatergic cotransmitter phenotypes of dopaminergic and noradrenergic neurons in the zebrafish brain. *J Comp Neurol* 2014; **522**: 2019–2037.
- Higashijima S, Mandel G, Fetcho JR. Distribution of prospective glutamatergic, glycinergic, and GABAergic neurons in embryonic and larval zebrafish. *J Comp Neurol* 2004; **480**: 1–18.
- Haug MF, Gesemann M, Mueller T, Neuhaus SC. Phylogeny and expression divergence of metabotropic glutamate receptor genes in the brain of zebrafish (*Danio rerio*). *J Comp Neurol* 2013; **521**: 1533–1560.
- Rico EP, de Oliveira DL, Rosemberg DB, Mussulini BH, Bonan CD, Dias RD *et al*. Expression and functional analysis of Na(+)-dependent glutamate transporters from zebrafish brain. *Brain Res Bull* 2010; **81**: 517–523.
- Brambilla P, Perez J, Barale F, Schettini G, Soares JC. GABAergic dysfunction in mood disorders. *Mol Psychiatry* 2003; **8**: 721–737, 715.
- Benes FM, Berretta S. GABAergic interneurons: implications for understanding schizophrenia and bipolar disorder. *Neuropsychopharmacology* 2001; **25**: 1–27.
- Edden RA, Crocetti D, Zhu H, Gilbert DL, Mostofsky SH. Reduced GABA concentration in attention-deficit/hyperactivity disorder. *Arch Gen Psychiatry* 2012; **69**: 750–753.
- Mueller T, Guo S. The distribution of GAD67-mRNA in the adult zebrafish (teleost) forebrain reveals a prosomeric pattern and suggests previously unidentified homologies to tetrapods. *J Comp Neurol* 2009; **516**: 553–568.
- Delgado L, Schmachtenberg O. Immunohistochemical localization of GABA, GAD65, and the receptor subunits GABAA α 1 and GABAB1 in the zebrafish cerebellum. *Cerebellum* 2008; **7**: 444–450.
- Doldan MJ, Prego B, Holmqvist BI, de Miguel E. Distribution of GABA-immunolabeling in the early zebrafish (*Danio rerio*) brain. *Eur J Morphol* 1999; **37**: 126–129.
- Mueller T, Vernier P, Wullmann MF. A phylotypic stage in vertebrate brain development: GABA cell patterns in zebrafish compared with mouse. *J Comp Neurol* 2006; **494**: 620–634.
- Kim YJ, Nam RH, Yoo YM, Lee CJ. Identification and functional evidence of GABAergic neurons in parts of the brain of adult zebrafish (*Danio rerio*). *Neurosci Lett* 2004; **355**: 29–32.
- McLean DL, Fetcho JR. Ontogeny and innervation patterns of dopaminergic, noradrenergic, and serotonergic neurons in larval zebrafish. *J Comp Neurol* 2004; **480**: 38–56.
- Rink E, Wullmann MF. The teleostean (zebrafish) dopaminergic system ascending to the subpallium (striatum) is located in the basal diencephalon (posterior tuberculum). *Brain Res* 2001; **889**: 316–330.
- Yamamoto K, Ruuskanen JO, Wullmann MF, Vernier P. Two tyrosine hydroxylase genes in vertebrates: New dopaminergic territories revealed in the zebrafish brain. *Mol Cell Neurosci* 2010; **43**: 394–402.
- Holzschuh J, Ryu S, Aberger F, Driever W. Dopamine transporter expression distinguishes dopaminergic neurons from other catecholaminergic neurons in the developing zebrafish embryo. *Mech Dev* 2001; **101**: 237–243.
- Yamamoto K, Ruuskanen JO, Wullmann MF, Vernier P. Differential expression of dopaminergic cell markers in the adult zebrafish forebrain. *J Comp Neurol* 2011; **519**: 576–598.
- Boehmler W, Obrecht-Pflumio S, Canfield V, Thisse C, Thisse B, Levenson R. Evolution and expression of D2 and D3 dopamine receptor genes in zebrafish. *Dev Dyn* 2004; **230**: 481–493.
- Li P, Shah S, Huang L, Carr AL, Gao Y, Thisse C *et al*. Cloning and spatial and temporal expression of the zebrafish dopamine D1 receptor. *Dev Dyn* 2007; **236**: 1339–1346.
- Tay TL, Ronneberger O, Ryu S, Nitschke R, Driever W. Comprehensive catecholaminergic projectome analysis reveals single-neuron integration of zebrafish ascending and descending dopaminergic systems. *Nat Commun* 2011; **2**: 171.
- Ma PM. Catecholaminergic systems in the zebrafish. I. Number, morphology, and histochemical characteristics of neurons in the locus coeruleus. *J Comp Neurol* 1994; **344**: 242–255.
- Ma PM. Catecholaminergic systems in the zebrafish. II. Projection pathways and pattern of termination of the locus coeruleus. *J Comp Neurol* 1994; **344**: 256–269.
- Schweitzer J, Lohr H, Filippi A, Driever W. Dopaminergic and noradrenergic circuit development in zebrafish. *Dev Neurobiol* 2012; **72**: 256–268.
- Peitsaro N, Sundvik M, Anichtchik OV, Kaslin J, Panula P. Identification of zebrafish histamine H1, H2 and H3 receptors and effects of histaminergic ligands on behavior. *Biochem Pharmacol* 2007; **73**: 1205–1214.
- Eriksson KS, Peitsaro N, Karlstedt K, Kaslin J, Panula P. Development of the histaminergic neurons and expression of histidine decarboxylase mRNA in the zebrafish brain in the absence of all peripheral histaminergic systems. *Eur J Neurosci* 1998; **10**: 3799–3812.
- Lucki I. The spectrum of behaviors influenced by serotonin. *Biol Psychiatry* 1998; **44**: 151–162.
- Norton WH, Folchert A, Bally-Cuif L. Comparative analysis of serotonin receptor (HTR1A/HTR1B families) and transporter (slc6a4a/b) gene expression in the zebrafish brain. *J Comp Neurol* 2008; **511**: 521–542.

- 45 Schneider H, Fritzy L, Williams J, Heumann C, Yochum M, Pattar K *et al*. Cloning and expression of a zebrafish 5-HT(2C) receptor gene. *Gene* 2012; **502**: 108–117.
- 46 Sallinen V, Sundvik M, Reenila I, Peitsaro N, Khrustalyov D, Anichtchik O *et al*. Hyperserotonergic phenotype after monoamine oxidase inhibition in larval zebrafish. *J Neurochem* 2009; **109**: 403–415.
- 47 Lillesaar C, Stigloher C, Tannhauser B, Wullmann MF, Bally-Cuif L. Axonal projections originating from raphe serotonergic neurons in the developing and adult zebrafish, *Danio rerio*, using transgenics to visualize raphe-specific pet1 expression. *J Comp Neurol* 2009; **512**: 158–182.
- 48 Drapeau P, Saint-Amant L, Buss RR, Chong M, McDearmid JR, Brustein E. Development of the locomotor network in zebrafish. *Prog Neurobiol* 2002; **68**: 85–111.
- 49 Rink E, Guo S. The too few mutant selectively affects subgroups of monoaminergic neurons in the zebrafish forebrain. *Neuroscience* 2004; **127**: 147–154.
- 50 Teraoka H, Russell C, Regan J, Chandrasekhar A, Concha ML, Yokoyama R *et al*. Hedgehog and Fgf signaling pathways regulate the development of tphR-expressing serotonergic raphe neurons in zebrafish embryos. *J Neurobiol* 2004; **60**: 275–288.
- 51 Bellipanni G, Rink E, Bally-Cuif L. Cloning of two tryptophan hydroxylase genes expressed in the diencephalon of the developing zebrafish brain. *Mech Dev* 2002; **119 Suppl 1**: S215–S220.
- 52 Wang Y, Takai R, Yoshioka H, Shirabe K. Characterization and expression of serotonin transporter genes in zebrafish. *Tohoku J Exp Med* 2006; **208**: 267–274.
- 53 Yamamoto K, Vernier P. The evolution of dopamine systems in chordates. *Front Neuroanat* 2011; **5**: 21.
- 54 Lillesaar C. The serotonergic system in fish. *J Chem Neuroanat* 2011; **41**: 294–308.
- 55 Hsieh DJ, Liao CF. Zebrafish M2 muscarinic acetylcholine receptor: cloning, pharmacological characterization, expression patterns and roles in embryonic bradycardia. *Br J Pharmacol* 2002; **137**: 782–792.
- 56 Papke RL, Ono F, Stokes C, Urban JM, Boyd RT. The nicotinic acetylcholine receptors of zebrafish and an evaluation of pharmacological tools used for their study. *Biochem Pharmacol* 2012; **84**: 352–365.
- 57 Klee EW, Schneider H, Clark KJ, Cousin MA, Ebbert JO, Hooten WM *et al*. Zebrafish: a model for the study of addiction genetics. *Hum Genet* 2012; **131**: 977–1008.
- 58 Arenzana FJ, Clemente D, Sanchez-Gonzalez R, Porteros n, Aijun J, Arevalo R. Development of the cholinergic system in the brain and retina of the zebrafish. *Brain Res Bull* 2005; **66**: 421–425.
- 59 Clemente D, Porteros A, Weruaga E, Alonso JR, Arenzana FJ, Aijun J *et al*. Cholinergic elements in the zebrafish central nervous system: Histochemical and immunohistochemical analysis. *J Comp Neurol* 2004; **474**: 75–107.
- 60 Postlethwait JH, Yan YL, Gates MA, Horne S, Amores A, Brownlie A *et al*. Vertebrate genome evolution and the zebrafish gene map. *Nat Genet* 1998; **18**: 345–349.
- 61 Fredriksson R, Schioth HB. The repertoire of G-protein-coupled receptors in fully sequenced genomes. *Mol Pharmacol* 2005; **67**: 1414–1425.
- 62 Panula P, Chen YC, Priyadarshini M, Kudo H, Semenova S, Sundvik M *et al*. The comparative neuroanatomy and neurochemistry of zebrafish CNS systems of relevance to human neuropsychiatric diseases. *Neurobiol Dis* 2010; **40**: 46–57.
- 63 Ruuskanen JO, Xhaard H, Marjamaki A, Salaneck E, Salminen T, Yan YL *et al*. Identification of duplicated fourth alpha2-adrenergic receptor subtype by cloning and mapping of five receptor genes in zebrafish. *Mol Biol Evol* 2004; **21**: 14–28.
- 64 Ruuskanen JO, Laurila J, Xhaard H, Rantanen VV, Vuoriluoto K, Wurster S *et al*. Conserved structural, pharmacological and functional properties among the three human and five zebrafish alpha 2-adrenoceptors. *Br J Pharmacol* 2005; **144**: 165–177.
- 65 Ruuskanen JO, Peitsaro N, Kaslin JV, Panula P, Scheinin M. Expression and function of alpha-adrenoceptors in zebrafish: drug effects, mRNA and receptor distributions. *J Neurochem* 2005; **94**: 1559–1569.
- 66 Doyon Y, McCammon JM, Miller JC, Faraji F, Ngo C, Katibah GE *et al*. Heritable targeted gene disruption in zebrafish using designed zinc-finger nucleases. *Nat Biotechnol* 2008; **26**: 702–708.
- 67 Moens CB, Donn TM, Wolf-Saxon ER, Ma TP. Reverse genetics in zebrafish by TILLING. *Brief Funct Genomic Proteomic* 2008; **7**: 454–459.
- 68 Peravali R, Gehrig J, Giselbrecht S, Lutjohann DS, Hadzhiev Y, Muller F *et al*. Automated feature detection and imaging for high-resolution screening of zebrafish embryos. *Biotechniques* 2011; **50**: 319–324.
- 69 Pardo-Martin C, Chang TY, Koo BK, Gilleland CL, Wasserman SC, Yanik MF. High-throughput in vivo vertebrate screening. *Nat Methods* 2010; **7**: 634–636.
- 70 Akerboom J, Chen TW, Wardill TJ, Tian L, Marvin JS, Mutlu S *et al*. Optimization of a GCaMP calcium indicator for neural activity imaging. *J Neurosci* 2012; **32**: 13819–13840.
- 71 Scott EK, Mason L, Arrenberg AB, Ziv L, Gosse NJ, Xiao T *et al*. Targeting neural circuitry in zebrafish using GAL4 enhancer trapping. *Nat Methods* 2007; **4**: 323–326.
- 72 Kawakami K, Takeda H, Kawakami N, Kobayashi M, Matsuda N, Mishina M. A transposon-mediated gene trap approach identifies developmentally regulated genes in zebrafish. *Dev Cell* 2004; **7**: 133–144.
- 73 Ahrens MB, Orger MB, Robson DN, Li JM, Keller PJ. Whole-brain functional imaging at cellular resolution using light-sheet microscopy. *Nat Methods* 2013; **10**: 413–420.
- 74 Kubo F, Hablitzel B, Dal Maschio M, Driever W, Baier H, Arrenberg AB. Functional architecture of an optic flow-responsive area that drives horizontal eye movements in zebrafish. *Neuron* 2014; **81**: 1344–1359.
- 75 Huang KH, Ahrens MB, Dunn TW, Engert F. Spinal projection neurons control turning behaviors in zebrafish. *Curr Biol* 2013; **23**: 1566–1573.
- 76 Tye KM, Deisseroth K. Optogenetic investigation of neural circuits underlying brain disease in animal models. *Nat Rev Neurosci* 2012; **13**: 251–266.
- 77 Pan YA, Livet J, Sanes JR, Lichtman JW, Schier AF. Multicolor Brainbow imaging in zebrafish. *Cold Spring Harb Protoc* 2011; **2011** prot5546.
- 78 Geschwind DH. Genetics of autism spectrum disorders. *Trends Cogn Sci* 2011; **15**: 409–416.
- 79 Weinberger DR. From neuropathology to neurodevelopment. *Lancet* 1995; **346**: 552–557.
- 80 Livet J, Weissman TA, Kang H, Draft RW, Lu J, Bennis RA *et al*. Transgenic strategies for combinatorial expression of fluorescent proteins in the nervous system. *Nature* 2007; **450**: 56–62.
- 81 Cai D, Cohen KB, Luo T, Lichtman JW, Sanes JR. Improved tools for the Brainbow toolbox. *Nat Methods* 2013; **10**: 540–547.
- 82 Pan YA, Freundlich T, Weissman TA, Schoppik D, Wang XC, Zimmerman S *et al*. Zebrow: multispectral cell labeling for cell tracing and lineage analysis in zebrafish. *Development* 2013; **140**: 2835–2846.
- 83 Gubernator NG, Zhang H, Staal RG, Mosharov EV, Pereira DB, Yue M *et al*. Fluorescent false neurotransmitters visualize dopamine release from individual presynaptic terminals. *Science* 2009; **324**: 1441–1444.
- 84 Rodriguez PC, Pereira DB, Borgkvist A, Wong MY, Barnard C, Sonders MS *et al*. Fluorescent dopamine tracer resolves individual dopaminergic synapses and their activity in the brain. *Proc Natl Acad Sci USA* 2013; **110**: 870–875.
- 85 Okubo Y, Sekiya H, Namiki S, Sakamoto H, Iinuma S, Yamasaki M *et al*. Imaging extrasynaptic glutamate dynamics in the brain. *Proc Natl Acad Sci USA* 2010; **107**: 6526–6531.
- 86 Marvin JS, Borghuis BG, Tian L, Cichon J, Harnett MT, Akerboom J *et al*. An optimized fluorescent probe for visualizing glutamate neurotransmission. *Nat Methods* 2013; **10**: 162–170.
- 87 Nguyen QT, Schroeder LF, Mank M, Muller A, Taylor P, Griesbeck O *et al*. An in vivo biosensor for neurotransmitter release and in situ receptor activity. *Nat Neurosci* 2010; **13**: 127–132.
- 88 White RM, Sessa A, Burke C, Bowman T, LeBlanc J, Ceol C *et al*. Transparent adult zebrafish as a tool for in vivo transplantation analysis. *Cell Stem Cell* 2008; **2**: 183–189.
- 89 Dodt H-U, Leischner U, Schierloh A, Jahrling N, Mauch CP, Deininger K *et al*. Ultramicroscopy: three-dimensional visualization of neuronal networks in the whole mouse brain. *Nat Methods* 2007; **4**: 331–336.
- 90 Erturk A, Mauch CP, Hellal F, Forstner F, Keck T, Becker K *et al*. Three-dimensional imaging of the unsectioned adult spinal cord to assess axon regeneration and glial responses after injury. *Nat Med* 2012; **18**: 166–171.
- 91 Hama H, Kurokawa H, Kawano H, Ando R, Shimogori T, Noda H *et al*. Scale: a chemical approach for fluorescence imaging and reconstruction of transparent mouse brain. *Nat Neurosci* 2011; **14**: 1481–1488.
- 92 Erturk A, Becker K, Jahrling N, Mauch CP, Hojer CD, Egen JG *et al*. Three-dimensional imaging of solvent-cleared organs using 3DISCO. *Nat Protoc* 2012; **7**: 1983–1995.
- 93 Chung K, Wallace J, Kim SY, Kalyanasundaram S, Andalman AS, Davidson TJ *et al*. Structural and molecular interrogation of intact biological systems. *Nature* 2013; **497**: 332–337.
- 94 Ke MT, Fujimoto S, Imai T. SeeDB: a simple and morphology-preserving optical clearing agent for neuronal circuit reconstruction. *Nat Neurosci* 2013; **16**: 1154–1161.
- 95 Kuwajima T, Sitko AA, Bhansali P, Jurgens C, Guido W, Mason C. ClearT: a detergent- and solvent-free clearing method for neuronal and non-neuronal tissue. *Development* 2013; **140**: 1364–1368.
- 96 Susaki EA, Tainaka K, Perrin D, Kishino F, Tawara T, Watanabe TM *et al*. Whole-brain imaging with single-cell resolution using chemical cocktails and computational analysis. *Cell* 2014; **157**: 726–739.
- 97 Seo HC, Saetre BO, Havik B, Ellingsen S, Fjose A. The zebrafish Pax3 and Pax7 homologues are highly conserved, encode multiple isoforms and show dynamic segment-like expression in the developing brain. *Mech Dev* 1998; **70**: 49–63.
- 98 Moore S, Ribes V, Terriente J, Wilkinson D, Relaix F, Briscoe J. Distinct regulatory mechanisms act to establish and maintain Pax3 expression in the developing neural tube. *PLoS Genet* 2013; **9**: e1003811.

- 99 Garnett AT, Square TA, Medeiros DM. BMP, Wnt and FGF signals are integrated through evolutionarily conserved enhancers to achieve robust expression of Pax3 and Zic genes at the zebrafish neural plate border. *Development* 2012; **139**: 4220–4231.
- 100 Fernandes AM, Beddows E, Filippi A, Driever W. Orthopedia transcription factor *otpa* and *otpb* paralogous genes function during dopaminergic and neuroendocrine cell specification in larval zebrafish. *PLoS ONE* 2013; **8**: e75002.
- 101 Ryu S, Mahler J, Acampora D, Holzschuh J, Erhardt S, Omodei D *et al*. Orthopedia homeodomain protein is essential for diencephalic dopaminergic neuron development. *Curr Biol: CB* 2007; **17**: 873–880.
- 102 Kastenhuber E, Kratochwil CF, Ryu S, Schweitzer J, Driever W. Genetic dissection of dopaminergic and noradrenergic contributions to catecholaminergic tracts in early larval zebrafish. *J Comp Neurol* 2010; **518**: 439–458.
- 103 Zhang P, Hirsch EC, Damier P, Duyckaerts C, Javoy-Agid F. *c-fos* protein-like immunoreactivity: distribution in the human brain and over-expression in the hippocampus of patients with Alzheimer's disease. *Neuroscience* 1992; **46**: 9–21.
- 104 Singewald N. Altered brain activity processing in high-anxiety rodents revealed by challenge paradigms and functional mapping. *Neurosci Biobehav Rev* 2007; **31**: 18–40.
- 105 Singewald N, Salchner P, Sharp T. Induction of *c-Fos* expression in specific areas of the fear circuitry in rat forebrain by anxiogenic drugs. *Biol Psychiatry* 2003; **53**: 275–283.
- 106 Baraban SC, Taylor MR, Castro PA, Baier H. Pentylene tetrazole induced changes in zebrafish behavior, neural activity and *c-fos* expression. *Neuroscience* 2005; **131**: 759–768.
- 107 Ellis LD, Seibert J, Soanes KH. Distinct models of induced hyperactivity in zebrafish larvae. *Brain Res* 2012; **1449**: 46–59.
- 108 Cachat J, Kyzar EJ, Collins C, Gaikwad S, Green J, Roth A *et al*. Unique and potent effects of acute ibogaine on zebrafish: the developing utility of novel aquatic models for hallucinogenic drug research. *Behav Brain Res* 2013; **236**: 258–269.
- 109 Riehl R, Kyzar E, Allain A, Green J, Hook M, Monnig L *et al*. Behavioral and physiological effects of acute ketamine exposure in adult zebrafish. *Neurotoxicol Teratol* 2011; **33**: 658–667.
- 110 Lau BY, Mathur P, Gould GG, Guo S. Identification of a brain center whose activity discriminates a choice behavior in zebrafish. *Proc Natl Acad Sci USA* 2011; **108**: 2581–2586.
- 111 von Trotha JW, Vernier P, Bally-Cuif L. Emotions and motivated behavior converge on an amygdala-like structure in the zebrafish. *Eur J Neurosci* 2014.
- 112 Virdee K, Cumming P, Caprioli D, Jupp B, Rominger A, Aigbirhio FI *et al*. Applications of positron emission tomography in animal models of neurological and neuropsychiatric disorders. *Neurosci Biobehav Rev* 2012; **36**: 1188–1216.
- 113 Arnsten AF, Rubia K. Neurobiological circuits regulating attention, cognitive control, motivation, and emotion: disruptions in neurodevelopmental psychiatric disorders. *J Am Acad Child Adolesc Psychiatry* 2012; **51**: 356–367.
- 114 Mullins MC, Hammerschmidt M, Haffter P, Nüsslein-Volhard C. Large-scale mutagenesis in the zebrafish: in search of genes controlling development in a vertebrate. *Curr Biol* 1994; **4**: 189–202.
- 115 Amsterdam A, Burgess S, Golling G, Chen W, Sun Z, Townsend K *et al*. A large-scale insertional mutagenesis screen in zebrafish. *Genes Dev* 1999; **13**: 2713–2724.
- 116 Petzold AM, Balciunas D, Sivasubbu S, Clark KJ, Bedell VM, Westcot SE *et al*. Nicotine response genetics in the zebrafish. *Proc Natl Acad Sci* 2009; **106**: 18662–18667.
- 117 Hwang WY, Fu Y, Reyon D, Maeder ML, Tsai SQ, Sander JD *et al*. Efficient genome editing in zebrafish using a CRISPR-Cas system. *Nat Biotechnol* 2013; **31**: 227–229.
- 118 Barkley RA, Fischer M, Smallish L, Fletcher K. Young adult outcome of hyperactive children: adaptive functioning in major life activities. *J Am Acad Child Adolesc Psychiatry* 2006; **45**: 192–202.
- 119 Schmidt S, Petermann F. Developmental psychopathology: Attention Deficit Hyperactivity Disorder (ADHD). *BMC Psychiatry* 2009; **9**: 58.
- 120 Berquin PC, Giedd JN, Jacobsen LK, Hamburger SD, Krain AL, Rapoport JL *et al*. Cerebellum in attention-deficit hyperactivity disorder: a morphometric MRI study. *Neurology* 1998; **50**: 1087–1093.
- 121 Arnsten AF. Catecholamine and second messenger influences on prefrontal cortical networks of "representational knowledge": a rational bridge between genetics and the symptoms of mental illness. *Cereb Cortex* 2007; **17 Suppl 1**: i6–15.
- 122 Bush G. Attention-deficit/hyperactivity disorder and attention networks. *Neuropharmacology* 2010; **35**: 278–300.
- 123 Rubia K. "Cool" inferior frontostriatal dysfunction in attention-deficit/hyperactivity disorder versus "hot" ventromedial orbitofrontal-limbic dysfunction in conduct disorder: a review. *Biol Psychiatry* 2011; **69**: e69–e87.
- 124 Arcos-Burgos M, Jain M, Acosta MT, Shively S, Stanescu H, Wallis D *et al*. A common variant of the latrophilin 3 gene, LPHN3, confers susceptibility to ADHD and predicts effectiveness of stimulant medication. *Mol Psychiatry* 2010; **15**: 1053–1066.
- 125 Lange M, Norton W, Coolen M, Chaminade M, Merker S, Proft F *et al*. The ADHD-linked gene *Lphn3.1* controls locomotor activity and impulsivity in zebrafish. *Mol Psychiatry* 2012; **17**: 855.
- 126 Lange M, Norton W, Coolen M, Chaminade M, Merker S, Proft F *et al*. The ADHD-susceptibility gene *Lphn3.1* modulates dopaminergic neuron formation and locomotor activity during zebrafish development. *Mol Psychiatry* 2012; **17**: 946–954.
- 127 Tay TL, Ronneberger O, Ryu S, Nitschke R, Driever W. Comprehensive catecholaminergic projectome analysis reveals single-neuron integration of zebrafish ascending and descending dopaminergic systems. *Nat Commun* 2011; **2**: 171.
- 128 Lange M, Norton W, Coolen M, Chaminade M, Merker S, Proft F *et al*. The ADHD-susceptibility gene *Lphn3.1* modulates dopaminergic neuron formation and locomotor activity during zebrafish development. *Mol Psychiatry* 2012; **17**: 946–954.
- 129 Wallis D, Hill DS, Mendez IA, Abbott LC, Finnell RH, Wellman PJ *et al*. Initial characterization of mice null for *Lphn3*, a gene implicated in ADHD and addiction. *Brain Res* 2012; **1463**: 85–92.
- 130 Hoffman K, Webster TF, Weisskopf MG, Weinberg J, Vieira VM. Exposure to polyfluoroalkyl chemicals and attention deficit/hyperactivity disorder in U.S. children 12–15 years of age. *Environ Health Perspect* 2010; **118**: 1762–1767.
- 131 Spulber S, Kilian P, Wan Ibrahim WN, Onishchenko N, Ulhaq M, Norrgren L *et al*. PFOS induces behavioral alterations, including spontaneous hyperactivity that is corrected by dexamfetamine in zebrafish larvae. *PLoS ONE* 2014; **9**: e94227.
- 132 Levin ED, Sledge D, Roach S, Petro A, Donerly S, Linney E. Persistent behavioral impairment caused by embryonic methylphenidate exposure in zebrafish. *Neurotoxicol Teratol* 2011; **33**: 668–673.
- 133 Gerlai R, Lahav M, Guo S, Rosenthal A. Drinks like a fish: zebra fish (*Danio rerio*) as a behavior genetic model to study alcohol effects. *Pharmacol Biochem Behav* 2000; **67**: 773–782.
- 134 Oliveira RF, Silva JF, Simoes JM. Fighting zebrafish: characterization of aggressive behavior and winner-loser effects. *Zebrafish* 2011; **8**: 73–81.
- 135 Kalueff AV, Gebhardt M, Stewart AM, Cachat JM, Brimmer M, Chawla JS *et al*. Towards a comprehensive catalog of zebrafish behavior 1.0 and beyond. *Zebrafish* 2013; **10**: 70–86.
- 136 Moretz JA, Martins EP, Robison BD. Behavioural syndromes and the evolution of correlated behavior in zebrafish. *Behav Ecol* 2007; **1**: 7–17.
- 137 Filby AL, Paull GC, Hickmore TF, Tyler CR. Unravelling the neurophysiological basis of aggression in a fish model. *BMC Genomics* 2010; **11**: 498.
- 138 Larson ET, O'Malley DM, Melloni RH Jr. Aggression and vasotocin are associated with dominant-subordinate relationships in zebrafish. *Behav Brain Res* 2006; **167**: 94–102.
- 139 Norton WH, Stumpfenhorst K, Faus-Kessler T, Folchert A, Rohner N, Harris MP *et al*. Modulation of *Fgfr1a* signaling in zebrafish reveals a genetic basis for the aggression-boldness syndrome. *J Neurosci* 2011; **31**: 13796–13807.
- 140 Sneddon LU, Schmidt R, Fang Y, Cossins AR. Molecular correlates of social dominance: a novel role for ependymin in aggression. *PLoS ONE* 2011; **6**: e18181.
- 141 Filby AL, Paull GC, Searle F, Ortiz-Zarragoitia M, Tyler CR. Environmental estrogen-induced alterations of male aggression and dominance hierarchies in fish: a mechanistic analysis. *Environ Sci Technol* 2012; **46**: 3472–3479.
- 142 Rohner N, Bercsenyi M, Orban L, Kolanczyk ME, Linke D, Brand M *et al*. Duplication of *fgfr1* permits *Fgf* signaling to serve as a target for selection during domestication. *Curr Biol* 2009; **19**: 1642–1647.
- 143 Thisse B, Thisse C. Functions and regulations of fibroblast growth factor signaling during embryonic development. *Dev Biol* 2005; **287**: 390–402.
- 144 Caramillo EM, Khan KM, Collier AD, Echevarria DJ. Modeling PTSD in the zebrafish: Are we there yet? *Behav Brain Res* 2014; pii **S0166-4328**: 00296–4.
- 145 Stewart AM, Yang E, Nguyen M, Kalueff AV. Developing zebrafish models relevant to PTSD and other trauma- and stressor-related disorders. *Prog Neuropsychopharmacol Biol Psychiatry* 2014; pii **S0278-5846**: 00153–00155.
- 146 Bass SL, Gerlai R. Zebrafish (*Danio rerio*) responds differentially to stimulus fish: the effects of sympatric and allopatric predators and harmless fish. *Behav Brain Res* 2008; **186**: 107–117.
- 147 Gerlai R. Zebrafish antipredatory responses: a future for translational research? *Behav Brain Res* 2010; **207**: 223–231.
- 148 Blank M, Guerim LD, Cordeiro RF, Vianna RM. A one-trial inhibitory avoidance task to zebrafish: rapid acquisition of an NMDA-dependent long-term memory. *Neurobiol Learn Mem* 2009; **92**: 529–534.
- 149 Morin C, de Souza Silva MA, Muller CP, Hardigan P, Spieler RE. Active avoidance learning in zebrafish (*Danio rerio*)—the role of sensory modality and inter-stimulus interval. *Behav Brain Res* 2013; **248**: 141–143.
- 150 Xu X, Scott-Scheiern T, Kempker L, Simons K. Active avoidance conditioning in zebrafish (*Danio rerio*). *Neurobiol Learn Mem* 2007; **87**: 72–77.

- 151 Pradel G, Schmidt R, Schachner M. Involvement of L1.1 in memory consolidation after active avoidance conditioning in zebrafish. *J Neurobiol* 2000; **43**: 389–403.
- 152 Agetsuma M, Aoki T, Aoki R, Okamoto H. Cued fear conditioning in zebrafish (*Danio rerio*). In: Kalueff AV, Stewart AM (eds). *Zebrafish Protocols for Neuro-behavioral Research* vol. 66. Humana Press: New York, NY, pp 257–264, 2012.
- 153 Egan RJ, Bergner CL, Hart PC, Cachat JM, Canavello PR, Elegante MF *et al*. Understanding behavioral and physiological phenotypes of stress and anxiety in zebrafish. *Behav Brain Res* 2009; **205**: 38–44.
- 154 Speedie N, Gerlai R. Alarm substance induced behavioral responses in zebrafish (*Danio rerio*). *Behav Brain Res* 2008; **188**: 168–177.
- 155 Wong K, Elegante M, Bartels B, Elkhayat S, Tien D, Roy S *et al*. Analyzing habituation responses to novelty in zebrafish (*Danio rerio*). *Behav Brain Res* 2010; **208**: 450–457.
- 156 Mahabir S, Chatterjee D, Buske C, Gerlai R. Maturation of shoaling in two zebrafish strains: a behavioral and neurochemical analysis. *Behav Brain Res* 2013; **247**: 1–8.
- 157 Miller N, Gerlai R. Quantification of shoaling behaviour in zebrafish (*Danio rerio*). *Behav Brain Res* 2007; **184**: 157–166.
- 158 Miller NY, Gerlai R. Oscillations in shoal cohesion in zebrafish (*Danio rerio*). *Behav Brain Res* 2008; **193**: 148–151.
- 159 Stewart AM, Nguyen M, Wong K, Poudel MK, Kalueff AV. Developing zebrafish models of autism spectrum disorder (ASD). *Prog Neuropsychopharmacol Biol Psychiatry* 2014; **50**: 27–36.
- 160 Green J, Collins C, Kyzar EJ, Pham M, Roth A, Gaikwad S *et al*. Automated high-throughput neurophenotyping of zebrafish social behavior. *J Neurosci Methods* 2012; **210**: 266–271.
- 161 Pham M, Raymond J, Hester J, Kyzar E, Gaikwad S, Bruce I *et al*. Assessing social behavior phenotypes in adult zebrafish: shoaling, social preference and mirror biting tests. In: Kalueff AV, Stewart AM (eds). *Zebrafish Protocols for Neuro-behavioral Research*. Humana Press: New York, NY, 2012.
- 162 Hageman I, Andersen HS, Jorgensen MB. Post-traumatic stress disorder: a review of psychobiology and pharmacotherapy. *Acta Psychiatr Scand* 2001; **104**: 411–422.
- 163 Alderman SL, Bernier NJ. Ontogeny of the corticotropin-releasing factor system in zebrafish. *Gen Comp Endocrinol* 2009; **164**: 61–69.
- 164 Alsop D, Vijayan M. The zebrafish stress axis: molecular fallout from the teleost-specific genome duplication event. *Gen Comp Endocrinol* 2009; **161**: 62–66.
- 165 Griffiths BB, Schoonheim PJ, Ziv L, Voelker L, Baier H, Gahtan E. A zebrafish model of glucocorticoid resistance shows serotonergic modulation of the stress response. *Front Behav Neurosci* 2012; **6**: 68.
- 166 Ziv L, Muto A, Schoonheim PJ, Meijnsing SH, Strasser D, Ingraham HA *et al*. An affective disorder in zebrafish with mutation of the glucocorticoid receptor. *Mol Psychiatry* 2013; **18**: 681–691.
- 167 Yehuda R. Biology of posttraumatic stress disorder. *J Clin Psychiatry* 2001; **62 Suppl 17**: 41–46.
- 168 Chakravarty S, Reddy BR, Sudhakar SR, Saxena S, Das T, Meghah V *et al*. Chronic unpredictable stress (CUS)-induced anxiety and related mood disorders in a zebrafish model: altered brain proteome profile implicates mitochondrial dysfunction. *PLoS ONE* 2013; **8**: e63302.
- 169 Gotovac K, Sabioncello A, Rabatic S, Berki T, Dekaris D. Flow cytometric determination of glucocorticoid receptor (GCR) expression in lymphocyte subpopulations: lower quantity of GCR in patients with post-traumatic stress disorder (PTSD). *Clin Exp Immunol* 2003; **131**: 335–339.
- 170 Schaaf MJ, Champagne D, van Laanen IH, van Wijk DC, Meijer AH, Meijer OC *et al*. Discovery of a functional glucocorticoid receptor β -isoform in zebrafish. *Endocrinology* 2008; **149**: 1591–1599.
- 171 Mehta D, Binder EB. Gene x environment vulnerability factors for PTSD: the HPA-axis. *Neuropharmacology* 2012; **62**: 654–662.
- 172 Alsop D, Vijayan M. The zebrafish stress axis: molecular fallout from the teleost-specific genome duplication event. *Gen Comp Endocrinol* 2009; **161**: 62–66.
- 173 Dutton MA, Lee EW, Zukowska Z. NPY and extreme stress: lessons learned from posttraumatic stress disorder. *EXS* 2006; **95**: 213–222.
- 174 Nguyen M, Yang E, Neelkantam N, Mikhaylova A, Arnold R, Poudel MK *et al*. Developing 'integrative' zebrafish models of behavioral and metabolic disorders. *Behav Brain Res* 2013; **256**: 172–187.
- 175 Tyrka A. Biomarkers of Risk for Post-Traumatic Stress Disorder (PTSD): DTIC Document 2011.
- 176 Alazizi A, Liu MY, Williams FE, Kurogi K, Sakakibara Y, Suiko M *et al*. Identification, characterization, and ontogenic study of a catechol O-methyltransferase from zebrafish. *Aquat Toxicol (Amsterdam, Netherlands)* 2011; **102**: 18–23.
- 177 Aldeco M, Arslan BK, Edmondson DE. Catalytic and inhibitor binding properties of zebrafish monoamine oxidase (zMAO): comparisons with human MAO A and MAO B. *Comp Biochem Physiol B Biochem Biol* 2011; **159**: 78–83.
- 178 Mann KD, Hoyt C, Feldman S, Blunt L, Raymond A, Page-McCaw PS. Cardiac response to startle stimuli in larval zebrafish: sympathetic and parasympathetic components. *Am J Physiol Regul Integr Comp Physiol* 2010; **298**: R1288–R1297.
- 179 Robinson TE, Berridge KC. The neural basis of drug craving: an incentive-sensitization theory of addiction. *Brain Res Rev* 1993; **18**: 247–291.
- 180 Rink E, Wullimann MF. The teleostean (zebrafish) dopaminergic system ascending to the subpallium (striatum) is located in the basal diencephalon (posterior tuberculum). *Brain Res* 2001; **889**: 316–330.
- 181 Rink E, Wullimann MF. Development of the catecholaminergic system in the early zebrafish brain: an immunohistochemical study. *Dev Brain Res* 2002; **137**: 89–100.
- 182 Klee EW, Ebbert JO, Schneider H, Hurt RD, Ekker SC. Zebrafish for the study of the biological effects of nicotine. *Nicotine Tob Res* 2011; **13**: 301–312.
- 183 Darland T, Dowling JE. Behavioral screening for cocaine sensitivity in mutant zebrafish. *Proc Natl Acad Sci USA* 2001; **98**: 11691–11696.
- 184 Ninkovic J, Folchert A, Makhankov YV, Neuhauss SC, Sillaber I, Straehle U *et al*. Genetic identification of AChE as a positive modulator of addiction to the psychostimulant D-amphetamine in zebrafish. *J Neurobiol* 2006; **66**: 463–475.
- 185 Kily LJ, Cowe YC, Hussain O, Patel S, McElwaine S, Cotter FE *et al*. Gene expression changes in a zebrafish model of drug dependency suggest conservation of neuro-adaptation pathways. *J Exp Biol* 2008; **211**: 1623–1634.
- 186 Lau B, Bretaud S, Huang Y, Lin E, Guo S. Dissociation of food and opiate preference by a genetic mutation in zebrafish. *Genes Brain Behav* 2006; **5**: 497–505.
- 187 Guo S, Wilson SW, Cooke S, Chitnis AB, Driever W, Rosenthal A. Mutations in the zebrafish unmask shared regulatory pathways controlling the development of catecholaminergic neurons. *Dev Biol* 1999; **208**: 473–487.
- 188 Levkowitz G, Zeller J, Sirotkin HI, French D, Schilbach S, Hashimoto H *et al*. Zinc finger protein too few controls the development of monoaminergic neurons. *Nat Neurosci* 2002; **6**: 28–33.
- 189 Lau B, Bretaud S, Huang Y, Lin E, Guo S. Dissociation of food and opiate preference by a genetic mutation in zebrafish. *Genes Brain Behav* 2006; **5**: 497–505.
- 190 Webb KJ, Norton WH, Trumbach D, Meijer AH, Ninkovic J, Topp S *et al*. Zebrafish reward mutants reveal novel transcripts mediating the behavioral effects of amphetamine. *Genome Biol* 2009; **10**: R81.
- 191 Mathur P, Lau B, Guo S. Conditioned place preference behavior in zebrafish. *Nat Protoc* 2011; **6**: 338–345.
- 192 Mathur P, Berberoglu MA, Guo S. Preference for ethanol in zebrafish following a single exposure. *Behav Brain Res* 2011; **217**: 128–133.
- 193 Bretaud S, Li Q, Lockwood B, Kobayashi K, Lin E, Guo S. A choice behavior for morphine reveals experience-dependent drug preference and underlying neural substrates in developing larval zebrafish. *Neuroscience* 2007; **146**: 1109–1116.
- 194 Ponzoni L, Braidia D, Pucci L, Andrea D, Fasoli F, Manfredi I *et al*. The cytosine derivatives, CC4 and CC26, reduce nicotine-induced conditioned place preference in zebrafish by acting on heteromeric neuronal nicotinic acetylcholine receptors. *Psychopharmacology* 2014; **1–13**.
- 195 Everitt BJ, Robbins TW. Neural systems of reinforcement for drug addiction: from actions to habits to compulsion. *Nat Neurosci* 2005; **8**: 1481–1489.
- 196 Pelloux Y, Everitt BJ, Dickinson A. Compulsive drug seeking by rats under punishment: effects of drug taking history. *Psychopharmacology* 2007; **194**: 127–137.
- 197 Kily LJM, Cowe YCM, Hussain O, Patel S, McElwaine S, Cotter FE *et al*. Gene expression changes in a zebrafish model of drug dependency suggest conservation of neuro-adaptation pathways. *J Exp Biol* 2008; **211**: 1623–1634.
- 198 Brennan CH, Parmar A, Kily LKM, Ananthathevan A, Doshi A, Patel S *et al*. Conditioned place preference models of drug dependence and relapse to drug seeking: studies with nicotine and ethanol. *NeuroMethods* 2011; **52**: 163–180.
- 199 Shaham Y, Hope BT. The role of neuroadaptations in relapse to drug seeking. *Nat Neurosci* 2005; **8**: 1437–1439.
- 200 Navarro HJ, Doran CM, Shakeshaft AP. Measuring costs of alcohol harm to others: a review of the literature. *Drug Alcohol Depend* 2011; **114**: 87–99.
- 201 Tran S, Gerlai R. Recent advances with a novel model organism: alcohol tolerance and sensitization in zebrafish (*Danio rerio*). *Prog Neuropsychopharmacol Biol Psychiatry* 2014; pii **S0278-5846**: 00038–4.
- 202 Miller N, Greene K, Dydinski A, Gerlai R. Effects of nicotine and alcohol on zebrafish (*Danio rerio*) shoaling. *Behav Brain Res* 2013; **240**: 192–196.
- 203 Gerlai R, Chatterjee D, Pereira T, Sawashima T, Krishnannair R. Acute and chronic alcohol dose: population differences in behavior and neurochemistry of zebrafish. *Genes Brain Behav* 2009; **8**: 586–599.
- 204 Tran S, Chatterjee D, Gerlai R. An integrative analysis of ethanol tolerance and withdrawal in zebrafish (*Danio rerio*). *Behav Brain Res* 2014; pii **S0166-4328**: 00109–0.
- 205 Fernandes Y, Tran S, Abraham E, Gerlai R. Embryonic alcohol exposure impairs associative learning performance in adult zebrafish. *Behav Brain Res* 2014; **265**: 181–187.

- 206 Pan Y, Kaiguo M, Razak Z, Westwood JT, Gerlai R. Chronic alcohol exposure induced gene expression changes in the zebrafish brain. *Behav Brain Res* 2011; **216**: 66–76.
- 207 Barkley RA. Behavioral inhibition, sustained attention, and executive functions: constructing a unifying theory of ADHD. *Psychol Bull* 1997; **121**: 65.
- 208 Dalley JW, Cardinal RN, Robbins TW. Prefrontal executive and cognitive functions in rodents: neural and neurochemical substrates. *Neurosci Biobehav Rev* 2004; **28**: 771–784.
- 209 Urcelay GP, Dalley JW. Linking ADHD, impulsivity, and drug abuse: a neuropsychological perspective. *Curr Top Behav Neurosci* 2012; **9**: 173–197.
- 210 Clayton IC, Richards JC, Edwards CJ. Selective attention in obsessive, compulsive disorder. *J Abnorm Psychol* 1999; **108**: 171.
- 211 Belin D, Mar AC, Dalley JW, Robbins TW, Everitt BJ. High impulsivity predicts the switch to compulsive cocaine-taking. *Science* 2008; **320**: 1352–1355.
- 212 Molander AC, Mar A, Norbury A, Steventon S, Moreno M, Caprioli D *et al*. High impulsivity predicting vulnerability to cocaine addiction in rats: some relationship with novelty preference but not novelty reactivity, anxiety or stress. *Psychopharmacology* 2011; **215**: 721–731.
- 213 Dalley JW, Everitt BJ, Robbins TW. Impulsivity, compulsivity, and top-down cognitive control. *Neuron* 2011; **69**: 680–694.
- 214 Robbins T. The 5-choice serial reaction time task: behavioural pharmacology and functional neurochemistry. *Psychopharmacology* 2002; **163**: 362–380.
- 215 Economidou D, Pelloux Y, Robbins TW, Dalley JW, Everitt BJ. High impulsivity predicts relapse to cocaine-seeking after punishment-induced abstinence. *Biol Psychiatry* 2009; **65**: 851–856.
- 216 Parker MO, Millington ME, Combe FJ, Brennan CH. Development and implementation of a three-choice serial reaction time task for zebrafish (*Danio rerio*). *Behav Brain Res* 2012; **227**: 73–80.
- 217 Parker M, Ife D, Ma J, Pancholi M, Smeraldi F, Straw C *et al*. Development and automation of a test of impulse control in zebrafish. *Front Syst Neurosci* 2013; **7**: 65.
- 218 Parker MO, Brock AJ, Sudwants A, Brennan CH. Atomoxetine reduces anticipatory responding in a 5-choice serial reaction time task for adult zebrafish. *Psychopharmacology* 2014; 1–9.
- 219 Robinson ES, Eagle DM, Mar AC, Bari A, Banerjee G, Jiang X *et al*. Similar effects of the selective noradrenaline reuptake inhibitor atomoxetine on three distinct forms of impulsivity in the rat. *Neuropsychopharmacology* 2008; **33**: 1028–1037.
- 220 Force A, Lynch M, Pickett FB, Amores A, Yan YL, Postlethwait J. Preservation of duplicate genes by complementary, degenerative mutations. *Genetics* 1999; **151**: 1531–1545.
- 221 Nguyen M, Yang E, Neelkantan N, Mikhaylova A, Arnold R, Poudel MK *et al*. Developing 'integrative' zebrafish models of behavioral and metabolic disorders. *Behav Brain Res* 2013; **256**: 172–187.
- 222 Ito H, Yamamoto N. Non-laminar cerebral cortex in teleost fishes? *Biol Lett* 2009; **5**: 117–121.
- 223 Aizawa H. Habenula and the asymmetric development of the vertebrate brain. *Anat Sci Int* 2013; **88**: 1–9.
- 224 Tomaiuolo M, Gonzalez C, Medina JH, Piriz J. Lateral Habenula determines long-term storage of aversive memories. *Front Behav Neurosci* 2014; **8**: 170.
- 225 Agetsuma M, Aizawa H, Aoki T, Nakayama R, Takahoko M, Goto M *et al*. The habenula is crucial for experience-dependent modification of fear responses in zebrafish. *Nat Neurosci* 2010; **13**: 1354–1356.
- 226 Amo R, Aizawa H, Takahoko M, Kobayashi M, Takahashi R, Aoki T *et al*. Identification of the zebrafish ventral habenula as a homolog of the mammalian lateral habenula. *J Neurosci* 2010; **30**: 1566–1574.
- 227 Okamoto H, Agetsuma M, Aizawa H. Genetic dissection of the zebrafish habenula, a possible switching board for selection of behavioral strategy to cope with fear and anxiety. *Dev Neurobiol* 2012; **72**: 386–394.
- 228 Shimada Y, Hirano M, Nishimura Y, Tanaka T. A high-throughput fluorescence-based assay system for appetite-regulating gene and drug screening. *PLoS ONE* 2012; **7**: e52549.
- 229 Chen TW, Wardill TJ, Sun Y, Pulver SR, Renninger SL, Baohan A *et al*. Ultra-sensitive fluorescent proteins for imaging neuronal activity. *Nature* 2013; **499**: 295–300.
- 230 Chatterjee D, Shams S, Gerlai R. Chronic and acute alcohol administration induced neurochemical changes in the brain: comparison of distinct zebrafish populations. *Amino acids* 2014; **46**: 921–930.
- 231 Hough LB. *Histamine*. Lippincott-Raven: Philadelphia, 1999.
- 232 Gerlai R. Antipredatory behavior of zebrafish: adaptive function and a tool for translational research. *Evol Psychol* 2013; **11**: 591–605.
- 233 Cachat J, Canavello P, Elegante M, Bartels B, Hart P, Bergner C *et al*. Modeling withdrawal syndrome in zebrafish. *Behav Brain Res* 2010; **208**: 371–376.
- 234 Chatterjee D, Gerlai R. High precision liquid chromatography analysis of dopaminergic and serotonergic responses to acute alcohol exposure in zebrafish. *Behav Brain Res* 2009; **200**: 208–213.
- 235 Pan Y, Chatterjee D, Gerlai R. Strain dependent gene expression and neurochemical levels in the brain of zebrafish: focus on a few alcohol related targets. *Physiol Behav* 2012; **107**: 773–780.

Uncorrected