



Simpson, S.L., Boyce, A.J., Lee, M.R., and Lindgren, P. (2014) *Stable isotope studies of post impact hydrothermal deposits within the sub-crater environment of the Rochechouart structure*. In: 77th Annual Meeting of the Meteoritical Society, 8-13 September 2014, Casablanca, Morocco.

Copyright © 2014 The Authors

<http://eprints.gla.ac.uk/97473/>

Deposited on: 25 September 2014

Enlighten – Research publications by members of the University of Glasgow  
<http://eprints.gla.ac.uk>

**Stable isotope studies of post impact hydrothermal deposits within the sub-crater environment of the Rochechouart structure.** S. L. Simpson, A. J. Boyce, M. R. Lee and P. Lindgren. School of Geographical and Earth Sciences, University of Glasgow, UK. E-mail: S.Simpson.1@Research.gla.ac.uk.

**Introduction:** Hypervelocity impacts into hydrous terrestrial targets can initiate hydrothermal circulation [1]. The longevity of these systems is highly variable, and depends on the amount of water available in the target (*i.e. meteoric, hydrous minerals*) and amount of heat generated (*i.e. impact melt, elevated geothermal gradient*) [1]. Approximately one-third of terrestrial impact structures have evidence for impact-hydrothermalism, and this process has recently been confirmed on Mars [1, 2]. These systems are ideal environments to search for life, as they provide an exogenic source of heat to tectonically “dead” planets.

**Purpose:** We have sought evidence for past life in the sub crater environment of the  $201 \pm 2$  Ma [3], 23km diameter [4], Rochechouart impact structure located in west-central France. The crater has been highly eroded, yet enough material remains to provide a full suite of impactites including the shocked basement directly below the transient crater floor. The target was primarily crystalline, located on the western edge of the Central Massif. Hettangian (199-196 Ma) sedimentary carbonates, crop out 17.5km west of the center of the structure [4]. These deposits may have influenced, possibly covered, Rochechouart during cooling, providing a water source to fuel hydrothermal activity. A strong hydrothermal overprint, primarily K-metasomatism, is a feature of Rochechouart [4, 5]. Of particular interest in this study is the shocked basement directly beneath the transient crater floor, which hosts a network of shock-related fractures, faults and lithic breccias ideal for fluid transport [1, 6].

**Methods and Results:** Carbonates and sulphides found within autochthonous and parautochthonous impactites were extracted and analyzed for  $\delta^{13}\text{C}$ ,  $\delta^{18}\text{O}$  and  $\delta^{34}\text{S}$ . Pyrite  $\delta^{34}\text{S}$  values range from -10‰ to -26‰ (VCDT), which is indicative of biological reduction of sulphate, and carbonate values reflect a mixture of organic and inorganic reservoirs, as well cool (< 100°C) depositional conditions [7].

In July 2014, a second field campaign is planned for further collection of impact-hydrothermal sulphides and carbonates from the shocked basement, as well as nearby Hettangian sedimentary deposits for comparative isotope work.

**Acknowledgements:** Funding has been kindly provided by the Barringer Family Fund for Meteorite Impact Research [8]. Thanks are also extended to Philippe Lambert for his guidance and support, and the Réserve Naturelle de l'Astroblème de Rochechouart-Chassenon for permission to sample [9].

**References:** [1] Osinski, G. R. et al. 2013. *Icarus* 224: 347–363. [2] Marzo, G. A. et al. 2010. *Icarus* 208: 667-683. [3] Schmeider, M. et al. 2010. *Meteoritics & Planetary Science* 45 Nr 8: 1225–1242. [4] Lambert, P. 2010. *The Geological Society of America Special Paper* 465 [5] Tagle, R. et al. 2009. *Geochimica et Cosmochimica Acta* 73: 4891–4906 [6] French, B. 1998. *LPI Contribution* 954. [7] Pirajno, F. 2009. *Springer, Geological Survey of Western Australia* ISBN 978-1-4020-8612-0 [8] Barringer Family Fund for Meteorite Impact Research. 2014. Barringer Crater Company, Lunar and Planetary Institute, Houston, TX. [9] Réserve Naturelle de l'Astroblème de Rochechouart-Chassenon, *Rochechouart, France*.