

PAPER • OPEN ACCESS

Casimir effect for perfect electromagnetic conductors (PEMCs): a sum rule for attractive/repulsive forces

To cite this article: Stefan Rode *et al* 2018 *New J. Phys.* **20** 043024

View the [article online](#) for updates and enhancements.

Related content

- [Macroscopic quantum electrodynamics in nonlocal and nonreciprocal media](#)
Stefan Yoshi Buhmann, David T Butcher and Stefan Scheel
- [Cavity-QED interactions of two correlated atoms](#)
Saeideh Esfandiarpour, Hassan Safari, Robert Bennett *et al.*
- [Damped vacuum states of light](#)
T G Philbin



PAPER

Casimir effect for perfect electromagnetic conductors (PEMCs): a sum rule for attractive/repulsive forces

 Stefan Rode¹, Robert Bennett¹ and Stefan Yoshi Buhmann^{1,2}
¹ Physikalisches Institut, Albert-Ludwigs-Universität Freiburg, Hermann-Herder-Str. 3, D-79104 Freiburg, Germany

² Freiburg Institute for Advanced Studies, Albert-Ludwigs-Universität Freiburg, Albertstr. 19, D-79104 Freiburg, Germany

 E-mail: stefan.rode511@web.de
Keywords: perfect electromagnetic conductor (PEMC), Casimir effect, nonreciprocal media, duality, Casimir repulsion

RECEIVED

4 October 2017

REVISED

20 December 2017

ACCEPTED FOR PUBLICATION

24 January 2018

PUBLISHED

16 April 2018

Original content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](https://creativecommons.org/licenses/by/4.0/).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.



Abstract

We discuss the Casimir effect for boundary conditions involving perfect electromagnetic conductors, which interpolate between perfect electric conductors and perfect magnetic conductors. Based on the corresponding reciprocal Green's tensor we construct the Green's tensor for two perfectly reflecting plates with magnetoelectric coupling (non-reciprocal media) within the framework of macroscopic quantum electrodynamics. We calculate the Casimir force between two arbitrary perfect electromagnetic conductor plates, resulting in a universal analytic expression that connects the attractive Casimir force with the repulsive Boyer force. We relate the results to a duality symmetry of electromagnetism.

1. Introduction

Nonvanishing zero-point energies are a pervasive feature of quantum mechanics and quantum field theory. The fact that energy fluctuations of the vacuum lead to physically observable macroscopic forces was first discovered by Hendrik Casimir in [1], who calculated the attractive force between two uncharged metallic plates due to the fluctuations of the electromagnetic field, which turned out to be given by the simple expression

$$f_{\text{attr}} = -\frac{\hbar c \pi^2}{240 d^4} \quad (1)$$

for plates separated by a distance d . The origin of this force is the non-vanishing expectation value of the squared electric and magnetic fields in the vacuum state, which is then modified by the presence of surfaces. These vacuum fluctuations give rise to various forms of matter-vacuum interaction. The inverse fourth-power distance-dependence leads to negligibly small forces on large distance scales. However, in the nanometre regime the Casimir effect and other vacuum fluctuation induced forces can become significant or even dominant. In particular, the Casimir force poses a challenge for constructing microelectromechanical systems [2]. It causes effects such as stiction [3, 4], which is the permanent adhesion of two nano-structural elements. In order to remove such impeding effects, possible ways of manipulating the Casimir force between bodies have been pursued.

Of particular interest are repulsive Casimir forces [5]. The first result in this field was obtained by Boyer in [6], who considered an assembly of two parallel plates, one of them perfectly conducting, the other one perfectly permeable. He found the Casimir force to be repulsive in this case and showed that the ratio of his result to the attractive force calculated by Casimir reads

$$f_{\text{rep}} = -\frac{7}{8} f_{\text{attr}}. \quad (2)$$

It has also been theoretically shown that the magnitude of the Casimir force between two plates of any magnetodielectric properties has to fall between the result of Casimir and the result of Boyer [7], which we shall confirm here.

Due to the difficulty of realising materials whose permeability is perfect or nearly perfect, other ways of implementing repulsive Casimir forces have been considered. Kenneth and Klich [8] have discussed the

opportunities of materials with non-trivial but finite magnetic susceptibilities for instance. As another approach the Casimir forces on materials with polarisation-twisting effects have been studied. In particular the vacuum interaction properties of topological insulators [9, 10] and of chiral metamaterials [11, 12], have been investigated for generalised boundary conditions [13]. For the case of a scalar field confined by Robin boundary conditions, the Casimir force has been obtained to be either repulsive or attractive [14], as is also the case for thin films described as Chern–Simons boundaries [15, 16]. Here we will study perfect electromagnetic conductors (PEMCs) as introduced by Lindell and Sihvola [17], which are an idealised class of nonreciprocal polarisation-mixing materials whose response is characterised by a single parameter M . We will calculate the Casimir force between two PEMC plates in terms of this parameter, which will allow us to continuously vary the Casimir force between the two extremal values.

From a more fundamental point of view, the Casimir force in PEMC media is of interest because of its close relation to duality invariance. It has been shown [18, 19] that a linear magnetodielectric medium breaks the duality invariance that holds for the free Maxwell equations, causing them to instead have a discrete \mathbb{Z}_4 -symmetry. Allowing material response that violates Lorentz-reciprocity restores duality invariance [20]—PEMC media provide these properties. For this reason we will determine the relation between the PEMC parameter M and the duality angle of a perfect conductor to obtain a coherent picture of the impact of duality transformations on Casimir forces.

2. The Casimir force on nonreciprocal bodies

The Green's tensor $\mathbb{G} = \mathbb{G}(\mathbf{r}, \mathbf{r}', \omega)$ of Maxwell's equations in a region with tensor-valued permittivity $\varepsilon = \varepsilon(\mathbf{r}, \omega)$, permeability $\mu = \mu(\mathbf{r}, \omega)$ and cross-polarisabilities $\zeta = \zeta(\mathbf{r}, \omega)$ and $\xi = \xi(\mathbf{r}, \omega)$ (discussed in detail in section 3) is defined to satisfy [20]:

$$\left[\nabla \times \frac{1}{\mu} \nabla \times - \frac{i\omega}{c} \nabla \times \frac{\zeta}{\mu} + \frac{i\omega}{c} \frac{\xi}{\mu} \nabla \times - \frac{\omega^2}{c^2} \left(\varepsilon - \frac{\xi\zeta}{\mu} \right) \right] \mathbb{G} = \mathbb{I} \delta(\mathbf{r} - \mathbf{r}') \quad (3)$$

subject to appropriate boundary conditions, and where \mathbb{I} is the three-dimensional identity matrix. Then, quantised electromagnetic fields can be constructed via [21]

$$\hat{\mathbf{E}}(\mathbf{r}, \omega) = i\mu_0\omega \int d^3\mathbf{r}' \mathbb{G}(\mathbf{r}, \mathbf{r}', \omega) \cdot \hat{\mathbf{j}}_{\mathbf{N}}(\mathbf{r}', \omega) \quad (4)$$

where $\hat{\mathbf{j}}_{\mathbf{N}}$ is a noise-current source, and here we have taken $\omega > 0$, but the corresponding negative frequency fields can be constructed by hermitian conjugation. The noise current $\hat{\mathbf{j}}_{\mathbf{N}}$ is given explicitly in terms of noise polarisation $\hat{\mathbf{P}}_{\mathbf{N}}$ and noise magnetisation $\hat{\mathbf{M}}_{\mathbf{N}}$ by

$$\hat{\mathbf{j}}_{\mathbf{N}}(\mathbf{r}, \omega) = -i\omega\hat{\mathbf{P}}_{\mathbf{N}} + \nabla \times \hat{\mathbf{M}}_{\mathbf{N}} \quad (5)$$

with

$$\begin{pmatrix} \hat{\mathbf{P}}_{\mathbf{N}} \\ \hat{\mathbf{M}}_{\mathbf{N}} \end{pmatrix} = \sqrt{\frac{\hbar}{\pi}} \mathcal{S} \begin{pmatrix} \hat{\mathbf{f}}_e \\ \hat{\mathbf{f}}_m \end{pmatrix}, \quad \mathcal{S}\mathcal{S}^\dagger = \begin{pmatrix} \varepsilon_0 \text{Im}(\varepsilon - \xi\mu^{-1}\zeta) & (2i\mu Z_0)^{-1}(\xi - \zeta^*) \\ (2i\mu Z_0)^{-1}(\zeta - \xi^*) & -\text{Im}(\mu^{-1})/\mu_0 \end{pmatrix}, \quad (6)$$

where $Z_0 = \sqrt{\mu_0/\varepsilon_0}$ denotes the impedance of free space. Here and throughout we use the convention that 3×3 matrices in position space (tensors) are represented by 'open-face' symbols (\mathbb{A} , \mathbb{B} etc), while 2×2 matrices acting in polarisation or duality space are represented by 'calligraphic' symbols (\mathcal{A} , \mathcal{B} etc). The quantities $\hat{\mathbf{f}}_\lambda$ are bosonic quasiparticle excitations satisfying:

$$[\hat{\mathbf{f}}_\lambda(\mathbf{r}, \omega), \hat{\mathbf{f}}_{\lambda'}^\dagger(\mathbf{r}', \omega')] = \delta_{\lambda\lambda'} \delta(\mathbf{r} - \mathbf{r}') \delta(\omega - \omega') \mathbb{I}, \quad (\lambda = e, m) \quad (7)$$

with all other commutators being zero. From a macroscopic point of view the Casimir force \mathbf{F} between arbitrary bodies can be interpreted as the ground-state expectation value of the Lorentz force, or equivalently by an integral over the Maxwell stress tensor $\hat{\mathbb{T}}$:

$$\mathbf{F} = \int_{\partial V} d\mathbf{A} \cdot \langle \hat{\mathbb{T}} \rangle \quad (8)$$

with ∂V being the boundary of a volume enclosing the body on which the force is to be calculated, and the stress tensor is

$$\hat{\mathbb{T}} = \varepsilon_0 \hat{\mathbf{E}} \otimes \hat{\mathbf{E}} + \frac{1}{\mu_0} \hat{\mathbf{B}} \otimes \hat{\mathbf{B}} - \frac{1}{2} \left(\varepsilon_0 \hat{\mathbf{E}}^2 + \frac{1}{\mu_0} \hat{\mathbf{B}}^2 \right) \mathbb{I}, \quad (9)$$

where the fields are obtained from equation (4) together with $\hat{\mathbf{B}} = (i\omega)^{-1} \nabla \times \hat{\mathbf{E}}$. We can now evaluate the expectation value in the vacuum state $|\{0\}\rangle$ of the noise current quanta $\hat{\mathbf{f}}_\lambda$ by using $\hat{\mathbf{f}}_\lambda|\{0\}\rangle = 0$ and the

commutator (7) above. We will also use an integral relation that can be derived from the definition (3) of the Green's tensor [20, 22]

$$\Im[\mathbb{G}(\mathbf{r}, \mathbf{r}', \omega)] = \mu_0 \int d^3s \mathbb{G}(\mathbf{r}, \mathbf{s}, \omega)(i\omega, \overleftarrow{\nabla}) \cdot \mathcal{S} \cdot \mathcal{S}^\dagger \cdot (i\omega, \nabla \times)^T \mathbb{G}^{*T}(\mathbf{r}', \mathbf{s}, \omega), \quad (10)$$

where

$$\Im[\mathbb{A}] \equiv \frac{1}{2i}(\mathbb{A} - \mathbb{A}^\dagger) \quad (11)$$

is the generalised imaginary part of a tensor. Employing equation (10) as well as the electric field given by equation (4), one obtains in agreement with [11] the vacuum expectation values of the dyadic products appearing in equation (9) in Fourier space:

$$\langle \hat{\mathbf{E}}(\mathbf{r}, \omega) \otimes \hat{\mathbf{E}}(\mathbf{r}', \omega') \rangle = \frac{\mu_0 \omega^2 \hbar}{\pi} \Im[\mathbb{G}(\mathbf{r}, \mathbf{r}', \omega)] \delta(\omega - \omega'). \quad (12)$$

This relation is essentially a form of the fluctuation–dissipation theorem, as first shown under very general conditions in [23]. Its role is to link the field correlations required for evaluation of the quantum stress tensor and the classical Green's function of the medium. Using similar relations for the remaining terms in equation (9), transforming back to position space and rotating to imaginary frequencies u yields

$$\begin{aligned} \langle \hat{\mathbb{T}} \rangle = & -\frac{\hbar}{2\pi} \int_0^\infty du \int_{\partial V} dA \cdot \left\{ \frac{\xi^2}{c^2} [\mathbb{G}^{(1)}(\mathbf{r}, \mathbf{r}, iu) + \mathbb{G}^{(1)T}(\mathbf{r}, \mathbf{r}, iu)] \right. \\ & + \overleftarrow{\nabla} \times [\mathbb{G}^{(1)}(\mathbf{r}, \mathbf{r}', iu) + \mathbb{G}^{(1)T}(\mathbf{r}', \mathbf{r}, iu)] \times \overleftarrow{\nabla}'|_{\mathbf{r}' \rightarrow \mathbf{r}} \\ & \left. - \text{tr} \left[\frac{\xi^2}{c^2} [\mathbb{G}^{(1)}(\mathbf{r}, \mathbf{r}, iu)] + \overleftarrow{\nabla} \times [\mathbb{G}^{(1)}(\mathbf{r}, \mathbf{r}', iu)] \times \overleftarrow{\nabla}'|_{\mathbf{r}' \rightarrow \mathbf{r}} \right] \mathbb{I} \right\}, \quad (13) \end{aligned}$$

from which the force can be computed by means of equation (8). In this formula the Green's tensor has been replaced by its scattering part $\mathbb{G}^{(1)}$ defined via

$$\mathbb{G} = \mathbb{G}^{(0)} + \mathbb{G}^{(1)}, \quad (14)$$

where $\mathbb{G}^{(0)}$ is the bulk part of the Green's tensor, which does not contribute to the Casimir force regardless of the system's geometry. In addition we exploit the fact that [20]

$$\lim_{|\omega| \rightarrow \infty} \mathbb{G}^{(0)}(\mathbf{r}, \mathbf{r}') = \lim_{|\omega| \rightarrow \infty} \mathbb{G}(\mathbf{r}, \mathbf{r}') = -\mathbb{I} \delta(\mathbf{r} - \mathbf{r}') \quad (15)$$

which functions as a cutoff for high frequencies, allowing one to obtain a finite result. Note in particular that we did not assume the validity of the Lorentz reciprocity condition $\mathbb{G}(\mathbf{r}, \mathbf{r}', \omega) = \mathbb{G}^T(\mathbf{r}', \mathbf{r}, \omega)$, which is connected with time reversal invariance [20]. We hence have derived an expression for the Casimir force of arbitrary nonreciprocal bodies, which can also be obtained as a particular case of results for general magneto-dielectrics (see, for example, [24, 25]).

3. Bi-isotropic media and PEMCs

In order to obtain a tuneable Casimir force we will consider a class of materials whose reflection behaviour is in some sense intermediate between the extreme cases of the perfect electric conductor (PEC) and perfect magnetic conductor (PMC), which are respectively characterised by infinite permittivity ε or infinite permeability μ . These materials are known as bi-isotropic (see, for example, [18]), and in macroscopic quantum electrodynamics the response of such a medium is conveniently described by four material constants; the familiar ε and μ , as well as two cross-polarizabilities ξ and ζ . In principle all these quantities are permitted to be tensor-valued, which leads to the more general case of bi-anisotropic media. We will confine ourselves to bi-isotropic media, in which the material response shows no direction-dependence. This means that the four material constants are scalar (or pseudo-scalar) valued and fulfil the constitutive relations

$$\hat{\mathbf{D}} = \varepsilon_0 \varepsilon \hat{\mathbf{E}} + \frac{1}{c} \xi \hat{\mathbf{H}}, \quad (16)$$

$$\hat{\mathbf{B}} = \mu_0 \mu \hat{\mathbf{H}} + \frac{1}{c} \zeta \hat{\mathbf{E}}, \quad (17)$$

where we have chosen our definitions in such a way that all four material constants are dimensionless. For a fundamental theory of linear material response see [26].

3.1. Duality transformation

By allowing for nonzero (or even infinite) cross-polarisabilities ξ and ζ we achieve an interpolation between PECs and PMCs. To do this we note that Maxwell's equations for classical fields in media in the absence of free charges of currents can be arranged in the following way

$$\nabla \cdot \begin{pmatrix} Z_0 \mathbf{D} \\ \hat{\mathbf{B}} \end{pmatrix} = 0, \quad (18)$$

$$\nabla \times \begin{pmatrix} \mathbf{E} \\ Z_0 \mathbf{H} \end{pmatrix} + i\omega \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} Z_0 \mathbf{D} \\ \mathbf{B} \end{pmatrix} = 0. \quad (19)$$

These equations are invariant under an $SO(2)$ transformation, i.e. they remain valid when the vectors of fields are multiplied with a matrix of the form

$$\mathcal{D} = \begin{pmatrix} \cos(\theta) & \sin(\theta) \\ -\sin(\theta) & \cos(\theta) \end{pmatrix}. \quad (20)$$

The fields forming a vector in this formalism are called dual partners. The constitutive relations for the quantised fields then read [19]

$$\begin{pmatrix} Z_0 \hat{\mathbf{D}} \\ \hat{\mathbf{B}} \end{pmatrix} = \frac{1}{c} \begin{pmatrix} \varepsilon & \xi \\ \zeta & \mu \end{pmatrix} \begin{pmatrix} \hat{\mathbf{E}} \\ Z_0 \hat{\mathbf{H}} \end{pmatrix} + \begin{pmatrix} 1 & \xi \\ 0 & \mu \end{pmatrix} \begin{pmatrix} Z_0 \mathbf{P}_N \\ \mu_0 \mathbf{M}_N \end{pmatrix}, \quad (21)$$

where the noise polarisation $\hat{\mathbf{P}}_N$ and magnetisation polarisation $\hat{\mathbf{M}}_N$ are related to the noise current $\hat{\mathbf{j}}_N$. Note that in case of reciprocal materials the reduced number of degrees of freedom stemming from having $\xi = -\zeta$ leads to the constraint that θ has to be a integer multiple of $\pi/2$, in which case the continuous symmetry of duality invariance hence reduces to a discrete \mathbb{Z}_4 -symmetry [19]. The consideration of polarisation of polarisation-mixing material constants ξ and ζ , however, restores the continuity of duality invariance [20].

3.2. Perfect electromagnetic conductors (PEMC)

We will now focus on PEMCs as a special case of bi-isotropic materials. The concept of PEMCs has been introduced by Lindell and Sihvola [17, 27, 28], finding applications in waveguide and antenna engineering [29]. At a boundary with normal vector \mathbf{n} the PEMC reflection properties are defined via

$$\mathbf{n} \cdot (Z_0 \mathbf{D} - M \mathbf{B}) = 0, \quad (22)$$

$$\mathbf{n} \times (Z_0 \mathbf{H} + M \mathbf{E}) = 0. \quad (23)$$

They show a transmission-free, polarisation-mixing reflection behaviour [17]. The pseudoscalar material parameter M interpolates between PEC ($M \rightarrow \infty$) and PMC ($M = 0$ boundaries). We can now relate M to the magnetoelectric material constants introduced in the previous section by comparing equations (22) and (23) with the general constitutive relations (21). One arrives at

$$\xi = \zeta = \pm \sqrt{\mu \varepsilon}, \quad (24)$$

$$M = \frac{\xi}{\mu} = \pm \sqrt{\frac{\varepsilon}{\mu}} \quad (25)$$

in the limit $\mu \rightarrow \infty$, $\varepsilon \rightarrow \infty$, with M being finite. In other words, a PEMC is a very specific limiting case of a bi-isotropic medium with a strong response. Though it is not obvious from equations (24) and (25), these equations are consistent with reciprocal media for the PEC ($\varepsilon \rightarrow \infty$, ζ , $\xi \ll \varepsilon$) and PMC ($\mu \rightarrow \infty$, ζ , $\xi \ll \mu$) limits, as detailed in [17]. As pointed out by Dzyaloshinskii [30] and further investigated in [31], Cr_2O_3 is a naturally occurring crystal with a weak nonreciprocal cross-polarisability. The close analogy of such an electromagnetic response with that of the PEMC as well as the Tellegen medium and the axion field in particle physics is also discussed in [32]

PEMC materials can be seen as the dual transform of a PEC by a finite duality transformation angle θ . Transforming the PEC-boundary conditions

$$\mathbf{n} \cdot \mathbf{B}^* = \mathbf{n} \cdot [-\sin(\theta) Z_0 \mathbf{D} + \cos(\theta) \mathbf{B}] = 0, \quad (26)$$

$$\mathbf{n} \times \mathbf{E}^* = \mathbf{n} \times [\sin(\theta) Z_0 \mathbf{H} + \cos(\theta) \mathbf{E}] = 0 \quad (27)$$

directly gives equations (22) and (23) if the identification

$$M = \cot(\theta) \quad (28)$$

is made and the positive sign in equations (24) and (25) is taken. This means that the PEC case corresponds to $\theta = 0$ and the PMC case to $\theta = \pi/2$, with all other cases appearing for intermediate angles in the range $(0, \pi/2)$. If the negative sign were included in equations (24) and (25), the angle θ would instead be equal to $-\pi/2$ for the PMC, running to $\theta = 0$ for the PEC.




Figure 1. Three-layer system with perfectly reflecting, cross-polarising boundary surfaces at $z = 0$ and $z = L$.

4. The Green's tensor of two PEMC plates

In order to apply our general result (13) to two PEMC plates, we first need to find the respective Green's tensor for the setup specified in figure 1.

4.1. General structure of the Green's tensor

The reflection properties of a nonreciprocal plate are described by four reflection coefficients r_{ss} , r_{sp} , r_{ps} , r_{pp} corresponding to all possible combinations of the polarisation directions (s or p) of the incoming and outgoing light. Here the index p denotes an electric field polarisation parallel to the plane of incidence (transverse-magnetic (TM) polarisation), while s indicates perpendicular polarisation (transverse-electric (TE) polarisation). The reflected wave \mathbf{v}_{refl} corresponding to a general incident wave \mathbf{v}_{inc} at a boundary described by these four coefficients can therefore be represented as a matrix multiplication:

$$\mathbf{v}_{\text{refl}} = \mathcal{R} \cdot \mathbf{v}_{\text{inc}} = \begin{pmatrix} r_{ss} & r_{sp} \\ r_{ps} & r_{pp} \end{pmatrix} \cdot \begin{pmatrix} v_s \\ v_p \end{pmatrix}. \quad (29)$$

A setup consisting of two plates is considered as a three-layer system, where we require the Green's tensor for all positions in the middle layer. This consists of waves travelling from \mathbf{r} to \mathbf{r}' and being reflected any number of times, which can be elegantly taken into account by means of a Neumann series, as is well-known (see, for example, [33]). For matrices \mathcal{R}^\pm representing two plates being located at $z = 0$ (\mathcal{R}^-) and $z = L$ (\mathcal{R}^+) respectively we define

$$\mathcal{D}_{\sigma_i \sigma_j}^\pm = \left[\sum_{n=0}^{\infty} (\mathcal{R}^\pm \cdot \mathcal{R}^\mp)^n \cdot (e^{-2ik^+L})^n \right]_{\sigma_i \sigma_j} = (\mathcal{I} - \mathcal{R}^\pm \cdot \mathcal{R}^\mp e^{-2ik^+L})_{\sigma_i \sigma_j}^{-1} \quad (30)$$

with σ_i, σ_j denoting the polarisation directions s and p and \mathcal{I} is the two-dimensional identity matrix. Using the general form of the Green's tensor we obtain the result

$$\begin{aligned} \mathbb{G}^{(1)}(\mathbf{r}, \mathbf{r}', \omega) &= \frac{1}{8\pi^2} \int \frac{d^2 k^\parallel}{k^\perp} e^{ik^\parallel(\mathbf{r}-\mathbf{r}')} \\ &\times [\mathbf{e}^+ \cdot \mathcal{R}^+ \cdot (\mathcal{D}^-)^{-1} \cdot \mathcal{R}^- \cdot \mathbf{e}^{+T} e^{ik^\parallel(2L+z-z')} + \mathbf{e}^- \cdot \mathcal{R}^- \cdot (\mathcal{D}^+)^{-1} \cdot \mathcal{R}^+ \cdot \mathbf{e}^{-T} e^{ik^\parallel(2L-z+z')} \\ &+ \mathbf{e}^- \cdot \mathcal{R}^- \cdot (\mathcal{D}^+)^{-1} \cdot \mathbf{e}^{+T} e^{ik^\parallel(z+z')} + \mathbf{e}^+ \cdot \mathcal{R}^+ \cdot (\mathcal{D}^-)^{-1} \cdot \mathbf{e}^{-T} e^{ik^\parallel(2L-z-z')}], \end{aligned} \quad (31)$$

where $\mathbf{e}^\pm = (\mathbf{e}_s^\pm, \mathbf{e}_p^\pm)$, with

$$\mathbf{e}_s^\mp = \mathbf{e}_s^\mp = \mathbf{e}_{k^\parallel} \times \mathbf{e}_z, \quad \mathbf{e}_p^\mp = 1/k(k^\parallel \mathbf{e}_z \mp k^\perp \mathbf{e}_{k^\parallel}) \quad \mathbf{k} = k^\parallel \mathbf{e}_{k^\parallel} + k^\perp \mathbf{e}_{k^\perp}. \quad (32)$$

Note that the matrix multiplication is performed in (s, p) -space. The Green's tensor's spatial components are obtained by the outer product of the respective polarisation vectors. In this expression the first two terms account for an even number of multiple reflections between \mathbf{r} and \mathbf{r}' , while odd numbers of reflections contribute to the final two terms. Similarly to the case of reciprocal materials [21], it turns out that the terms representing an odd number of reflections do not contribute to the Casimir force.

4.2. PEMC reflection matrices

The boundary conditions (22) and (23) for the fields lead to polarisation-mixing effects at a PEMC boundary. In terms of the magnetoelectric constants these reflection coefficients for radiation incident from medium 1 onto medium 2 are given by [10]

$$r_{ss} = \frac{(k_1^\perp - \mu k_2^\perp)\Omega_\varepsilon - k_1^\perp k_2^\perp \xi^2}{(k_1^\perp + \mu k_2^\perp)\Omega_\varepsilon + k_1^\perp k_2^\perp \xi^2}, \quad (33)$$

$$r_{ps} = \frac{-2\mu k_1^\perp k_2^\perp \xi}{(k_1^\perp + \mu k_2^\perp)\Omega_\varepsilon + k_1^\perp k_2^\perp \xi^2} = r_{sp}, \quad (34)$$

$$r_{pp} = \frac{\left[k_1^\perp - \left(\varepsilon - \frac{\xi^2}{\mu} \right) k_2^\perp \right] \Omega_\mu - k_1^\perp k_2^\perp \xi^2}{\left[k_1^\perp + \left(\varepsilon - \frac{\xi^2}{\mu} \right) k_2^\perp \right] \Omega_\mu + k_1^\perp k_2^\perp \xi^2}, \quad (35)$$

with $\Omega_\mu = \mu(k_1^\perp + \mu k_2^\perp)$, $\Omega_\varepsilon = \mu[k_1^\perp + (\varepsilon - \xi^2/\mu)k_2^\perp]$ and k_i^\perp representing the component of the wave vector perpendicular to the interface. In the PEMC-limit, with all response functions going to infinity and $M = \sqrt{\varepsilon/\mu}$, one obtains in matrix form:

$$\mathcal{R} = \begin{pmatrix} r_{ss} & r_{sp} \\ r_{ps} & r_{pp} \end{pmatrix} = \frac{1}{1 + M^2} \begin{pmatrix} 1 - M^2 & -2M \\ -2M & M^2 - 1 \end{pmatrix} \quad (36)$$

which is independent of the incoming wave vector.

Introducing the corresponding duality transformation angle θ via equation (28), one obtains for two plates:

$$\mathcal{R}^\pm = \begin{pmatrix} -\cos(2\theta^\pm) & \sin(2\theta^\pm) \\ \sin(2\theta^\pm) & \cos(2\theta^\pm) \end{pmatrix} \quad (37)$$

with θ^\pm being the respective duality transformation angle that defines the properties of each plate. We can now also calculate the corresponding multiple-reflection contributions to obtain

$$(\mathcal{D}^\pm)^{-1} = \frac{b}{1 - 2b \cos(2\delta) + b^2} \begin{pmatrix} b - \cos(2\delta) & \sin(2\delta) \\ -\sin(2\delta) & b - \cos(2\delta) \end{pmatrix} \quad (38)$$

with $b = e^{-2ik^\perp L}$ and $\delta = \theta^+ - \theta^-$.

5. Casimir force between two PEMC plates

In order to solve the Green's tensor integral we introduce polar coordinates (k^\parallel, φ) for the two-dimensional integral over \mathbf{k}^\parallel . This simplifies the calculation considerably because the reflection matrices as well as D^\pm and D^\mp do not depend on φ , the angular dependence appears only in the dyadic product of the polarisation vectors, which may be straightforwardly integrated.

We can compute a force $d\mathbf{F}/dA$ per unit area from the stress tensor via equation (8). Making use of the fact that $d\mathbf{A} \parallel \mathbf{e}_z$, we have


$$\begin{aligned} \mathbf{f} &= \frac{d\mathbf{F}}{dA} = -\frac{1}{A} \int_{\partial V} d\mathbf{A} \cdot \langle \mathbb{T} \rangle \\ &= -\frac{\hbar}{2\pi} \int_0^\infty du \sum_{j=x,y,z} \left(\langle \mathbb{T} \rangle_{jj} \mathbf{e}_j \right) \Bigg|_{z=L}, \end{aligned} \quad (39)$$

where $\langle \mathbb{T} \rangle$ is given by (13). The symmetry of the problem requires that \mathbf{f} has no x - or y - components which can indeed be seen from the fact that no combination of the polarisation vectors yields a x - z - or y - z -component when integrated over φ . We will hence suppress the fact that \mathbf{f} is a vector and just calculate its absolute value.

We now insert our obtained Green's tensor (31) into (13) and observe that the contributions from the terms containing curls equal the contributions from those without. After setting $\kappa = ik^\perp$, transforming to polar co-ordinates $k^\parallel = \kappa \cos(\phi)$, $u/c = \kappa \sin(\phi)$ and carrying out the trivial angular integration we get

$$f = \frac{\hbar c}{4\pi^2} \int_0^\infty d\kappa \kappa^3 e^{-2\kappa L} \text{Tr}[\mathcal{R}^+ \cdot (\mathcal{D}^-)^{-1} \cdot \mathcal{R}^- + \mathcal{R}^- \cdot (\mathcal{D}^+)^{-1} \cdot \mathcal{R}^+]. \quad (40)$$

This generalisation of Lifshitz's formula for planar systems [34] agrees with results for reciprocal polarisation-mixing plates such as gratings [35–37], which is itself a consequence of the general validity of the fluctuation-dissipation theorem. Remarkably, the result is hence insensitive to the fact that the plates are non-reciprocal at this level.



Substituting $x = \kappa L$ and performing the matrix multiplications we find the following integral

$$\begin{aligned}
 f &= -\frac{\hbar c}{\pi^2 L^4} \int_0^\infty dx x^3 \frac{e^{2x} \cos(2\delta) - 1}{1 - 2e^{2x} \cos(2\delta) + e^{4x}} \\
 &= -\frac{\hbar c}{\pi^2 L^4} \int_0^\infty dx x^3 \left(\frac{\frac{1}{2}e^{2x}(e^{2i\delta} + e^{-2i\delta})}{(1 - e^{2x}e^{2i\delta})(1 - e^{2x}e^{-2i\delta})} - \frac{1}{(1 - e^{2x}e^{2i\delta})(1 - e^{2x}e^{-2i\delta})} \right) \tag{41}
 \end{aligned}$$

which can be analytically integrated to finally obtain our main result

$$f(\theta^+, \theta^-) = -\frac{3\hbar c}{8\pi^2 L^4} \text{Re}(\text{Li}_4[e^{2i(\theta^+ - \theta^-)}]), \tag{42}$$

where we have made use of the polylogarithm function $\text{Li}_n(z) = \sum_{k=1}^\infty z^k/k^n$. Our result (42) immediately demonstrates an invariance of the Casimir force under duality transforms of material parameters; it only depends on the difference of the PEMC angles, so a simultaneously applied duality transformation does not change the Casimir force. Thus we may write $f(\theta^+, \theta^-) = f(\theta^+ - \theta^-) = f(\delta)$. We can easily check that this is indeed compatible with the results of Casimir and Boyer via

$$\text{Re Li}_4(e^{i\phi}) = \sum_{k=1}^\infty \frac{\cos^k(k\phi)}{k^4} = \frac{\pi^4}{90} - \frac{\pi^2 \phi^2}{12} + \frac{\pi \phi^3}{12} - \frac{\phi^4}{48}, \tag{43}$$

where we used de Moivre’s identity followed by formula (27.8.6(3)) of [38] (see also [39], (25.12(ii))), giving

$$f(\delta) = -\frac{\hbar c}{8\pi^2 L^4} \left[\frac{\pi^4}{30} - \delta^2(\pi - \delta)^2 \right]. \tag{44}$$

Then we obtain the special cases of Casimir ($\delta = 0$, corresponding to any choice of identical plates)

$$f(0) = -\frac{\hbar c}{240\pi^2 L^4} \tag{45}$$

and Boyer ($\delta = \pi/2$)

$$f(\pi/2) = \frac{7}{8} \cdot \frac{\hbar c}{240\pi^2 L^4}. \tag{46}$$

We show the results for intermediate angles in figure 2. It is seen that there is some value δ_{crit} for which there is no Casimir force, solving equation (44) for this gives

$$\delta_{\text{crit}} = \frac{\pi}{2} \left(1 - \sqrt{1 - 2\sqrt{\frac{2}{15}}} \right) \approx 0.96 \cdot \frac{\pi}{4}. \tag{47}$$

In the case of a scalar field under Robin boundary conditions, it has previously been shown numerically that such a ‘zero-force’ parameter exists between the extreme cases of attraction and repulsion [14], here we extend this to the electromagnetic field as well as finding an analytic value for δ_{crit} . It is also interesting to notice that the following holds

$$\int_0^{\pi/2} d\delta f(\delta) = 0, \quad (48)$$

so even though the force is not symmetric around the central angle $\delta = \pi/4$, the enclosed areas to the left and the right of the zero-force angle δ_{crit} are equal. Thus our result represents a sum rule for the Casimir force for PEMCs; the sum of Casimir forces over the entire PEMC parameter space is zero.

6. Conclusion

In order to calculate the Casimir force between two PEMC plates we have constructed the Green's tensor for two nonreciprocal plates in terms of their reflection properties. The result is duality invariant as well as it is compatible with the theorem derived by Kenneth and Klich that the Casimir force between identical bodies is always attractive [8]. It also verifies for a certain class of local nonreciprocal media described in section 2 the prediction that the Casimir force between two plates of any possible material will fall in between the results of Casimir and Boyer [7], which had thus far only been shown for magnetodielectrics. The derived Green's tensor is hence also applicable for different lossless material classes. In particular the focus might go to perfectly reflecting chiral materials ($\xi = -\zeta$ in terms of material constants) to explore the full parameter space of the Casimir effect.

For more realistic scenarios of course the corrections due to imperfect reflection or non-zero temperature are of high interest. For these cases the derived PEMC case can be viewed as a theoretical upper limit for the Casimir force since we assumed the reflection coefficients as well as the PEMC parameter to be frequency independent, and calculated the force at zero temperature. In less idealised cases one would expect the resulting Casimir force to lie somewhere 'under the curve' for the respective value of θ .

Acknowledgments

We would like to thank F Hehl for inspiring this work and S Fuchs, F Lindel, A Sihvola, C Henkel and M Bordag for stimulating discussions. The authors would like to thank an anonymous referee for a large number of constructive comments. We acknowledge financial support from the Deutsche Forschungsgemeinschaft via grants BU1803/3-1 and GRK 2079/1. RB and SYB additionally acknowledge support from the Alexander von Humboldt foundation, and SYB acknowledges support from the Freiburg Institute of Advanced Studies (FRIAS).

References

- [1] Casimir H B G 1948 On the attraction between two perfectly conducting plates *Proc. K. Ned. Akad.* **360** 793–5
- [2] Serry F M, Walliser D and Maclay G J 1995 The anharmonic Casimir oscillator (ACO)-the Casimir effect in a model microelectromechanical system *J. Microelectromech. Syst.* **4** 193–205
- [3] Tas N, Sonnenberg T, Jansen H, Legtenberg R and Elwenspoek M 1996 Stiction in surface micromachining *J. Micromech. Microeng.* **6** 385–97
- [4] Buks E and Roukes M L 2001 Stiction, adhesion energy, and the Casimir effect in micromechanical systems *Phys. Rev. B* **63** 033402
- [5] Woods L M, Dalvit D A R, Tkatchenko A, Rodriguez-Lopez P, Rodriguez A W and Podgornik R 2016 Materials perspective on Casimir and van der Waals interactions *Rev. Mod. Phys.* **88** 045003
- [6] Boyer T H 1974 Van der Waals forces and zero-point energy for dielectric and permeable materials *Phys. Rev. A* **9** 2078–84
- [7] Henkel C and Joulain K 2005 Casimir force between designed materials: what is possible and what not *Europhys. Lett.* **72** 929–35
- [8] Kenneth O, Klich I, Mann A and Revzen M 2002 Repulsive casimir forces *Phys. Rev. Lett.* **89** 033001
- [9] Grushin A G and Cortijo A 2011 Tunable Casimir repulsion with three-dimensional topological insulators *Phys. Rev. Lett.* **106** 020403
- [10] Fuchs S, Crosse J A and Buhmann S Y 2017 Casimir–Polder shift and decay rate in the presence of nonreciprocal media *Phys. Rev. A* **95** 023805
- [11] Butcher D T, Buhmann S Y and Scheel S 2012 Casimir–Polder forces between chiral objects *New J. Phys.* **14** 113013
- [12] Rosa F S S, Dalvit D A R and Milonni P W 2008 Casimir–Lifshitz theory and metamaterials *Phys. Rev. Lett.* **100** 183602
- [13] Asorey M and Muñoz-Castañeda J M 2013 Attractive and repulsive Casimir vacuum energy with general boundary conditions *Nucl. Phys. B* **874** 852–76
- [14] Romeo A and Saharian A A 2002 Casimir effect for scalar fields under Robin boundary conditions on plates *J. Phys. A: Math. Gen.* **35** 1297–320
- [15] Markov V N and Pis'mak Y M 2006 Casimir effect for thin films in QED *J. Phys. A: Math. Gen.* **39** 6525–32
- [16] Marachevsky V N 2017 Casimir effect for Chern–Simons layers in the vacuum *Theor. Math. Phys.* **190** 315–20
- [17] Lindell I V and Sihvola A H 2005 Perfect electromagnetic conductor *J. Electromagn. Waves Appl.* **19** 861–9
- [18] Serdyukov A, Semchenko I, Tretyakov S and Sihvola A 2001 *Electromagnetics of Bi-Anisotropic Materials: Theory and Applications* (Amsterdam: Gordon and Breach Science)
- [19] Buhmann S Y and Scheel S 2009 Macroscopic quantum electrodynamics and duality *Phys. Rev. Lett.* **102** 140404
- [20] Buhmann S Y, Butcher D T and Scheel S 2012 Macroscopic quantum electrodynamics in nonlocal and nonreciprocal media *New J. Phys.* **14** 083034
- [21] Buhmann S Y 2012 *Dispersion Forces I—Macroscopic Quantum Electrodynamics and Ground-state Casimir, Casimir–Polder and van der Waals Forces* (Springer Tracts in Modern Physics vol 247) (Berlin: Springer)
- [22] Eckhardt W 1982 First and second fluctuation–dissipation-theorem in electromagnetic fluctuation theory *Opt. Commun.* **41** 305–9

- [23] Agarwal G S 1975 Quantum electrodynamics in the presence of dielectrics and conductors: I. Electromagnetic-field response functions and black-body fluctuations in finite geometries *Phys. Rev. A* **11** 230–42
- [24] Kheirandish F, Soltani M and Sarabadani J 2010 Casimir force in the presence of a medium *Phys. Rev. A* **81** 052110
- [25] Philbin T G 2011 Casimir effect from macroscopic quantum electrodynamics *New J. Phys.* **13** 063026
- [26] Hehl F W and Obukhov Y N 2003 *Foundations of Classical Electrodynamics: Charge, Flux, and Metric* (Basel: Birkhäuser)
- [27] Lindell I V and Sihvola A H 2005 Transformation method for problems involving perfect electromagnetic conductor (PEMC) structures *IEEE Trans. Antennas Propag.* **53** 3005–11
- [28] Lindell I V and Sihvola A H 2005 Realization of the PEMC boundary *IEEE Trans. Antennas Propag.* **53** 3012–8
- [29] Sihvola A H and Lindell I V 2008 Perfect electromagnetic conductor medium *Ann. Phys.* **17** 787–802
- [30] Dzyaloshinskii I E 1960 On the magneto-electrical effect in antiferromagnets *J. Exp. Theor. Phys.* **37** 881
- [31] Hehl F W, Obukhov Y N, Rivera J-P and Schmid H 2008 Relativistic analysis of magnetoelectric crystals: extracting a new 4-dimensional P odd and T odd pseudoscalar from Cr₂O₃ data *Phys. Lett. A* **372** 1141–6
- [32] Zhang R Y, Zhai Y W, Lin S R, Zhao Q, Wen W and Ge M L 2015 Time circular birefringence in time-dependent magnetoelectric media *Sci. Rep.* **5** 13673
- [33] Chew W 1995 *Waves and Fields in Inhomogeneous Media* (Piscataway, NJ: IEEE)
- [34] Lifshitz E M 1956 The theory of molecular attractive forces between solids *J. Exp. Theor. Phys.* **2** 73–83
- [35] Lambrecht A and Marachevsky V N 2008 Casimir interaction of dielectric gratings *Phys. Rev. Lett.* **101** 160403
- [36] Contreras-Reyes A M, Guérout R, Maia Neto P A, Dalvit D A R, Lambrecht A and Reynaud S 2010 Casimir–Polder interaction between an atom and a dielectric grating *Phys. Rev. A* **82** 052517
- [37] Bender H, Stehle C, Zimmermann C, Slama S, Fiedler J, Scheel S, Buhmann S Y and Marachevsky V N 2014 Probing atom-surface interactions by diffraction of Bose–Einstein condensates *Phys. Rev. X* **4** 011029
- [38] Abramowitz M and Stegun I A 1970 *Handbook of Mathematical Functions: With Formulas, Graphs, and Mathematical tables* (New York: Dover)
- [39] Olver F W J et al *NIST Digital Library of Mathematical Functions*, <http://dlmf.nist.gov/>, (Release 1.0.16 of 2017-09-18)