

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18

Leaf traits related to productivity in *Populus deltoides* during the post-flooding period.

MARÍA E. RODRÍGUEZ¹, FABIO G. ACHINELLI¹², VIRGINIA M. C. LUQUEZ¹³

1 - Instituto de Fisiología Vegetal (INFIVE), CCT CONICET La Plata – FCAyF UNLP, CC 327, 1900

La Plata, Argentina.

2 – CIC Buenos Aires, Argentina.

3: Author for correspondence. vluquez@agro.unlp.edu.ar. Tel: +54-221-423-6618.

FAX: +54-221-423-3698.

19 **Abstract**

20 Flooding stress induces changes in trees at plant and leaf level that can reduce growth and
21 productivity. In this work, we explored changes in leaf traits related to productivity during the post-
22 flooding period in three poplar clones with different degrees of flooding sensibility. Our hypothesis was
23 that changes in leaf traits could lead to a higher photosynthetic activity in the post-flooding period to
24 compensate for the reduction in carbon fixation under flooding.

25 Plants were grown in pots in a greenhouse. Flooding was induced by filling the pots with tap
26 water up to 5 cm over the surface soil for 28 days. After this period, flooding ended and plant recovery
27 was followed for 42 days.

28 Flooding caused changes at plant and leaf level, not only during flooding but also after the
29 stress ended. During this post-flooding period, the formerly flooded plants of all clones produced
30 leaves with increased area and thickness compared to the control plants, but the photosynthetic rate
31 was not increased. The plants compensated for the reduced growth under flooding by substituting the
32 leaf area loss instead of increasing the photosynthetic activity.

33

34

35 **Key words:** *Populus deltoides* – flooding – leaf traits – photosynthesis

36

37

38 **Key Message:**

39 After a flooding period, *Populus deltoides* plants compensate for the reduced growth under flooding by
40 substituting the leaf area loss instead of increasing the leaf photosynthetic activity.

41

42 Introduction

43 The tolerance to flooding of woody plants varies according to species and genotypes, the age
44 of the plant, the degree of covering by water, the flood duration and the conditions of the floodwater
45 (Kozłowski 1997, Glenz et al. 2006). Among the most conspicuous responses to flooding, we can find
46 growth reduction, development of hypertrophied lenticels, adventitious roots and aerenchyma
47 formation; accelerated leaf senescence and abscission; changes in the absorption and availability of
48 mineral nutrients; and several metabolic changes caused by hypoxic or anoxic conditions (Kozłowski
49 1997, Bailey-Serres and Voeselek 2008). During root hypoxia, photosynthetic activity can be reduced
50 by stomatal closure in different poplar clones (Bejaoui et al. 2006, Gong et al. 2007, Guo et al. 2011).

51 In *Populus*, several morphological leaf traits are related to productivity: total leaf area (Rae et
52 al. 2004, Monclus et al. 2005, Marron et al. 2005), number of leaves on the main stem (Rae et al.
53 2004), individual leaf area (Monclus et al. 2005, Marron et al. 2005), specific leaf area (Marron et al.
54 2005), and stomatal density (Al Afas et al. 2006). Some of these traits are affected by flooding: in
55 *Populus trichocarpa x deltoides*, root hypoxia reduces leaf growth rate and final leaf size through the
56 reduction of both cell size and cell number (Smit et al. 1989); in *Populus angustifolia*, flooding reduces
57 leaf number and size (Rood et al. 2010); and in *Populus* plants with flooded roots, specific leaf weight
58 increases (i.e., specific leaf area decreases, Liu and Dickman 1992).

59 These flood-induced leaf modifications will probably affect plant productivity. Under flooding,
60 the combination of a reduced rate of leaf expansion and an acceleration of leaf senescence and
61 abscission can reduce the photosynthetically active leaf area, thus decreasing plant growth (Luquez et
62 al. 2012). This combined with a reduction in the photosynthesis rate due to stomatal closure results in
63 a reduced availability of photosynthates for growth. In addition to that, there are changes in dry matter
64 partitioning and a decrease in the root/shoot ratio (Kozłowski 1997).

65 In spite of the well-documented changes induced by flooding in leaf morphology and
66 physiology, little is known about the effects of these modifications in the post-flooding period, although
67 they are likely to affect growth recovery. These alterations cannot be neglected in a climate change
68 scenario, where areas with extensive poplar plantations like the Lower Paraná River Delta will
69 experience flooding events more frequently (Barros et al. 2006). Even when these flooding episodes
70 do not cause plant death, they may alter plant and leaf traits, with potentially lasting effects on forest
71 growth and productivity.

72 In a previous work, we identified three *Populus deltoides* clones planted in the Paraná Delta
73 area with different degrees of growth reduction under flooding. The degree of growth reduction
74 correlated with the overall reduction in total leaf area, individual leaf size and leaf expansion rate
75 (Luquez et al. 2012). In the present work, we explored more extensively the changes experienced by
76 these clones in the post-flooding period. We analyzed the changes induced by flooding in leaf traits
77 that affect productivity by comparing three cohorts of leaves: the first cohort -L1- expanded before
78 flooding induction, the second -L2- expanded during flooding, and the third -L3- expanded after the
79 flooding episode. Our hypothesis was that changes in leaf architecture and biochemistry could lead to
80 a higher net photosynthetic rate in the post-flooding period to compensate for the reduction in carbon
81 fixation under flooding.

82

83

84 **Material and Methods**

85 *Plant material, experimental design and stress treatment*

86 The *Populus deltoides* W. Bartram ex Marshall clones used in this work were Alton, Stoneville
87 67 (ST67) and 149-82. These clones were selected because they showed different degrees of growth
88 reduction under flooding in a previous experiment: Alton was tolerant, 149-82 was sensitive, and ST67
89 was sensitive but to a lesser degree than 149-82 (Luquez et al. 2012).

90 Two experiments were carried out. In the 2009 experiment, one-year-old cuttings of 60 cm
91 long were planted in 7 L pots filled with clay loam soil on August 7, 2009. The pots were placed in a
92 greenhouse in a completely randomized design, with 10 replicates for each clone and treatment.
93 Irradiance inside the greenhouse on clear days reached a maximum value of $1282 \mu\text{moles m}^{-2} \text{s}^{-1}$. Bud
94 flush occurred between August 20 and August 31, 2009. A slow-release commercial fertilizer (NPK
95 12:5:14 plus Mg, S, Ca, Zn, Fe, Mo and B) was added to the pots to ensure an adequate nutrient
96 availability. The dose was 1 g of fertilizer per pot, and the fertilization treatment was repeated twice
97 before the beginning of the flooding treatment. To avoid fungal diseases, the trees were treated once
98 a week with two commercial fungicides (Benomyl 50% WP and Carbendazim 50% SC). Before the
99 treatment, trees were pruned and only one shoot was kept, in order to minimize the variability induced
100 by several shoots per tree. Flooding started when the shoots were 2 months old, and was induced by
101 placing the potted trees inside a sealed 10 L pot filled with tap water up to approximately 5 cm above

102 soil level; water was added when necessary to keep this level. The control plants were watered
103 regularly to field capacity. The flooding stress treatment started on October 28, 2009 and lasted for 35
104 days.

105 In the 2011-2012 experiment, one-year-old cuttings of 20 cm long were planted in 4.5 L pots
106 filled with a 1:1 soil-sand mix. The plants were treated as described above, except for fertilization. Pots
107 were watered weekly with 50 ml of complete Hoagland solution (Legget and Frere 1971). Flooding
108 was induced as described above, by placing the potted trees inside a sealed 6 L pot. The flooding
109 stress treatment started on November 2, 2011 and lasted for 28 days. After that, the formerly flooded
110 plants were removed from the sealed pots, water was allowed to drain, and the plants were measured
111 for 44 days.

112 In the 2011-2012 experiment, three leaves were tagged in each plant, as described in Luquez
113 et al. (2012): one leaf expanded before flooding (L1), one leaf expanded during the period of flooding
114 (L2) and one leaf expanded after flooding ended (L3). Morphological, physiological and biochemical
115 measurements were carried out on these leaves (see below). Unless otherwise stated, all data
116 presented were measured in the 2011-2012 experiment.

117

118 *Growth measurements and microscopic observations*

119 Total shoot height was measured with a graduated stick. At the end of the experiment, all
120 leaves were scanned and the total leaf area (TLA) was determined with the Image J software
121 (<http://rsbweb.nih.gov/ij/>, Schneider et al. 2012). The individual leaf area (ILA) of leaves L1, L2 and L3
122 were determined in the same way. Dry mass was determined after drying leaves, shoots and roots at
123 65°C to constant weight. Specific leaf area (SLA, $\text{cm}^2 \text{g}^{-1}$) was determined by taking a leaf disc of
124 known area (2.27 cm^2) from each cohort and drying them to constant weight as described above. The
125 Relative Growth Rate (RGR) for stem height growth was calculated according to Whitehead and
126 Myerscough (1962).

127 Imprints were taken from the abaxial surface of leaves L1, L2 and L3 using clear lacquer and
128 transparent tape. The imprints were fixed on glass slides, observed at 20x and photographed with a
129 digital camera (Olympus Evolt E-330). Four pictures were taken for each imprint, each representing
130 one observation field. The number of stomata per field (stomatal density) and the total number of
131 epidermal cells per field (epidermal cell density) were counted using the Image J software

132 (<http://rsbweb.nih.gov/ij/>, Schneider et al. 2012), and the stomatal index (SI) was calculated according
133 to Masle et al. (2005):

134

$$135 \quad SI = (100 \times \text{stomatal density}) / (\text{stomatal density} + \text{epidermal cell density})$$

136

137 To determine leaf thickness, a piece of leaf around the main vein of leaves L1, L2 and L3 was
138 fixed in FAA (formalin-alcohol-acetic acid). The leaves were cut by hand with a razor blade; seven
139 cuttings were made of each sample. The cuttings were observed at 10x and photographed with a
140 digital camera (Olympus Evolt E-330) and three measurements of thickness were performed on each
141 side of the vein every 0.05 mm. Leaf thickness was calculated as an average of the six measurements
142 made in all seven cuttings.

143

144 *Gas exchange measurements*

145 Photosynthetic activity (A), transpiration and stomatal conductance (gs) were measured with
146 an IRGA CIRAS II, PP Systems in the experiment on the latest fully expanded leaf. Water Use
147 Efficiency (WUE) was measured as the ratio between A and transpiration. The measurements were
148 carried out between 10:00 am and 3:00 pm, under an irradiance of 1500 $\mu\text{moles m}^{-2} \text{s}^{-1}$.

149

150 *Chlorophyll and Rubisco content*

151 One 5-mm-diameter leaf disc (chlorophyll) and two 10-mm-diameter leaf discs (Rubisco) were
152 frozen in liquid nitrogen and stored at -80 °C until the determinations were carried out.

153 Chlorophyll content was determined using N,N-Dimethylformamide according to the method of
154 Inskeep and Bloom (1985).

155 Rubisco content was determined by SDS-PAGE according to Laemmli (1970). Two 1-cm-
156 diameter leaf discs were homogenized in 1X sample buffer (62.5mM Tris pH 6.8; 5% w/v SDS, 5% v/v
157 glycerol, 5% v/v β -mercaptoethanol) and centrifuged at 10,000 rpm for 8 min at 4 °C. For SDS-PAGE
158 analysis, proteins in the supernatant were separated in 1.5 mm thick minigels with 12% of acrylamide
159 concentration as in Laemmli (1970). A volume equivalent to 2.62 mm² of leaf area was loaded in each
160 lane. Proteins were visualized by staining with Coomassie Brilliant Blue R-250. Gels were digitized
161 and analyzed for background subtraction and banding density using the Image J software

162 (<http://rsbweb.nih.gov/ij/>). Three or four replicates per treatment were analyzed. The amount of
163 Rubisco large Sub-unit (LSU) was calculated as a percentage of the initial content.

164

165 *Statistical Analysis*

166 The statistical analysis was carried out with R software version 2.8.1 (R Development Core
167 Team, 2010). ANOVA and mean test were carried out using the **agricolae R** package.

168

169

170 **Results**

171 Dry matter partitioning was measured in the 2009 experiment (Fig. 1). Total dry weight was
172 significantly reduced only in 149-82, but flooding altered dry matter partitioning in all clones. Root
173 biomass was reduced in all clones and root/shoot ratio decreased in flooded plants compared to
174 controls (Fig. 1, in italics). However, the loss of root biomass in Alton was lower than in the other
175 clones. Its root biomass under flooding was reduced by 25% compared to control plants, while the
176 reduction in the other clones was of 52% (ST67) and 66% (149-82). Consequently, the root/shoot ratio
177 decreased by 35% in Alton flooded plants compared to controls, whereas it decreased by 50% in the
178 other clones.

179 The periodical growth in height was similar in both experiments; therefore, only data from 2011
180 are presented. During the first two weeks of flooding, there were no differences in height between
181 control and flooded plants, but marked differences began to appear among clones after three weeks
182 (Fig. 2). Flooding did not reduce height in Alton, with no differences in RGR between both treatments
183 (Fig.2, left hand side of the arrow). Flooded plants of 149-82 and ST67 (Fig. 2) reduced their height
184 after the third week of flooding, but RGR was only significantly reduced in 149-82 (Fig.2, left hand side
185 of the arrow). After four weeks, the flooding episode was ended and the plants were allowed to
186 recover and measured for another 42 days. At the end of the recovery period, there were no
187 differences in height between formerly flooded and non-flooded plants in Alton and ST67, while the
188 levels of formerly flooded plants in 149-82 were still significantly lower than those of non-flooded
189 plants. The RGR in the post-flooding period was significantly higher in formerly flooded plants of Alton
190 and ST67, but not in 149-82 (Fig.4 right hand side).

191 Total leaf area (TLA) was measured discriminating the area developed in the post-flooding
192 period from the area previously expanded (before/during flooding) (Fig. 3). After the 42-day-period of
193 recovery, there were no significant differences in TLA between control and formerly flooded plants for
194 any of the clones, but the relative number of leaves expanded before/during and after flooding was
195 different (data not shown). There were no significant differences in leaf area expanded before/during
196 flooding between control and flooded Alton, but it was significantly smaller in formerly flooded plants of
197 149-82 and ST67. The expanded area after the flooding period was significantly larger in formerly
198 flooded plants than in the control treatment in all clones.

199 A and g_s were measured throughout the flooding and the recovery periods (Fig. 4). Both
200 variables were reduced by flooding in all clones, but the reduction was less marked in Alton. After the
201 end of the stress, g_s of formerly flooded plants recovered to similar values as control plants. There
202 was a significant correlation between g_s and A in control and flooded plants, but the relation was
203 weaker in the post-flooding period, remaining significant only in ST67.

204 A, g_s and WUE were measured in leaves L1, L2 and L3 when they reached their full
205 expansion (Table 2); in the case of L2 and L3 it happened after the end of flooding. In the cohort
206 expanded during flooding (L2), A did not differ between treatments. g_s was significantly higher only in
207 149-82 flooded plants, while WUE decreased in all clones but only significantly in Alton. In the cohort
208 expanded in the post-flooding period (L3), there were no differences in A, g_s or WUE.

209 We determined ILA, SI, SLA and leaf thickness on the three cohorts, L1, L2 and L3 (Table 1).
210 On leaf L2, ILA and SLA were not significantly affected by flooding in any of the clones. Flooding
211 reduced SI in ST67 and increased leaf thickness in 149-82. In the cohort expanded during the post-
212 flooding period (L3, Table 1), LAI increased in all clones, albeit not significantly in 149-82. There was
213 no change in SLA, but leaf thickness increased significantly in all clones. SI decreased only in ST67.

214 We measured the chlorophyll and Rubisco content in all three cohorts of leaves (Table 3). We
215 did not find significant differences between flooded and control plants in any of the clones.

216

217

218 **Discussion**

219 In *Populus* and other species, flooding causes the root system to die back, and the most
220 tolerant genotypes develop new adventitious roots with aerenchyma (Kozłowski 1997, Cao and
221 Conner 1999). Our results confirm this, since the genotype with more tolerance -i.e., less growth
222 reduction under flooding- was Alton, which had a greater root biomass, newly developed roots with
223 aerenchyma, and a root/shoot ratio less affected by flooding. The most sensitive clone, 149-82,
224 developed neither hypertrophied lenticels nor adventitious roots (see additional figure 1). The variation
225 in root biomass seems to be related to the growth recovery capability after flooding. The extensive root
226 loss in 149-82 is the likely cause for the slow growth recovery in the post-flooding period. More roots
227 imply a higher capability for water transport and nutrient absorption, allowing for the maintenance of a
228 larger leaf area during flooding. In poplar, total leaf area often correlates with biomass accumulation
229 (Rae et al. 2004, Monclus et al. 2005, Marron et al. 2005). In our experiment, 42 days after the end of
230 the stress episode, TLA was not significantly different between control and formerly flooded plants.
231 However, when discriminating between the areas developed before/during the flooding and post-
232 flooding periods, a clear difference emerged. The formerly flooded plants developed a greater leaf
233 area than the controls during the recovery period, thus compensating for the area loss under flooding
234 due to an increased abscission. There was no difference in the number of leaves expanded after the
235 end of the flooding stress period (data not shown); hence, the difference is due to the increase in the
236 area of leaves expanded in the post-flooding period.

237 Growth rate depends ultimately on the carbon fixing capacity, and this can be reduced by
238 flooding stress (Bejaoui et al. 2006, Gong et al. 2007, Guo et al. 2011). We found a significant
239 correlation between g_s and A during flooding, suggesting that the main cause for carbon fixation
240 reduction is stomatal closure. But the correlation is weaker in the post-flooding period, suggesting that
241 other factors could have an influence on A . Several leaf traits that correlate with biomass accumulation
242 in poplar (Rae et al. 2004, Monclus et al. 2005, Marron et al. 2005) can be altered by different
243 environmental factors and stresses, like root hypoxia (Smit et al. 1989) and increased CO_2
244 concentration (Ceulemans et al. 1995). There are also differences among genotypes, leaf side and
245 leaf position in the canopy (Al Afas et al 2006, Dillen et al. 2008). It has been shown that a higher
246 stomatal density can enhance photosynthetic capacity in *Arabidopsis* (Tanaka et al. 2013). These
247 morphological and biochemical alterations of leaves could increase photosynthetic activity in the post-
248 flooding period, thus compensating for the reduction of leaf carbon fixation under flooding due to leaf

249 area reduction and stomatal closure. To answer this question, we measured several leaf traits related
250 to productivity in cohorts of leaves expanded before, during and after the flooding period (L1, L2 and
251 L3, respectively), and measured gas exchange when these leaves reached their full expansion. The
252 gas exchange measurements in L2 and L3 were taken after the end of the flooding period, when g_s
253 reached similar values as those of control plants. Consequently, any differences in photosynthetic
254 activity will be caused by alterations in the leaf architecture induced by flooding but not by a reduction
255 of g_s .

256 The area of leaf L2 decreased but not to the same extent as in our previous work (Luquez et
257 al. 2012). The cause of this difference may lie on the length of the flooding period, which was shorter
258 than in the previous experiment. Regarding SLA, there were differences only at clonal level but not
259 between treatments. In those experiments with longer flooding periods, we found a reduction in SLA
260 on these same clones (data not shown), as reported by Liu and Dickman (1992) for hybrid poplar. As
261 for ILA, it is likely that the length of the flooding period influenced SLA, as it does to other plant
262 responses to this stress (Kozłowski 1997, Glenz et al. 2006). The lack of a clear trend of change in the
263 morphological data, mirrored what happened with gas exchange, Rubisco and chlorophyll data for L2,
264 i.e., it did not show any differences caused by flooding.

265 The leaf expanded in the post-flooding period (L3) showed clear trends regarding leaf area
266 and thickness, since both increased in the formerly flooded plants. SLA did not change, possibly
267 because both area and width increased at the same time. SLA modulates maximum photosynthetic
268 rate (A_{max}) and nitrogen use efficiency on leaves of an ample range of species: leaves with higher SLA
269 have a higher A_{max} per unit leaf N (Reich et al. 1998). Our results seems to fit in this broader pattern,
270 since the lack of change in SLA was accompanied with no change in the photosynthetic rate or the
271 fraction of leaf N involved directly in the photosynthesis, represented by Rubisco and chlorophyll
272 content. *P. deltoides* plants growing under different combinations of water and nitrogen availability,
273 shows moderate plasticity in leaf traits (Funk et al. 2007) and this seems to be the case in our results
274 as well. There were changes in leaf thickness and ILA, but most of the leaf traits did not change.

275 Contrary to our hypothesis, there was no compensatory increase of the photosynthetic rate in
276 the post-flooding period. It seems that *Populus deltoides* plants increase their growth rate after
277 flooding by an increase in leaf area rather than by a higher photosynthetic capacity.

278

279 **Conflicts of interest:** the authors declare that they have no conflicts of interest.

280

281

282 Author contribution statement: MER carried out the most part of the experiments and the
283 statistical analysis, FGA helped with the experimental part, VMCL did part of the statistical
284 analysis and wrote the paper.

285

286

287 **Acknowledgements:**

288 This work was supported by grants from Agencia Nacional de Promoción Científica y Tecnológica
289 (ANPCyT PICT 487) and Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET PIP
290 269) to VMCL. Thanks to S. Martínez for helping with the microscopy work and to S. Cortizo for
291 providing clone 149-82. VMCL is a researcher from CONICET. MER held fellowships from ANPCyT
292 and CONICET.

293

294 **References**

295 Al Afas N, Marron N, Ceulemans R (2006) Clonal variation in stomatal characteristics related to
296 biomass production of 12 poplar (*Populus*) clones in a short rotation coppice culture. *Env Exp Bot* 58:
297 279-286.

298

299 Bailey–Serres J, Voisenek LACJ (2008) Flooding Stress: Acclimations and Genetic diversity. *Ann Rev*
300 *Plant Biol* 59: 313-339.

301

302 Barros V, Menéndez A, Natenzon C, Kokot R, Codignotto J, Re M, Bronstein P, Camilloni I, Ludueña
303 S, González S, Ríos D (2006) Vulnerability to floods in the metropolitan area of Buenos Aires under
304 future climate change. *AIACC Working Papers No 26*.
305 http://www.aiaccproject.org/working_papers/working_papers.html. Accessed on April 3, 2014.

306

- 307 Bejaoui Z, Albouchi A, Abassi M, El Aouni MH (2006) Influence d'une hydromorphie modérée ou
308 severe sur la production de biomasse et les échanges gazeux de plants de peuplier euraméricain. Can
309 J For Res 36: 2654-2665.
- 310
- 311 Cao FL, Conner WH (1999) Selection of flood tolerant *Populus deltoides* clones for reforestation
312 projects in China. For Ecol Manag 117: 211-220.
- 313
- 314 Ceulemans R, Van Praet L, Jiang XN (1995) Effects of CO₂ enrichment, leaf position and clone on
315 stomatal index and epidermal cell density in poplar (*Populus*). New Phytol 131: 99-107.
- 316
- 317 Dillen SY, Marron N, Koch B, Ceulemans R (2008) Genetic variation of stomatal traits and carbon
318 isotope discrimination in two hybrid poplar families (*Populus deltoides* S9-2 x *Populus nigra* Ghoy and
319 *P. deltoides* S9-2 x *P. trichocarpa* V24). Ann Bot 102: 399-407.
- 320
- 321 Funk JL, Jones CG, Lerdau MT (2007) Leaf and shoot level plasticity in response to different nutrient
322 and water availabilities. Tree Phys 27: 1731-1739.
- 323
- 324 Glenz C, Schlaepfer R, Iorgulescu I, Kienast F (2006) Flooding tolerance of Central European tree and
325 shrub species. For Ecol Manag 235: 1-13.
- 326
- 327 Gong JR, Zhang XS, Huang YM, Zhang CL (2007) The effects of flooding on several hybrid poplars
328 clones in Northern China. Agroforestry Syst 69: 77-88.
- 329
- 330 Guo XY, Huang Z, Xu A, Zhang X (2011) A comparison of physiological, morphological and growth
331 responses of 13 hybrid poplars clones under flooding. Forestry 84: 1-12.
- 332
- 333 Inskeep WP, Bloom PR (1985) Extinction coefficients of chlorophyll a and b in N,N-Dimethylformamide
334 and 80% acetone. Plant Physiol 77: 483-485.

- 335
- 336 Kozłowski TT (1997) Responses of woody plants to flooding and salinity. Tree Physiology Monograph
337 No 1. <http://www.pucrs.br/fabio/fisiovegetal/Encharcamento.pdf>. Accessed August 5, 2013.
- 338
- 339 Laemmli UK (1970) Cleavage of structural proteins during the assembly of the head of bacteriophage
340 T4. Nature 227: 680-685.
- 341
- 342 Leggett JE, Frere MH (1971) Growth and nutrient uptake by soybean plants in nutrient solutions of
343 graded concentrations. Plant Physiol 41: 457-460.
- 344
- 345 Liu Z, Dickmann DI (1992) Responses of two hybrid *Populus* clones to flooding, drought and nitrogen
346 availability. I. Morphology and growth. Can J Bot 70: 2265 – 2270.
- 347
- 348 Luquez VMC, Achinelli F, Cortizo S (2012) Evaluation of flooding tolerance in cuttings of *Populus*
349 clones used for forestation at the Paraná River Delta, Argentina. Southern Forests 74: 61-70.
- 350
- 351 Marron N, Villar M, Dreyer E, Delay D, Boudouresque E, Petit JM, Delmotte FM, Guehl JM, Brignolas
352 F (2005) Diversity of leaf traits related to productivity in 31 *Populus deltoides* x *Populus nigra* clones.
353 Tree Physiol 25: 425-435.
- 354
- 355 Masle J, Gilmore SR, Farquhar GD (2005) The ERECTA gene regulates plant transpiration in
356 Arabidopsis. Science 436: 866 – 869.
- 357
- 358 Monclus R, Dreyer E, Delmotte FM, Villar M, Delay D, Boudouresque E, Petit J, Marron N, Bréchet C,
359 Brignolas F (2005) Productivity, leaf traits and carbon isotope discrimination in 29 *Populus deltoides* x
360 *Populus nigra* clones. New Phytol 167: 53 – 62.
- 361
- 362 R Development Core Team (2010) R: A language and environment for statistical computing. R
363 Foundation for Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0. <http://www.R-project.org>.
364 Accessed November 9 2010.

365

366 Rae AM, Robinson KM, Street N, Taylor G (2004) Morphological and physiological traits influencing
367 biomass productivity in short-rotation coppice poplar. *Can J For Res* 34: 1488 – 1498.

368

369 Reich PB, Ellsworth DS, Walters MB (1998) Leaf structure (specific leaf area) modulates
370 photosynthesis-nitrogen relations: evidences from within and across species and functional groups.
371 *Functional Ecol* 12: 948-958.

372

373 Rood SB, Nielsen JL, Shenton L, Gill KM, Letts MG (2010) Effects of flooding on leaf development,
374 transpiration and photosynthesis in narrowleaf cottonwood, a willow-like poplar. *Photosynthesis Res*
375 104: 31-39.

376

377 Schneider CA, Rasband WS, Eliceiri KW (2012) NIH image to Image J: 25 years of image analysis.
378 *Nature Methods* 9 (7) 671-675.

379

380 Smit B, Stachowiak M, Van Volkenburgh E (1989) Cellular process limiting leaf growth in plants under
381 hypoxic root stress. *J Exp Bot* 40: 89-94.

382

383 Tanaka Y, Sugano S, Shimada T, Hara-Nishimura I (2013) Enhancement of leaf photosynthetic
384 capacity through increased stomatal density in *Arabidopsis*. *New Phytol* 198: 757-764.

385

386 Whitehead FH, Myerscough PJ (1962) Growth Analysis of plants. *New Phytol* 61: 314-321.

387

388 Table 1 - Individual Leaf Area (ILA, cm²), Stomatal Index (SI), Specific Leaf Area (SLA, cm² g⁻¹) and
 389 Leaf Thickness (μm) in three cohorts of poplar leaves. The first cohort (L1) completed its expansion
 390 before flooding induction, the second cohort (L2) expanded during the period of flooding, and the third
 391 cohort (L3) expanded after the end of the stress treatment. Means followed by the same letter do not
 392 differ significantly (p<0.05 LSD). C: control, F: flooded.

393

394

Treatment	Cohort	ILA	SI	SLA	Thickness
Alton C	L1	66.3 a	8.2 a	175 b	265 a
149-82 C	L1	89.3 b	9.0 b	181 b	221 b
ST67 C	L1	68.2 a	8.8 ab	222 a	221 b
Alton C	L2	102.3 a	8.8 a	95 b	309 b
Alton F	L2	98.1 a	9.0 a	94 b	315 b
149-82 C	L2	106.8 a	8.5 a	112 a	278 a
149-82 F	L2	98.5 a	8.9 a	106 a	295 c
ST67 C	L2	95.1 a	10.1 b	114 a	272 a
ST67 F	L2	103.1 a	8.8 a	113 a	278 a
Alton C	L3	78.9 c	7.7 a	99 b	310 d
Alton F	L3	112.5 ab	8.3 ab	107 ab	323 b
149-82 C	L3	102.8 ab	8.1 ab	110 ab	297 a
149-82 F	L3	116.2 a	7.7 a	114 a	326 b
ST67 C	L3	87.9 c	10.1 c	117 a	277 a
ST67 F	L3	127.6 b	8.9 b	118 a	289 c

395

396

397

398

399

400 Table 2 – Net Photosynthesis (A , $\mu\text{moles CO}_2 \text{ m}^{-2} \text{ s}^{-1}$), Stomatal Conductance (g_s , $\text{mmoles H}_2\text{O m}^{-2} \text{ s}^{-1}$) and instantaneous Water Use Efficiency (WUE) in three cohorts of poplar leaves. The first cohort
 401 (L1) completed its expansion before flooding induction, the second cohort (L2) expanded during the
 402 period of flooding, and the third cohort (L3) expanded after the end of the stress treatment. Means
 403 followed by the same letter do not differ significantly ($p < 0.05$ LSD). C: control, F: flooded.
 404

405

406

<i>Clone /</i>		<i>Leaf</i>		
<i>Treatment</i>	<i>Cohort</i>	<i>A</i>	<i>g_s</i>	<i>WUE</i>
Alton C	L1	17.3 a	322 a	3.78 a
149-82 C	L1	14.2 a	311 a	2.93 b
ST67 C	L1	16.4 a	262 a	3.93 ab
Alton C	L2	12.6 ab	144 ab	3.12 ab
Alton F	L2	14.2 b	196 b	2.65 bc
149-82 C	L2	11.2 a	123 a	2.74 ab
149-82 F	L2	10.7 a	191 b	2.16 c
ST67 C	L2	12.3 ab	116 a	3.20 a
ST67 F	L2	12.2 ab	157 ab	2.96 ab
Alton C	L3	15.0 a	98 b	6.85 a
Alton F	L3	13.7 a	89 ab	5.07 a
149-82 C	L3	16.3 a	94 ab	6.52 a
149-82 F	L3	13.1 a	77 ab	5.27 a
ST67 C	L3	14.4 a	71 a	6.15 a
ST67 F	L3	13.6 a	81 ab	5.43 a

407

408

409

410

411

412 Table 3 - Chlorophyll (Chl, $\mu\text{g cm}^{-2}$) and Rubisco content (as percentage of the initial content) in three
 413 cohorts of poplar leaves. The first cohort (L1) completed its expansion before flooding induction, the
 414 second cohort (L2) expanded during the period of flooding, and the third cohort (L3) expanded after
 415 the end of the stress treatment. Means followed by the same letter do not differ significantly ($p < 0.05$
 416 LSD). C: control, F: flooded.

417

418

<i>Clone / Treatment</i>	<i>Leaf Cohort</i>	<i>Chl a</i>	<i>Chl b</i>	<i>Total Chl</i>	<i>Rubisco LSU</i>
Alton C	L1	31.6 a	11.6 a	44.5 a	100
149-82 C	L1	29.7 a	11.6 a	41.3 a	100
ST67 C	L1	32.5 a	33.4 a	44.5 a	100
Alton C	L2	25.2 ab	10.1 ab	35.2 ab	73 a
Alton F	L2	26.5 a	10.3 a	36.8 a	82 a
149-82 C	L2	24.8 ab	9.9 ab	34.8 ab	88 a
149-82 F	L2	26.6 a	10.4 a	37.0 a	91 a
ST67 C	L2	23.5 bc	9.9 ab	33.3 bc	72 a
ST67 F	L2	21.4 c	9.4 b	30.8 c	66 a
Alton C	L3	27.1 ab	10.8 ab	37.9 ab	82 a
Alton F	L3	27.1 ab	10.9 ab	38.0 ab	76 a
149-82 C	L3	28.7 bc	11.2 bc	39.4 bc	100 a
149-82 F	L3	30.2 c	11.8 c	42.0 c	108 a
ST67 C	L3	25.0 a	10.3 a	35.3 a	85 a
ST67 F	L3	27.4 ab	11.3 bc	38.7 abc	87 a

419

420

421

422

423 **Legends to the figures**

424 Fig. 1 – Dry matter partitioning between roots, stem and leaves in three *Populus deltoides* clones -
425 Alton, 149-82 and ST67-, in the 2009 experiment. The root system of the plants was flooded (F) for 35
426 days, while the control plants (C) were maintained under well-drained conditions. Means with the
427 same letter do not differ significantly ($p < 0.05$ LSD) for total dry matter. In italics: root/shoot ratio for
428 each treatment and clone (shoot = stem + leaves).

429
430 Fig. 2 – Growth in height of three *Populus deltoides* clones: Alton, 149-82 and ST67. The treatments
431 were control (well-drained, black circles) and flooded (white circles). The arrows indicate the end of
432 the flooding treatment. The asterisks indicate statistically significant differences between control and
433 flooded plants of the same clone. Relative Growth Rate (RGR) values are multiplied by 10^3 . c: control,
434 f: flooding; pf: plants previously flooded.

435
436 Fig.3 - Total leaf area and area expanded after the end of flooding of three *Populus deltoides* clones:
437 Alton, 149-82 and ST67. The treatments were control (C) and flooded (F). In the 2011 experiment and
438 after 28 days of flooding, the plants were allowed to drain and their recovery was followed for 42 days.
439 Means with the same letter do not differ significantly ($p < 0.05$ LSD). Vertical bars: standard error of the
440 mean.

441
442 Fig. 4 - Net Photosynthesis (A , $\mu\text{moles CO}_2 \text{ m}^{-2} \text{ s}^{-1}$) and Stomatal Conductance (g_s , $\text{mmoles H}_2\text{O m}^{-2}$
443 s^{-1}) of three *Populus deltoides* clones: Alton, 149-82 and ST67. The treatments were control (well-
444 drained, black circles), flooded (white circles) and plants flooded after the end of the stress treatment
445 (grey circles). r : Pearson correlation coefficient. The asterisk indicates statistically significant
446 differences ($p < 0.05$).

447

FIG.1




FIG.2






FIG.3








FIG.4



Supplementary Figure 1

A - Cuttings of the clones used, showing that Alton (a) and ST67 (b) developed hypertrophied lenticels (HL) and adventitious roots (AL), while 149-82 (c) did not.



B - The adventitious roots had aerenchyma (marked with an arrow) that developed only in flooded plants. Length of the bar: 100 μm . C: Control. F: Flooded.

