

## Nota

Cuad. herpetol. 33(1): 29-31 (2019)

**New observations on diet of the South American two-striped forest-pitviper *Bothrops bilineatus smaragdinus* (Hoge, 1966)**Pablo J. Venegas<sup>1</sup>, Juan C. Chávez-Arribasplata<sup>1</sup>, Eduardo Almora<sup>1</sup>, Pablo Grilli<sup>2</sup>, Vilma Duran<sup>1</sup><sup>1</sup> Centro de Ornitología y Biodiversidad (CORBIDI). Calle Santa Rita 105, Urb. Los Huertos de San Antonio, Surco, Lima 33, Perú.<sup>2</sup> Cátedra de Ecología General y Recursos Naturales, Universidad Nacional Arturo Jauretche. Av. Calchaquí 6299, Florencio Varela (CP 1888) Provincia de Buenos Aires, Argentina.

Recibida: 14 Noviembre 2017

Revisada: 28 Febrero 2018

Aceptada: 29 Enero 2019

Editor Asociado: J. Goldberg

doi: 10.31017/CdH.2019.(2017-031)

## ABSTRACT

The arboreal, nocturnal and widely distributed snake *Bothrops bilineatus smaragdinus* (Squamata, Viperidae) is known as a generalist whose diet items have rarely been identified to species level. In this work we report three new items for its diet: an adult frog *Dendropsophus marmoratus* (Anura, Hylidae), an adult lizard *Thecadactylus solimoensis* (Squamata, Phyllodactylidae), and an adult bat *Carollia brevicauda* (Chiroptera, Phyllostomidae), the latter being the first record of predation of a Chiropteran by this species.

Key Words: Anurophagy; *Carollia brevicauda*; *Dendropsophus marmoratus*; Diet; *Thecadactylus solimoensis*; Saurophagy.

The venomous snake, *Bothrops bilineatus smaragdinus* (Hoge, 1966), is widely distributed across the western side of the Amazon basin, encompassing Amazonian Colombia and southern Venezuela, northern and western Brazil, eastern Ecuador, Peru and Bolivia (Campbell and Lamar, 2004). With mostly nocturnal habits, it is an arboreal snake with preference of low forest strata and upland *terra firme* forests (Duellman, 1978; Campbell and Lamar, 2004; Turci *et al.*, 2009; Valencia *et al.*, 2016). According to the preys recorded by some authors, *B. b. smaragdinus* feeds on frogs, lizards, birds and rodents (Carrillo de Espinoza, 1983; Cunha and Nascimento, 1993; Martins *et al.*, 2002; Campbell and Lamar, 2004). However, specific identified items in the diet of *B. b. smaragdinus* are rarely reported. Dixon and Soini (1986) found tree frogs *Osteocephalus* and *Hyla* (now *Boana* or *Dendropsophus*) in specimens from near Iquitos, Peru. Valencia *et al.* (2016) found remnants of *Anolis fuscoauratus* and rodents in the stomachs of Ecuadorian specimens.

Herein, we report three new items for the diet of *B. b. smaragdinus*, and the first record of predation of a Chiropteran by this species. The first predation event was observed in February, 2007 by P. Grilli. In this case (Fig. 1), a juvenile of *B. b. smaragdinus* was observed swallowing an adult *Dendropsophus*

*marmoratus* (Laurenti, 1768) at 1.4 m above the ground on a branch at Cashiriari (11°51'53.4" S; 72°46'45.4" W, 588 m a.s.l.), La Convención, Cusco department, Peru. Both individuals, snake and frog, were not collected.

For the other two records, we examined the stomach contents of two specimens of *B. b. smaragdinus* from the herpetological collection of the Centro de Ornitología y Biodiversidad (CORBIDI), Lima, Peru. In one of these, an adult male (CORBIDI 4799) collected in March 2008 by V. Duran at Corrientes river (03°03'54" S, 75°49'35" W, 2000 m a.s.l.), Andoas, Loreto department, Peru. We found an adult *Thecadactylus solimoensis* (Bergmann and Russell, 2007) (CORBIDI 5077). The prey's measurements were: Snout-vent-length (SVL): 136 mm, Head Width (HW): 19.9 mm and Head length (HL): 27.8 mm. Snake's measurements were: SVL 510 mm, TL 91 mm, HL 21.66 mm and HW 15.46 mm. The other specimen was an adult female (CORBIDI 11169), collected on 7 May 2012 by V. Duran at Malvinas (11°39'16.1" S; 73°1'5.8" W, 400 m a.s.l.), La Convención, Cusco department, Peru. Its stomach contained a recently swallowed adult of *Carollia brevicauda*. The bat measurements were SVL: 55 mm, HW: 9 mm. Snake's measurements were SVL: 657 mm, TL: 64 mm, HW: 12.2 mm.

Author for correspondence: [juancarlos.chavez@corbidi.org](mailto:juancarlos.chavez@corbidi.org)



**Figure 1.** A juvenile of *B. bilineatus smaragdinus* swallowing an adult *Dendropsophus marmoratus* (Anura, Hylidae) at Cashiriari, La Convención, Cusco department, Peru.

From the dietary items reported here, the bat represents the most interesting record. From the several events of bat consumption by snakes reported for Central and South America, only a small portion belongs to species of the Viperidae family. Previous reports are: *Bothriechis schlegelii* and *Bothrops asper* feeding on the bat *Glossophaga soricina* in Honduras and Mexico respectively (Groves, 1961; Villa and Lopez-Forment, 1966); *Crotallus pusillus* feeding on the bat *Eptesicus fuscus* (Duellman, 1961); and *Bothrops pauloensis* feeding on an undetermined species of bat in Central Brazil (Valdujo *et al.*, 2002). However, Esbérard and Vrcibradic (2007) have pointed out that it is not possible to determine if these records are from predation of live bats or from necrophagy, especially those records obtained from stomach contents.

Although, the bat *Carollia brevicauda* has been

previously reported as prey of another viper species, *Bo. schlegelii*, from western Ecuador (Meza-Ramos *et al.*, 2010), and the broad distribution of this bat species, as reported by Gardner (2007) (known from Panama to Peru, Bolivia and Southeastern Brazil), overlaps with that of *B. b. smaragdinus*; we concluded, that this is the first report of this kind of interaction between both species. The described nocturnal behavior for the bat includes resting in tree holes and hollow roots and having a preference for fruits of the Piperaceae family (Estrada-Villegas *et al.*, 2010; Ramos Pereira *et al.*, 2010). This behavior may indicate a particular predation activity pattern similar to that suggested for *Bo. schlegelii*, being this the selection of ambush sites for success in predation (Sorrel, 2009). If vipers waited near the Piperaceae fruit or near the bats' dens and resting places they would have the opportunity to ambush these bats. We suggest this may be a probable strategy of predation extended to other semi-arboreal pitvipers, such as *B. b. smaragdinus* and a topic worthy of further research.

#### Acknowledgements

We are indebted with K. Siu-Ting by the early review of the manuscript. Specimens were collected with the following permits: 72-2008-INRENA-IFFS-DCB and 594-2012-AG-DGFFS-DGEFFS.

#### Literature cited

- Campbell, J. A. & Lamar, W. W. 2004. The Venomous Reptiles of the Western Hemisphere. Volume II. Cornell University Press. Ithaca and London.
- Carrillo de Espinoza, N. 1983. Contribución al conocimiento de las serpientes venenosas del Perú de las familias Viperidae, Elapidae e Hydrophiidae (Ophidia: Reptilia). *Publico Museo Historia Natural "Javier Prado", Universidad Nacional Mayor San Marcos, Lima; (Ser. A, Zool.)* 30: 1-55
- Cunha, O. R. & Nascimento, F. P. 1993. Ofídios da Amazônia: as cobras da região Leste do Pará. *Boletim do Museu Paraense Emilio Goeldi. Serie Zoologia* 9: 1-191.
- Dixon, J. R. & Soini, P. 1986. The reptiles of the Upper Amazon Basin, Iquitos Region, Peru. II Crocodilians, turtles and snakes. Milwaukee Public Museum. Milwaukee.
- Duellman, W. E. 1961. The amphibians and reptiles of Michoacán, México. *Publications of the University of Kansas Museum of Natural History* 15: 1-148.
- Duellman, W. E. 1978. The Biology of an Equatorial Herpetofauna in Amazonian Ecuador. Miscellaneous Publication N°65: University of Kansas. Lawrence.
- Esbérard, C. E. & Vrcibradic, D. 2007. Snakes preying on bats: new records from Brazil and a review of recorded cases in the Neotropical Region. *Revista Brasileira de Zoologia* 24: 848-853.

- Estrada-Villegas, S.; Pérez-Torres, J. & Stevenson, P. R. 2010. Ensamblaje de murciélagos en un bosque subandino colombiano y análisis sobre la dieta de algunas especies. *Mastozoología Neotropical* 17: 31-41.
- Gardner, A. L. 2007. Mammals of South America. volume 1: marsupials, xenarthrans, shrews, and bats. University of Chicago Press. Chicago, Illinois, and London.
- Groves, F. 1961. A feeding record of the palm viper, *Bothrops schlegelii*. *Herpetologica* 17: 277.
- Martins, M.; Marques, O. A. & Sazima, I. 2002. Ecological and phylogenetic correlates of feeding habits in Neotropical pitvipers of the genus *Bothrops*: 307-328. In: Schuett, G.W.; Höggren, M.; Douglas, M.E. & Greene, H.W. (eds.), *Biology of the Vipers* Eagle Mountain Publishing. Eagle Mountain.
- Meza-Ramos, P.; Almendáriz, A. L. & Yáñez-Muñoz, M. H. 2010. Datos sobre la dieta de *Bothriechis schlegelii* (Berthold, 1846) (Serpentes-Viperidae) en el occidente del Ecuador. *Boletín técnico 9, Serie Zoológica* 6: 15-18.
- Ramos Pereira, M. J.; Marques, J. T. & Palmeirim, J. M. 2010. Ecological responses of frugivorous bats to seasonal fluctuation in fruit availability in Amazonian forests. *Biotropica* 42: 680-687.
- Schuett, G. W.; Höggren, M.; Douglas, M. E. & Greene H. W. 2002. *Biology of the vipers*. Eagle Mountain Publishing. Eagle Mountain, UT.
- Sorrell, G. G. 2009. Diel movement and predation activity patterns of the eyelash palm-pitviper (*Bothriechis schlegelii*). *Copeia* 2009: 105-109.
- Turci, L. C. B.; Albuquerque, S.; Bernarde, P. S. & Miranda, D. B. 2009. Uso do hábitat, atividade e comportamento de *Bothriopsis bilineatus* e de *Bothrops atrox* (Serpentes: Viperidae) na floresta do Rio Moa, Acre – Brasil. *Biota Neotropica* 9: 197-206.
- Valencia, J. H.; Garzón-Tello, K. & Barragán-Paladines, M. E. 2016. Serpientes Venenosas del Ecuador: Sistemática, Taxonomía, Historia Natural, Conservación, Envenenamiento y Aspectos Antropológicos. Fundación Herpetológica Gustavo Orcés, Universidad de Texas. Fondo Ambiental Nacional. Quito.
- Valdujo, P. H., C. Nogueira & Martins, M. 2002. Ecology of *Bothrops neuwiedii pauloensis* (Serpentes: Viperidae: Crotalinae) in the Brazilian Cerrado. *Journal of Herpetology* 36: 169-176.
- Villa, B. & Lopez-Forment, W. 1966. Cinco casos de depredación de pequeños vertebrados en murciélagos de México. *Anales del Instituto de Biología de la Universidad de México* 37: 187-193.

© 2019 por los autores, licencia otorgada a la Asociación Herpetológica Argentina. Este artículo es de acceso abierto y distribuido bajo los términos y condiciones de una licencia Atribución-No Comercial 2.5 Argentina de Creative Commons. Para ver una copia de esta licencia, visite <http://creativecommons.org/licenses/by-nc/2.5/ar/>

