

# A new species of *Zamischus* (Hymenoptera, Cynipoidea, Figitidae) from Brazil

Norma Díaz<sup>1</sup>, Fabiana Gallardo<sup>1</sup>, Cecilia Dorfey<sup>2</sup> & Andreas Köhler<sup>2</sup>

<sup>1</sup>División Entomología, Facultad de Ciencias Naturales y Museo, UNLP, Paseo del Bosque s/n, 1900 La Plata, Buenos Aires, Argentina. [ndiaz@fcnym.unlp.edu.ar](mailto:ndiaz@fcnym.unlp.edu.ar)

<sup>2</sup>Laboratório de Entomologia, Universidade de Santa Cruz do Sul, Av. Independência 2293, Bairro Universitário, 96815-900 Santa Cruz do Sul-RS, Brazil. [andreas@unisc.br](mailto:andreas@unisc.br)

**ABSTRACT.** A new species of *Zamischus* (Hymenoptera, Cynipoidea, Figitidae) from Brazil. A new species of *Zamischus* from Brazil is described and diagnosed. The species is closely related to *Z. brasiliensis* but differs in the morphology of antennae and scutellum. *Z. aquilesi* **sp. nov.** was collected via Malaise trapping in a tobacco field in Santa Cruz do Sul, Rio Grande do Sul.

**KEYWORDS.** Brazil; Cynipoidea; Eucilinae; Figitidae.

**RESUMO.** Uma nova espécie de *Zamischus* (Hymenoptera, Cynipoidea, Figitidae) do Brasil. Uma nova espécie de *Zamischus* do Brasil é descrita e diagnosticada. A espécie é intimamente relacionada a *Z. brasiliensis*, mas difere na morfologia da antena e do escutelo. *Z. aquilesi* **sp. nov.** foi coletada com armadilha de Malaise em uma lavoura de tabaco em Santa Cruz do Sul, Rio Grande do Sul.

**PALAVRAS-CHAVE:** Brasil; Cynipoidea; Eucilinae; Figitidae.

Species of *Zamischus* Ashmead are minute, stick-like wasps, which belong to the parasitic Eucilinae, the richest and most diverse subfamily of Figitidae in the Neotropical Region (Díaz *et al.* 2008). This genus was proposed by Ashmead (1903a) with the type-species, *Z. brasiliensis* Ashmead, 1903, designated, but not described. Weld (1921) described and illustrated the type-species, which he later added to his key to the world genera of Cynipoidea (Weld 1952). Yoshimoto (1971) described *Z. elongatus* Yoshimoto, 1971 and the male of *Z. brasiliensis*. Later Díaz (1974) recorded *Z. brasiliensis* from Argentina. The two known species of *Zamischus* are from South America. The purpose of this study is to describe a new species of this genus collected in a tobacco field in Santa Cruz do Sul, RS, Brazil, with Malaise trap.

## MATERIAL AND METHODS

One specimen (female) was studied. It is housed at Coleção Zoológica do Departamento de Biologia, Universidade de Santa Cruz do Sul, Rio Grande do Sul, Brazil. Terminology used in descriptions follows Fontal-Cazalla *et al.* (2002). The photographs were taken with a Canon Powershot A 520 adapted to a Leica stereomicroscope (S8APO).

## RESULTS

### *Zamischus* Ashmead

*Zamischus* Ashmead 1903a: 221, 1903b: 60, 68; Dalla Torre & Kieffer 1910: 101; Rohwer & Fagan 1917: 378; Weld 1921: 437, 1952: 103, 189; Yoshimoto 1971: 979; Díaz 1974: 18.

Type species. *Zamischus brasiliensis* Ashmead.

**Diagnosis.** Species of *Zamischus*, as well as *Perischus* Weld, 1931 possess a propodeum that is produced posteriorly into a long tapering neck and the metasoma is attached to the propodeum via an elongate slender petiole; the result is an overall long and slender body. In *Zamischus*, the pronotal plate has two teeth on posterior margin; mesoscutum is smooth, marginal cell of fore wing is open at base as well as on margin, and the petiole is sulcate. In *Perischus* the posterior margin of the pronotal plate is shallowly emarginate, mesoscutum is transversely striate, marginal cell is closed, and the petiole dull and striate.


**Remarks.** *Zamischus* + *Perischus* lineage which is characterized by a series of striking autapomorphies have few characters providing reliable indications of their affinities to other eucilines and is possible that many of the Neotropical grade taxa, such as these, represent relictual, morphologically isolated South American lineages and that many of the intermediate taxa are now extinct, making it difficult to resolve relationships correctly (Fontal-Cazalla *et al.* 2002; Buffington *et al.* 2007)

**Distribution.** Brazil and Argentina.

### *Zamischus aquilesi* Díaz & Gallardo **sp. nov.**

(Figs. 1–7)

Female (Fig. 1). Total length 5 mm. General coloration black; mandible reddish brown; fore leg yellowish brown, base of femur and apical tarsal segment darker; mid leg yellowish brown, base of coxa, trochanter, femur and apical tarsal segment darker; tarsal segments of hind leg, except last,



Figs. 1–7. *Zamischus aquilesei* Díaz & Gallardo **sp. nov.** Female: 1, habitus (lateral view) (scale bar, 1mm); 2, head (lateral view); 3, antenna; 4, flagellomeres 1 and 2; 5, mesosoma (lateral view); 6, mesoscutum and scutellum (dorsal view); 7, fore wing. (scale bar, 0.5 mm).

yellowish brown; veins of wing yellow; base of syntergum and hypopygium reddish brown.

Head (Figs. 2–4). Smooth and polished with scattered setiferous punctures on face and genae, broader than thorax, slightly tilted forward and ovate in profile. Gena not margined; malar groove present. Antennae with 11 flagellomeres, filiform, inserted on prominence about middle of face; first flagellomere longer than scape plus pedicel, shorter than second flagellomere, slightly curved, distally swollen, smooth; flagellomeres 2–11 subequal in length, cylindrical, ridged longitudinally, with rhinaria.

Mesosoma (Figs. 5–7). In lateral view longer than high. Pronotal plate smooth, posterior margin with two triangular teeth, slanting anteriorly in diagonal plane (as seen in profile). Sides of pronotum smooth and polished, depression of anterodorsal corner finely striate; highly setose behind genae. Mesoscutum (Fig. 6) smooth, in dorsal view longer than wide; notauli absent; with two pairs of setiferous punctures and a longitudinal depression in the posterior half; parascutal impressions present. Scutellum (Fig. 6) rounded behind; disk minutely punctate; scutellar plate large, dorsal surface slightly convex, densely punctate, each puncture with single short seta, tapering anteriorly into septum between scutellar fovea, glandular release pit on posterior-facing slope; lateral bars longitudinally striate, length half of scutellum. Mesopleuron smooth, mesopleural carina present. Metapleuron longer than high, with two longitudinal carinae, area between carinae concave, anteroventral cavity conspicuous, setose. Hind coxa cylindrical. Fore wing (Fig. 7) with a single longitudinal vein and basal cross vein; marginal cell long, narrow, open at base and margin; surface sparsely pubescent, margin ciliate. Propodeum as long as the half of petiole; propodeal carinae running along dorsal margin, auxiliary propodeal carinae present; area between propodeal carinae minutely punctate.

Metasoma (Fig. 1). Petiole very long (more than twice the length of the propodeum, 23:10), longitudinally ridged, with two dorsal longitudinal carinae, lesser prominent carinae laterally; gaster short, ovate, laterally compressed, glabrous at base; syntergum present.

Male. Unknown.

Material examined. Holotype. 1 female, UNISC, Brazil: Rio Grande do Sul, Santa Cruz do Sul, 14XII-2009, Malaise trap, Dorfey Coll.

Etymology. Dedicated to Dr. Carlos Aquiles Darrieu (Facultad de Ciencias Naturales y Museo, Universidad Nacional de La Plata, Buenos Aires, Argentina)

Distribution. Brazil: Rio Grande do Sul.

Biology. Host unknown.

Remarks. Female of *Z. aquilesi* **sp. nov.** is more similar to *Z. brasiliensis* than to *Z. elongatus*. In both species, the petiole is very long and the gaster is short, but the species

can be separated from one another by the antennae and scutellar morphology; *Z. aquilesi* **sp. nov.** possesses antenna filiform and large densely punctate scutellar plate (antenna widened toward apex and small scutellar plate with peripheral row of setiferous punctures in *Z. brasiliensis*). The female of *Z. elongatus* is separated from *Z. brasiliensis* and *Z. aquilesi* by the presence of a short petiole and elongate gaster.

#### ACKNOWLEDGEMENTS

We thank Marta Loiácono, Matthew Buffington and an anonymous reviewer for the critical review of the manuscript, Mariano Lucía for to take the photographs, Cecilia Gorretta and Paulina Hernández for technical support (CIC). Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Comisión de Investigaciones Científicas de la Provincia de Buenos Aires (CIC), Universidad Nacional de La Plata (UNLP) – Argentina and Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPQ) – Brazil, for their constant support.

#### REFERENCES

- Ashmead, W. H. 1903a. Classification of the gall-wasps and parasitic cynipoids, of the Superfamily Cynipoidea. II. *Psyche* **10**: 59–72.
- Ashmead, W. H. 1903b. Some new genera in the Cynipoidea. *Proceedings of the Entomological Society of Washington* **5**: 222.
- Buffington, M. L.; J. A. A. Nylander & J. M. Heraty. 2007. The phylogeny and evolution of Figitidae (Hymenoptera: Cynipoidea). *Cladistics* **23**: 403–431.
- Dalla Torre K. & J. Kieffer. 1910. Cynipoidea. *Das Tierreich* **24**: 1–891.
- Díaz, N. B. 1974. Anotaciones sobre cinipoideos argentinos. I. (Hymenoptera). *Neotrópica* **20**: 17–20.
- Díaz, N. B.; F. E. Gallardo; A. L. Gaddi; M. Jiménez; P. Ros-Farré; J. Paretas-Martínez & J. Pujade-Villar. 2008. Avances en el conocimiento de las Figitidae Neotropicales (Hymenoptera, Cynipoidea), p. 141–158. *In*: J. Llorente-Bousquet, J. & A. Lanteri (eds.). *Contribución taxonómica en órdenes de insectos hiperdiversos*. Mexico D. F., Las Prensas de Ciencias, UNAM, viii+221 p.
- Fontal-Cazalla, F. M.; M. L. Buffington; G. Nordlander; J. Liljeblad; P. Ros-Farré; J. L. Nieves-Aldrey; J. Pujade-Villar & F. Ronquist. 2002. Phylogeny of the Eucoilinae (Hymenoptera: Cynipoidea: Figitidae). *Cladistics* **18**: 154–199.
- Rohwer, S. & M. Fagan. 1917. The type-species of the genera of the Cynipoidea or the gall wasps and the parasitic cynipoids. *Proceedings of the United States National Museum* **53**: 357–380.
- Weld, L. H. 1921. Notes on certain genera of parasitic Cynipidae proposed by Ashmead with descriptions of genotypes. *Proceedings of the United States National Museum* **59**: 433–451.
- Weld, L. H. 1931. Additional notes on types with description of a new genus (Hymenoptera: Cynipidae). *Proceedings of the Entomological Society of Washington* **33**: 220–227.
- Weld, L. H. 1952. *Cynipoidea (Hymenoptera) 1905–1950*. Ann Arbor, Privately printed. 351 p.
- Yoshimoto, C. M. 1971. A new species of the genus *Zamischus* and description of the male *Z. brasiliensis* (Hymenoptera: Cynipidae: Eucoilinae). *The Canadian Entomologist* **103**: 979–982.