

Auxotrophy Accounts for Nodulation Defect of Most *Sinorhizobium meliloti* Mutants in the Branched-Chain Amino Acid Biosynthesis Pathway

Maria de las Nieves Peltzer,¹ Nicolas Roques,² V erena Poinso, ^{3,4} O. Mario Aguilar,¹ Jacques Batut,² and Delphine Capela²

¹Instituto de Biotecnolog a y Biolog a Molecular, CCT-La Plata, Universidad Nacional de La Plata, Facultad de Ciencias Exactas, Calles 47 y 115, 1900-La Plata, Argentina; ²Laboratoire des Interactions Plantes-Microorganismes, UMR CNRS-INRA 2594/441, F-31320 Castanet Tolosan, France; ³Universit  Paul Sabatier, UMR 5623, Laboratoire des IMRCP, F-31062 Toulouse, France; ⁴CNRS, UMR 5623, Laboratoire des IMRCP, F-31062 Toulouse, France

Submitted 24 February 2008. Accepted 1 June 2008.

Some *Sinorhizobium meliloti* mutants in genes involved in isoleucine, valine, and leucine biosynthesis were previously described as being unable to induce nodule formation on host plants. Here, we present a reappraisal of the interconnection between the branched-chain amino acid biosynthesis pathway and the nodulation process in *S. meliloti*. We characterized the symbiotic phenotype of seven mutants that are auxotrophic for isoleucine, valine, or leucine in two closely related *S. meliloti* strains, 1021 and 2011. We showed that all mutants were similarly impaired for nodulation and infection of the *Medicago sativa* host plant. In most cases, the nodulation phenotype was fully restored by the addition of the missing amino acids to the plant growth medium. This strongly suggests that auxotrophy is the cause of the nodulation defect of these mutants. However, we confirmed previous findings that *ilvC* and *ilvD2* mutants in the *S. meliloti* 1021 genetic background could not be restored to nodulation by supplementation with exogenous amino acids even though their Nod factor production appeared to be normal.

Rhizobia are soil bacteria which are able to fix nitrogen in symbiotic association with leguminous plants. The fixation process occurs in a specialized organ, the nodule, which is the end point of a complex developmental program. The process starts with a molecular recognition system that enables specific rhizobia to enter root hairs. Most rhizobia penetrate the root through the formation of infection threads that cross several cell layers in the root. Simultaneously, root cortical cells divide and differentiate to form the nodule. Then, bacteria are released from the infection threads into the plant cells and, in some legumes, differentiate into nitrogen-fixing bacteroids.

Signal molecules involved in the early recognition process were previously identified. Flavonoid compounds present in root exudates induce activation of what are collectively called

nodulation genes (*nod*, *nol*, and *noe* genes), which are present in most rhizobia. The activity of nodulation proteins yields *N*-acylated oligomers of *N*-acetyl-D-glucosamine, known as Nod factors, which play a role in both nodule organogenesis and nodule colonization. The common nodulation genes *nodABC* are unique to rhizobia and are found in most rhizobia. Specific nodulation genes (*nodMnolFGnodN*, *nodIJ*, *nodQPGEF*, *nodH*, and *nodLnoeAB* in *Sinorhizobium meliloti*) determine the specific structure of Nod factors and are characteristic of rhizobium species. However, only mutations in *nodABC* and *nodH* genes in *S. meliloti* lead to a strict Nod⁻ phenotype (no nodule formation) on *Medicago* spp. host plants (Roche et al. 1991). Inactivation of other nodulation genes just delays nodulation or affects infection, sometimes in a plant-specific way (Ardourel et al. 1995; Debelle  et al. 1986; Demont et al. 1993). For some mutants, the absence of nodulation phenotype might be due to gene redundancy in the genome (e.g., *nodM* and *nodPQ*). In *S. meliloti*, expression of the nodulation genes is regulated by three LysR-type NodD regulators. NodD1 and NodD2 are transcriptional activators that depend on plant inducers, flavonoids, and betaines, respectively. NodD3 activates *nod* genes independently of plant compounds and its expression depends on a complex regulation involving the regulatory proteins SyrM and NodD1 and the NtrBC two-component system which responds to the nitrogen status of the cell (Dusha and Kondorosi 1993; Maillet et al. 1990).

Apart from mutations in nodulation genes, very few mutations leading to a Nod⁻ phenotype (no nodule formed) have been described in *S. meliloti*. Auxotrophic mutations in a number of amino acid biosynthetic pathways such as histidine, asparagine, cysteine, or arginine biosynthesis do not impair the ability of *S. meliloti* to initiate nodule formation (Randhawa and Hassani 2002), whereas several mutants that affect the biosynthesis of branched-chain amino acids (BCAA) isoleucine, valine, and leucine have repeatedly been shown to be defective in the initiation of nodule formation on host legumes (Aguilar and Grasso 1991; Hassani et al. 2002; Lopez et al. 2001; Pobigaylo et al. 2008; Sanjuan-Pinilla et al. 2002; Truchet et al. 1980). This suggested a specific and important role of the BCAA biosynthesis pathway in the nodulation process in *S. meliloti*.

Aguilar and Grasso (1991) first characterized a mutant in the *ilvC* gene encoding a key enzyme in the isoleucine and valine biosynthesis which was unable to form nodules on alfalfa even when isoleucine and valine were added to the plant culture

Corresponding author: D. Capela; Telephone: +1 33 5 61 28 54 54; Fax: +1 33 5 61 28 50 61; E-mail: Delphine.Capela@toulouse.inra.fr

*The e-Xtra logo stands for "electronic extra" and indicates that three supplementary tables and two supplementary figures are published online.

This article is in the public domain and not copyrightable. It may be freely reprinted with customary crediting of the source. The American Phytopathological Society, 2008.

medium. Lopez and associates (2001) further demonstrated that both induction of nodulation genes by addition of the plant flavone luteolin and Nod factor production were strongly affected in this mutant. More recently, Hassani and associates (2002) showed that not only an *ilvC* mutant but also an *ilvI/B/G* mutant in the *S. meliloti* Rmd201 strain were impaired for nodulation and infection processes on *Medicago sativa*. As in the Aguilar and Grasso study (1991), the Nod⁻ phenotype of both mutants was not rescued by addition of exogenous isoleucine and valine. Another *S. meliloti* auxotrophic strain defective for host legume nodulation was identified by Sanjuan-Pinilla and associates (2002). This strain was mutated in *leuA1*, the gene encoding the first enzyme of the leucine biosynthesis pathway; nodulation gene expression was also affected in this mutant. In this case, addition of leucine to the plant growth medium allowed the mutant to nodulate alfalfa roots, although with a slower kinetics than the wild-type strain. Truchet and associates (1980) also reported a leucine auxotrophic mutant in *S. meliloti* strain L5-30 that formed ineffective nodules, inside which mutants were sequestered in infection threads. It was also found that addition of leucine or leucine precursors restored the capacity of the mutants to invade plant cells and fix nitrogen. In contrast, two independent *ilvD* mutants that were auxotrophic for isoleucine and valine amino acids in strains 1021 and Rmd201 were found to be symbiotically active on *Medicago* plants (Aguilar and Grasso 1991; Hassani et al. 2002).

Hence, no clear picture emerged from these different studies regarding the link between BCAA synthesis and nodulation in *S. meliloti* (Supplementary Table S1 provides a summary of previously published data). In particular, it was not clear whether the entire isoleucine, valine, and leucine biosynthetic pathway is necessary for the nodulation process in *S. meliloti* or whether only certain enzymes or intermediary products of the pathway are required. Another question that remained unanswered was whether all the Nod⁻ auxotrophic strains are impaired for Nod factor production and whether this could account for the nodulation defect of these strains.

Here, we present a complete reappraisal of the role of the BCAA biosynthesis pathway in the nodulation process in *S. meliloti*. In two different *S. meliloti* genetic backgrounds, we constructed or identified a set of mutants that were auxotrophic for isoleucine, valine, or leucine and that affected almost all enzymatic steps of the biosynthetic pathway. We then systematically and concomitantly assessed their symbiotic phenotype on host plants in the absence or presence of the required amino acids. We also systematically monitored their ability to express nodulation genes in response to the flavone, luteolin.

RESULTS

Identification of the *S. meliloti* genes involved in the BCAA biosynthesis pathway.

The biosynthesis of BCAA valine, isoleucine, and leucine occurs through a parallel set of reactions beginning with pyruvate and α -ketobutyrate (Fig. 1). Four common enzymes are required for the biosynthesis of isoleucine and valine: an acetohydroxy acid synthase (AHAS), an acetohydroxy acid isomeroreductase, a dihydroxy-acid dehydratase, and a BCAA aminotransferase. In *Escherichia coli*, these enzymes are encoded by the *ilvBN* (AHAS isozyme I), *ilvGM* (AHAS isozyme II), *ilvIH* (AHAS isozyme III), *ilvC*, *ilvD*, and *ilvE* genes, respectively. Leucine is formed from α -ketoisovalerate, an intermediate in valine biosynthesis. Leucine biosynthesis requires three additional enzymes: α -isopropylmalate synthase, α -isopropylmalate isomerase, and β -isopropylmalate dehydrogenase. These enzymes are encoded by *leuA*, *leuC*, *leuD*, and

leuB genes in *E. coli*. Likely orthologs of all *E. coli* genes have been identified on the *S. meliloti* genome, except for genes encoding the two regulatory subunits of AHAS-I and -II, *ilvN* and *ilvM*. Some of the *ilv* and *leu* genes have paralogs in *S. meliloti*, including 3 *ilvB*, 5 *ilvD*, 2 *leuA*, and 2 *ilvE* genes. To identify the *S. meliloti* genes involved in BCAA biosynthesis, we tested the auxotrophy or prototrophy of mutants of most of these genes on minimal medium supplemented or not with either isoleucine and valine or leucine. Mutants were either available from Tn5 mutant libraries in 2011 or 1021 *S. meliloti* strains (Milcamps et al. 1998; Pobigaylo et al. 2006) or constructed during the course of this study. The results of auxotrophy assays are listed in Table 1. Mutation in *ilvI* (SMc01431), which affects the first common step of the isoleucine and valine biosynthesis pathway, led to auxotrophy, whereas single mutants in *ilvB1* (SMc02263), *ilvB2* (SMb21530), and *ilvG* (SMc04455) were prototrophs. Mutation in *ilvH* (SMc01530), the only homologue of the regulatory subunit of the AHAS complex, was a prototroph, which is consistent with the fact




Fig. 1. Predicted branched-chain amino acid biosynthesis pathway in *Sinorhizobium meliloti*. The figure is adapted from Hassani and associates (2002). AHAS, acetohydroxy acid synthase; AHAI, acetohydroxy acid isomeroreductase; DHAD, dihydroxy acid dehydratase; BCAA AT, branched-chain amino acid aminotransferase; IPMS, α -isopropylmalate synthase; IPMI, α -isopropylmalate isomerase; IPMDH, β -isopropylmalate dehydrogenase. *S. meliloti* genes involved in each enzymatic reaction are indicated based on auxotrophic mutations. The *ilvIH* genes putatively form an operon. Mutations in *ilvIH*, *ilvC*, and *ilvD2* are prototrophic for leucine because α -ketoisovalerate can be produced by IlvE from valine and 2-oxoglutarate.

that, in enterobacteria, IlvH has no enzymatic activity per se but is required for full IlvI activity as well as for the inhibition of IlvI activity by valine (Mendel et al. 2001). All the *ilvD* mutants, except *ilvD2* (SMc04045), were prototrophs, indicating that only *ilvD2* is involved in the synthesis of isoleucine and valine. Similarly, a single mutation in the *leuA1* gene (SMc02117) led to leucine auxotrophy, indicating that the *leuA2* copy (SMc02546) may not be required for the pathway. We did not obtain auxotrophic mutations affecting the BCAA aminotransferase IlvE, the last step of the BCAA biosynthesis pathway. The two single *ilvE1* (SMc02896) and *ilvE2* (SMc00042) mutants were prototrophs, suggesting that the two *ilvE* copies may be functionally redundant.

All auxotrophic mutations in the BCAA biosynthetic pathway led to a defect in nodulation.

We inoculated *M. sativa* plants with auxotrophic mutants that were affected in the different steps of the BCAA biosynthesis pathway and subsequently monitored nodule formation for approximately 40 days. Mutants were obtained from two closely related strains, 1021 and 2011. We observed that all mutants exhibited a severe nodulation defect with, however, slight differences in phenotype between the two genetic backgrounds, 1021 and 2011 (Table 2). In strain 2011, nodules induced by *ilvI*, *ilvC*, *ilvD2*, *leuA1*, and *leuB* mutants appeared on roots approximately 20 days after those induced by the wild-type strain. The nodules formed did not fix nitrogen and cytological observations showed that these nodules were not invaded by bacteroids, although a few infection threads could be observed in the leucine auxotrophs 2011 *leuA1* and 2011 *leuB* (not shown). In strain 1021, the phenotype observed for *ilvC* and *ilvD2* was more pronounced than for *ilvI*, *leuA1*, *leuB*, *leuC*, and *leuD* because these two mutants exhibited a strong delay in nodulation (20 days after the wild-type strain) and the few nodules formed were mainly pseudonodules (root deformations such as bumps due to few cortical cell divisions) or abnormally developed nodules showing multiple or central vascular strands. None of these nodules or pseudonodules was invaded. Nodulation was also delayed in the 1021 *ilvI*, *leuA1*, *leuB*, *leuC*, and *leuD* mutants but the delay was shorter than in strain 2011 (approximately 7 days after the wild-type strain). However, we observed that the symbiotic phenotypes of 1021 *ilvI*, *leuA1*, and *leuD* mutants were biased by the presence of a number of nitrogen-fixing nodules induced by mutant revertants. Isolation of bacteria from pink nodules showed that these nodules were invaded by prototrophic neomycin-sensitive bacteria. These mutants were constructed by plasmid insertions that revert at a higher frequency in plants, where they are not subjected to selection pressure, than the Tn5 mutations which are present in the other mutants (Table 1).

We concluded from this first set of plant tests that all mutations in the BCAA biosynthesis pathway led to a defect in nodulation in *S. meliloti* strains 2011 and 1021, contrary to what was previously shown for the *ilvD* (Nod⁺ Fix⁺), *leuC/D*, and *leuB* (Nod⁺ Fix⁻) mutants (Aguilar and Grasso 1991; Hassani et al. 2002). All the delayed nodules formed were not invaded and, therefore, displayed no nitrogen fixation.

Addition of exogenous amino acids restored both prototrophy and nodulation in strain 2011.

Addition of relevant amino acids (isoleucine and valine for *ilv* mutants or leucine for *leu* mutants) to the growth medium of *M. sativa* plants completely restored the nodulation ability of all 2011 auxotrophs (Table 2). In these conditions, mutant-induced nodules appeared at the same time as wild-type-induced nodules or with a delay of only a few days. However,

the nodules formed by *ilv* mutants in the presence of isoleucine and valine were not effective and cytological studies showed that, even though some infection threads were formed, nodule cells were not at all or only weakly invaded by bacteroids. This suggests that isoleucine and valine concentrations inside the plant cells may limit bacterial growth and, hence, the infection capacity of corresponding auxotrophs. In contrast, nodules induced by *leu* mutants in the presence of exogenous leucine were invaded in the normal way and fixed nitrogen, indicating that leucine may be provided by the plant to the bacteria during the infection process as well as in the symbiosome.

Strains 2011 and 1021 behaved differently with respect to the symbiotic phenotype of *ilvC* and *ilvD2* mutants.

Supplementation experiments performed with 1021 auxotrophs showed that the nodulation ability of *ilvI*, *leuA1*, *leuB*, *leuC*, and *leuD* strains could be fully restored by addition of exogenous amino acids in a way similar to that observed with 2011 mutants. In contrast, the nodulation ability of the *ilvD2* mutant was only partially rescued by addition of isoleucine and valine and no effect was observed on the nodulation ability of *ilvC* when isoleucine and valine were added to the plant growth medium (Fig. 2). In the presence of isoleucine and valine, nodules induced by the 1021 *ilvD2* mutant appeared with a marked delay (approximately 10 days) compared with wild-type nodules. These nodules were devoid of bacteria and, thus, ineffective. In the case of the 1021 *ilvC* mutant, very few nodules were formed even in the presence of amino acids and most were only pseudonodules (cortical cell divisions that were not invaded), as in the absence of exogenous amino acids. Cytological observations of roots inoculated with the 1021 *ilvC* and *ilvD2* strains in the presence of amino acids showed some root hair curling and the formation of very few infection threads that aborted rapidly (not shown).

In order to rule out the possible occurrence of secondary mutations affecting nodulation in these two strains, we complemented the 1021 *ilvC* and 1021 *ilvD2* mutants with the wild-type *ilvC* and *ilvD2* genes constitutively expressed in trans from a plasmid. Complementation of the 1021 *ilvC* and 1021 *ilvD2* mutants fully restored the prototrophy, nodulation, and nitrogen-fixing abilities of these strains (Fig. 2).

To try to understand the phenotypic difference between the 1021 and 2011 *ilvC* and *ilvD2* mutants, we compared the growth rates of the four mutants in minimal medium liquid cultures containing different isoleucine and valine concentrations (0.04, 0.4, and 0.8 mM). Because the growth rates of mutants were similar between the two genetic backgrounds (data not shown), we inferred that the 1021 *ilvC* and *ilvD2* nodulation behavior was not due to a less efficient import of BCAA in strain 1021.

We also ruled out the possibility that the mutation itself could account for the phenotypic difference between the 1021 *ilvC* (GMI11498) and 2011 *ilvC* (GMI11523) by reconstructing the same *ilvC* mutation (Tn5 insertion 744 bp after the start codon) using the pQB2 plasmid in the 2011 genetic background (strain GMI11544). The nodulation ability of this strain was fully restored by addition of isoleucine and valine into the plant growth medium like the former 2011 *ilvC* (GMI11523) mutant (data not shown).

Nod factor production in the *ilvC* and *ilvD2* mutants was not affected.

Two mutations that led to BCAA auxotrophy, *ilvC* and *leuA1*, have previously been shown to have impaired nodulation gene expression in *S. meliloti* (Lopez et al. 2001; Sanjuan-Pinilla et al. 2002). We wanted to know whether all auxotrophic mutations in the BCAA biosynthesis pathway were similarly impaired for

nodulation gene expression. Consequently, we assessed the expression of the key nodulation gene, *nodC*, in the absence and presence of the plant inducer luteolin in the 1021 and 2011 auxotrophs overexpressing the *nod* gene activator NodD1 on the pRmM57 plasmid (Mulligan and Long 1989). NodC expression was determined by measuring β -galactosidase activity of the *nodC-lacZ* translational fusion also carried by the pRmM57

plasmid (Table 3). In these experiments, auxotrophic strains were cultured in the presence of 0.8 mM isoleucine, valine, or leucine to allow normal growth of the strains.

Again, strains 1021 and 2011 behaved differently with respect to *nod* gene expression. Previous work demonstrated that *nod* gene expression in the presence of luteolin is higher in strain 1021 than in strain 2011, probably due to a different amount of

Table 1. Bacterial (*Sinorhizobium meliloti*) strains and plasmids

Strains, plasmids	Description ^a	Auxotrophy	Reference
Strains			
GMI11495	2011, Str ^r derivative of 2011	...	J. Dénarié
GMI11496	1021, Str ^r derivative of 2011	...	Meade et al. 1982
GMI11497	1021 <i>ilvB1</i> ::pVO155 (881 bp), Str ^r , Neo ^f	...	This work
GMI11498	1021 <i>ilvC</i> ::Tn5 (744 bp), Str ^r , Neo ^f	Ile, Val	This work
GMI11499	1021 <i>ilvD2</i> ::Tn5 (\approx 900 bp), Str ^r , Neo ^f	Ile, Val	This work
GMI11500	1021 <i>ilvE1</i> ::pVO155 (491 bp), Str ^r , Neo ^f	...	This work
GMI11501	1021 <i>ilvH</i> ::Tn5-luxAB (167 bp), Str ^r , Neo ^f	...	Milcamps et al. 1998
GMI11502	1021 <i>ilvI</i> ::pVO155 (962 bp), Str ^r , Neo ^f	Ile, Val	This work
GMI11503	1021 <i>leuA1</i> ::pVO155 (709 bp), Str ^r , Neo ^f	Leu	This work
GMI11504	1021 <i>leuB</i> ::Tn5-luxAB (372 bp), Str ^r , Neo ^f	Leu	Milcamps et al. 1998
GMI11505	1021 <i>leuC</i> ::Tn5-luxAB (742 bp), Str ^r , Neo ^f	Leu	Milcamps et al. 1998
GMI11506	1021 <i>leuD</i> ::pVO155 (324 bp), Str ^r , Neo ^f	Leu	This work
GMI11507	1021 SMc04348::pVO155 (398 bp), Str ^r , Neo ^f	Ile, Val	This work
GMI11508	1021 pGMI50122, Str ^r , Gen ^f	...	This work
GMI11509	GMI11498 pGMI50120, Str ^r , Neo ^f , Gen ^f	...	This work
GMI11511	GMI11499 pGMI50121, Str ^r , Neo ^f , Gen ^f	...	This work
GMI11512	GMI11507 pGMI50120, Str ^r , Neo ^f , Gen ^f	...	This work
GMI11513	GMI11507 pGMI50122, Str ^r , Neo ^f , Gen ^f	Ile, Val	This work
GMI11540	1021 <i>sitD</i> ::pK19mob2HMB Ω (844 bp), Str ^r , Neo ^f	...	A. Becker
GMI11541	1021 Tn5, Str ^r , Neo ^f	...	This work
GMI11514	1021 pRmM57	...	Capela et al. 2005
GMI11515	GMI11498 pRmM57, Str ^r , Neo ^f , Tet ^f	Ile, Val	This work
GMI11516	GMI11499 pRmM57, Str ^r , Neo ^f , Tet ^f	Ile, Val	This work
GMI11517	GMI11502 pRmM57, Str ^r , Neo ^f , Tet ^f	Ile, Val	This work
GMI11518	GMI11503 pRmM57, Str ^r , Neo ^f , Tet ^f	Leu	This work
GMI11519	GMI11504 pRmM57, Str ^r , Neo ^f , Tet ^f	Leu	This work
GMI11520	GMI11505 pRmM57, Str ^r , Neo ^f , Tet ^f	Leu	This work
GMI11521	GMI11506 pRmM57, Str ^r , Neo ^f , Tet ^f	Leu	This work
GMI11542	GMI11540 pRmM57, Str ^r , Neo ^f , Tet ^f	...	This work
GMI11543	GMI11541 pRmM57, Str ^r , Neo ^f , Tet ^f	...	This work
GMI11522	2011 <i>ilvB2</i> ::mTn5 (693 bp), Str ^r , Neo ^f	...	Pobigaylo et al. 2006
GMI11523	2011 <i>ilvC</i> ::mTn5 (391 bp), Str ^r , Neo ^f	Ile, Val	Pobigaylo et al. 2006
GMI11524	2011 <i>ilvD1</i> ::mTn5 (512 bp), Str ^r , Neo ^f	...	Pobigaylo et al. 2006
GMI11525	2011 <i>ilvD2</i> ::mTn5 (583 bp), Str ^r , Neo ^f	Ile, Val	Pobigaylo et al. 2006
GMI11526	2011 <i>ilvD3</i> ::mTn5 (406 bp), Str ^r , Neo ^f	...	Pobigaylo et al. 2006
GMI11527	2011 <i>ilvD4</i> ::mTn5 (19 bp), Str ^r , Neo ^f	...	Pobigaylo et al. 2006
GMI11528	2011 <i>ilvD5</i> ::mTn5 (931 bp), Str ^r , Neo ^f	...	Pobigaylo et al. 2006
GMI11529	2011 <i>ilvE2</i> ::mTn5 (239 bp), Str ^r , Neo ^f	...	Pobigaylo et al. 2006
GMI11530	2011 <i>ilvG</i> ::mTn5 (719 bp), Str ^r , Neo ^f	...	Pobigaylo et al. 2006
GMI11531	2011 <i>ilvI</i> ::mTn5 (966 bp), Str ^r , Neo ^f	Ile, Val	Pobigaylo et al. 2006
GMI11532	2011 <i>leuA1</i> ::mTn5 (1379 bp), Str ^r , Neo ^f	Leu	Pobigaylo et al. 2006
GMI11533	2011 <i>leuB</i> ::mTn5 (765 bp), Str ^r , Neo ^f	Leu	Pobigaylo et al. 2006
GMI11544	2011 <i>ilvC</i> ::Tn5 (744 bp), Str ^r , Neo ^f	Ile, Val	This work
GMI11534	2011 pRmM57	...	This work
GMI11535	GMI11523 pRmM57, Str ^r , Neo ^f , Tet ^f	Ile, Val	This work
GMI11536	GMI11525 pRmM57, Str ^r , Neo ^f , Tet ^f	Ile, Val	This work
GMI11537	GMI11531 pRmM57, Str ^r , Neo ^f , Tet ^f	Ile, Val	This work
GMI11538	GMI11532 pRmM57, Str ^r , Neo ^f , Tet ^f	Leu	This work
GMI11539	GMI11533 pRmM57, Str ^r , Neo ^f , Tet ^f	Leu	This work
Plasmids			
pBBR1MCS-5	Cloning vector, Gen ^f	...	Kovach et al. 1995
pGMI50120	pBBR1MCS-5 containing the <i>ilvC</i> gene, Gen ^f	...	This work
pGMI50121	pBBR1MCS-5 containing the <i>ilvD2</i> , Gen ^f	...	This work
pGMI50122	pBBR1MCS-5 containing the SMc04348 gene, Gen ^f	...	This work
pK19mob2HMB Ω	Suicide plasmid, Kan ^f	...	A. Becker
pMA6B	7.5-kb <i>EcoRI</i> fragment from MA6-B (1021 <i>ilvD2</i> ::Tn5) strain in pSUP102, Neo ^f , Tet ^f	...	Grasso, unpublished
pRmM57	pSymA <i>nod</i> region, <i>nodC-lacZ</i> translational fusion, Tet ^f	...	Mulligan and Long 1989
pRK600	Helper plasmid, Chl ^f	...	Finan et al. 1986
pQB2	11-kb <i>EcoRI</i> fragment from strain 1028 (1021 <i>ilvC</i> ::Tn5) in pSUP202, Neo ^f , Tet ^f	...	Aguilar and Grasso 1991
pVO155	Suicide plasmid, Kan ^f , Amp ^f	...	Oke and Long 1999
pXLGD4	<i>hemA-lacZ</i> translational fusion, Tet ^f	...	Leong et al. 1985

^a The location of the Tn5 and plasmid insertions from the start codon of genes are indicated in brackets. Str^r = streptomycin resistance, Neo^f = neomycin resistance, Gen^f = gentamicin resistance, Tet^f = tetracyclin resistance, Chl^f = chloramphenicol resistance, Kan^f = kanamycin resistance, Amp^f = ampicillin resistance, Ile = isoleucine, Val = valine, Leu = leucine.

active NodD proteins in the cells (Wais et al. 2002). Here, we confirmed that *nod* gene expression in the presence of luteolin is higher (approximately two- to threefold) in the 1021 wild-type strain than in the 2011 wild-type strain.

The *ilvC* mutants derived from both strains 1021 and 2011 presented a reduced level of β -galactosidase activity which reached approximately 66 and 50%, respectively, of the level of the wild-type strains after 4 h of incubation with luteolin. However, these differences were no longer observed 24 h after induction, in contrast to previous work (Lopez et al. 2001). In addition, the *ilvD2* mutant derived from strain 2011 also displayed a significant reduction (50%) in *nodC* induction by luteolin compared with the wild-type strain 4 h after addition of luteolin. Again, this reduction disappeared 24 h after induction.

Surprisingly, we observed that the *nodC-lacZ* translational fusion carried by the pRmM57 plasmid was highly susceptible to the presence of the antibiotic neomycin in the culture medium. Indeed, this fusion was no more inducible by luteolin when neomycin was added to the medium in all the neomycin-resistant mutant strains we tested, independently of the mutation itself (Supplementary Fig. S1A and B). Moreover, this defect was observed at the protein level (β -galactosidase dosage of the *nodC-lacZ* translational fusion) but not at the RNA level (quantitative reverse-transcriptase polymerase chain reaction [qRT-PCR] experiments). In our experiments, the neomycin antibiotic was not added to the cultures of the mutants. To check if the β -galactosidase activities measured in these conditions for the mutants were not due to mutant revertants, we estimated the mutation reversion rate by counting prototroph colonies on minimal medium plates at the end of each experiment and found that it was 10^{-8} for the Tn5 mutants and 10^{-6} for the pVO155 plasmid insertion mutants.

In order to directly assess Nod factor production by the *ilvC* and *ilvD2* mutants, we performed biochemical analyses of supernatants of 24-h luteolin-induced cultures from *ilvC*, *ilvD2*, and wild-type strains in the two genetic backgrounds, 1021 and 2011. Consistently with the β -galactosidase dosages performed 24 h after addition of luteolin, both *ilvC* and *ilvD2* mutants were shown to produce Nod factors in similar quantity and of similar quality to the wild-type strains in both 1021 and 2011 backgrounds (Fig. 3).

From these results, we concluded that Nod factor biosynthesis is not the limiting factor that prevents nodulation of 1021 *ilvC* and 1021 *ilvD2* strains, at least in the presence of 0.8 mM isoleucine and valine.

The TetR-type regulator SMC04348 adjacent to *ilvC* did not contribute to the symbiotic defect.

In order to assess the consequences of *ilvC* and *ilvD2* inactivation on global gene expression in the *S. meliloti* strain 1021, we analyzed the transcriptomes of these two mutants compared with the isogenic wild-type strain grown in minimal medium containing ammonium and succinate as nitrogen




Fig. 2. Nodulation kinetics of the 1021 *ilvC* and 2011 *ilvD2* auxotrophs and complemented strains. Results of a typical nodulation experiment. **A**, GMI11496: 1021 wild-type strain; GMI11498: 1021 *ilvC* mutant; GMI11509: 1021 *ilvC* complemented strain. The difference in the number of nodules between the wild type and the 1021 *ilvC* complemented strains is not statistically significant. **B**, GMI11495: 2011 wild-type strain; GMI11499: 2011 *ilvD2* mutant; GMI11511: 2011 *ilvD2* complemented strain. IV: 0.2 mM isoleucine and valine were added to the plant growth medium.

Table 2. Symbiotic phenotype of auxotrophic mutants on *Medicago sativa*

Strains	Parental strain	Genotype	Auxotrophy	Plant growth medium ^a	
				Without amino acids	With amino acids
GMI11531	2011	<i>ilvI</i>	Ile, Val	Nod ^D (20 days) Fix ^{-b}	Nod ⁺ Fix ⁻
GMI11523	2011	<i>ilvC</i>	Ile, Val	Nod ^D (20 days) Fix ^{-b}	Nod ⁺ Fix ⁻
GMI11525	2011	<i>ilvD2</i>	Ile, Val	Nod ^D (20 days) Fix ^{-b}	Nod ⁺ Fix ⁻
GMI11532	2011	<i>leuA1</i>	Leu	Nod ^D (20 days) Fix ^{-b}	Nod ⁺ Fix [±]
GMI11533	2011	<i>leuB</i>	Leu	Nod ^D (20 days) Fix ^{-b}	Nod ⁺ Fix [±]
GMI11502	1021	<i>ilvI</i>	Ile, Val	Nod ^D (7 days) Fix ^{±c}	Nod ⁺ Fix ^{±c}
GMI11498	1021	<i>ilvC</i>	Ile, Val	Nod ^D (20 days) Fix ^{-b}	Nod ^D (20 days) Fix ⁻
GMI11499	1021	<i>ilvD2</i>	Ile, Val	Nod ^D (20 days) Fix ^{-b}	Nod ^D (10 days) Fix ⁻
GMI11503	1021	<i>leuA1</i>	Leu	Nod ^D (7 days) Fix ^{±c}	Nod ⁺ Fix [±]
GMI11505	1021	<i>leuC</i>	Leu	Nod ^D (7 days) Fix ⁻	Nod ⁺ Fix [±]
GMI11506	1021	<i>leuD</i>	Leu	Nod ^D (7 days) Fix ^{±c}	Nod ⁺ Fix [±]
GMI11504	1021	<i>leuB</i>	Leu	Nod ^D (7 days) Fix ⁻	Nod ⁺ Fix [±]

^a Nodulation phenotypes were determined in the absence or presence of 0.2 mM of the missing amino acids in the plant growth medium. Nod^D = nodulation delayed, Nod⁺ = wild type nodulation kinetics, Fix⁺ = nitrogen-fixing nodules, Fix⁻ = non-nitrogen-fixing nodules, Fix[±] = partial impairment of nitrogen fixation. The number of days of delay before nodulation compared with the wild-type strains was averaged from at least three independent experiments.

^b Number of induced nodules was low.

^c Pink nitrogen-fixing nodules were due to mutant revertants.

and carbon sources and supplemented with isoleucine and valine. We found very few genes whose expression varied between mutants and the wild type (not shown). Among them, only one gene, SMc04348, was consistently induced in the *ilvC* and *ilvD2* mutants. SMc04348 encodes a TetR-type regulator and is located immediately upstream from the *ilvC* gene in the genome. We validated the induction of this gene by qRT-PCR in the two mutants (Supplementary Fig. S2A). We wanted to know whether overexpression of SMc04348 could affect the symbiotic properties of the wild-type strain. Consequently, we constructed a strain (GMI11508) overexpressing the TetR regulator SMc04348 on a plasmid and tested its symbiotic performances on *M. sativa*. We found

that overexpression of SMc04348 in strain 1021 influenced neither the nodulation nor the nitrogen-fixation ability of the strain. We also constructed a SMc04348 null mutant by plasmid insertion (GMI11507). The 1021 SMc04348 mutant was auxotrophic for isoleucine and valine, like the *ilvC* mutant, and displayed the same nodulation phenotype as the 1021 *ilvC* strain. We showed that the auxotrophy and the nodulation phenotype of this strain was due to the polarity of the SMc04348 mutation on the *ilvC* gene because in trans expression of *ilvC* on a plasmid (GMI11512) restored both the prototrophy and the nodulation ability of the SMc04348 mutant whereas in trans expression of SMc04348 on a plasmid (GMI11513) did not reverse the phenotype.

Table 3. Expression of *nodC* in auxotrophic mutants

Strains	Parental strain	Genotype	β-Galactosidase activity ^a		
			-Lut	+Lut (4 h)	+Lut (24 h)
GMI11534	2011	Wild type	20 ± 1	472 ± 70	251 ± 40
GMI11537	2011	<i>ilvI</i>	21 ± 2	352 ± 63	nd
GMI11535	2011	<i>ilvC</i>	15 ± 2	204 ± 38*	266 ± 56
GMI11536	2011	<i>ilvD2</i>	15 ± 1	233 ± 24*	256 ± 30
GMI11538	2011	<i>leuA1</i>	20 ± 1	452 ± 10	nd
GMI11539	2011	<i>leuB</i>	15 ± 1	361 ± 54	nd
GMI11514	1021	Wild type	27 ± 1	1,254 ± 93	785 ± 71
GMI11517	1021	<i>ilvI</i>	27 ± 1	1,415 ± 94	nd
GMI11515	1021	<i>ilvC</i>	22 ± 5	834 ± 75*	1,074 ± 232
GMI11516	1021	<i>ilvD2</i>	26 ± 2	1,287 ± 74	nd
GMI11518	1021	<i>leuA1</i>	34 ± 3	1,328 ± 125	nd
GMI11520	1021	<i>leuC</i>	28 ± 1	1,183 ± 67	nd
GMI11521	1021	<i>leuD</i>	24 ± 2	971 ± 83*	nd
GMI11519	1021	<i>leuB</i>	29 ± 4	1,048 ± 197	nd

^a β-Galactosidase activities were measured in Miller units in the absence or presence of 10 μM luteolin over a period of 4 or 24 h. Average results and standard deviation of three independent experiments are indicated. An asterisk (*) indicates that the difference between the mutant and the wild type was statistically significant (*t* test, *P* < 0.05); nd = not determined.




Fig. 3. Nod factor production in the *ilvC* and *ilvD2* mutants and in the respective wild-type strains. High-performance liquid chromatography UV traces recorded at 206 nm of the XAD extract eluted with 50% acetonitrile/water fraction and injected on an Xterra reversed-phase column. **A**, 1021 *ilvC* (blue), 1021 *ilvD2* (pink), and 1021 wild type (black). **B**, 2011 *ilvC* (blue), 2011 *ilvD2* (pink), and 2011 wild type (black). Asterisk (*) indicates low molecular weight polyethylene glycol present in water of cultures that were enriched during extraction.

DISCUSSION

Apart from studies on mutants in the nodulation genes, reports in the literature on rhizobial mutants that are completely deficient in nodulation are rare. However, three independent studies reported several *S. meliloti* mutants which were defective in host legume nodulation and unable to synthesize the BCAA isoleucine, valine, and leucine. This led us to reexamine the role in nodulation of this biosynthesis pathway as a whole in *S. meliloti*.

Based on the *E. coli* biosynthesis pathway model, we searched for homologous *ilv* and *leu* genes on the *S. meliloti* genome and checked their requirement for the synthesis of isoleucine, valine, and leucine by auxotrophy tests of mutants of these genes. These assays enabled the reconstruction of most of the biosynthesis pathway in *S. meliloti*. Only one AHAS isozyme was identified, the AHAS isozyme III (IlvIH) ubiquitously found in bacteria, whereas two BCAA aminotransferases (IlvE1 and IlvE2) catalyzing the last step of the pathway may be functional in *S. meliloti*.

Here, we characterized the symbiotic phenotype of three mutants auxotrophic for isoleucine and valine (*ilvI*, *ilvC*, and *ilvD2*) and four mutants auxotrophic for leucine (*leuA1*, *leuC*, *leuD*, and *leuB*) in two closely related *S. meliloti* strains, 1021 and 2011. We found that all auxotrophic mutants in both strains presented a similar phenotype on plants in the absence of amino acids. Indeed, the mutants were severely deficient in their nodulation and infection abilities on *M. sativa* host plants. Nodules were formed after a delay of several days compared with nodules induced by the wild-type strains, and were few in number, often abnormally developed and not invaded by bacteria. These results indicate that BCAA are essential for both nodulation and infection in *S. meliloti*.

Unexpectedly, we also observed that the nodulation deficiency of most *ilv* and *leu* mutants was essentially restored by addition of BCAA to the plant growth medium. For restoration to be successful, amino acids had to be added to the plant culture medium and not to the inoculum itself. Other than the noticeable exceptions of the *ilvC* and *ilvD2* mutants in the 1021 background, our results demonstrated that the nodulation deficiency of the mutants was entirely linked to their auxotrophy. This suggests that the concentration of BCAA in alfalfa root exudates is limiting and does not satisfy the requirements for the normal metabolic activity of isoleucine, valine, and leucine auxotrophs. However, unlike leucine auxotrophs, supplementation with exogenous amino acids restored neither the nitrogen fixation nor the normal infection process in isoleucine and valine auxotrophs. Therefore, isoleucine and valine amino acids may also be limiting inside the alfalfa root tissues.

Some of our data confirmed results obtained in previous studies while other data disagreed. We confirmed that supplementation with amino acids does not restore the symbiotic phenotype of the *ilvC* mutant in strain 1021 although the *ilvC* nodulation phenotype is background dependent. We also confirmed that the 1021 *leuA1* mutant can nodulate alfalfa in the presence of leucine. One striking discrepancy between our results and previous studies mainly concerns the *ilvD2* phenotype. Although this mutant was found to be symbiotically effective in two independent studies in strains 1021 and Rmd201 (Aguilar and Grasso 1991; Hassani et al. 2002), our *ilvD2* mutants in strains 1021 and 2011 were strongly impaired for nodulation even in the presence of amino acids in strain 1021. Very recently, Pobigaylo and associates (2008) confirmed the nodulation phenotype of the 2011 *ilvD2* mutant using the same strain as the one used in this study.

Another major difference between our study and previous studies was the ability to produce Nod factors by the *ilv* and

leu auxotrophs. In our study, we found that all auxotrophic strains were able to induce a *nodC-lacZ* fusion, contrary to previous works on the *ilvC* and *leuA1* mutants (Aguilar and Grasso 1991; Sanjuan-Pinilla et al. 2002). We only observed a partial reduction of *nodC-lacZ* expression in the *ilvC* and *ilvD2* mutants a few hours after incubation with luteolin. However, this reduction disappeared after a longer incubation with luteolin. In addition, we demonstrated by biochemical analyses that the 1021 and 2011 *ilvC* mutants normally produce Nod factors in 24-h luteolin-induced cultures. Interestingly, we found that the *nodC-lacZ* translational fusion carried by the pRmM57 plasmid was very weakly expressed at the protein level in the presence of neomycin in the culture medium even upon induction by luteolin. Neomycin blocks translation initiation; therefore, we suspect that the translation of the *nodC* mRNA is particularly sensitive to this mechanism even in strains that are resistant to this antibiotic. This sensitivity could explain results reported by Aguilar and Grasso (1991) and Sanjuan-Pinilla and associates (2002) concerning the lack of expression of this fusion in the neomycin-resistant *ilvC* and *leuA1* mutants. Our *nod* gene induction and Nod factor production assays clearly demonstrated that the irreversibility of the nodulation phenotype of the 1021 *ilvC* and *ilvD2* auxotrophs by supplementation with amino acids was not due to a defect in Nod factor synthesis.

In this study, we found that the *ilvC* and *ilvD2* mutations had an impact on nodulation which depended on the genetic background of the strain. Although the *S. meliloti* 1021 and 2011 strains are very close to each other, because they are two independent spontaneous streptomycin-resistant derivatives of the same parental strain, they display many genetic differences and many phenotypic differences. Differences include *nod* gene induction level by NodD1 and luteolin and the calcium-spiking kinetic response in host plants (Wais et al. 2002), the nitrogen fixation efficiency on some ecotypes of *M. truncatula* (Terpolilli et al. 2008), their ability to form biofilms (Fujishige et al. 2006), and the differential expression of hundreds of genes, including phosphate starvation responsive genes (Krol and Becker 2004; *personal observation*). The symbiotic phenotype of *ilvC* and *ilvD2* mutants is possibly another difference between the two strains. Alternatively, we cannot exclude that secondary mutations have been selected during the construction of these two particular mutants in strain 1021, either because strain 1021 is more mutator than strain 2011 or because BCAA biosynthesis genes are essential genes of central metabolism, or both.

Preliminary analyses of the 1021 *ilvC* and 1021 *ilvD2* transcriptomes in free-living cultures did not provide any clues regarding genes deregulated in these mutants that may be involved in their symbiotic defect. We investigated the role of the TetR-type transcription regulator SMc04348, located upstream from *ilvC* on the genome, whose expression was found to be higher in both mutants than in the wild-type strain. Induction of this gene in the mutants was observed in both 1021 and 2011 genetic backgrounds. However, we found that the level of expression of SMc04348 in the mutants depended on the source of nitrogen in the culture medium, expression of this gene being higher in the presence of 15 mM ammonia than in the presence of 6 mM glutamate, suggesting that this regulator may connect BCAA biosynthesis and nitrogen metabolism. SMc04348 was found to be linked to *ilvC* in the genome of several α proteobacteria such as *Rhizobium elii*, *Mesorhizobium loti*, *Agrobacterium tumefaciens* or *Brucella* spp. This conserved synteny suggests that the regulator may, thus, be functionally linked to *ilvC* or to the isoleucine and valine biosynthetic pathway, although we do not have further evidence for this. Preliminary transcriptome analyses of strains overexpressing SMc04348 or inactivated in SMc04348

cultivated in minimal medium did not help identify the SMc04348 target genes (not shown). Thus, the role of this regulator remains unknown. The characterization of the symbiotic phenotypes of the SMc04348 overexpressing strain (GMI11508) and the SMc04348 mutant strains (GMI11507 and GMI11512) indicated that this gene does not play an essential role in the nodulation process in *S. meliloti*.

In rhizobia, isoleucine, valine, or leucine auxotrophic mutations do not always lead to a nodulation defect. For example, leucine auxotrophs of *Bradyrhizobium japonicum* form effective nodules on soybean (Kummer and Kuykendall 1989). In contrast, auxotrophic mutations in leucine biosynthesis genes prevent nodulation of photosynthetic *Bradyrhizobium* spp. whose genomes lack classical *nod* genes. In the same photosynthetic *Bradyrhizobium* spp., other auxotrophic mutations (in particular, for proline amino acid) also lead to a nodulation defect on *Aeschynomene* host plants. In the two latter cases, the phenotype was rescued by addition of the missing amino acids to the plant nutrient medium (Giraud et al. 2007). Taken together, these data and our results indicate that the nodulation phenotype of auxotrophic mutants in rhizobia is likely determined by the availability of amino acids and their derivatives in the root exudates. Hosts with different profiles of exudates may explain the different impacts of rhizobial auxotrophy on the symbiotic interaction. In natural environments, the intrinsic metabolic capacity of rhizobia to synthesize amino acids, which may be limiting in the rhizosphere of their host plants, is essential for both the establishment of an effective symbiosis and their competitiveness.

In conclusion, we believe that this work clarifies the role of the BCAA biosynthesis in the nodulation process of *S. meliloti*. Indeed, we have shown that all mutations that are auxotrophic for BCAA lead to a severe nodulation defect on *M. sativa* host plants, demonstrating that the entire biosynthesis pathway is necessary for nodulation. We also demonstrated that auxotrophy is likely the primary cause of the nodulation defect in most of these mutants. However, the nodulation defect of two noticeable auxotroph mutants, *ilvC* and *ilvD2* in the specific 1021 genetic background, remains to be elucidated.

MATERIALS AND METHODS

Bacterial strains and growth conditions.

Bacterial strains and plasmids used in this study are listed in Table 1. Auxotrophic strains were grown either in rich medium (Luria-Bertani [LB] with 2.5 mM CaCl₂ and 2.5 mM MgSO₄) or in Vincent minimal medium modified as previously described (Becker et al. 2004) supplemented with isoleucine and valine or leucine at a final concentration of 0.8 mM. The concentrations of antibiotics used for *S. meliloti* cultures were 200 µg/ml for streptomycin, 100 µg/ml for neomycin, 10 µg/ml for tetracycline, and 30 µg/ml for gentamycin.

Constructions of strains and plasmids.

All *S. meliloti* 2011 mTn5 mutants were obtained from the mTn5 random mutant library (Pobigaylo et al. 2006). The *S. meliloti* 1021 *ilvH*, *leuB*, and *leuC* mutants were obtained from a Tn5 random mutant library (Milcamps et al. 1998). The 2011 *ilvC* (GMI11544), 1021 *ilvC* (GMI11498), and 1021 *ilvD2* (GMI11499) mutants were constructed by site-specific mutagenesis using the suicide plasmids pQB2 and pMA6B carrying an *ilvC*::Tn5 and *ilvD2*::Tn5 cassette, respectively (Aguilar and Grasso 1991). Both plasmids were transferred to *S. meliloti* by biparental conjugation using *E. coli* S17-1 as donor strain. The double event of recombination was selected by searching for tetracycline-sensitive streptomycin- and neomycin-resistant colonies. The mutants were then checked by both Southern blot

and PCR analyses. The prototrophic 1021 Tn5 (GMI11541) mutant was obtained by a nonspecific Tn5 insertion from the pMA6B plasmid. The 1021 *ilvB1*, *ilvI*, *leuA1*, and *leuD* mutants were constructed by site-specific insertion mutagenesis using the nonreplicative plasmid pVO155. The pVO155 plasmid was digested by *Bam*HI and *Xba*I. A DNA fragment of each gene to be inactivated was amplified by PCR with primers containing a *Bam*HI and *Xba*I restriction site at their 5' end. Primers used for mutagenesis are listed in Supplementary Table S2. The PCR products were digested with *Bam*HI and *Xba*I, ligated to the pVO155 plasmid, and transformed into DH5α *E. coli*-competent cells. The plasmid constructions were checked by sequencing and then conjugating into *S. meliloti* strain 1021 using the pRK600 as helper plasmid. Transconjugants were selected three times on LB plates containing 2.5 mM CaCl₂, 2.5 mM MgSO₄, streptomycin at 200 µg/ml, and neomycin at 100 µg/ml. Correct plasmid integration was checked by Southern blot analysis.

For the construction of the complementation plasmids pGMI50120 (*ilvC*), pGMI50121 (*ilvD2*), and pGMI50122 (SMc04348), the wild-type *ilvC*, *ilvD2*, and SMc04348 genes were amplified by PCR using *S. meliloti* 1021 genomic DNA as template and ODC161-ODC162, ODC163-ODC164, and OCD177-ODC178 primer pairs, respectively, which carry a *Xba*I and *Bam*HI restriction site at their 5' end (Table S2). The three fragments were digested with *Bam*HI and *Xba*I and ligated into a *Bam*HI-*Xba*I-digested pBBR1MCS-5 plasmid. The resulting plasmids were transformed into *E. coli* DH5α-competent cells. Clones were screened by PCR and checked by DNA sequencing prior to introduction into *S. meliloti* *ilvC*, *ilvD2*, and SMc04348 auxotrophic mutants by triparental mating using the pRK600 as helper plasmid.

The pXLGD4 and pRmM57 replicative plasmids were introduced into all auxotrophs and wild-type strains by triparental mating using the pRK600 as helper plasmid.

Plant assays.

Seed of *Medicago sativa* cv. Europe were surface sterilized, germinated, and allowed to grown in 12-cm² plates containing slanting nitrogen-free Fahraeus agar medium for 3 days at 22°C with day and night cycles of 16 and 8 h, respectively. The plants were then inoculated with 2 × 10³ bacteria/plant. For supplementation experiments, either isoleucine and valine or leucine were added to the plant medium at 0.2 mM because higher amino acid concentrations inhibited the growth of plant roots. Nodulation kinetics were monitored for approximately 40 days.

Cytological studies.

For cytological observations, plants were inoculated with strains carrying the pXLGD4 plasmid that expresses the *hemA-lacZ* fusion at a constitutive level. For observations at early stages, entire roots were collected 7 days after inoculation, fixed with 2% glutaraldehyde solution for 1.5 h under vacuum, rinsed three times in Z' buffer (0.1 M potassium phosphate buffer, pH 7.4, 1 mM MgSO₄, and 10 mM KCl), and stained overnight at 37°C in Z' buffer containing 0.08% X-gal (5-bromo-4-chloro-3-indolyl-β-D-galactoside), 5 mM K₃Fe(CN)₆, and 5 mM K₄Fe(CN)₆. For observations at late stages, 35-day-old nodules were harvested, fixed with 2% glutaraldehyde in Z' buffer, and then sliced into 80-µm-thick longitudinal sections using a vibrating-blade microtom (Leica VT1000S, Heidelberg, Germany) before staining. Entire roots or cut nodules were observed with an optical microscope.

nodC gene induction assays.

S. meliloti strains carrying the pRmM57 plasmid were grown at 28°C in Vincent minimal medium containing 6 mM glutamate as sole nitrogen source and 0.8 mM isoleucine and

valine or leucine. Cultures were diluted in fresh medium at an optical density at 600 nm of 0.08 and left to grow for 3 h. The cultures were then incubated with or without 10 μ M luteolin (Sigma) for 4 or 24 more hours. β -Galactosidase activities were measured using 0.5 ml of culture as previously described (Miller 1972).

Nod factor production.

S. meliloti strains carrying the pRmM57 plasmid were grown at 28°C in Vincent minimal medium containing 6 mM glutamate as sole nitrogen source and 0.8 mM isoleucine and valine or leucine. Cultures were diluted in fresh medium at an optical density at 600 nm of 0.1 in the presence of 10 μ M luteolin (Sigma) and grown for 24 h. The cultures were then centrifuged and supernatants were stored at 4°C until analyses.

Nod factors were extracted from filtered culture supernatants by XAD4 bed adsorption (50 g of phase per liter). After 2 h of stirring, the bed was retained by filtration and washed with pure water. Nod factors were eluted using a 50% water/acetonitrile solution (200 ml). Purification was performed on a high-performance liquid chromatography (HPLC) system Dionex P680-170U (Dionex Corporation, Sunnyvale, CA, U.S.A.) with a narrowbore analytical C18 reverse-phase column (2.1 by 150 mm, 3.5 μ m; Xterra, Waters, Milford, MA, U.S.A.) for 8 min in isocratic solvent A (water/acetonitrile, 80:20 [vol/vol]), followed by linear gradient for 25 min from solvent A to solvent B (100% acetonitrile) and another isocratic step at 100% acetonitrile for 5 min. The return to the initial condition occurred within 10 min and an equilibration time of 10 min was necessary. The flow rate was of 0.2 ml min⁻¹ and the UV absorption was monitored at 206 nm.

The XAD extracts were dried under a nitrogen flux and redissolved in 500 μ l of 80:20 water/acetonitrile solution and 10 μ l was injected on the HPLC column.

Each HPLC fraction produced was analyzed using an ESI-QqToF Ultima apparatus (Waters) using direct infusion. Spectra were recorded in both the positive and negative mode. Peaks detected in the expected molecular weight range (m/z 1,000 to 1,500 for the simple charged species or 500 to 700 for the double-charged ones) were subjected to dual mass spectrometry (MS/MS) experiment to confirm their lipo-chito oligosaccharide nature and structure.

Energies were the following: probe, 3 kV; cone, 100 V; Rf, 70 V; and collision cell, 15 V for MS and 30 V for MS/MS. Collision gas was argon and direct inlet was solvent AcCN/H₂O 1:1 and 1% acetic acid; rate, 10 μ l min⁻¹. Concentrations were approximately 10⁻⁴ M, source temperature was 80°C, and drying gas temperature was 120°C.

RNA preparation.

Filtered bacteria were resuspended in 3 ml of lysis solution (1.4% sodium dodecyl sulfate, 4 mM EDTA, and proteinase K at 0.4 mg/ml) and incubated at 65°C for 20 min. Proteins were precipitated by adding 1.5 ml of NaCl 5M at 4°C for 10 min. Nucleic acids were precipitated from the supernatant by addition of one volume of isopropanol and the pellet was resuspended in RNase-free water. DNA was removed from RNA preparations using RNeasy columns (clean-up procedure with DNase I treatment) (Qiagen, Germany).

Transcriptome analyses.

RNA (20 μ g) was reverse transcribed using Superscript II reverse transcriptase (Invitrogen, Carlsbad, CA, U.S.A.) and random hexamers as primers. Cy3- and Cy5-labeled cDNAs were prepared according to DeRisi and associates (1997). The microarrays used in this study are described in Krol and Becker (2004) and were purchased from A. Becker (University

of Bielefeld, Germany). Hybridizations were performed in DIG Easy Hyb buffer (Roche, Mannheim, Germany) in Corning chambers at 42°C for 16 h. Data were acquired on a Genepix 4000 scanner (Axon Instruments, Foster City, CA, U.S.A.) and quantification of median signal was performed using Genepix pro 3.0 (Axon Instruments). Data analyses were carried out using Genesight 3.5 (Biodiscovery, Inc., El Segundo, CA, U.S.A.). At least two independent experiments were performed for each transcriptome comparison.

qRT-PCR.

Reverse transcriptions were performed from 1 μ g of RNA using Superscript II reverse transcriptase (Invitrogen) and random hexamers as primers. cDNAs were used for running real-time PCR on a LightCycler system (Roche) using the FastStart DNA Master^{PLUS} SYBRGreen I kit (Roche) according to the manufacturer's instructions. The *rplM* gene was used as reference gene for data normalization. Primers used are listed in Supplementary Table S3.

ACKNOWLEDGMENTS

We thank A. Becker (University of Bielefeld) and IIT GmbH (Bielefeld, Germany) for providing all the *S. meliloti* 2011 mTn5 mutants; and F. de Bruijn and L. Sauviac (Laboratoire Interactions Plantes Micro-Organismes Toulouse) for providing the *S. meliloti* 1021 *ilvH*, *leuB*, and *leuC* Tn5 mutants. The work was supported by the France-Argentina ECOS-Sud program (A04B01).

LITERATURE CITED

- Aguilar, O. M., and Grasso, D. H. 1991. The product of the *Rhizobium meliloti* *ilvC* gene is required for isoleucine and valine synthesis and nodulation of alfalfa. *J. Bacteriol.* 173:7756-7764.
- Ardoirel, M., Lortet, G., Maillet, F., Roche, P., Truchet, G., Promé, J. C., and Rosenberg, C. 1995. In *Rhizobium meliloti*, the operon associated with the nod box n5 comprises *nodL*, *noeA* and *noeB*, three host-range genes specifically required for the nodulation of particular *Medicago* species. *Mol. Microbiol.* 17:687-699.
- Becker, A., Berges, H., Krol, E., Bruand, C., Ruberg, S., Capela, D., Lauber, E., Meilhoc, E., Ampe, F., de Bruijn, F. J., Fourment, J., Francez-Charlot, A., Kahn, D., Kuster, H., Liebe, C., Puhler, A., Weidner, S., and Batut, J. 2004. Global changes in gene expression in *Sinorhizobium meliloti* 1021 under microoxic and symbiotic conditions. *Mol. Plant-Microbe Interact.* 17:292-303.
- Capela, D., Carrere, S., and Batut, J. 2005. Transcriptome-based identification of the NodD1 regulon of *Sinorhizobium meliloti*. *Appl. Environ. Microbiol.* 71:4910-4913.
- Debellé, F., Rosenberg, C., Vasse, J., Maillet, F., Martinez, E., Dénarié, J., and Truchet, G. 1986. Assignment of symbiotic developmental phenotypes to common and specific nodulation (nod) genetic loci of *Rhizobium meliloti*. *J. Bacteriol.* 168:1075-1086.
- Demont, N., Debellé, F., Aurelle, H., Dénarié, J., and Promé, J. C. 1993. Role of the *Rhizobium meliloti* *nodF* and *nodE* genes in the biosynthesis of lipo-oligosaccharidic nodulation factors. *J. Biol. Chem.* 268:20134-20142.
- DeRisi, J. L., Iyer, V. R., and Brown, P. O. 1997. Exploring the metabolic and genetic control of gene expression on a genomic scale. *Science* 278:680-686.
- Dusha, I., and Kondorosi, A. 1993. Genes at different regulatory levels are required for the ammonia control of nodulation in *Rhizobium meliloti*. *Mol. Gen. Genet.* 240:435-444.
- Finan, T. M., Kunkel, B., De Vos, G. F., and Signer, E. R. 1986. Second symbiotic megaplasmid in *Rhizobium meliloti* carrying exopolysaccharide and thiamine synthesis genes. *J. Bacteriol.* 167:66-72.
- Fujishige, N. A., Kapadia, N. N., De Hoff, P. L., and Hirsch, A. M. 2006. Investigations of *Rhizobium* biofilm formation. *FEMS (Fed. Eur. Microbiol. Soc.) Microbiol. Ecol.* 56:195-206.
- Giraud, E., Moulin, L., Vallenet, D., Barbe, V., Cytryn, E., Avarre, J. C., Jaubert, M., Simon, D., Cartieux, F., Prin, Y., Bena, G., Hannibal, L., Fardoux, J., Kojadinovic, M., Vuillet, L., Lajus, A., Cruveiller, S., Rouy, Z., Mangenot, S., Segurens, B., Dossat, C., Franck, W. L., Chang, W. S., Saunders, E., Bruce, D., Richardson, P., Normand, P., Dreyfus, B., Pignol, D., Stacey, G., Emerich, D., Vermeglio, A., Medigue, C., and

- Sadowsky, M. 2007. Legumes symbioses: Absence of Nod genes in photosynthetic bradyrhizobia. *Science* 316:1307-1312.
- Hassani, R., Prasad, C. K., Vineetha, K. E., Vij, N., Singh, P., Sud, R., Yadav, S., and Randhawa, G. S. 2002. Symbiotic characterization of isoleucine+valine and leucine auxotrophs of *Sinorhizobium meliloti*. *Indian J. Exp. Biol.* 40:1110-1120.
- Kovach, M. E., Elzer, P. H., Hill, D. S., Robertson, G. T., Farris, M. A., Roop, R. M., 2nd, and Peterson, K. M. 1995. Four new derivatives of the broad-host-range cloning vector pBRR1MCS, carrying different antibiotic-resistance cassettes. *Gene* 166:175-176.
- Krol, E., and Becker, A. 2004. Global transcriptional analysis of the phosphate starvation response in *Sinorhizobium meliloti* strains 1021 and 2011. *Mol. Genet. Genomics* 272:1-17.
- Kummer, R. M., and Kuykendall, L. D. 1989. Symbiotic properties of amino acid auxotrophs of *Bradyrhizobium japonicum*. *Soil Biol. Biochem.* 21:779-782.
- Leong, S. A., Williams, P. H., and Ditta, G. S. 1985. Analysis of the 5' regulatory region of the gene for delta-aminolevulinic acid synthetase of *Rhizobium meliloti*. *Nucleic Acids Res.* 13:5965-5976.
- Lopez, J. C., Grasso, D. H., Frugier, F., Crespi, M. D., and Aguilar, O. M. 2001. Early symbiotic responses induced by *Sinorhizobium meliloti* *ilvC* mutants in alfalfa. *Mol. Plant-Microbe Interact.* 14:55-62.
- Maillet, F., Debelle, F., and Dénarié, J. 1990. Role of the *nodD* and *syrM* genes in the activation of the regulatory gene *nodD3*, and of the common and host-specific nod genes of *Rhizobium meliloti*. *Mol. Microbiol.* 4:1975-1984.
- Meade, H. M., Long, S. R., Ruvkun, G. B., Brown, S. E., and Ausubel, F. M. 1982. Physical and genetic characterization of symbiotic and auxotrophic mutants of *Rhizobium meliloti* induced by transposon Tn5 mutagenesis. *J. Bacteriol.* 149:114-122.
- Mendel, S., Elkayam, T., Sella, C., Vinogradov, V., Vyazmensky, M., Chipman, D. M., and Barak, Z. 2001. Acetohydroxyacid synthase: A proposed structure for regulatory subunits supported by evidence from mutagenesis. *J. Mol. Biol.* 307:465-477.
- Milcamps, A., Ragatz, D. M., Lim, P., Berger, K. A., and de Bruijn, F. J. 1998. Isolation of carbon- and nitrogen-deprivation-induced loci of *Sinorhizobium meliloti* 1021 by Tn5-*luxAB* mutagenesis. *Microbiology* 144:3205-3218.
- Miller, J. H. 1972. Pages 352-355 in: *Experiments in Molecular Genetics*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, U.S.A.
- Mulligan, J. T., and Long, S. R. 1989. A family of activator genes regulates expression of *Rhizobium meliloti* nodulation genes. *Genetics* 122:7-18.
- Oke, V., and Long, S. R. 1999. Bacterial genes induced within the nodule during the *Rhizobium-legume* symbiosis. *Mol. Microbiol.* 32:837-849.
- Pobigaylo, N., Wetter, D., Szymczak, S., Schiller, U., Kurtz, S., Meyer, F., Nattkemper, T. W., and Becker, A. 2006. Construction of a large signature-tagged mini-Tn5 transposon library and its application to mutagenesis of *Sinorhizobium meliloti*. *Appl. Environ. Microbiol.* 72:4329-4337.
- Pobigaylo, N., Szymczak, S., Nattkemper, T. W., and Becker, A. 2008. Identification of Genes relevant to symbiosis and competitiveness in *Sinorhizobium meliloti* using signature-tagged mutants. *Mol. Plant-Microbe Interact.* 21:219-231.
- Randhawa, G. S., and Hassani, R. 2002. Role of rhizobial biosynthetic pathways of amino acids, nucleotide bases and vitamins in symbiosis. *Indian J. Exp. Biol.* 40:755-764.
- Roche, P., Debelle, F., Maillet, F., Lerouge, P., Faucher, C., Truchet, G., Dénarié, J., and Promé, J. C. 1991. Molecular basis of symbiotic host specificity in *Rhizobium meliloti*: *nodH* and *nodPQ* genes encode the sulfation of lipo-oligosaccharide signals. *Cell* 67:1131-1143.
- Sanjuan-Pinilla, J. M., Munoz, S., Nogales, J., Olivares, J., and Sanjuan, J. 2002. Involvement of the *Sinorhizobium meliloti* *leuA* gene in activation of nodulation genes by NodD1 and luteolin. *Arch. Microbiol.* 178:36-44.
- Terpolilli, J. J., O'Hara, G. W., Tiwari, R. P., Dilworth, M. J., and Howieson, J. G. 2008. The model legume *Medicago truncatula* A17 is poorly matched for N(2) fixation with the sequenced microsymbiont *Sinorhizobium meliloti* 1021. *New Phytol.* 14:14.
- Truchet, G., Michel, M., and Dénarié, J. 1980. Sequential analysis of the organogenesis of lucerne (*Medicago sativa*) root nodules using symbiotically-defective mutants of *Rhizobium meliloti*. *Differentiation* 16:163-172.
- Wais, R. J., Wells, D. H., and Long, S. R. 2002. Analysis of differences between *Sinorhizobium meliloti* 1021 and 2011 strains using the host calcium spiking response. *Mol. Plant-Microbe Interact.* 15:1245-1252.