

Lethal and sublethal effects after treatment with conventional and biorational insecticides on eggs of *Chrysoperla externa* (Neuroptera: Chrysopidae)

Efectos letales y subletales luego de exposición a insecticidas convencionales y biorracionales sobre huevos de *Chrysoperla externa* (Neuroptera: Chrysopidae)

MARINA HARAMBOURE*

Abstract: The neuropteran *Chrysoperla externa* is a generalist predator of many important agricultural pests in the Neotropical region. It occurs on several agricultural crops, and therefore it can be affected by pesticides. Here, the lethal and sublethal effects of conventional and biorational insecticides were assessed against *C. externa* eggs. The exposure method consisted on immersion of eggs in solutions with the maximum field recommended concentration of 25 mg a.i./L cypermethrin, 200 mg a.i./L acetamiprid, 40 mg a.i./L azadirachtin and 75 mg a.i./L pyriproxyfen. The viability of the eggs was not affected, but 100 % of the neonates from eggs treated with cypermethrin died after 48 h. Larvae from eggs treated with acetamiprid, azadirachtin and pyriproxyfen survived to the adult stage, with survival rates of 75 %, 70.6 % and 84.6 %, respectively. Regarding the reproductive parameters, the preoviposition period was not affected but the fecundity and fertility were lower with acetamiprid and pyriproxyfen, showing the sublethal effects of the insecticides acting in the adult stage.

Key words: IPM, neuropterans, Neotropical region, pesticides.

Resumen: El neuróptero *Chrysoperla externa* es un depredador neotropical generalista de plagas importantes en la agricultura. Se encuentra en campos tratados con insecticidas pudiendo ser afectada por los mismos. Los efectos letales y subletales de insecticidas convencionales y biorracionales fueron evaluados sobre huevos expuestos. Los huevos se trataron por inmersión en soluciones de 25 mg i.a./L cipermetrina, 200 mg i.a./L acetamiprid, 40 mg i.a./L azadiractina y 75 mg i.a./L piriproxifén a la máxima concentración recomendada de campo. La viabilidad de los huevos no afectada, pero el 100 % de las neonatas provenientes de tratamientos con cipermetrina murieron luego de 48 h de emergidas. Las larvas de huevos tratados con acetamiprid, azadiractina y piriproxifén sobrevivieron hasta el estado adulto en un 75 %, 70,6 % y 84,6 %, respectivamente. Con respecto a los parámetros reproductivos, el período de pre-oviposición no fue afectado, pero la fecundidad y fertilidad disminuyó en los tratamientos con acetamiprid y piriproxifén, demostrando que en estado adulto los individuos son afectados por tratamientos en el estado de huevo.

Palabras clave: MIP, neurópteros, región Neotropical, pesticidas.

Introduction

The Neotropical predator *Chrysoperla externa* Hagen (Neuroptera: Chrysopidae) is associated with several agricultural crops. This species exhibits a high degree of adaptability to different environmental conditions as well as a robust reproductive potential, an efficient predatory capability, and a ubiquitous distribution from the Antilles in the Caribbean down to the north of the Patagonia in Argentina (Adams and Penny 1987). Larvae of *C. externa* are effective natural enemies of aphids (Hemiptera: Aphididae), white flies (Hemiptera: Aleyrodidae), thrips (Thysanoptera: Thripidae), various families of lepidopterans (Lepidoptera), and acari (Arachnida: Acari) (Carvalho *et al.* 2002; Oliveira *et al.* 2009). Therefore, *C. externa* has been identified as a good candidate for being an effective biological control agent. During recent years the massive rearing of this species has been promoted along with its release in the field in South-American countries (Carvalho *et al.* 2002; Pappas *et al.* 2011).

Biological control and selective pesticides have proved to be compatible with Integrated Pest Management (IPM) (Kogan 1998; Galvan *et al.* 2005). Although chemical control should be the final option in an IPM program, several

agroecosystems depend on pesticide applications, and then their lethal and sublethal effects against beneficial organisms must be known (Shinde *et al.* 2009). Moreover, conventional insecticides in Argentina continue to be the first choice due to the low cost and, sometimes, the lack of information about alternative measures (Villaamil Lepori *et al.* 2013; Pórfido *et al.* 2014). In order to maximize the compatibility of natural enemies with chemicals, studies about effects of pesticides on non-target insects are necessary (Medina *et al.* 2003; Desneux *et al.* 2007).

Studies of pesticides on different stages of *C. externa* have been evaluated in the last decades, most of all on larvae, the predator stage. Eggs are not mobile and could be laid on unprotected spaces and therefore be exposed to pesticides (Soares *et al.* 2002). The importance of the toxicity assessment on beneficial insects' eggs resides in the fact that the establishment of the population depends on them and could be affected by pesticides applications (Castilhos *et al.* 2014). Among the groups of chemicals that have been analyzed on *C. externa* eggs are pyrethroids, neonicotinoids, IGR's, botanicals, organophosphates, organochlorines (Iannacone and Lamas 2003; Bueno and Freitas 2004; Godoy *et al.* 2004), and chrysopids have shown a differential susceptibility against them.

* Ph.D. Centro de Estudios Parasitológicos y de Vectores (CEPAVE) CONICET UNLP. Bvd. 120 e/60 y 64 S/N – La Plata, Buenos Aires, Argentina. Tel: +54 (0221) 4232140 Int 58. Fax: +54 (0221) 4232140 Int 12. marinaharamboure@gmail.com; marina.haramboure@conicet.gob.ar.

The pyrethroid cypermethrin is a broad-spectrum insecticide, with a long-term residual effect. It has been used indiscriminately since its synthesis in the 60's, with negative effects as the development of insecticide resistance (Zhong *et al.* 2013). In the 90's, neonicotinoids like acetamiprid started to be commercialized, classified as biorational insecticides by the United States-Environmental Protection Agency (US-EPA) due to their low toxicity in non-target organisms. However, subsequent studies have demonstrated a high toxicity to pollinators (Laycock *et al.* 2012; Whitehorn *et al.* 2012; Rundlöf *et al.* 2015), and nowadays the classification is being reconsidered (United States-Environmental Protection Agency 2016). The botanical insecticide azadirachtin has been commonly reported as a biorational insecticide with low toxicity to vertebrates, high selectivity and short-term residuality; however, its mode of action and biosynthesis are not well known (Mordue 2004). Pyriproxyfen is an insecticide commonly used in Argentina, classified as an insect growth regulator (IGR) that mimics the action of the juvenile hormone (Pórfido *et al.* 2014) and whose insecticidal activity occurs upon either contact or ingestion.

The aim of this study was to assess both lethal and sublethal effects of the conventional and biorational insecticides cypermethrin, acetamiprid, azadirachtin, and pyriproxyfen after application on eggs of *C. externa* under laboratory conditions.

Materials and methods

Insects. *Chrysoperla externa* colony was maintained in a rearing chamber under controlled conditions (25 °C ± 2; 70 % ± 5 RH; 16:8 L:D). Adults were fed on artificial diet according to Vogt *et al.* (2000) and reared in plastic containers ventilated in the upper side with a fine mesh. Containers were covered inside with black cardboard as oviposition substrate. Eggs were removed from the container and larvae were fed on nymphs and adults of *Rhopalosiphum padi* L. (Hemiptera: Aphididae) and a dietary supplement (Haramboure *et al.* 2016).

Insecticides. The following insecticides were used: Glextrin 25® (cypermethrin 25 % w/v, EC, Gleba S.A.), Mospilan® (acetamiprid 20 % w/w, WP, Summit-Agro S.A.), Neem-Azal® (azadirachtin 1.2 % w/v, EC, Agristar S.A.) and Epingle® (pyriproxyfen 10 % w/v, EC, Summit-Agro S.A.). Maximum field recommended concentrations (MFRC) were evaluated for each treatment: 25 mg a.i./L cypermethrin, 200 mg a.i./L acetamiprid, 40 mg a.i./L azadirachtin and 75 mg a.i./L pyriproxyfen. Dilutions were made using distilled water and controls were treated with distilled water alone. The term **conventional insecticide** will be used for cypermethrin, and **biorational insecticide** for acetamiprid, azadirachtin and pyriproxyfen, throughout the text.

Bioassays. Eggs of the same cohort of less than 24 h laid were treated. Little pieces of cardboard with eggs were cut and immerse for 5 s in a 10 mL glass beaker with the dilutions. Three replicates of 20 eggs were treated for each chemical and followed during the life cycle until the adult stage. After that, eggs were maintained in Petri dishes under the controlled conditions before mentioned; recently hatched larvae were kept individually in 2 cm diameter x 1 cm height plastic vials until pupation. Individuals that survived to adults

were identified by gender through the external genitalia, then paired in five couples and kept in 4 cm diameter x 6 cm height cylindrical vials.

Eggs viability and survival of their offspring were evaluated, as well as the developmental time, preoviposition period (time from adult emergence to first oviposition), and five-days-accumulated fecundity (total number of eggs laid during five days) and fertility (total number of hatched larvae from those eggs).

Statistics. Kaplan-Meier curves were calculated to compare probability of death between the treatments, from egg to adult stage. Survival curves were compared through long-rank tests with χ^2 . Those individuals who reached the adult stage were censored data.

Egg viability, developmental time of eggs, larvae, and pupa, fecundity and fertility were analyzed by ANOVA. When normality and homoscedasticity assumptions were not accomplished they were analyzed by the non-parametric Kruskal-Wallis test. Means or medians, respectively, were compared using LSD or Box and Whisker Plot tests. $P \leq 0.05$ was considered significant. SPSS Software was used.

Results and discussion

Viability and survival curves. The viability of eggs treated with the four insecticides did not differ from the control, which was of 100 % ($F = 2.83$; $df = 4, 10$; $P = 0.082$). The chorion of chrysopids eggs has a proteic, crystalline component arranged in layers (Furneaux and Mackay 1972); this natural barrier which encloses the embryo could lead to mechanical resistance, being this stage one of the most tolerant to insecticides (Silva *et al.* 2012; Rugno *et al.* 2015). Similar results were obtained after exposition of eggs from the same species to different insecticides (Rimoldi *et al.* 2008; Silva *et al.* 2012). On the other hand, Rugno *et al.* (2015) recorded a high mortality of the chrysopid *Ceraeochrysa cubana* Hagen (Neuroptera: Chrysopidae) treated with the neonicotinoid imidacloprid. Even though they belong to the same family, chorion of *C. cubana* eggs could be formed by a different structure from those of *C. externa*, conferring a higher susceptibility. As Castilhos *et al.* (2014) have hypothesized, oil-based formulations could form a layer around the egg, blocking the micropile not allowing air interchange between the embryo and the environment causing asphyxia and dead. In the present study, oil solvent based insecticides were cypermethrin, azadirachtin and pyriproxyfen, and the wettable powder one was acetamiprid. Even so, the eclosion was of 100 % in all treatments and the developmental period between eggs to larvae was the same, showing that asphyxia did not happen and neither did the penetration/absorption of the insecticides.

Larvae from eggs treated with cypermethrin died within the first 48 h, whilst those treated with acetamiprid, azadirachtin and pyriproxyfen survived to the adult stage in percentages of 75 %, 70.6 % and 84.6 %, respectively (Fig. 1 and Table 1). During the eclosion neonates had to contact the insecticides in the chorion surface, exposing themselves to their toxicity. The deleterious effect of cypermethrin agrees with that of lambda-cyhalothrin on *C. cubana* (Rugno *et al.* 2015). On the contrary, results obtained by Silva *et al.* (2012) have suggested that the viability of eggs of *C. externa* would not vary after exposition to betacyfluthrin. Although both

insecticides belong to the same chemical group, pyrethroids are formed by mixtures of *cis* and *trans* isomers, with more or less insecticide effect, respectively. Depending on the percentage of the isomers in the given pyrethroid, the insecticidal effect is different (Aldridge 2013; Casida 2013). In this case, cypermethrin has 48-50 % of *cis* isomers and betacyfluthrin 30-40 % (Vodeb and Petanovska-Ilievska 2006); consequently, cypermethrin must be more toxic. It is also important to take into account the exposition method; in Silva *et al.* (2012) the eggs were sprayed by a Potter's tower Residual tests on *Chrysoperla carnea* Stephens (Neuroptera: Chrysopidae) eggs with different groups of pesticides have shown a differential viability according with the treatment, being acetamiprid and imidacloprid highly toxic compared to IGR's, which did not differ to the control (Nasreen *et al.* 2005). Medina *et al.* (2003) have hypothesized that *C. carnea* performs a rapid excretion of pyriproxifen once inside the body, so it could explain the low toxicity. Neonates mortality within the first 24 h of treated was related to the insecticide residual in the chorion when the egg hatches, in the same line as the present study. Also, Cloyd and Bethke (2011) concluded that acetamiprid residues were toxic to newly hatched nymphs of the plant bug *Deraeocoris brevis* Uhler (Hemiptera: Miridae), but did not affect the egg hatch.

Developmental time. Developmental time from eggs to neonates did not differ between all treatments ($F = 0.85$; $df = 4, 84$; $P = 0.49$) and was of 4.78 (± 0.1) days on average. However, developmental time of larvae was shortened to approximately 3 days with acetamiprid, in first and third larval stages (Fig. 2) (L1-L2: $H = 26.56$; $df = 3$; $P \leq 0.001$. L2-L3: $F = 0.96$; $df = 3, 58$; $P = 0.418$. L3-P: $F = 3.14$; $df = 3, 58$; $P = 0.032$). Azadirachtin and pyriproxifen shortened the first stage as well, but did not affect the total larval stage. Pupal stage lasted for approximately 10 days in all treatments ($H = 1.57$; $df = 3$; $P = 0.66$). Studies with different insecticides on *C. externa* eggs and the Palearctic *C. carnea* did not reveal differences in the developmental time of eggs and pupae as our results, but neither in the larval stage (Silva *et al.* 2012; Rugno *et al.* 2015).

Reproductive parameters. Preoviposition period lasted 7.55 ± 0.27 days ($F = 1.69$; $df = 3, 16$; $P = 0.21$). No differences

have been found between controls and all treatments, as reported by Silva *et al.* (2012) in *C. externa* eggs treated with several other pesticides, and by Gontijo *et al.* (2014) in larvae of *C. carnea* treated with chlorantranilprole and thiamethoxam. Apparently, the development of the ovaries is not affected by these insecticides. Cumulative fecundity and fertility during the first five ovipositions was higher in the controls and azadirachtin treatment, whereas exposition to pyriproxifen and acetamiprid reduced significantly these parameters (Fecundity: $F = 36.35$; $df = 3, 16$; $P \leq 0.001$. Fertility: $F = 25.29$; $df = 3, 16$; $P \leq 0.001$) (Fig. 3). These two insecticides reduced the reproductive capacity of the females, and this was in accordance with treatments on eggs from the same species with the IGR methoxyfenozide, although the lower fecundity was observed within the 24 h of the oviposition period, besides fertility was not affected (Rimoldi *et al.* 2008). Pyriproxifen is a juvenile hormone analogue that inhibits the vitellogenesis and cause abnormal development of the ovary and subsequent sterility of the insect (Ohashi *et al.* 2012). The lower values of both fecundity and fertility provoked by a neonicotinoid were also observed in other beneficial organisms as pollinators (e.g. *Bombus terrestris* L., *Osmia bicornis* L., *Apis mellifera* L.) (Laycock *et al.* 2012; Whitehorn *et al.* 2012; Rundlöf *et al.* 2015) and coccinellids (e.g. *Serangium japonicum* Chapin, *Harmonia axyridis* Pallas, *Coleomegilla maculate* DeGeer, *Hippodamia convergens* Guérin-Méneville) (He *et al.* 2012; Awasthi *et al.* 2013; Moscardini *et al.* 2015). The physiology of an insect is affected by its nervous system; therefore, a neurotoxin should interfere in these processes. On the other hand, the innocuous effect of azadirachtin on reproductive parameters of *C. externa* was also recorded with the bioinsecticide spinosad (Rimoldi *et al.* 2008). This could be due to an enhanced metabolism of azadirachtin that was applied in eggs, through the life time of the insect, leading to adults with no effect of this insecticide. Moreover, Medina *et al.* (2004) tested azadirachtin on adults of *C. carnea* and found that they were not affected irrespective of the mode of exposure; authors stated that presence of this insecticide must be constant to produce any effect. This could explain that adults of *C. externa* from treated eggs did not show deleterious effects in reproductive parameters due to the absence of the insecticide in the insect body.

The importance of evaluating the effects of pesticides on natural enemies relies on the occurrence of these beneficial arthropods in the agroecosystems, and the potential exposure to chemical applications. Predators could be exposed to a given insecticide indirectly by feeding on contaminated prey, or directly by insecticide spray and by residual contact with treated plants (Medina *et al.* 2003).




Figure 1. Insecticide effect on the survival of *Chrysoperla externa* offspring. Kaplan-Meier, χ^2 , log rank test, $P \leq 0.05$.

Table 1. Effect of cypermethrin, acetamiprid, azadirachtin and pyriproxifen on the survival of *Chrysoperla externa* offspring. Kaplan-Meier, χ^2 log rank test, $P \leq 0.05$.

	χ^2	df	P
Global comparison	85.08	4	< 0.001
Control - Cypermethrin	39.22	1	< 0.001
Control - Acetamiprid	5.75	1	0.016
Control - Azadirachtin	5.86	1	0.015
Control - Pyriproxifen	5.92	1	0.015




Figure 2. Development time of *Chrysoperla externa* with different treatments. C: control; Cy: cypermethrin; Ac: acetamiprid; A: azadirachtin; P: pyriproxyfen. Bars are means \pm SE. Asterisks denote significant differences between treatments and controls. ANOVA, Fisher LSD, $P \leq 0.05$.

The present work has presented the effect of 25 ppm of cypermethrin in *C. externa* eggs; with this concentration and without any surfactant, the insecticide resulted highly toxic. There are no previous immersion tests in *C. externa* eggs with acetamiprid, pyriproxyfen or azadirachtin so far, in any concentration; the sublethal effects are variable among those products. It is remarkable that the eggs survived to all the insecticides applied, but the residual effect did last as long as to affect the neonates in contact with the chorion. Future studies on the ultrastructure of the chorion of *C. externa* eggs should be done in order to explain their tolerance to insecticides.

Conclusions

The viability of *C. externa* eggs treated by immersion with cypermethrin, acetamiprid, azadirachtin and pyriproxyfen were not affected, but neonates hatched from eggs treated with cypermethrin died after 48 h. On the other hand, reproductive parameters were affected by pyriproxyfen and acetamiprid which reduced the fecundity and fertility of adults from treated eggs. These results show the importance of evaluate the compatibility between insecticides and natural enemies, especially in situations where chemical control is predominant and conventional insecticides are still used. In an IMP program with *C. externa* acting as a bio-controller, cypermethrin should be avoided and pyriproxyfen and acetamiprid should be used with caution. Azadirachtin proved to be innocuous. Future studies on the other stages of this species treated with these insecticides should be done; also, the ultrastructure of the chorion should be characterized to explain the lack of penetration.

Acknowledgements

The author is grateful to Dr. Santiago Plischuk for his disinterested help and improvement of the final manuscript. She also wants to thank the anonymous reviewers for their valuable comments.




Figure 3. Fecundity (number of eggs/female) and fertility (number of larvae/eggs) of *Chrysoperla externa* with the different treatments. Bars are means \pm SE. Different letters denote significant differences between treatments. ANOVA, Fisher LSD, $P \leq 0.05$.

Literature cited

- ADAMS, P. A.; PENNY, N. D. 1985. Neuroptera of the Amazon basin. Part 11a. Acta Amazonica 15 (3-4): 413-480. doi:10.1590/1809-43921985153479.
- ALDRIDGE, W. N. 2013. Toxicology of pyrethroids. In: Miyamoto J, Kearney PC (Eds.). Pesticide chemistry: human welfare and the environment. Pergamon Press, Oxford, pp: 485-490.
- AWASTHI, N. S.; BARKHADE, U. P.; PATIL, S. R.; LANDE, G. K. 2013. Comparative toxicity of some commonly used insecticides to cotton aphid and their safety to predatory coccinellids. Bioscan 8 (3): 1007-1010.
- BUENO, A. F., FREITAS, S. 2004. Effect of the insecticides abamectin and lufenuron on eggs and larvae of *Chrysoperla externa* under laboratory conditions. BioControl 49 (3): 277-283.
- CARVALHO, G. A.; CARVALHO, C. F.; SOUZA, B.; ULHÔA, J. L. 2002. Selectivity of insecticides to *Chrysoperla externa* (Hagen) (Neuroptera: Chrysopidae). Neotropical Entomology 31 (4): 615-621. doi:10.1590/S1519-566X2002000400015.
- CASIDA, J. E. 2013. Novel aspects of metabolism of pyrethroids. In: Miyamoto J, Kearney PC (Eds.). Pesticide chemistry: human welfare and the environment. Pergamon Press, Oxford, pp: 187-192.
- CASTILHOS, R. V.; GRÜTZMACHER, A. D.; SIQUEIRA, P. R. B.; DE MORAES, Í. L.; GAUER, C. J. 2014. Seletividade de agrotóxicos utilizados em pessegueiro sobre ovos e pupas do predador *Chrysoperla externa*. Ciência Rural 44 (11): 1921-1928.
- CLOYD, R. A.; BETHKE, J. A. 2011. Impact of neonicotinoid insecticides on natural enemies in greenhouse and interior scape environments. Pest Management Science 67 (1): 3-9. doi:10.1002/ps.2015.
- DESNEUX, N., DECOURTYE, A.; DELPUECH, J. M. 2007. The sublethal effects of pesticides on beneficial arthropods. Annual Review of Entomology 52: 81-106. doi:10.1146/annurev.ento.52.110405.091440.
- FURNEAUX, P. J. S.; MACKAY, A. L. 1972. Crystalline protein in the chorion of insect egg shells. Journal of Ultrastructure and Molecular Structure Research 38 (3): 343-359. doi:10.1016/S0022-5320(72)90010-X.
- GALVAN, T. L.; KOCH, R. L.; HUTCHISON, W. D. 2005. Toxicity of commonly used insecticides in sweet corn and soybean to multicolored Asian lady beetle (Coleoptera:

- Coccinellidae). *Journal of Economic Entomology* 98 (3): 780-789. doi:10.1603/0022-0493-98.3.780.
- GODOY, M. S.; CARVALHO, G. A.; MORAES, J. C.; JÚNIOR, M. G.; MORAIS, A. A.; COSME, L. V. 2004. Selectivity of insecticides used in citrus crops to eggs and larvae of *Chrysoperla externa* (Hagen) (Neuroptera: Chrysopidae). *Neotropical Entomology* 33 (5): 639-646.
- GONTIJO, P. C.; MOSCARDINI, V. F.; MICHAUD, J. P.; CARVALHO, G. A. 2014. Non-target effects of chlorantraniliprole and thiamethoxam on *Chrysoperla carnea* when employed as sunflower seed treatments. *Journal of Pest Science* 87 (4): 711-719.
- HARAMBOURE, M.; MIRANDE, L.; SCHNEIDER, M. I. 2016. Improvement of the mass rearing of larvae of the Neotropical lacewing *Chrysoperla externa* through the incorporation of a new semiliquid artificial diet. *BioControl* 61 (1): 69-78. doi:10.1007/s10526-015-9699-7.
- HE, Y.; ZHAO, J.; ZHENG, Y.; DESNEUX, N.; WU, K. 2012. Lethal effect of imidacloprid on the coccinellid predator *Serangium japonicum* and sublethal effects on predator voracity and on functional response to the whitefly *Bemisia tabaci*. *Ecotoxicology* 21 (5): 1291-1300. doi:10.1007/s10646-012-0883-6.
- IANNACONE, O.; LAMAS, M. 2003. Efectos toxicológicos de extractos de molle (*Schinus molle*) y lantana (*Lantana camara*) sobre *Chrysoperla externa* (Neuroptera: Chrysopidae), *Trichogramma pintoi* (Hymenoptera: Trichogrammatidae) y *Copidosoma koehleri* (Hymenoptera: Encyrtidae) en el Perú. *Agricultura Técnica* 63 (4): 347-360.
- KOGAN, M. 1998. Integrated pest management: historical perspectives and contemporary developments. *Annual Review of Entomology* 43 (1): 243-270. doi:10.1146/annurev.ento.43.1.243.
- LAYCOCK, I.; LENTHALL, K. M.; BARRATT, A. T.; CRESSWELL, J. E. 2012. Effects of imidacloprid, a neonicotinoid pesticide, on reproduction in worker bumble bees (*Bombus terrestris*). *Ecotoxicology* 21 (7): 1937-1945. doi:10.1007/s10646-012-0974-4.
- MEDINA, P.; SMAGGHE, G.; BUDIA, F.; TIRRY, L.; VIÑUELA, E. 2003. Toxicity and absorption of azadirachtin, diflubenzuron, pyriproxyfen, and tebufenozide after topical application in predatory larvae of *Chrysoperla carnea* (Neuroptera: Chrysopidae). *Environmental Entomology* 32 (1): 196-203.
- MEDINA, P.; BUDIA, F.; DEL ESTAL, P.; VIÑUELA, E. 2004. Influence of azadirachtin, a botanical insecticide, on *Chrysoperla carnea* (Stephens) reproduction: toxicity and ultrastructural approach. *Journal of Economic Entomology* 97 (1): 43-50.
- MORDUE, A. J. 2004. Present concepts of the mode of action of azadirachtin from neem. In: Koul, O.; Wahab, S. (Eds.). *Neem: today and in the new millennium*. Springer, Netherlands, pp: 229-242.
- MOSCARDINI, V. F.; GONTIJO, P. C.; MICHAUD, J. P.; CARVALHO, G. A. 2015. Sublethal effects of insecticide seed treatments on two Nearctic lady beetles (Coleoptera: Coccinellidae). *Ecotoxicology* 24 (5): 1152-1161. doi:10.1007/s10646-015-1462-4.
- NASREEN, A.; MUSTAFA, G.; ASHFAQ, M. 2005. Mortality of *Chrysoperla carnea* (Stephens) (Neuroptera: Chrysopidae) after exposure to some insecticides; laboratory studies. *South Pacific Studies* 26 (1): 1-6.
- OHASHI, K.; NAKADA, K.; ISHIWATARI, T.; MIYAGUCHI, J.; SHONO, Y.; LUCAS, J. R.; MITO, N. 2012. Efficacy of pyriproxyfen-treated nets in sterilizing and shortening the longevity of *Anopheles gambiae* (Diptera: Culicidae). *Journal of Medical Entomology* 49 (5): 1052-1058. doi:10.1603/ME12006.
- OLIVEIRA, S. A.; AUAD, A. M.; SOUZA, B.; CARVALHO, C. A.; SOUZA, L. S.; AMARAL, R. L.; SILVA, D. M. 2009. Benefits of honey and pollen of forage on the biological parameters of *Chrysoperla externa* (Hagen, 1861) (Neuroptera: Chrysopidae). *Arquivos do Instituto Biológico (São Paulo)* 76 (4): 583-588.
- PAPPAS, M. L.; BROUFAS, G. D.; KOVEOS, D. S. 2011. Chrysopid predators and their role in biological control. *Journal of Entomology* 8 (3): 301-326. doi: 10.3923/je.2011.301.326.
- PÓRFIDO, O. D.; BUTLER, E.; DE TITTO, E.; ISSALY, P.; BENÍTEZ, R. 2014. Los Plaguicidas en la República Argentina. Ministerio de Salud de la Nación, Argentina.
- RIMOLDI, F.; SCHNEIDER, M. I.; RONCO, A. E. 2008. Susceptibility of *Chrysoperla externa* eggs (Neuroptera: Chrysopidae) to conventional and biorational insecticides. *Environmental Entomology* 37 (5): 1252-1257. doi:10.1093/ee/37.5.1252.
- RUGNO, G. R.; ZANARDI, O. Z.; YAMAMOTO, P. T. 2015. Are the pupae and eggs of the lacewing *Ceraeochrysa cubana* (Neuroptera: Chrysopidae) tolerant to insecticides?. *Journal of Economic Entomology* 108 (6): 2630-2639. doi:10.1093/jee/tov263.
- RUNDLÖF, M.; ANDERSSON, G. K.; BOMMARCO, R.; FRIES, I.; HEDERSTRÖM, V.; HERBERTSSON, L.; JONSSON, O.; KLATT, B. K.; PEDERSEN, T. R.; YOURSTONE, J.; SMITH, H. G. 2015. Seed coating with a neonicotinoid insecticide negatively affects wild bees. *Nature* 521(7550): 77-80. doi:10.1038/nature14420.
- SHINDE, C. U.; PATEL, M. B.; MEHENDALE, S. K. 2009. Relative toxicity of different insecticides to the larvae of *Chrysoperla carnea* (Stephens) under laboratory conditions. *Pest Management In Horticultural Ecosystems* 15 (2): 161-164.
- SILVA, R. A.; CARVALHO, G. A.; CARVALHO, C. F.; BASTOS SILVA, D. 2012. Effects of pesticides on eggs of *Chrysoperla externa* (Neuroptera: Chrysopidae) and consequences on subsequent development. *Revista Colombiana de Entomología* 38 (1): 58-63.
- SOARES, J. J. 2002. Fatores que afetam a predação de *Chrysoperla externa* (Neuroptera: Chrysopidae). *Campina Grande: Embrapa*, 30 p.
- US EPA - United States-Environmental Protection Agency. 2016. (March 29, 2016) Groups of pesticides in registration review. (Web page) <https://www.epa.gov/pesticide-reevaluation/groups-pesticides-registration-review>. [November 12, 2016].
- VILLAAMIL LEPORI, E. C.; MITRE, G. B.; NASSETTA, M. 2013. Situación actual de la contaminación por plaguicidas en Argentina. *Revista Internacional de Contaminación Ambiental* 29: 25-43.
- VODEB, L.; PETANOVSKA-ILIEVSKA, B. 2006. HPLC-DAD with different types of column for determination of beta-cyfluthrin in pesticide formulations. *Acta Chromatography* 17: 188-201.
- VOGT, H.; BIGLER, F.; BROWN, K.; CANDOLFI, M. P.; KEMMETER, F.; KÜHNER, C.; MOLL, M.; TRAVIS, A.; UFER, A.; VIÑUELA, E.; WLADBURGER, M.; WALTERSDORFER, A. 2000. Laboratory method to test effects of plant protection products on larvae of *Chrysoperla carnea* (Neuroptera: Chrysopidae). pp: 27-44. In: Candolfi, M. P.; Blümel, S.; Forster, R.; Bakker, F. M.; Grimm, C.; Hassan, S. A.; Heimbach, U.; Mead-Briggs, M. A.; Raber, B.; Schmuck, R.; Vogt, H. (Eds.). *Guidelines to evaluate side-effects of plant protection products to non-target arthropods*. International Organization for Biological and Integrated Control of Noxious Animals and Plants. West Palearctic Regional Section (IOBC/WPRS). Ghent, Belgium. 158 p.
- WHITEHORN, P. R.; O'CONNOR, S.; WACKERS, F. L.; GOULSON, D. 2012. Neonicotinoid pesticide reduces bumble bee colony growth and queen production. *Science* 336 (6079): 351-352. doi:10.1126/science.1215025.
- ZHONG, D.; CHANG, X.; ZHOU, G.; HE, Z.; FU, F.; YAN, Z.; ZHU, G.; XU, T.; BONIZZONI, M.; WANG, M-H.; CUI, L.; ZHENG, B.; CHEN, B.; YAN, G. 2013. Relationship between

knock-down resistance, metabolic detoxification and organismal resistance to pyrethroids in *Anopheles sinensis*. PLoS One 8 (2):e55475. doi:10.1371/journal.pone.0055475.

Received: 26-mar-2016 • Accepted: 24-jun-2017

Suggested citation:

HARAMBOURE, M. 2017. Lethal and sublethal effects after treatment with conventional and biorational insecticides on eggs of *Chrysoperla externa* (Neuroptera: Chrysopidae). Revista Colombiana de Entomología 43 (2): 161-166. Julio - Diciembre 2017. ISSN 0120-0488.