



This document is a postprint version of an article published in Diseases Of Aquatic Organisms © Inter-Research after peer review. To access the final edited and published work see <https://doi.org/10.3354/dao03336>

Document downloaded from:



1 **Baseline pathological data of the wedge clam *Donax trunculus* from the Tyrrhenian Sea**
2 **(Mediterranean Basin)**

3

4 F. Carella^{1,*}, N. Carrasco², G. De Vico¹

5 ¹ Department of Biology, University of Naples Federico II, MSA, 80126 Naples, Italy

6 ²IRTA-Sant Carles de la Rapita and Catalonia's Aquaculture R&D and Innovation Reference
7 Network (XRAq), Ctra. Poblenou Km 5, 43540 Sant Carles de la Rapita, Spain

8 ABSTRACT: In recent years, a collapse in *Donax trunculus* fishing yields has occurred in the
9 Tyrrhenian Sea (Mediterranean Basin). There is little information available on the impact disease
10 may have had on *D. trunculus* populations. For the first time, a pathological survey was performed
11 on the natural beds of the bivalve on the Campania and Lazio coasts, western Italy. Detected
12 pathogens and related diseases were analysed, and their prevalence and mean intensity values
13 were calculated. Viral particles, *Chlamydia-like* organisms, ciliates, coccidians, microcells and
14 trematodes were observed. An unknown ciliate was linked to severe inflammatory and necrotic
15 lesions in the digestive gland. Metacercariae of the trematode *Postmonorchis* sp. were also
16 strongly represented in almost all samples, reaching high levels of infection; however, none of
17 the pathogens described required the World Organisation for Animal Health to be notified. Initial
18 results indicated that further surveys related to environmental data are necessary in order to assess
19 the relevance of these early observations in managing the declining *D. trunculus* population in
20 the Tyrrhenian Sea.

21 KEY WORDS: Wedge clam • Protozoa • Bivalve disease • Trematodes

22

23 1. INTRODUCTION

24 *Donax trunculus* (Bivalvia: Donacidae) is an Atlantic-Mediterranean warm-temperate bivalve
25 species found in the Mediterranean Sea, the Black Sea and in the Atlantic Ocean in coastal regions
26 from Senegal to France. In these environments, *D. trunculus* populations are distributed in sandy
27 beds between depths of 0-2 m in the Mediterranean, and from 0-6 m on the Atlantic coasts, with
28 the highest densities at 0-3 m (Deval 2009). This bivalve represents an important fishery resource
29 due to its high economic value. Landings in Europe over the last 12 yr (since 2006) were 11202
30 t, with a maximum yield of 1355 t in 2005, followed by a steady decline, with only 516 t in 2011
31 (FAO 2013). A recent collapse in yields has occurred in the Tyrrhenian Sea of the Mediterranean
32 Basin (de la Huz et al. 2002, Rambaldi et al. 2010, Marie et al. 2016). The causes of this collapse
33 are unclear. Abiotic factors resulting from anthropogenic activity—such as beach nourishment
34 and chemical pollutants—along with extreme meteorological events related to global warming,
35 have been advocated as causing significant persistent changes to the environment of the sandy
36 beds of the Tyrrhenian coast, with serious consequences for the bivalve resources (Rambaldi et
37 al. 2010). However, other environmental stressors associated with climate change may also have
38 facilitated the spread of diseases into previously unaffected shellfish populations, exacerbated
39 disease where it already exists or facilitated the emergence of novel pathogens, which also may
40 have caused severe losses. There is very little information available on the prevalence and
41 distribution of disease in *D. trunculus* populations in the Mediterranean Sea. Although pathogens
42 such as *Rickettsia/Chlamydia-like* organisms (RLOs/CLOs), nematodes and trematodes have
43 been described as parasites of *Donax* spp. (Sindermann & Rosenfield 1967, Comps & Raimbault
44 1978, Ansell 1983, de Montaudouin et al. 2014), few studies have reported specifically on *D.*
45 *trunculus*, and little is known of the pathogens that may adversely affect the health of these
46 populations. In general, trematodes are the most often described parasites of the family
47 Donacidae, including Fellosto- midae and Monorchidae (Palombi 1933a,b, Carella et al. 2013, de
48 Montaudouin et al. 2014).

49 In the present study, we sampled *D. trunculus* from the Tyrrhenian Sea over several years and
50 describe the pathogens and diseases found using histopathology and molecular techniques to
51 provide baseline data on the health status of this species.

52 2. MATERIALS AND METHODS

53 2.1 Sampling

54 From 2008 to 2015, 330 wedge clams *D. trunculus* were collected during the summer or autumn
55 from different geographical areas. The sampling sites in the Campania region were the Volturno
56 River mouth (41° 0' 57" N, 13° 56' 50" E) from July 2008 to 2010 (90 clams) and Litorale Domitio
57 (41°4'8"N, 13° 51'4"E) from June and July 2012 (60 clams) and June 2014 and 2015 (60 clams).
58 In September 2012, samples were also collected at Formia (41° 14'21"N, 13° 36' 43" E) (30 clams)
59 in the Lazio region. In October 2013, clams were collected in 3 areas of Salerno Province
60 including Foce Sele (30 clams) (40° 29' 02" N, 14° 56' 29" E), Foce Tusciano (30 clams) (40° 34'
61 50" N, 14° 52' 58" E) and Solofrone (30 clams) (40° 22' 10" N, 15°00'0"E) (Fig. 1). The clams
62 were sampled from the shore using a hand-held semi-circular dredge as used by the local
63 fishermen.

64 2.2 Animal tissue processing

65 The animals were opened in the laboratory and examined for external signs of pathology under a
66 dissecting microscope (Nikon SMZ-10). The bivalves were then fixed in Davidson's fixative for
67 24 to 48 h and processed for routine histopathology. Additionally, pieces of digestive gland, gills
68 and gonads were also preserved from each animal for further molecular and transmission electron
69 microscopy (TEM) analysis. For histopathology, 2 transverse sections, approx. 5 mm thick,
70 including the mantle, gonad, digestive gland, gills, kidney and foot, were excised from each clam
71 and placed into histological cassettes. The tissue samples were embedded in paraffin wax, sliced
72 in 5 µm sections and subsequently stained with haematoxylin and eosin (H&E). For better
73 characterization of the tissue, additional stains, such as PAS-BA and Masson's Trichrome (Mazzi
74 1977), were also used on selected slides from every sampled area. Furthermore, to improve the
75 characterization of the prokaryote-like inclusions, additional staining techniques were used:
76 Gram's method for Gram-positive and Gram-negative bacteria, Macchiavello stain (Mazzi 1977)
77 and Giemsa (Howard & Smith 1983). All micrographs were captured using a Nikon DS-Fi1 video
78 camera mounted on a Nikon 50i microscope connected to a computer. Histological sections were
79 examined for the presence of parasites and pathological alterations. Pathogen prevalence (*P*) and
80 mean intensity (*I*) of the different parasites were calculated. For evaluation of the infection
81 intensity of trematodes such as *Postmonorchis* sp. present in gill tissue, data were obtained by
82 histological examination (histological infection density, HID) as described by Carella et al.
83 (2013).

84 The percentage of clams affected by each parasite and their pathological condition were
85 determined for each sampled bed included in the survey. The overall prevalence (and 95% CI) of
86 each parasite and pathological condition were calculated for the whole study.

87 2.3 TEM

88 After histological examination, small pieces of digestive gland parasitized by protozoa and
89 individuals with digestive glands that had a foamy aspect at the level of A cells under light
90 microscopy were selected from the stored tissues for ultrastructural analysis by TEM.

91 Tissues were fixed in 2% glutaraldehyde in filtered seawater for 24 h at 4°C, post-fixed in 2%
92 tetroxide osmium (2 h at 4°C) and embedded in EPON. Ultrathin sections (70–90 nm) were cut,
93 the tissues were stained with uranyl acetate and lead citrate and were observed using a JEOL 100
94 CX2 transmission electron microscope at the Centro Interdipartimentale di Servizio per la
95 Microscopia Elettronica (CISME), University of Naples.

96

97 2.4 Molecular identification of RLOs/CLOs

98 For RLO/CLO identification, DNA isolation was performed from pieces of digestive gland
99 previously preserved in 96% EtOH. DNA extractions were performed employing the DNeasy
100 Blood and Tissue Kit (Qiagen) following the manufacturer's instructions. The DNA concentration
101 was measured using a Nanodrop 2000c spectrophotometer (ThermoScientific), and DNA quality
102 was verified by electrophoresis on a 1 % agarose gel stained with ethidium bromide.

103 The 16S primers RCF/RCR described by Costa et al. (2012) for *Rickettsia/Chlamydia-like*
104 bacteria were used. The PCR mixture consisted of 0.4 pM of each primer RCF/RCR (Sigma),
105 PCR reactions were carried out using PCR buffer at 1x concentration, 1.5 mM MgCl₂, 200 pM of
106 each deoxynucleotide (dNTP mixture; Takara Bio) and 1 unit of *Taq* DNA Polymerase
107 (Invitrogen), 2 pl of template DNA was added to the mixture to make a final volume of 25 pl.
108 After a denaturation step at 94°C for 5 min, 30 cycles were run with an Eppendorf thermal cycle
109 as follows: denaturation at 94°C for 1 min, annealing at 51°C for 1 min and elongation at 72°C
110 for 1 min. A final elongation step at 72°C for 10 min was performed.

111 The PCR products were run on 1.2% agarose gel. The amplified fragments were gel-eluted and
112 directly sequenced. The sequencing reactions were run on a 310 Automated Sequencer (Applied
113 Biosystems). The best BLAST hits for the obtained nucleotide sequences were downloaded and
114 aligned with their corresponding sequences using ClustalW. The alignment files were employed
115 to construct neighbour-joining trees using the software MEGA v.7 (Kumar et al. 2016).

116 3. RESULTS

117 A total of 330 wedge clams from the Lazio and Campania regions were examined. Histopathologi-
118 cal analysis revealed different symbionts and pathological conditions affecting *Donax trunculus*.
119 Observed pathogens and their tropism are shown in Fig. 2; see also Table 1 & Figs. 3-9.

120 3.1 Viral infection

121 Under light microscopy, in specimens from Litorale Domitio from 2014 and 2015, some A cells
122 of the digestive epithelium were vacuolated and foamy. TEM examination of the digestive tubules
123 from 3 cases from 2014 and 2 from 2015 showed the presence of viral particles 60 nm in
124 dimension in A cells. The virions exhibited an icosahedral symmetry (Fig. 3). Moreover,
125 cytoplasmic viral factories (where viral replication and assembly likely take place) constituted by
126 both electron-dense cytoplasmic inclusions and membrane vesicles were visible (Fig. 3 insert).

127 3.2 Prokaryote infection by CLOs

128 During histology, light microscopy observations of H&E-stained tissues revealed rounded
129 intracellular inclusions of prokaryote-like colonies. The inclusions were (mean ± SD) 7.65 ± 1.79
130 pm in length and were found in the epithelium of the digestive gland in clams from all localities.
131 These colonies occupied the cytoplasm of the A cells of the digestive tubules, in some cases
132 causing hypertrophy or destruction (Fig. 4A,B). In some cases, an infiltrative mild/strong
133 haemocytic reaction was detected. The colonies were positive for *Rickettsia/Chlamydia* by
134 Macchiavello and Giemsa staining (Fig. 4C,D). Most infected clams had 8 to 20 inclusions
135 section⁻¹, but a few clams had 20 to 30 inclusions section⁻¹. The maximum *P* and mean *I* were
136 recorded in Litorale Domitio in 2015 (83%), with 30 colonies per histological section. No
137 prokaryote-like colonies were observed at Foce Sele and Solofrone in October 2013 (Table 1).
138 PCR amplification was run on 3 samples from Volturmo (2008 and 2009), 3 samples from Litorale
139 Domitio (2012) and 2 samples from Formia (2012) on a 350 bp region produced by the partial
140 16SrRNA. BLAST analysis revealed 100% identity among all the sequences, and the best
141 nucleotide identity score (92 %) was obtained for various strains of *Clamydia-* like bacteria, such
142 as *Chlamydia pecorum*, *C. ibidis* and *Estrella lausannensis* (Fig. 5). The sequences were deposited
143 in Genebank (accession no. MH492627).

144 3.3 Protists

145 Two different types of ciliates were observed: one located in the gills and another in the digestive
146 gland. Moreover, oocytes of the gregarine *Nematopsis* sp. were also detected in the connective
147 tissue of the digestive gland and gills (Fig. 6). These oocysts were oval in shape, had a maximum
148 dimension of (mean \pm D) 9.49 ± 1.24 μ m (n = 20) and usually contained 1 sporozoite oocyst⁻¹.
149 Ciliated protozoa were observed adjacent to the gills (Fig. 6A). They were oval-shaped, had dense
150 ciliature and measured 10 to 12 μ m in length. They may have been *Hypocomella* sp., which have
151 been described in other bivalve species, but no reports have occurred in the wedge clam. The
152 ciliates had 2 nuclei (macro- and micronucleus). They did not appear to cause any specific host
153 response. The maximum P were 36% in Foce Sele and 40% in Tusciano in 2013.

154 In 2 samples from 2014 and 1 sample from 2015, the presence of small protozoans similar to
155 haplosporidian microcell parasites were detected in different tissues of *D. trunculus* from Litorale
156 Domitio (June 2015), particularly in muscular and connective tissues and the intracellular space
157 within haemocytes (Fig. 6C,D). The infection was focal and low for any given organ. They mostly
158 resembled the *Mikrocytos*-like parasite previously described by Garcia et al. (2018), while for the
159 immune cell tropism they were similar to other microcells belonging to *Bonamia* species.

160
161 An unidentified ciliate endoparasite was detected in the interstitial spaces of digestive tubules in
162 the digestive gland (Fig. 7). The parasite was present in every sampled area, showing a high
163 prevalence and infection intensity, with an overall P of 219 (Fig. 7A). The parasite was always
164 accompanied by an intense inflammatory reaction, and in some cases it was associated with focal
165 necrosis of the digestive tubules and intestinal epithelia (Fig. 7B). Generally, the parasites were
166 grouped with others and were often present at the haemo-lymphatic sinus level, eating host
167 haemocytes (Fig. 7C,D). Morphologically, the ciliates were ovoid with a tapered anterior end
168 and round in crosssection. They had an estimated maximum body width
169

170 3.4 Metazoa

171 Metacercariae of the trematode *Postmonorchis* sp., as previously described by Carella et al. (2013),
172 were also detected (Fig. 8A-C). They were located at the gill level and were distributed across the
173 upper, middle and lower part of filaments, in many cases reaching high infection intensity and
174 macroscopic visibility (Fig. 8A). The mean diameter of the cysts was (mean \pm SD) $180-220 \pm$
175 23.82 μ m (n = 50). In a few cases, the parasite was detected in the foot, the stomach and the
176 kidney-pericardium system. The inflammatory response included encapsulation and ranged from
177 mild to intense in gill tissues, eliciting hypertrophy and hyperplasia of the gill epithelia.

178 Different phases of development of the metacercariae were observed. In fact, some of them had
179 a smaller dimension (~160 μ m) and a thinner cyst wall compared to other metacercariae (Fig. 8B).
180 In these cases, the pathogen appeared to be more susceptible to the host inflammatory response.
181 Other histopathological findings included mild hypertrophy or hyperplasia of mucus cells and
182 the connective tissue of the gills, as demonstrated by PAS-BA and Trichrome staining. A higher
183 prevalence was detected in the northern part of the Campania region, at the Volturno River mouth
184 in 2008 and 2009.

185 Sporocysts of the Fellodistomidae trematode *Bacciger bacciger* (Rudolphi 1819) were observed
186 in *D. trunculus* in all localities except for the areas of Salerno Province. Sporocysts containing
187 developing cercariae were found mostly in the gonad, but also in the foot, mantle and/or the whole
188 visceral mass (Fig. 8D). The fully developed cercaria showed a setiferous tail, eosinophilic
189 staining and an ovoid body of 310 ± 52 μ m (n = 10) (Fig. 8E). When present, the parasite generally
190 showed heavy infection intensity. The tissues of the gonad and digestive gland of these clams
191 were replaced by sporocysts and cercariae that caused intense damage and host castration.
192 Animals did not show a haemocyte reaction to the pathogen. Within the sporocyst, the cercariae
193 were typically in the same range of developmental stages.

194
195

197 This study revealed, for the first time, the occurrence of different pathogens and related
198 pathological conditions affecting the wedge clam *Donax trunculus* along the Campania and Lazio
199 coasts.

200 The presence of viral particles in A cells of the digestive gland were observed in our study.
201 Previous reports described 2 types of birnavirus in the bivalve *Tellina tenuis*, tellina virus1 (TV-
202 1) and tellina virus2 (TV-2) (Chung & Paetzel 2013, Nobiron et al. 2008), in digestive tubule
203 cells. TV-1 was described in connection with a population of diseased *T. tenuis* with watery flesh.
204 Infected animals had atrophic digestive tissues and were markedly thinner than healthier
205 specimens (Renault & Novoa 2004, Nobiron et al. 2008). The virus showed morphological and
206 physio- chemical characteristics similar to infectious pancreatic necrosis virus (IPNV), an
207 aquabirnavirus that infects salmonid fishes (Hill 1976, Underwood et al. 1977). On the basis of
208 their location and dimension, the virions observed in our case were in concordance with the
209 above-mentioned reports; however, further studies are needed to definitely assign them to this
210 group.

211 RLOs and CLOs are obligate, intracellular bacterial parasites associated with a variety of
212 vertebrate and invertebrate hosts (Gollas-Galvan et al. 2014). Several authors have documented
213 the presence of RLOs/CLOs in many aquatic animals, including fish, molluscs and crustaceans
214 (Gulka & Chang 1984, Wang & Gu 2002, Sun & Wu 2004), with most of the literature focused
215 on teleost fishes and bivalve molluscs. In some cases, these pathogens are causative agents of
216 severe mortality outbreaks in farmed aquatic species such as the *Rickettsia* *Candidatus*
217 *Xenohaliotis californiensis* that causes mass mortalities of *Haliotis* spp. in many countries
218 (Crosson et al. 2014). However, little is known about the pathogen's life cycle or host range
219 (Ferrantini et al. 2009). In this study, the observed intracellular bacteria were recorded within the
220 cytoplasmic vacuoles of the digestive gland epithelium of *D. trunculus*. In many cases, the
221 immune response of the host was mild. Only some specimens had digestive epithelia destruction
222 as reported by Comps & Raimbault (1978). Our molecular study revealed that the prokaryotic
223 inclusion in the digestive gland belonged to *Chlamydiales*, as also reported by Costa et al. (2012)
224 in *Ruditapes decussatus*. Interestingly, in contrast to RLOs, which have been described in
225 different tissues in bivalves, CLOs have been reported only in digestive tubules. In particular, a
226 study by Renault & Cochen- nec (1995) reported the presence of *Chlamydia* sp. in the Japanese
227 oyster *Crassostrea gigas*, where indirect fluorescent and peroxidase conjugated antibody tests
228 suggested that this agent might share common antigens with the prokaryotic agent *Chlamydia*
229 *psittaci*, strain *ovis*. Infections by members of the genera *Chlamydia* have been described in a
230 variety of adult bivalve molluscs, with reports in bivalves in Chesapeake Bay by Harshbarger et
231 al. (1977), in *Mercenaria mercenaria* from the Great South Bay (Meyers 1979) in *Argopecten*
232 *irradians* in Canada (Morrison & Shum 1982) and in mussels from Basque Country (Cajarville
233 & Angulo 1991).

234 Regarding protists, the gregarine *Nematopsis* sp. was not highly represented, with no possible
235 species identification. This parasite has not been previously reported in *D. trunculus*, but
236 *Nematopsis* sp. has been reported in different bivalve species such as *Mytilus galloprovincialis*
237 (*Carella* 2010), *R. decussatus* and *Cerastoderma glaucum* (*Longshaw & Malham* 2013).

238 Ciliate parasites are very common organisms in bivalves (Bower et al. 1994). In this study, 2 types
239 of ciliates were observed. One was present adjacent to the gills and caused evident damage. An
240 unidentified ciliate was observed in the digestive tissues of specimens in all the sampled areas
241 and had high prevalence and infection intensity. The protozoa were characterized by the presence
242 of cilia, a rounded nucleus and an elongated, tube-shaped body structure. The cytoplasm of the
243 ciliates was typically full of food vacuoles, which appeared to contain debris. Our study showed
244 that ciliate infections can cause significant morbidity related to intense tissue damage. Other
245 studies reported the detrimental effects of ciliated protozoa on the digestive tissue of bivalves.
246 Elston et al. (1999) described mortality episodes linked to an invasive opportunistic ciliate
247 belonging to Scuticociliatida. The same group of protozoa has been described in other bivalve
248 species and in crustaceans where infection is fatal (Morado & Small 1994, 1995, Cawthorn et al.
249 1996). We still do not have a specific taxonomic affiliation for the observed protozoa in *D.*
250 *trunculus* from the Tyrrhenian area. Further molecular analyses are needed to identify the
251 observed pathogen.

252 Microcell parasites are small intracellular protozoans mostly detected in molluscs, and they can
253 be associated with high mortality rates. They are essentially mollusc parasites represented by 2
254 genera: *Bonamia* and *Mikrocytos* (Carnegie & Cochenec- Laureau 2004). Microcell-like
255 parasites were detected at low prevalence in our samples in the muscles and connective tissues,
256 and the infection was focal and low for any given organ. This was the first time that a microcell
257 parasite has been detected in Italy. Further molecular analyses are required in order to classify
258 the organism.

259 Metazoan parasites detected in this survey included different phases of development of digenean
260 trematodes. *B. bacciger* was reported by Ramon et al. (1999), with adult phases originally
261 discovered by Rudolphi (1819) in marine fishes. In bivalves, the literature reports that heavy
262 sporocyst infiltration of *B. bacciger* can cause complete castration and depletion of body
263 energy, with host soft tissues becoming empty and watery (Lauckner 1983). On the French
264 Atlantic and Italian Mediterranean coasts, the parasite caused large-scale fluctuations in the
265 abundance of *Ruditapes* spp. and *D. vittatus* populations. It was also previously reported in the
266 Campania region by Palombi (1933a,b) at Fusaro Lagoon, with higher prevalence during the
267 spring from March to May. In our study, the samples were all collected during the summer
268 season, with a higher infection prevalence in Litorale Domitio in 2014 and 2015.

269 Metacercariae of *Postmonorchis* sp. have already been described by Carella et al. (2013). The
270 prevalence of the pathogen ranged from 76 to 100%, while the infection intensity fluctuated in
271 the study areas. An inflammatory response was always recorded in parasitized animals and was
272 characterized by inflammatory capsules that adhered to and enclosed the foreign body; such
273 responses were, in some cases, ineffective against completely formed metacercariae with a
274 thicker covering, as already reported (Carella et al. 2013).

275 Young (1953) described *Postmonorchis donacis* from the Pacific coast of the United States. The
276 adult phases were found in teleosts belonging to the families Sciaenidae and Embiotocidae, and
277 cercariae and metacercariae were detected in the bean clam *D. gouldii*. The reported
278 metacercariae of *Postmonorchis* sp. in *D. trunculus* showed a combination of features of the 3
279 described species of *Postmonorchis* present in the literature: *P. variabilis*, *P. orthopristsis* and *P.*
280 *donacis*.

281 In conclusion, the present study was the first attempt to identify pathogens of *D. trunculus* col-
282 lected from a natural shellfish bed on the Tyrrhenian coast, combining histological and
283 molecular analyses that should be useful as a reference in future health surveys. Gill ciliates,
284 *Nematopsis* sp., and *Chlamydia*-like colonies seemed to have limited pathogenicity and light
285 infection intensity. On the other hand, based on their pathogenic potential, the observed viral
286 particles, ciliated protozoa, bucephalid spore-cysts and monorchidae metacercariae caused
287 heavy infections.

288 Among the protozoa, the unidentified ciliate protozoan in the digestive gland was significant for
289 its high prevalence and was possibly detrimental to the wedge clam populations. None of the
290 observed parasites are on the list of obligatory notification of the World Organization for
291 Animal Health (OIE).

292 In this study, infections with microcell parasites were observed in a few specimens of *D.*
293 *trunculus*. In France, these parasites caused animal mortality (Garcia et al. 2018). Based on
294 morphological characteristics and its detection in the wedge clam, we consider that the pathogen
295 probably belongs to the novel *Mikrocytos* species detected in France. This underlines the
296 possible importance of including microcells on the list of potential disease agents.

297 Our data is limited to one season with corresponding environmental parameters. However, these
298 initial results show that further surveys related to environmental data and seasons are necessary
299 in order to assess the relevance of the pathogens and diseases observed in this declining *D.*
300 *trunculus* population in the Tyrrhenian Sea.

301
302 *Acknowledgements.* The authors acknowledge Dr. Grazia Villari for technical assistance with
303 histology. The authors also thank Dr. Francesco Bolinesi who provided technical assistance in
304 sampling in Salerno Province. The authors acknowledge the Nucleo Carabinieri Subacquei of
305 Napoli for field and diving support in the area of Volturmo and Litorale Domitio. Sample
306 processing and examination with TEM were performed by the CISME (University of Naples-
307 Department of Biology) with technical assistance from Dr. Sergio Sorbo.

308 LITERATURE CITED

309 Ansell AD (1983) The biology of the genus *Donax*. In: McLachlan A, Erasmus T (eds) Sandy beaches as ecosystems. Dr. W. Junk

310 Publishers, The Hague, p 607–635
311
312 Bower SM, McGladdery SE, Price IM (1994) Synopsis of infectious diseases and parasites of commercially exploited shellfish. *Annu*
313 *Rev Fish Dis* 4:1–199
314
315 Cajaraville MP, Angulo E (1991) Chlamydia-like organisms in digestive and duct cells of mussels from the Basque coast. *J Invertebr*
316 *Pathol* 58:381–386
317
318 Carella F (2010) Biotechnologies to evaluate the environmental status: new test organisms in ecotoxicology and histopathological and
319 molecular biomarkers in natural population. PhD thesis, University of Naples Federico II
320
321 Carella F, Culurgioni J, Aceto S, Fichi G, Pretto T, Luise D, Gustinelli A (2013) Postmonorchis sp. inq. (Digenea: Monorchidae)
322 metacercariae infecting natural beds of wedge clam *Donax trunculus* in Italy. *Dis Aquat Org* 106: 163–172
323
324 Carnegie RB, Cochenne-Laureau N (2004) Microcell parasites of oysters: recent insights and future trends. *Aquat Living Resour*
325 17:519–528
326
327 Cawthorn RJ, Lynn DH, Despres B, MacMillan R, Maloney R, Loughlin M, Bayer R (1996) Description of *Anophryoides haemophila*
328 n. sp. (Scuticociliatida: Orchitophryidae), a pathogen of American lobsters *Homarus americanus*. *Dis Aquat Org* 24:143–148
329
330 Chung IY, Paetzel M (2013) Tellina virus 1 VP4 peptidase. In: Rawlings ND, Galvesen G (eds) *Handbook of proteolytic enzymes*,
331 Vol 3. Academic Press, Amsterdam, p 3523–3527
332
333 Comps M, Raimbault R (1978) Infection rickettsienne de la glande digestive de *Donax trunculus* Linné. *Sci Pêche* 281:11–12
334
335 Costa PM, Carreira S, Lobo J, Costa MH (2012) Molecular detection of prokaryote and protozoan parasites in the commercial bivalve
336 *Ruditapes decussatus* from southern Portugal. *Aquaculture* 370-371:61–67
337
338 Crosson LM, Wight N, Van Blaricom GR, Kiryu I, Moore JD, Friedman CS (2014) Abalone withering syndrome: distribution,
339 impacts, current diagnostic methods and new findings. *Dis Aquat Org* 108:261–270
340
341 de la Huz R, Lastra M, López J (2002) The influence of grain size on burrowing, growth and metabolism of *Donax trunculus* L.
342 (*Bivalvia*: *Donacidae*). *J Sea Res* 47: 85–95
343
344 de Montaudouin X, Bazairi H, Ait Mlik K, Gonzalez P (2014) Bacciger bacciger (Trematoda: Fellodistomidae) infection effects on
345 wedge clam *Donax trunculus* condition. *Dis Aquat Org* 111:259–267
346
347 Deval MC (2009) Growth and reproduction of the wedge clam (*Donax trunculus*) in the Sea of Marmara, Turkey. *J Appl Ichthyol*
348 25:551–558
349
350 Elston RA, Cheney D, Frelie P, Lynn D (1999) Invasive orchitophryid ciliate infections in juvenile Pacific and Kumamoto oysters,
351 *Crassostrea gigas* and *Crassostrea sikamea*. *Aquaculture* 174:1–14
352
353 FAO (2013) Fisheries Global Information System (FIGIS). www.fao.org/fishery/statistics/global-capture-production/query/en, 2013
354
355 Ferrantini F, Fokini SI, Modeo L, Andreoli I, Dini F, Gortz HD, Petroni G (2009) ‘*Candidatus Cryptoprodotis polytropus*’, a novel
356 Rickettsia-like organism in the ciliated protist *Pseudomicrothorax dubius* (Ciliophora, Nassophorea). *J Eukaryot Microbiol*
357 56:119–129
358
359 Garcia C, Haond C, Chollet B, Nerac M and others (2018) Descriptions of *Mikrocytos veneroides* n. sp. and *Mikrocytos donaxi* n. sp.
360 (*Ascetosporea*: *Mikrocytida*: *Mikrocytiidae*), detected during important mortality events of the wedge clam *Donax trunculus* Linnaeus
361 (*Veneroida*: *Donacidae*), in France between 2008 and 2011. *Parasit Vectors* 11:119
362
363 Gollas-Galvan T, Avila-Villa LA, Martinez-Porchas M, Hernandez-Lopez J (2014) Rickettsia-like organisms from cultured aquatic
364 organisms, with emphasis on necrotizing hepatopancreatitis bacterium affecting penaeid shrimp: an overview on an emergent concern.
365 *Rev Aquacult* 6:256–269
366
367 Gulka G, Chang PW (1985) Pathogenicity and infectivity of a rickettsia-like organism in the sea scallop, *Placopecten magellanicus*. *J*
368 *Fish Dis* 8:309–318
369
370 Harshbarger JC, Chang SC, Otto SV (1977) Chlamydiae (with phages), mycoplasmas, and rickettsia in Chesapeake Bay bivalves.
371 *Science* 196:666–668
372
373 Hill BJ (1976) Properties of a virus isolated from the bivalve mollusc *Tellina tenuis* (Da Costa). In: Page LA (ed) *Wildlife diseases*.
374 Plenum Press, New York, NY, p 445–452
375
376 Howard DW, Smith CS (1983) Histological techniques for marine bivalve mollusks. NOAA Tech Memo NMFSF/NEC-25. National
377 Oceanic and Atmospheric Administration, Woods Hole, MA
378
379 Kumar S, Stecher G, Tamura K (2016) MEGA7: molecular evolutionary genetics analysis version 7.0 for bigger datasets. *Mol Biol*
380 *Evol* 33:1870–1874
381
382 Lauckner G (1983) Diseases of Mollusca. *Bivalvia*. In: Kinne O (ed) *Diseases of marine animals*, Vol 11. Biologische Anstalt
383 Helgoland, Hamburg, p 477–961
384
385 Longshaw M, Malham SK (2013) A review of the infectious agents, parasites, pathogens and commensals of European cockles

386 (Cerastoderma edule and C. glaucum). J Mar Biol Assoc UK 93:227–247
387
388 Marie AD, Lejeune C, Karapatsiou E, Cuesta JA and others (2016) Implications for management and conservation of the population
389 genetic structure of the wedge clam Donax trunculus across two biogeographic boundaries. Sci Rep 6:39152
390
391 Mazzi V (1977) Manuale di tecniche istologiche e istochimiche. Piccin, Rome Meyers TR (1979) Preliminary studies on a chlamydial
392 agent in the digestive diverticular epithelium of hard clams Mercenaria mercenaria (L.) from Great South Bay, New
393 York. J Fish Dis 2:179–189
394
395 Morado JF, Small EB (1994) Morphology and stomatogenesis of Mesanophrys pugettensis n. sp. (Scuticociliatida: rchitophryidae), a
396 facultative parasitic ciliate of the Dungeness crab, Cancer magister (Crustacea: Decapoda). Trans Am Microsc Soc 113:343–364
397
398 Morado JF, Small E (1995) Ciliate parasites and related diseases of Crustacea: a review. Rev Fish Sci 3:275–354 Morrison C, Shum
399 C (1982) Chlamydia like organisms in the digestive diverticula of the bay scallop, Argopecten irradians (Lmk). J Fish Dis 5:173–184
400
401 Nobiron I, Galloux M, Henry C, Torhy C and others (2008) Genome and polypeptides characterization of Tellina virus 1 reveals a
402 fifth genetic cluster in the Bimaviridae family. Virology 371:350–361
403
404 Palombi A (1933a) Bacciger bacciger (Rud.) Nicoll, 1914, forma adulta di Cercaria pectinata Huet, 1891. Boll Soc Nat Napoli
405 44:217–219
406
407 Palombi A (1933b) Cercaria pectinata Huet et Bacciger bacciger (Rud.) Rapporti genetici e biologia. Boll Zool 4:1–11
408
409 Rambaldi E, Lanni L, Pelusi P, Binda F and others (2010) Valutazione dei banchi naturali di molluschi bivalvi eduli (telline, Donax
410 trunculus e cannolicchi, Ensis siliqua) lungo la fascia costiera della provincia di Latina e indicazioni gestionali per una pesca
411 sostenibile. Biol Mar Mediterr 17:328–329
412
413 Ramòn M, Gracenea M, Gonzales-Moreno O (1999) Bacciger bacciger (Trematoda, Fellodistomidae) infection in commercial clams
414 Donax trunculus (Bivalvia, Donacidae) from the sandy beaches of the Western Mediterranean. Dis Aquat Org 35:37–46
415
416 Renault T, Cochenec N (1995) Chlamydia-like organisms in ctenidia and mantle cells of the Japanese oyster Crassostrea gigas from
417 the French Atlantic coast. Dis Aquat Org 23:153–159
418
419 Renault T, Novoa B (2004) Viruses infecting bivalve molluscs. Aquat Living Resour 17:397–409
420
421 Rudolphi CA (1819) Entozoorum synopsis cui accedunt mantissa duplex et indices locupletissimi. Riicker, Berlin
422
423 Sindermann CJ, Rosenfield A (1967) Principal diseases of commercially important marine bivalve Mollusca and Crustacea. Fish Bull
424 66:335–385
425
426 Sun J, Wu X (2004) Histology, ultrastructure, and morphogenesis of a rickettsia-like organism causing disease in the oyster,
427 Crassostrea ariakensis Gould. J Invertebr Pathol 86:77–86
428
429 Underwood BO, Smale CJ, Brown F, Hill BJ (1977) Relationship of a virus from Tellina tenuis to infectious pancreatic necrosis virus.
430 J Gen Virol 36:93–109
431
432 Wang W, Gu Z (2002) Rickettsia-like organism associated with tremor disease and mortality of the Chinese mitten crab Eriocheir
433 sinensis. Dis Aquat Org 48:149–153
434
435 Young RT (1953) Postmonorchis donacis, a new species of monorchid trematode from the Pacific coast, and its life history. J Wash
436 Acad Sci 43:88–93