

1
2
3
4
5 **Compound-Specific Carbon Isotope Compositions of Aldehydes and Ketones in the**
6 **Murchison Meteorite**
7

8 Danielle N. Simkus^{1,2,3*} (danielle.n.simkus@nasa.gov), José C. Aponte^{3,4}
9 (jose.c.aponte@nasa.gov), Robert W. Hiltz⁵ (hiltzr@macewan.ca), Jamie E. Elsil³
10 (jamie.e.cook@nasa.gov), Christopher D. K. Herd¹ (herd@ualberta.ca)

11 ¹Department of Earth and Atmospheric Sciences, University of Alberta, Edmonton, AB, Canada

12 ²*Current affiliation:* NASA Postdoctoral Program at NASA Goddard Space Flight Center,
13 Greenbelt, MD, USA

14 ³Solar System Exploration Division, Code 691, NASA Goddard Space Flight Center, Greenbelt,
15 MD, USA

16 ⁴Catholic University of America, Washington, D.C., USA

17 ⁵Department of Physical Sciences, MacEwan University, Edmonton, AB, Canada

18
19 Accepted to:

20 Meteoritics & Planetary Science
21
22
23

24 *Corresponding author:

25 Danielle N. Simkus

26 *Formerly at:*

27 Department of Earth & Atmospheric Sciences, University of Alberta, Edmonton, AB, Canada

28 *Present address:*

29 Solar System Exploration Division, Code 691

30 NASA Goddard Space Flight Center

31 Greenbelt, MD 20771

32 Phone: (301) 614-5887

33 danielle.n.simkus@nasa.gov
34
35

ABSTRACT


Compound-specific carbon isotope analysis ($\delta^{13}\text{C}$) of meteoritic organic compounds can be used to elucidate the abiotic chemical reactions involved in their synthesis. The soluble organic content of the Murchison carbonaceous chondrite has been extensively investigated over the years, with a focus on the origins of amino acids and the potential role of Strecker-cyanohydrin synthesis in the early solar system. Previous $\delta^{13}\text{C}$ investigations have targeted α -amino acid and α -hydroxy acid Strecker products and reactant HCN; however, $\delta^{13}\text{C}$ values for meteoritic aldehydes and ketones (Strecker precursors) have not yet been reported. As such, the distribution of aldehydes and ketones in the cosmos and their role in prebiotic reactions have not been fully investigated. Here, we have applied an optimized *O*-(2,3,4,5,6-pentafluorobenzyl)hydroxylamine (PFBHA) derivatization procedure to the extraction, identification and $\delta^{13}\text{C}$ analysis of carbonyl compounds in the Murchison meteorite. A suite of aldehydes and ketones, dominated by acetaldehyde, propionaldehyde and acetone, were detected in the sample. $\delta^{13}\text{C}$ values, ranging from -10.0% to $+66.4\%$, were more ^{13}C -depleted than would be expected for aldehydes and ketones derived from the interstellar medium, based on interstellar $^{12}\text{C}/^{13}\text{C}$ ratios. These relatively ^{13}C -depleted values suggest that chemical processes taking place in asteroid parent bodies (e.g. oxidation of the IOM) may provide a secondary source of aldehydes and ketones in the solar system. Comparisons between $\delta^{13}\text{C}$ compositions of meteoritic aldehydes and ketones and other organic compound classes were used to evaluate potential structural relationships and associated reactions, including Strecker synthesis and alteration-driven chemical pathways.

1. INTRODUCTION

Carbonaceous chondrite meteorites are some of the most primitive materials in our solar system, and their organic inventory contains a wealth of information about the chemistry of the presolar cloud, the protoplanetary disk, and the subsequent chemical processes that took place on and within parent body asteroids. Investigating the origins of organic compounds in meteorites may provide insights into the chemical processes that led to the origin of life on Earth, as well as the potential for life to exist elsewhere in our solar system. The Murchison meteorite, a CM type 2 meteorite that fell in southeastern Australia in 1969, is one of the most widely studied carbonaceous chondrites due to the large mass of sample material available, the relatively pristine nature of the meteorite and the structural diversity of its organic content. The findings from ~50 years of research on the chemistry of organic compounds of the Murchison meteorite have built a valuable reference data set for all other carbonaceous chondrites. The soluble organic inventory of the Murchison meteorite includes amino acids, carboxylic acids, polycyclic aromatic hydrocarbons (PAHs), amines, polyols, aldehydes, ketones, and many more compounds (see reviews and references therein: Botta and Bada, 2002; Pizzarello et al. 2006; Sephton, 2002, 2014).

Aldehydes and ketones (collectively referred to as “carbonyl compounds”) may play an important role as precursors to many biologically relevant organic compounds in the solar system. One of the most widely studied chemical reactions involving aldehyde and ketone precursors is Strecker-cyanohydrin synthesis, a proposed source of α -amino acids and α -hydroxy acids in meteorites (Scheme 1; Peltzer and Bada, 1978). The α -amino acid products of Strecker synthesis are of particular astrobiological interest as they are monomers of proteins and enzymes in living systems and their delivery by meteorites may hold clues to the origins of prebiotic organic compounds on the early Earth (Kvenvolden et al., 1970). For this reason, the distribution and stable isotopic compositions of amino acids in meteorites have been extensively analyzed to elucidate their synthetic pathways (e.g., Strecker synthesis (Peltzer et al., 1984), Michael addition (Miller et al., 1957), carbon dioxide (CO₂) addition to amines (Hudson et al., 2009), and reductive amination of keto acids (Huber and Wächtershäuser, 2003)). Despite the potential significance of Strecker synthesis and the relevance of aldehydes and ketones in prebiotic organic chemistry, the stable carbon isotopic compositions ($\delta^{13}\text{C}$) of aldehydes and ketones in meteorites have not yet been

1 measured. Thus, the synthetic relationships between meteoritic aldehydes, ketones and amino
 2 acids have not been fully investigated, representing a significant gap in the literature.



Scheme 1. The Strecker-cyanohydrin synthesis reaction (illustration adapted from Elsila et al., 2016), yielding α -amino acids in the presence of NH_3 and α -hydroxy acids in the absence of NH_3 (Peltzer et al., 1984). Colors are used to trace the path of different carbon atoms from precursor to products.

3

4 In addition to the Strecker reaction, aldehydes and ketones are potentially structurally
 5 related to several other prebiotic organic compounds (Figure 1), though some of these reactions
 6 have yet to be evaluated under astrochemically-relevant conditions. Carbonyl compounds and
 7 alcohols may be interconverted via oxidation/hydrogenation reactions (Bisschop et al., 2007).
 8 Aldehydes can be further oxidized to produce carboxylic acids (Corey et al., 1968), the chemical
 9 building blocks of biological membranes (Deamer et al., 2002). Aldehydes and ketones may also
 10 be converted to amines via reductive amination (Burk et al., 1994; Abdel-Magid et al., 1996), and
 11 these amines may be subsequently converted to amino acids via CO_2 addition from photochemical
 12 and/or ion-irradiation reactions (Holtom et al., 2005; Hudson et al., 2009; Bossa et al., 2009; Lee
 13 et al., 2009). The simplest aldehyde, formaldehyde, is thought to play a particularly important role
 14 in the synthesis of prebiotic organic compounds (Cleaves, 2008). In addition to the reactions
 15 summarized above, formaldehyde has been proposed as a precursor to meteoritic sugars, sugar-
 16 alcohols and insoluble organic matter (IOM) via formose reactions (Cooper et al., 2001; Cody et
 17 al., 2011; Meinert et al., 2016).




Figure 1. Schematic of potential synthetic relationships between carbonyl compounds⁽¹⁾ and other structurally-related organic compounds. Illustrated are the amino acid⁽²⁾ and hydroxy acid⁽³⁾ products of Strecker synthesis, carboxylic acids⁽⁴⁾ produced via oxidation of aldehydes, alcohol⁽⁵⁾ precursors to aldehydes/ketones, and amines⁽⁶⁾ formed through reductive amination of aldehydes/ketones, decarboxylation of amino acids, and nucleophilic substitution of alcohols. **A.** The simplest aldehyde (formaldehyde) and structurally-related organics. **B.** The simplest ketone (acetone) and structurally-related organics.

1

2 Formaldehyde is ubiquitous in the interstellar medium (ISM) and comets (Irvine, 1999;

3 Cleaves, 2008; Mumma and Charnley, 2011). Proposed mechanisms for the formation of

4 interstellar formaldehyde include successive hydrogenation of carbon monoxide (CO) on ice/grain

5 surfaces (Tielens and Whittet, 1997; Watanabe and Kouchi, 2002; Awad et al., 2005), gas-phase

6 reactions (Shalabiea and Greenberg, 1994), and UV photolysis of H₂O-CO ices (Allamandola et

7 al., 1988; Schutte et al., 1996). Higher molecular weight carbonyl compounds in the interstellar

8 medium may have been synthesized by further addition of carbon from CO (Charnley et al., 2004)

9 and then become incorporated into comets and asteroids during early solar system formation (Botta

10 and Bada, 2002).

11 Aldehydes and ketones have been previously investigated in Murchison meteorite samples

12 via colorimetric analysis, gas chromatography (GC), and gas chromatography-mass spectrometry


13 (GC-MS) analysis of water extracts and headspace gas (Jungclaus et al., 1976) and by applying

14 the 1998 Environmental Protection Agency (EPA) *O*-(2,3,4,5,6-pentafluorobenzyl)

15 hydroxylamine (PFBHA) derivatization method (EPA Method #556 (1998); Pizzarello and

16 Holmes (2009); Monroe and Pizzarello (2011); Pizzarello et al., 2012)). Derivatization of volatile

1 aldehydes and ketones produces less polar, more thermally stable oxime derivatives that are more
 2 amenable to GC-MS analysis. Furthermore, derivatization of the carbonyl compounds allows for
 3 chromatographic separation of these species from alcohols and other unknown compounds which
 4 would otherwise coelute (Jungclaus et al., 1976). The PFBHA-derivatization method (Scheme 2)
 5 is commonly used for the analysis of carbonyl compounds in environmental and biological samples
 6 and it has been modified in the literature for various suites of terrestrial compounds (e.g. Spaulding
 7 and Charles, 2002; Rodigast et al., 2015; see Table S1); however, this procedure has not previously
 8 been optimized specifically for the analysis of carbonyl compounds extracted from carbonaceous
 9 chondrites. Method optimization is particularly important for maximizing yields for isotopic
 10 analyses when working with limited quantities of extraterrestrial samples of low organic content.
 11 The previous reports of aldehydes and ketones in Murchison identified similar suites of
 12 compounds, dominated by low molecular weight acetaldehyde, formaldehyde and acetone;
 13 however, the explanation of their origin was limited by the absence of stable isotopic
 14 measurements. Compound-specific $\delta^{13}\text{C}$ analysis of meteoritic aldehydes and ketones is a key next
 15 step for understanding the origins of these carbonyl compounds in the solar system.



Scheme 2. Derivatization of carbonyl compounds. PFBHA reacts with an aldehydes or ketones to produce the (*E*)- and (*Z*)-PFBHA derivatives.

16
 17 The objectives of the present study were: (a) to optimize the EPA PFBHA derivatization
 18 method specifically for the analysis of aldehydes and ketones in astromaterials in order to
 19 maximize both yields and reproducibility for the detection of low molecular weight aldehydes and
 20 ketones in samples with organic contents in nmol/g of meteorite range; (b) to identify, quantify,
 21 and, for the first time, measure the compound-specific $\delta^{13}\text{C}$ values of aldehydes and ketones in the
 22 Murchison meteorite; and (c) to compare the measured $\delta^{13}\text{C}$ values of indigenous aldehydes and
 23 ketones with previously measured $\delta^{13}\text{C}$ values of other meteoritic organics including amino acids,

1 hydroxy acids, HCN, amines, carboxylic acids, and IOM in Murchison, in order to investigate
2 potential synthetic relationships between these compound classes.

3 **2. MATERIALS AND METHODS**

4 **2.1. Chemicals and reagents**

5 Standards and reagents were purchased from Sigma Aldrich and Fisher Scientific. Five
6 aldehydes and five ketones previously identified in carbonaceous chondrites (Jungclaus et al.,
7 1976; Yabuta et al., 2007; Pizzarello and Holmes, 2009; Monroe and Pizzarello, 2011; Pizzarello
8 et al., 2012) were used for the method development experiments and as the standards:
9 formaldehyde, acetaldehyde, propionaldehyde, butyraldehyde, benzaldehyde, acetone, 2-
10 butanone, 2-pentanone, 2-hexanone, and acetophenone. Ultrapure water (Millipore Direct Q3 UV,
11 18.2 M Ω , 3 ppb total organic carbon; hereafter referred to as “water”), HPLC grade
12 dichloromethane (DCM), double-distilled 6 M HCl, and *O*-(2,3,4,5,6-pentafluorobenzyl)
13 hydroxylamine hydrochloride of $\geq 99.0\%$ purity were used. All glassware and tools were wrapped
14 in aluminum foil and heated at 450°C for a minimum of 6 hours before use. All vials were capped
15 with PTFE-lined lids.

16 **2.2. Murchison sample extraction**

17 An interior chip of the Murchison carbonaceous chondrite (CM2, USNM 54512; extracted
18 mass: 0.5075 g) was provided by the Smithsonian National Museum of Natural History,
19 Washington, D.C. The sample was powdered using a porcelain mortar and pestle and extracted
20 with 1 mL of water in a flame-sealed glass ampule at 100°C for 24 hours. After extraction, the
21 sample was centrifuged, and the supernatant was transferred to a glass vial. The residual meteorite
22 solid was rinsed three more times using 0.5 mL of water for each rinse. The aqueous supernatant
23 and rinses were combined into one fraction and filtered through quartz wool to remove any
24 remaining solid material from the solution. The quartz wool filter was rinsed two more times using
25 0.5 mL of water for each rinse. The entire extraction procedure was carried out in parallel with a
26 procedural solvent blank and a serpentine mineral analogue blank (0.5013 g; powdered and
27 combusted at 500°C for 16 hours before extraction).

28 **2.3. Derivatization of aldehydes and ketones**

1 Carbonyl compounds were derivatized using an optimized EPA Method #556 for PFBHA
2 derivatization (see Supplementary Information), as follows: 1 mL of 0.2 mg/mL PFBHA solution
3 was added to the 3.5 mL of combined water extract and rinses. The solution was agitated for 5
4 minutes and then left to react for 24 hours at room temperature to allow the derivatization reaction
5 to go to completion. The reaction was quenched by adding 100 μ L of 0.4 M HCl solution. Two
6 mL of dichloromethane were then added to the solution to extract the derivatized carbonyl
7 compounds. The resulting mixture was agitated for 5 minutes and then left undisturbed for 30
8 minutes to allow the dichloromethane and water layers to settle. The dichloromethane layer was
9 separated from the aqueous layer and brought through an acid-wash step (3 mL of 0.4 M HCl).
10 The extraction of the aqueous layer was repeated with another 2 mL of dichloromethane and the
11 isolated dichloromethane layer was washed with 3 mL of 0.4 M HCl. The 2 x 2 mL
12 dichloromethane extracts containing the PFBHA derivatives were combined, the volume was
13 reduced to 200 μ L under a stream of nitrogen, and the resulting concentrated solution was analyzed
14 by gas chromatography coupled to mass spectrometry and isotope ratio mass spectrometry (GC-
15 MS/IRMS; section 2.4).

16 **2.4. Identification, quantification and compound-specific $\delta^{13}\text{C}$ analysis of aldehydes and** 17 **ketones**

18 The analysis of derivatized carbonyl compounds was performed using GC-MS/IRMS,
19 which provides compound-specific identification and stable carbon isotopic ratios in parallel
20 (Elsila et al., 2012; Aponte et al. 2014). The GC separation was accomplished using a Thermo
21 Trace GC equipped with a 5 m base-deactivated fused silica guard column (Restek, 0.25 mm ID)
22 and three 30 m length \times 0.25 mm I.D. \times 0.5 μ m film thickness Rxi-5ms capillary columns (Restek)
23 connected using Press-Tight[®] connectors (Restek). The oven program was set as follows: initial
24 temperature was 40 $^{\circ}\text{C}$, ramped at 10 $^{\circ}\text{C}/\text{min}$ to 160 $^{\circ}\text{C}$, ramped at 5 $^{\circ}\text{C}/\text{min}$ to 190 $^{\circ}\text{C}$, ramped at
25 10 $^{\circ}\text{C}/\text{min}$ to 290 $^{\circ}\text{C}$ and held for 7 min. The carrier gas used was UHP helium (5.0 grade) at 2.6
26 mL/min flow rate. Triplicate injections of PFBHA derivatives were made in splitless mode in
27 aliquots of 1 μ L; splitless mode was used to maximize sensitivity and minimize potential isotopic
28 fractionation during injection. The mass spectrum was used to identify and quantify the meteoritic
29 carbonyl compounds by comparison to reference standards and application of calibration curves.
30 Five-point external calibration curves were prepared for each individual isomer for ten carbonyl

1 standards (Section 2.1). These calibration curves were obtained immediately prior to the
 2 Murchison sample analysis to limit temporal variations of response factors, in lieu of introducing
 3 internal standards which could result in chromatographic coelutions. Selected ion mass-to-charge
 4 ratio ($m/z = 181.0$) was used to identify and quantify compounds (see Table 1 for compound
 5 identifications). Concentrations were calculated using quadratic equations (average $R^2 = 0.995$)
 6 derived from the calibration curves of each individual isomer. Approximately 10% of the sample
 7 eluting from the GC column was directed into a Thermo DSQII electron-impact quadrupole mass
 8 spectrometer (ion source set at 200 °C and 70 eV). The remaining 90% of each eluting compound
 9 was directed through a Thermo GC-C III interface for oxidation of the compounds to carbon
 10 dioxide; the carbon stable isotopic measurement was then made on a Thermo MAT 253 IRMS.
 11 The $\delta^{13}\text{C}$ values of the eluting compounds were obtained after injection of three pulses of
 12 precalibrated CO_2 ($\delta^{13}\text{C} = -24.23\text{‰}$ VPDB) into the IRMS and computation using Thermo Isodat
 13 2.5 software. In order to correct for the isotopic contribution from carbon added by the
 14 derivatization reagent, $\delta^{13}\text{C}$ values were also determined for both derivatized carbonyl standards
 15 and underivatized carbonyl standards. The underivatized standards were analyzed on a Costech
 16 ECS 4010 combustion elemental analyzer (EA) connected to the IRMS. The final $\delta^{13}\text{C}$ values of
 17 the meteoritic aldehydes and ketones (carbonyls) were calculated using equation 1 (derived from
 18 equation 1 of Docherty et al., 2001).

$$19 \quad \delta^{13}\text{C}_{\text{sample carbonyl}} = \left[\left(\frac{n_{\text{carbonyl}} + n_d}{n_{\text{carbonyl}}} \right) \times (\delta^{13}\text{C}_{\text{derivatized sample carbonyl}} - \delta^{13}\text{C}_{\text{derivatized std carbonyl}}) \right] + \delta^{13}\text{C}_{\text{underivatized std carbonyl}} \quad (1)$$

20 where n_{carbonyl} = number of carbon atoms in underivatized carbonyl and n_d = number of carbons
 21 added by derivatizing reagent. The precision (standard deviation) of the $\delta^{13}\text{C}$ values was obtained
 22 using equation 2 (Docherty et al., 2001).

$$23 \quad \sigma^2_{\text{sample carbonyl}} = \sigma^2_{\text{underivatized std}} + \left[\sigma^2_{\text{derivatized std}} \times \left(\frac{n_{\text{carbonyl}} + n_d}{n_{\text{carbonyl}}} \right)^2 \right] + \left[\sigma^2_{\text{derivatized sample}} \times \left(\frac{n_{\text{carbonyl}} + n_d}{n_{\text{carbonyl}}} \right)^2 \right] \quad (2)$$

24
 25
 26
 27
 28

3. RESULTS

3.1. Identification, quantification and compound-specific $\delta^{13}\text{C}$ analysis of aldehydes and ketones in the Murchison meteorite

A suite of aldehydes and ketones were identified and quantified in the Murchison meteorite and the $\delta^{13}\text{C}$ values were measured for those compounds present in sufficient abundance. Table 1 lists the concentrations and $\delta^{13}\text{C}$ values for the measurable aldehydes and ketones in the Murchison sample. Figure 2 illustrates the selected ion ($m/z = 181.0$) GC-MS chromatogram of representative injections from the derivatized Murchison meteorite extract, the procedural blanks, and a mixture of derivatized aldehydes and ketone standards. Two chromatographic peaks are observed for most compounds, as (*E*)- and (*Z*)-isomers (two molecules with the same molecular formula but different stereometric configurations) are produced during derivatization for those carbonyl compounds with asymmetrical chemical structures. Abundances are reported as sums of the (*E*)- and (*Z*)-isomers, whereas the $\delta^{13}\text{C}$ values for the two isomers are reported individually. The elution order of the (*E*)- and (*Z*)-isomer peaks reported here is based on peak identities from previous studies that have used similar GC columns (EPA Method #556, 1998; Cancho et al., 2001). The Murchison sample was dominated by the low molecular weight aldehyde acetaldehyde (273.4 nmol/g), followed by propionaldehyde (149.8 nmol/g), acetone (87.5 nmol/g), and formaldehyde and higher molecular weight carbonyl compounds at lower abundances. Formaldehyde exhibited the most ^{13}C -enriched carbon isotope composition (+66.4‰). The aldehyde $\delta^{13}\text{C}$ values (+20.3 to +66.4‰) were more ^{13}C -enriched in comparison to the ketone $\delta^{13}\text{C}$ values (−10.0‰ and +11.8‰). The $\delta^{13}\text{C}$ values for the acetaldehyde (*E*)- and (*Z*)-isomers fell within one standard deviation of one another. A trace amount of formaldehyde was observed in the procedural blank (0.5 nmol) and in the combusted serpentine blank (0.9 nmol), and trace amounts of acetaldehyde (2.1 nmol) and acetone (0.7 nmol) were observed in the combusted serpentine blank. The Murchison formaldehyde, acetaldehyde and acetone abundances were corrected using these serpentine blank measurements (Table 1).

Table 1. Concentrations (nmol/g) and $\delta^{13}\text{C}$ values of aldehydes and ketones identified in the Murchison meteorite.

Peak label	Compound	Abundance (nmol/g)		$\delta^{13}\text{C}$ (‰)
		This study ^a (\pm S.D. ^b)	Pizzarello and Holmes (2009)	
Aldehydes				
1	Formaldehyde ^c	19.2 \pm 2.8	10.0	+66.4 \pm 3.2
2	(<i>E</i>)-Acetaldehyde ^c	273.4 \pm 48.7	24.0	+25.7 \pm 0.9
3	(<i>Z</i>)-Acetaldehyde			+27.0 \pm 0.9
5	(<i>E</i>)-Propionaldehyde ^c	149.8 \pm 22.8	23.5	<i>n.d.</i>
6	(<i>Z</i>)-Propionaldehyde			+41.8 \pm 1.3
9	(<i>E</i>)-Butyraldehyde ^c	42.2 \pm 6.6	32.2	+20.3 \pm 1.3
10	(<i>Z</i>)-Butyraldehyde			<i>n.d.</i>
16	(<i>E</i>)-Benzaldehyde	<i>n.d.</i>	7.6	<i>n.d.</i>
17	(<i>Z</i>)-Benzaldehyde			<i>n.d.</i>
Ketones				
4	Acetone	87.5 \pm 15.4	47.3	+11.8 \pm 1.5
7	(<i>E</i>)-2-Butanone	36.0 \pm 5.7	6.9	<i>n.d.</i>
8	(<i>Z</i>)-2-Butanone ^c			-10.0 \pm 1.2
11	(<i>E</i>)-2-Pentanone	12.3 \pm 1.8	1.5	<i>n.d.</i>
12	(<i>Z</i>)-2-Pentanone			<i>n.d.</i>
13	(<i>E</i>)-2-Hexanone	3.5 \pm 0.9	25.0	<i>n.d.</i>
14	(<i>Z</i>)-2-Hexanone			<i>n.d.</i>
15	(<i>E</i>)-Acetophenone	1.9 \pm 0.6	<i>n.d.</i>	<i>n.d.</i>
18	(<i>Z</i>)-Acetophenone			<i>n.d.</i>
Total abundance		626.2 \pm 105.3	200	

^aAbundances represent the sum of the two (*E*) and (*Z*) isomers, where applicable.

^bS.D. – Standard deviation of three sequential injections.

^cPartially co-eluting with an unidentified compound

n.d. – not determined.




Figure 2. GC-MS chromatograms of derivatized aldehyde and ketone standards, the serpentine mineral analogue blank, and the Murchison meteorite extract, and a GC-IRMS chromatogram of carbonyl compounds in the Murchison sample measured for $\delta^{13}\text{C}$. GC-MS peaks in the upper three chromatograms represent PFBHA derivatives, containing the characteristic ion of $m/z = 181.0$. The lower GC-MS

chromatogram is a total ion chromatogram (TIC). The GC-IRMS chromatogram shows the $m/z = 44$ ($^{12}\text{CO}_2$ peak) measured during carbon compound-specific isotope analysis. Peak *R* represents unreacted PFBHA derivatization reagent. The identities of the labeled peaks are presented in Table 1.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27

4. DISCUSSION

4.1. Method optimization for PFBHA derivatization and analysis of aldehydes and ketones in astromaterials

4.1.1. Modifications to the EPA Method #556

We have optimized the EPA Method #556 (Scheme 2), used previously for the analysis of the Murchison meteorite (Pizzarello and Holmes, 2009), with an aim to maximize yields and reproducibility for a set of ten aldehyde and ketone standards previously identified in carbonaceous chondrites (Section 2.1). The EPA Method #556 has been similarly optimized for atmospherically relevant dialdehydes, diketones, and unsaturated and aromatic aldehydes (Rodigast et al., 2015). Our method optimization work, though focused on relatively lower molecular weight target analytes, produced results that are consistent with those obtained by Rodigast et al. (2015). Derivatization times reported in the literature vary widely, from 10 minutes to 96 hours at room temperature (see Table S1, adapted from Rodigast et al., 2015). While most aldehydes are expected to derivatize completely within a few hours, ketones are known to require much longer derivatization times and are generally given 24 hours for the reaction to reach completion (Kobayashi et al., 1980; Glaze et al., 1989; Yamada and Somiya, 1989). In the present study, we tested four derivatization times (2, 8, 16 and 24 hours) to confirm the influence of reaction time on our suite of standards (Figure 3a). While the total yields of the lower molecular weight aldehydes appear to plateau within 2 hours, the ketones required at least 16 hours in order to reach their highest relative yields, as indicated by the similar (within error) GC-MS peak areas for the 16-hour and 24-hour samples. As such, we have chosen a 24-hour derivatization time to maximize our yields and reproducibility. The EPA Method #556 recommends using hexane to extract the carbonyl derivatives from aqueous solution; however, dichloromethane is also commonly used in the literature for this method (e.g. Spaulding and Charles, 2002; de Marcellus et al., 2015; Rodigast et al., 2015; see Table S1). Thus, we compared the extraction efficiencies of dichloromethane and hexane for our suite of derivatized standards (Figure 3b). The use of dichloromethane resulted in

1 relatively higher yields and greater reproducibility for all our compounds of interest and was,
2 therefore, selected as the extraction solvent for this study.




Figure 3. **A.** Total yields (GC-MS peak areas) of aldehyde and ketone PFBHA derivatives from 2-, 8-, 16- and 24-hour derivatization reactions (standard abundances: 1 mL of 50 $\mu\text{g/mL}$ solution). Each individual bar represents 3 replicate samples and error bars represent one standard deviation from the mean. All PFBHA derivatives were isolated using dichloromethane (DCM) as the extraction solvent. **B.** Extraction efficiency of DCM vs. hexane for aldehyde and ketone PFBHA derivatives. Each bar represents the average of 12 replicate samples and the error bars represent one standard deviation from the mean. All derivatization reactions were carried out for 24 hours.

3 Other changes to the method included: (1) using a lower concentration of PFBHA solution
4 (0.2 mg/mL), as the expected yields of aldehydes and ketones in astromaterials are much lower
5 than terrestrial environmental samples and excess PFBHA reagent in solution can result in less
6 effective extraction of the derivatives into the organic phase (Rodigast et al., 2015); (2) replacing
7 H_2SO_4 with HCl, as HCl can be double-distilled for higher purity; (3) carrying out a 30 minute
8 extraction time, to allow for a complete extraction of the target analytes, as recommended by
9 Rodigast et al. (2015); and (4) storing samples in solution, at low temperature (4°C), to avoid
10 decomposition of the PFBHA derivatives (see Section 4.1.2).

11 4.1.2. Stability of the PFBHA derivatives

12 The stability of the aldehyde and ketone derivatives during sample work-up and long-term
13 storage is particularly important when working with astromaterials, as sample abundances are

1 generally small and limited, and degradation of the compounds may induce stable isotopic and
2 enantiomeric fractionations. Our experiments revealed a notable drawback of the PFBHA
3 derivatization method; the PFBHA derivatives appear to degrade when the sample solutions are
4 evaporated to dryness, and the decomposition reaction appears to be accelerated when the samples
5 are stored dry at room temperature, as opposed to in low temperature conditions (4°C) (see Figure
6 S1). A decomposition product, identified by the NIST Mass Spectral Library as PFBHA, was
7 observed in the GC-MS chromatograms of dried samples (see Figure S2), suggesting that the
8 aldehyde and ketone derivatives decomposed back to their original constituents (i.e., PFBHA and
9 volatile aldehydes and ketones) during or after the solvent evaporation step. This decomposition
10 reaction, involving hydrolysis of the C-N bond (Kalia and Raines, 2008), could be taking place as
11 a result of a small amount of residual acid in the vials, which becomes difficult to remove as the
12 sample is concentrated. As a preventative measure, we stored the PFBHA derivatives in solution,
13 at low temperatures (<4°C). Several studies report using amber vials and bottles for derivatizing
14 and storing aldehyde and ketone samples, implying that the PFBHA derivatives may be light-
15 sensitive (Glaze et al., 1989; Spaulding and Charles, 2002; Hudson et al., 2007; Serrano et al.,
16 2013). We have ruled out photolysis as the cause of the decomposition reaction observed here by
17 limiting sample exposure to light. The decomposition of the derivatives in dried samples persisted
18 despite carrying out all derivatization reactions, clean-up steps, and storage in amber vials.

19 **4.2. Distribution and abundances of aldehydes and ketones in Murchison**

20 The total yield of aldehydes and ketones reported here is substantially higher (3-fold larger)
21 than total abundances reported for Murchison by previous studies (Table 1; Jungclaus et al., 1976,
22 Pizzarello and Holmes, 2009), likely resulting from increased extraction efficiencies and
23 derivatization yields. This variation between studies may also be partially attributable to sample
24 heterogeneity, as has been reported for other compound classes (Cronin and Pizzarello, 1983;
25 Krishnamurthy et al., 1992; Pizzarello et al., 2003; Aponte et al., 2014). Considering the ubiquitous
26 presence of formaldehyde in the ISM, the abundance of formaldehyde in the Murchison meteorite
27 (19.2 nmol/g) is lower than expected. A relatively low abundance of formaldehyde in Murchison
28 was similarly observed by Pizzarello and Holmes (2009) and Jungclaus et al. (1976) and may be
29 partially attributable to its low boiling point (-19°C), as the Murchison meteorite samples have
30 been stored long-term in room temperature conditions, potentially allowing volatile compounds to

1 be lost. Its low abundance relative to other carbonyl species in Murchison may also be a result of
2 its high reactivity and involvement in abiotic synthetic reactions inside the parent body, including
3 for example the formose reaction; it is possible that formaldehyde originally contained in
4 Murchison may have converted into more complex molecules, including insoluble
5 macromolecules, during secondary processing on the asteroid parent body (Cooper et al., 2001;
6 Cody et al., 2011; Meinert et al., 2016).

7 **4.3. Potential origins for the aldehydes and ketones in Murchison**

8 **4.3.1. Isotopic interpretation of Murchison aldehyde and ketone $\delta^{13}\text{C}$ compositions**

9 The positive $\delta^{13}\text{C}$ values measured here for individual aldehydes and ketones in Murchison
10 (+11.8‰ to +66.4‰) are attributed to an extraterrestrial origin, as terrestrial aldehydes and ketones
11 generally exhibit negative $\delta^{13}\text{C}$ values (−28.3 to −17.0‰; Goldstein and Shaw, 2003). The negative
12 $\delta^{13}\text{C}$ value (−10.0‰) measured for 2-butanone in Murchison is not unusual for meteoritic organic
13 compounds (Sephton and Gilmour, 2001) and does not necessarily indicate terrestrial input; it may
14 reflect a distinct abiotic origin for this type of compound in the parent body asteroid.

15 In order to make direct comparisons between $\delta^{13}\text{C}$ values of different meteoritic compound
16 classes, some assumptions need to be made regarding the original isotopic compositions of the
17 compounds, the alteration history of the asteroid parent body and the associated chemical reactions
18 (Aponte et al., 2017). The carbon isotope compositions of early solar system aldehydes and ketones
19 prior to their incorporation into the asteroid parent body are assumed to have been highly ^{13}C -
20 enriched ($\delta^{13}\text{C} > +117\text{‰}$), based on recent $^{12}\text{C}/^{13}\text{C}$ measurements for interstellar formaldehyde
21 ($\delta^{13}\text{C} = 117\text{-}4933\text{‰}$, $^{12}\text{C}/^{13}\text{C} = 15\text{-}80$; Wirström et al., 2012) and the average $^{12}\text{C}/^{13}\text{C}$ measurement
22 for nearby molecular clouds ($\delta^{13}\text{C} = 117\text{-}392\text{‰}$, $^{12}\text{C}/^{13}\text{C} = 64\text{-}80$; Henkel et al., 1980, 1982). If the
23 aldehydes and ketones detected in the Murchison meteorite are relict, unreacted Strecker synthesis
24 precursors, we would expect them to be relatively enriched in ^{13}C compared to their corresponding
25 α -amino acid products. This prediction assumes that the Strecker synthesis reaction involves
26 significant kinetic isotope fractionations (i.e. that isotopically lighter (^{12}C -enriched) molecules
27 react faster than isotopically heavier (^{13}C -enriched) molecules at a rate that significantly affects
28 the relative $\delta^{13}\text{C}$ values of the products and reactants). This prediction also assumes that the amino
29 acid products have not been isotopically fractionated via subsequent chemical reactions during

1 thermal/aqueous alteration (i.e. that the pool of amino acid products has not been shifted towards
2 a relatively ^{13}C -enriched composition due to the preferential reaction of isotopically lighter amino
3 acids). The isotopic composition of the unreacted carbonyl pool, which would gradually shift
4 towards a more ^{13}C -enriched composition as isotopically light carbonyl molecules are
5 preferentially consumed depends on what proportion of the aldehydes and ketones have been
6 converted into amino acids or vice versa, which remains unknown. Lastly, it is assumed that any
7 variation observed between studies primarily reflects differences in methodologies and that the
8 Murchison meteorite does not exhibit large heterogeneity between subsamples.

9 The $\delta^{13}\text{C}$ measurements for the aldehydes and ketones in Murchison are less ^{13}C -enriched
10 than would be expected for a primordial pool of unreacted Strecker synthesis precursors, assuming
11 that early solar system aldehydes and ketones incorporated into the Murchison parent body asteroid
12 exhibited highly ^{13}C -enriched isotopic signatures and that this signature either remained unaltered
13 until our analyses or was amplified as Strecker synthesis shifts the reactant pool towards a more
14 ^{13}C -enriched composition. The discrepancy between the ^{13}C -isotopic signatures for the carbonyl
15 compounds we found in Murchison and the ^{13}C -enriched values expected from interstellar
16 observations may indicate that all of the early solar system aldehydes and ketones in the asteroid
17 parent body were consumed early on, either via Strecker amino acid synthesis or through alternate
18 chemical reactions, and were later resynthesized from reactions occurring through aqueous and
19 thermal processing inside the asteroid parent body. This is a reasonable possibility as aldehydes
20 and ketones are highly reactive species and are likely to polymerize and react with other chemical
21 species fairly readily in asteroidal conditions (Cleaves, 2008; Cody et al. 2011; Kebukawa and
22 Cody, 2015; Kebukawa et al., 2017). Our observations suggest that the aldehydes and ketones
23 identified in this study were synthesized at a later stage, potentially during aqueous alteration
24 reactions after accretion of the parent body asteroid. A secondary generation of aldehydes and
25 ketones on asteroid parent bodies may imply that these compounds are more readily available in
26 the solar system than previously thought.

27 **4.3.2. Chemical oxidation of insoluble organic matter as a potential source of carbonyl** 28 **compounds in Murchison**

29 Low-temperature aqueous alteration of IOM has been proposed as a potential source of
30 water-soluble organic compounds in carbonaceous chondrites (Cody and Alexander, 2005). The

1 oxidation of IOM in aqueous solution has been shown to convert aliphatic carbon to CO moieties,
2 in addition to organic acids and CO₂ (Cody and Alexander, 2005) and this mechanism may
3 partially explain the contrasting isotopic signatures for the carbonyl compounds in Murchison and
4 those seen in interstellar environments. The more labile, aliphatic moieties of the IOM are expected
5 to be relatively ¹³C-enriched compared to the aromatic portions (Kerridge et al., 1987), as the
6 aromatic components of the IOM are considered to reflect a longer history of chemical processing
7 in the cold ISM and/or during early solar system formation, resulting in progressive ¹³C-depletion
8 over time. Consistent with this theory, the carbonyl δ¹³C compositions reported here (−10.0 to
9 +66.4‰; Figure 4) are generally more ¹³C-enriched in comparison to measurements of IOM-
10 derived aromatics in Murchison (−25‰ to −1‰; Figure 4; Sephton et al., 1998; Sephton and
11 Gilmour, 2001). The Murchison ketone δ¹³C values (−10.0‰ and +11.8‰) are the most ¹³C-
12 depleted of the carbonyl suite, suggesting that they are derived from a more ¹³C-depleted source
13 (e.g. IOM) in comparison to the aldehydes. IOM-derived CO moieties are discussed as primarily
14 representing ketone compounds, rather than aldehydes, based on their solid-state ¹H and ¹³C
15 Nuclear Magnetic Resonance (NMR) spectra (Cody and Alexander, 2005). Likewise, pyrolysis of
16 IOM from the Murchison meteorite has been shown to release several aromatic ketones (Remusat
17 et al., 2005).

18 **4.3.3. Alteration-driven decomposition of soluble organics**

19 Another possible origin for the carbonyl compounds detected in Murchison may be
20 alteration-driven degradation of other soluble organic compound classes (e.g. amino acids, keto
21 acids) in the parent body asteroid. Similarities or trends observed between the compound-specific
22 carbon isotope compositions of the carbonyl compounds in Murchison and other indigenous
23 organics may be indicative of precursor-product synthetic relationships. While these values and
24 data comparisons are not definitive indicators for specific chemical reactions, due to unknowns
25 regarding the original budget of primordial soluble organic compounds and the effects of aqueous
26 alteration on isotopic compositions (Section 4.3), they do provide an informative basis for
27 elucidating the reaction history of aldehydes and ketones in the asteroid parent body.

28 **4.3.3.1. Strecker-cyanohydrin synthesis precursors and products**

1 The $\delta^{13}\text{C}$ values of Strecker α -amino acid and α -hydroxy acid products depend on a
2 combination of: (1) the carbon isotope composition of the precursor carbonyl compound, (2) the
3 carbon isotope composition of the reactant HCN, and (3) the kinetic isotope fractionation effects
4 associated with each step of the reaction (Scheme 1; Elsila et al., 2012; Aponte et al., 2017). The
5 α -carbon and the alkyl (R) groups of the amino acid and hydroxy acid products are derived from
6 the precursor aldehyde or ketone, while the carboxyl carbon is derived from HCN (Scheme 1). The
7 relative influence of the aldehyde or ketone $\delta^{13}\text{C}$ value on the amino acid/hydroxy acid $\delta^{13}\text{C}$ value
8 depends on the number of carbon atoms contained in, and thus contributed from, the R groups. For
9 example, among the Strecker amino acid products, meteoritic HCN ($\delta^{13}\text{C} = +1.4$ to $+7.3\%$;
10 Pizzarello et al., 2004) will have a substantial influence on the glycine isotopic composition and
11 will play an increasingly lesser role on amino acid $\delta^{13}\text{C}$ values as the carbonyl R groups increase
12 in chain length. Comparing the $\delta^{13}\text{C}$ values of carbonyl compounds, amino acids, hydroxy acids
13 and HCN can be used to evaluate a potential Strecker precursor-product relationship between these
14 compounds.

15 In this study, we focus primarily on isotopic comparisons between aldehydes, ketones and
16 their structurally analogous α -amino acids, due to the large data set of stable isotopic measurements
17 previously attained for amino acids in Murchison (Engel et al., 1990; Pizzarello et al., 2004; Elsila
18 et al., 2012). Strecker amino acid products are expected to be relatively depleted in ^{13}C compared
19 to their aldehyde and ketone precursors (Elsila et al., 2012; Aponte et al. 2014, 2017), unless the
20 chemical reactions do not involve significant carbon isotope fractionations, in which case the $\delta^{13}\text{C}$
21 values for the reactants and products would be approximately equal. Amino acids with $\delta^{13}\text{C}$ values
22 that are relatively enriched in ^{13}C compared to their structurally analogous aldehydes and ketones
23 can be interpreted as having been either isotopically fractionated to some extent via subsequent
24 parent body processing-driven chemical reactions following Strecker synthesis, or as having been
25 derived, at least partially, from a carbon source that does not involve Strecker starting materials.




Figure 4. Carbon stable isotope compositions of aldehydes and ketones (circles, this study), their structurally analogous amines (Aponte et al., 2014), amino acids (Engel et al., 1990; Pizzarello et al., 2004; Elsila et al., 2012), monocarboxylic acids (Yuen et al., 1984; Huang et al., 2005), hydroxy acids (Pizzarello et al., 2010) and aromatics derived from macromolecular insoluble organic material (Sephton et al., 1998; Sephton and Gilmour, 2001) in the Murchison meteorite. Individual (*E*) and (*Z*) isomers of derivatized aldehydes and ketones are labeled where applicable. Upper indices assigned to individual amines, carboxylic acids and hydroxy acids in the legend denote the corresponding data points for each compound in the figure.

1 The carbon isotope compositions of the Murchison aldehydes and ketones (−10.0‰ to
 2 +66.4‰) and their structurally-analogous Strecker amino acids (+5.0‰ to +51.7‰) fall within a
 3 similar wide range of values (Figure 4). The aldehyde $\delta^{13}\text{C}$ values are generally more ^{13}C -enriched
 4 than their corresponding Strecker products; the formaldehyde, propionaldehyde, and
 5 butyraldehyde values are relatively enriched in ^{13}C compared to their structurally analogous amino
 6 acids, and all four aldehyde $\delta^{13}\text{C}$ values are more ^{13}C -enriched compared to the structurally
 7 analogous hydroxy acid $\delta^{13}\text{C}$ values (Figure 4). This relative ^{13}C -enrichment observed for the
 8 aldehydes relative to Strecker products suggests that a precursor-product relationship could relate
 9 these compound classes in Murchison, as kinetic isotope effects involved in the reaction would
 10 produce isotopically light (^{13}C -depleted) amino acid and hydroxy acid products, relative to the
 11 carbonyl reactants. In contrast, the ketone $\delta^{13}\text{C}$ values are ^{13}C -depleted relative to both their
 12 structurally-analogous amino acids and hydroxy acids, implying that other factors likely
 13 influenced the isotopic compositions of Murchison ketones. We consider the possibility that the

1 aldehydes, ketones and amino acids may be intimately connected, not solely by Strecker synthesis,
2 but also by decomposition of the amino acids during aqueous alteration on the parent body asteroid.
3 An episodic decomposition or dissolution of α -amino acids in between intermittent Strecker
4 synthesis could explain the similar $\delta^{13}\text{C}$ values amongst the aldehydes, ketones and amino acids,
5 although this is currently only speculative, as the temperatures experienced during aqueous
6 alteration on the Murchison asteroid are arguably insufficient for amino acid degradation. Aqueous
7 solutions of α -amino acids can undergo decomposition at elevated temperatures, yielding aldehyde
8 and ketone products (Schonberg and Moubacher, 1951); however, these reactions generally take
9 place at temperatures $> 220^\circ\text{C}$ (Lien and Nawar, 1974) and temperature estimates for aqueous
10 alteration on the Murchison parent body asteroid are much lower, ranging from $0\text{-}80^\circ\text{C}$ (Clayton
11 and Mayeda, 1999; Baker et al., 2002; Guo and Eiler, 2007). Whether or not long-duration (several
12 million years; Krot et al., 2015) aqueous alteration, catalyzed by mineral interactions, would be
13 sufficient to induce amino acid degradation at these low temperatures has not been evaluated
14 experimentally; however, laboratory simulations of aqueous alteration in the presence of mineral
15 surfaces at high temperatures ($>156^\circ\text{C}$) have shown that amino acid degradation via
16 decarboxylation and deamination is greatly accelerated under these conditions (McCollom, 2013).

17 In addition to possible decomposition of water-soluble amino acids, long-duration parent
18 body aqueous alteration could also result in the release of carbonyl compounds from the insoluble
19 organic matter (Cody and Alexander, 2005). The relative depletion in ^{13}C observed for the ketones
20 in Murchison may reflect this secondary source of carbonyl compounds (Section 4.3.2; Figure 4).

21 **4.3.3.2. $\delta^{13}\text{C}$ comparisons with keto acids, amines, and carboxylic acids**

22 The similar $\delta^{13}\text{C}$ compositions of aldehydes, ketones and amino acids in Murchison could
23 also indicate that they are all partially derived from a common source of soluble organic carbon.
24 For example, some of the aldehydes, ketones and amino acids could be linked by reactions
25 involving keto acids; aldehydes and ketones could be derived from degradation of keto acids
26 during aqueous alteration, while the amino acids may be synthesized from the α -keto acids via
27 reductive amination (Huber and Wächtershäuser, 2003). For instance, thermal decomposition of
28 pyruvic acid yields acetaldehyde (Taylor, 1987), while reductive amination of pyruvic acid yields
29 its structurally analogous amino acid, alanine (Yanagawa et al., 1982). Keto acids, including
30 pyruvic acid, acetoacetic acid, oxaloacetic acid, and citric acid, have been detected in carbonaceous

1 chondrites (Cooper et al., 2005; Cooper et al., 2011); however, their stable isotopic compositions
2 have not yet been measured, limiting our ability to investigate this theory. Furthermore, these types
3 of chemical reactions have only been described for terrestrial samples and have not yet been
4 evaluated for cosmochemical conditions.

5 Amines and carboxylic acids are also potential structural relatives of carbonyl compounds
6 as they can be synthesized via reductive amination of aldehydes and ketones and oxidation of
7 aldehydes, respectively (Figure 1). The amine $\delta^{13}\text{C}$ values reported by Aponte et al. (2014) are
8 generally much more ^{13}C -enriched than the aldehydes and ketones reported here, suggesting that
9 the amines in Murchison were not produced via reductive amination of the aldehydes and ketones
10 identified. Murchison carboxylic acid $\delta^{13}\text{C}$ values, on the other hand, are ^{13}C -depleted relative to
11 the aldehyde values (Figure 4), consistent with synthesis via either oxidation of the aldehydes
12 (Corey et al., 1968) or oxidation of the macromolecular organic matter in Murchison (Oba and
13 Naraoka, 2006).

14 5. CONCLUSIONS

15 The aldehyde and ketone content of the Murchison meteorite and their carbon isotope
16 compositions provide a snapshot of the post-alteration organic chemistry of the parent body
17 asteroid, as well as some important information about the history of the asteroid and the synthetic
18 reactions involved. Our carbon isotope analyses of aldehydes and ketones in Murchison revealed
19 a distinct composition, dominated by acetaldehyde, propionaldehyde, and acetone, with carbon
20 isotopic compositions covering a range of ^{13}C -depleted values (-10.0‰ to $+66.4\text{‰}$), similar to
21 other previously analyzed soluble organics in Murchison. This is in contrast to the formaldehyde-
22 dominated, highly ^{13}C -enriched, carbonyl content of the interstellar medium. This large disparity
23 in composition between the two environments suggests that the organic chemistry and alteration
24 history of the parent body asteroid may play a major role in generating a secondary source of
25 carbonyl compounds in the solar system. Given the long history of the Murchison parent body
26 asteroid and the complexity of its organic content, this compilation of data demonstrates that the
27 aldehydes, ketones, and other compound classes are interconnected by a dynamic series of
28 chemical reactions.

29

ACKNOWLEDGEMENTS

1
2
3
4
5
6
7
8
9
10

D.N.S. and R.W.H. would like to thank Drs. Matthew Ross and Aaron Skelhorne, MacEwan University, for their assistance with the GC-MS analyses during method optimization. J.C.A. and J.E.E. acknowledge support from NASA's Emerging Worlds program for this work, as well as from the NASA Astrobiology Institute and the Goddard Center for Astrobiology, and a grant from the Simons Foundation (SCOL award 302497). Funding for this study was also provided by Natural Sciences and Engineering Research Council (NSERC) Grant 261740 to C.D.K.H., and NSERC CGS D3 and Izaak Walton Killam Memorial scholarships to D.N.S. Two anonymous reviewers are thanked for their careful review of this manuscript.

REFERENCES

- 1
- 2 Abdel-Magid A. F., K. G. Carson, B. D. Harris, C. A. Maryanoff, and R. D. Shah (1996) Reductive
3 amination of aldehydes and ketones with sodium triacetoxyborohydride. Studies on direct
4 and indirect reductive amination procedures. *Journal of Organic Chemistry* **61**, 3849-3862.
- 5 Allamandola, L. J., S. A. Sandford, and G. J. Valero (1988) Photochemical and thermal evolution
6 of interstellar/precometary ice analogs. *Icarus* **76**, 225-252.
- 7 Aponte, J. C., J. P. Dworkin, and J. E. Elsila (2014) Assessing the origins of aliphatic amines in
8 the Murchison meteorite from their compound-specific carbon isotopic ratios and
9 enantiomeric composition. *Geochimica et Cosmochimica Acta* **141**, 331-345.
- 10 Aponte, J. C., J. E. Elsila, D. P. Glavin, S. N. Milam, S. B. Charnley, and J. P. Dworkin (2017)
11 Pathways to meteoritic glycine and methylamine. *ACS Earth and Space Chemistry*, DOI:
12 10.1021/acsearthspacechem.6b00014
- 13 Awad, Z., T. Chigai, Y. Kimura, O. M. Shalabiea, and T. Yamamoto (2005) New rate constants
14 of hydrogenation of CO on H₂O-CO ice surfaces. *The Astrophysical Journal* **626**, 262-271.
- 15 Baker, L., I. A. Franchi, I. P. Wright, and C. T. Pillinger (2002) The oxygen isotopic composition
16 of water from Tagish Lake: Its relationship to low-temperature phases and to other
17 carbonaceous chondrites. *Meteoritics & Planetary Science* **37**, 977-985.
- 18 Bisschop, S. E., G. W. Fuchs, E. F. van Dishoeck, and H. Linnartz (2007) H-atom bombardment
19 of CO₂, HCOOH, and CH₃CHO containing ices. *Astronomy & Astrophysics* **474**, 1061-
20 1071.
- 21 Bossa, J.-B. F. Duvernay, P. Theulé, F. Borget, L. d'Hendecourt, T. Chiavassa (2009)
22 Methylammonium methylcarbamate thermal formation in interstellar ice analogs: a glycine
23 salt precursor in protostellar environments. *Astronomy & Astrophysics* **506**, 601-608.
- 24 Botta, O. and J. L. Bada (2002) Extraterrestrial organic compounds in meteorites. *Surveys in*
25 *Geophysics* **23**, 411-467.
- 26 Burk M. J., J. P. Martinez, J. E. Feaster, and N. Cosford (1994) Catalytic asymmetric reductive
27 amination of ketones via highly enantioselective hydrogenation of the C=N double bond.
28 *Tetrahedron* **50**, 4399-4428.

- 1 Cancho, B., F. Ventura and M. T. Galceran (2001) Determination of aldehydes in drinking water
2 using pentafluorobenzylhydroxylamine derivatization and solid-phase microextraction.
3 *Journal of Chromatography* **943**, 1-13.
- 4 Charnley, S. B., P. Ehrenfreund, T. J. Millar, A. C. A. Boogert, A. J. Markwick, H. M. Butner, R.
5 Ruitkamp, and S. D. Rodgers (2004) Observational tests for grain chemistry: Posterior
6 Isotopic labelling. *Monthly Notices of the Royal Astronomical Society* **347**, 157-162.
- 7 Clayton, R. N., and T. K. Mayeda (1999) Oxygen isotope studies in carbonaceous chondrites.
8 *Geochimica et Cosmochimica Acta* **63**, 2089-2104.
- 9 Cleaves, H. J. (2008) The prebiotic geochemistry of formaldehyde. *Precambrian Research*, **164**,
10 111-118.
- 11 Cody, G. D., and C. M. O'D. Alexander (2005) NMR studies of chemical structural variation of
12 insoluble organic matter from different carbonaceous chondrite groups. *Geochimica et*
13 *Cosmochimica Acta* **69**, 1085-1097.
- 14 Cody, G. D., E. Heying, C. M. O'D. Alexander, L. R. Nittler, A. L. D. Kilcoyne, S. A. Sandford,
15 and R. M. Stroud (2011) Establishing a molecular relationship between chondritic and
16 cometary organic solids. *PNAS* **108**, 19171-19176.
- 17 Cooper, G., N., N. Kimmich, W. Belisle, J. Sarinana, K. Brabham, and L. Garrel (2001)
18 Carbonaceous meteorites as a source of sugar-related organic compounds for the early
19 Earth. *Nature* **414**, 879-883.
- 20 Cooper, G., A. Dugas, A. Byrd, P. M. Chang and N. Washington (2005) Keto-acids in
21 carbonaceous meteorites. *36th Lunar and Planetary Science Conference*, #2381.
- 22 Cooper, G., C. Reed, D. Nguyen, M. Carter, and Y. Wang (2011) Detection and formation scenario
23 of citric acid, pyruvic acid, and other possible metabolism precursors in carbonaceous
24 meteorites. *PNAS* **108**, 14015-14020.
- 25 Corey, E. J., N. W. Gilman, and B. E. Ganem (1968) New methods for the oxidation of aldehydes
26 to carboxylic acids and esters. *Journal of the American Chemical Society* **90**, 5616-5617.
- 27 Cronin, J. R. and S. Pizzarello (1983) Amino acids in meteorites. *Advances in Space Research* **3**,
28 5-18.
- 29 Deamer, D., J. P. Dworkin, S. A. Sandford, M. P. Bernstein, and L. J. Allamandola (2002) The
30 first cell membranes. *Astrobiology* **2**, 371-381.

1 de Marcellus, P., C. Meinert, I. Myrgorodska, L. Nahon, T. Buhse, L. Le Sergeant D'Hendecourt,
2 and U. J. Meierhenrich (2015) Aldehydes and sugars from evolved precometary ice
3 analogs: Importance of ices in astrochemical and prebiotic evolution. *PNAS* **112**, 965-970.

4 Docherty, G., V. Jones, and R. P. Evershed (2001) Practical and theoretical considerations in the
5 gas chromatography/combustion/isotope ratio mass spectrometry $\delta^{13}\text{C}$ analysis of small
6 polyfunctional compounds. *Rapid Communications in Mass Spectrometry* **15**, 730-738.

7 Elsila, J. E., S. B. Charnley, A. S. Burton, D. P. Glavin, and J. P. Dworkin (2012) Compound-
8 specific carbon, nitrogen, and hydrogen isotopic ratios for amino acids in CM and CR
9 chondrites and their use in evaluating potential formation pathways. *Meteoritics &*
10 *Planetary Science* **47**, 1517-1536.

11 Engel, M. H., S. A. Macko, and J. A. Silfer (1990) Carbon isotope composition of individual amino
12 acids in the Murchison meteorite. *Nature* **348**, 47-49.

13 EPA Method 556: Determination of carbonyl compounds in drinking water by pentafluorobenzyl-
14 hydroxylamine derivatization and capillary gas chromatography with electron capture
15 detection (1998) National exposure research laboratory office of research and
16 development. US Environmental Protection Agency, Cincinnati, Ohio, 45268.

17 Glaze, W. H., M. Koga and D. Cancilla (1989) Ozonation byproducts. 2. Improvement of an
18 aqueous-phase derivatization method for the detection of formaldehyde and other carbonyl
19 compounds formed by the ozonation of drinking water. *Environmental Science &*
20 *Technology* **23**, 838-847.

21 Goldstein, A. H. and S. L. Shaw (2003) Isotopes of volatile organic compounds: An emerging
22 approach for studying budgets and chemistry. *Chemical Reviews* **103**, 5025-5048.

23 Guo, W., and J. M. Eiler (2007) Temperatures of aqueous alteration and evidence for methane
24 generation on the parent bodies of the CM chondrites. *Geochimica et Cosmochimica Acta*
25 **71**, 5565-5575.

26 Henkel, C., T. L. Wilson, and J. Biegging (1982) Further ($^{12}\text{C}/^{13}\text{C}$) ratios from formaldehyde: A
27 variation with distance from the galactic center. *Astronomy & Astrophysics* **109**, 344-351.

28 Holtom, P. D., C. J. Bennett, Y. Osamura, N. J. Mason, and R. I. Kaiser (2005). A combined
29 experimental and theoretical study on the formation of the amino acid glycine
30 ($\text{NH}_2\text{CH}_2\text{COOH}$) and its isomer (CH_3NHCOOH) in extraterrestrial ices. *The Astrophysical*
31 *Journal* **626**, 940.

- 1 Huang, Y., Y. Wang, M. R. Alexandre, T. Lee, C. Rose-Petruck, M. Fuller, and S. Pizzarello
2 (2005) Molecular and compound-specific isotopic characterization of monocarboxylic
3 acids in carbonaceous chondrites. *Geochimica et Cosmochimica Acta* **69**, 1073-1084.
- 4 Huang, Y., J. C. Aponte, J. Zhao, R. Tarozo, and C. Hallmann (2015) Hydrogen and carbon
5 isotopic ratios of polycyclic aromatic compounds in two CM2 carbonaceous chondrites
6 and implications for prebiotic synthesis. *Earth and Planetary Science Letters* **426**, 101-
7 108.
- 8 Huber, C., and G. Wächtershäuser (2003) Primordial reductive amination revisited. *Tetrahedron*
9 *Letters* **44**, 1695-1697.
- 10 Hudson, E. D., K. Okuda, and P. A. Ariya (2007) Determination of acetone in seawater using
11 derivatization solid-phase microextraction. *Analytical and Bioanalytical Chemistry* **388**,
12 1275-1282.
- 13 Hudson, R. L., Lewis, A. S., Moore, M. H., Dworkin, J. P., and M. P. Martin (2009) Enigmatic
14 isovaline: investigating the stability, racemization and formation of a non-biological
15 meteoritic amino acid. *ASP Conference Series* **420**, 157-162.
- 16 Irvine (1999) The composition of interstellar molecular clouds. *Space Science Reviews* **90**, 203-
17 218.
- 18 Jungclaus, G. A., G. U. Yuen, C. B. Moore, J. G. Lawless (1976) Evidence for the presence of low
19 molecular weight alcohols and carbonyl compounds in the Murchison meteorite.
20 *Meteoritics* **11**, 231-237.
- 21 Kalia, J. and R. T. Raines (2008) Hydrolytic stability of hydrazones and oximes. *Angewandte*
22 *Chemie* **47**, 7523-7526.
- 23 Kebukawa, Y., and G. D. Cody (2015) A kinetic study of the formation of organic solids from
24 formaldehyde: Implications for the origin of extraterrestrial organic solids in primitive
25 Solar System objects. *Icarus* **248**, 412-423.
- 26 Kebukawa, Y., Q. H. S. Chan, S. Tachibana, K. Kobayashi, and M. E. Zolensky (2017) One-pot
27 synthesis of amino acid precursors with insoluble organic matter in planetesimals with
28 aqueous activity. *Science Advances* **3**, 1-7.
- 29 Kobayashi, K., M. Tanaka, and S. Kawai (1980) Gas chromatographic determination of low-
30 molecular-weight carbonyl compounds in aqueous solution as their O-(2,3,4,5,6-
31 pentafluorobenzyl) oximes. *Journal of Chromatography* **187**, 413-417.

- 1 Krishnamurthy, R. V., S. Epstein, J. R. Cronin, S. Pizzarello, and G. U. Yuen (1992) Isotopic and
2 molecular analyses of hydrocarbons and monocarboxylic acids of the Murchison meteorite.
3 *Geochimica et Cosmochimica Acta* **56**, 4045-4058.
- 4 Krot, A. N., K. Nagashima, C. M. O'D Alexander, F. J. Ciesla, W. Fujiya, and L. Bonal (2015)
5 Sources of water and aqueous activity on the chondrite parent asteroids. In *Asteroids IV*, P.
6 Michel et al., Eds. University of Arizona Press, Tuscon, 635-660.
- 7 Kvenvolden, K., J. Lawless, K. Pering, E. Peterson, J. Flores, C. Ponnampereuma, I. R. Kaplan, C.
8 Moore (1970) Evidence for extraterrestrial amino-acids and hydrocarbons in the
9 Murchison meteorite. *Nature* **228**, 923-926.
- 10 Lee C.-W., J.-K. Kim, E.-S. Moon, Y. C. Minh, and H. Kang (2009) Formation of glycine on
11 ultraviolet-irradiated interstellar ice-analog films and implications for interstellar amino
12 acids. *The Astrophysical Journal* **697**, 428-435.
- 13 Lien, Y. C. and W. W. Nawar (1974) Thermal degradation of some amino acids. *Journal of Food*
14 *Science* **39**, 911-913.
- 15 McCollom, T. M. (2013) The influence of minerals on decomposition of the n-alkyl- α -amino acid
16 norvaline under hydrothermal conditions. *Geochimica et Cosmochimica Acta* **104**, 330-
17 357.
- 18 Meinert, C., I. Myrgorodska, P. de Marcellus, T. Buhse, L. Nahon, S. V. Hoffmann, L. Le Sergeant
19 d'Hendecourt, U. J. Meierhenrich (2016) Ribose and related sugars from ultraviolet
20 irradiation of interstellar ice analogs. *Science* **352**, 208-212.
- 21 Miller, S. (1957) The mechanism of synthesis of amino acids by electric discharges. *Biochimica*
22 *et Biophysica Acta* **23**, 480-489.
- 23 Monroe, A. A. and S. Pizzarello (2011) The soluble organic compounds of the Bells meteorite:
24 Not a unique or unusual composition. *Geochimica et Cosmochimica Acta* **75**, 7585-7595.
- 25 Mumma, M. J., and S. B. Charnley (2011) The chemical composition of comets. *Annual Review*
26 *of Astronomy and Astrophysics* **49**, 471-524.
- 27 Oba, Y. and H. Naraoka (2006) Carbon isotopic composition of acetic acid generated by hydrous
28 pyrolysis of macromolecular organic matter from the Murchison meteorite. *Meteoritics &*
29 *Planetary Science* **41** (8), 1175-1181.
- 30 Peltzer, E. T. and J. L. Bada (1978) A-Hydroxycarboxylic acids in the Murchison meteorite.
31 *Nature* **272**, 443-444.

- 1 Peltzer, E. T., J. L. Bada, G. Schlesinger, and S. L. Miller (1984) The chemical conditions on the
2 parent body of the Murchison meteorite: Some conclusions based on amino, hydroxy and
3 dicarboxylic acids. *Advances in Space Research* **4**, 69–74.
- 4 Pizzarello, S., M. Zolensky, and K. A. Turk (2003) Non racemic isovaline in the Murchison
5 meteorite: chiral distribution and mineral association. *Geochimica et Cosmochimica Acta*
6 **67**, 1589-1595.
- 7 Pizzarello, S., Y. S. Huang, and M. Fuller (2004) The carbon isotopic distribution of Murchison
8 amino acids. *Geochimica et Cosmochimica Acta* **68**, 4963-4969.
- 9 Pizzarello, S., G. W. Cooper, G. J. Flynn (2006) The nature and distribution of the organic material
10 in carbonaceous chondrites and interplanetary dust particles. *Meteorites and the Early*
11 *Solar System II*, D. S. Lauretta and H. Y. McSween, Eds. (University of Arizona Press,
12 2006), p. 625-650.
- 13 Pizzarello, S. and W. Holmes (2009) Nitrogen-containing compounds in two CR2 meteorites: ¹⁵N
14 composition, molecular distribution and precursor molecules. *Geochimica et*
15 *Cosmochimica Acta* **73**, 2150-2162.
- 16 Pizzarello, S., Y. Wang, and G. M. Chaban (2010) A comparative study of the hydroxy acids from
17 the Murchison, GRA 95229 and LAP 02342 meteorites. *Geochimica et Cosmochimica*
18 *Acta* **74**, 6206-6217.
- 19 Pizzarello, S., D. L. Schrader, A. A. Monroe, and D. S. Lauretta (2012) Large enantiomeric
20 excesses in primitive meteorites and the diverse effects of water in cosmochemical
21 evolution. *Proceedings of the National Academy of Sciences* **109**, 11949-11954.
- 22 Remusat, L., S. Derenne, F. Robert, and H. Knicker (2005) New pyrolytic and spectroscopic data
23 on Orgueil and Murchison insoluble organic matter: A different origin than soluble?
24 *Geochimica et Cosmochimica Acta* **69**, 3919-3932.
- 25 Rodigast, M., A. Mutzel, Y. Iinuma, S. Haferkorn, and H. Herrmann (2015) Characterisation and
26 optimization of a sample preparation method for the detection and quantification of
27 atmospherically relevant carbonyl compounds in aqueous medium. *Atmospheric*
28 *Measurement Technique* **8**, 2409-2416.
- 29 Schonberg, A. and R. Moubacher (1951) The Strecker degradation of α -amino acids. *Chemical*
30 *Reviews* **50**, 261-277.

- 1 Schutte, W. A., P. A. Gerakines, T. R. Geballe, E. F. vanDishoeck, and J. M. Greenberg (1996)
2 Discovery of solid formaldehyde toward the protostar GL 2136: Observations and
3 laboratory simulation. *Astronomy & Astrophysics* **309**, 633-647.
- 4 Seaman, V. Y., M. J. Charles, and T. M. Cahill (2006) A sensitive method for the quantification
5 of acrolein and other volatile carbonyls in ambient air. *Analytical Chemistry* **78**, 2405-
6 2412.
- 7 Sephton, M. A. (2002) Organic compounds in carbonaceous meteorites. *Natural Product Reports*
8 **19**, 292-311.
- 9 Sephton, M. A. (2014) Organic geochemistry of meteorites. *Treatise on Geochemistry 2nd Edition*,
10 H.D. Holland and K. K. Turekian, Eds. (Elsevier, 2014) vol. 12, p. 1-31.
- 11 Serrano, M., M. Silva, and M. Gallego (2013) Development of an environment-friendly
12 microextraction method for the determination of aliphatic and aromatic aldehydes in water.
13 *Analytica Chimica Acta* **784**, 77-84.
- 14 Shalabiea, O. M., and J. M. Greenberg (1994) Two key processes in dust/gas chemical modeling
15 – photoprocessing of grain mantles and explosive desorption. *Astronomy & Astrophysics*
16 **290**, 266-278.
- 17 Spaulding, R. S., and M. J. Charles (2002) Comparison of methods for extraction, storage, and
18 silylation of pentafluorobenzyl derivatives of carbonyl compounds and multi-functional
19 carbonyl compounds. *Analytical and Bioanalytical Chemistry* **372**, 808-816.
- 20 Taylor, R. (1987) The mechanism of thermal eliminations Part XXIII: [1] The thermal
21 decomposition of pyruvic acid. *International Journal of Chemical Kinetics* **19**, 709-713.
- 22 Tielens, A. G. G. M. and D. C. B. Whittet (1997) Ice in star forming regions. *Molecules in*
23 *Astrophysics: Probe and Processes*, E. F. van Dishoeck, Ed. (Springer Science & Business
24 Media), 45-60.
- 25 Watanabe, N. and A. Kouchi (2002) Efficient formation of formaldehyde and methanol by the
26 addition of hydrogen atoms to CO in H₂O-CO ice at 10 K. *The Astrophysical Journal* **571**,
27 173-176.
- 28 Wirström, E. S., S. B. Charnley, W. D. Geppert, and C. M. Persson (2012) Observations of carbon
29 isotopic fractionation in interstellar formaldehyde. *43rd Lunar and Planetary Science*
30 *Conference*, #1611.

1 Yamada, H., and I. Somiya (1989) The determination of carbonyl compounds in ozonated water
2 by the PFBOA method. *Ozone Science & Engineering* **11**, 127-141.

3 Yanagawa, H., Y. Makino, K. Sato, M. Nishizawa, and F. Egami (1982) Novel formation of α -
4 amino acids and their derivatives from oxo acids and ammonia in an aqueous medium. *The*
5 *Journal of Biochemistry* **91**, 2087-2090.

6 Yuen, G., N. Blair, D. J. Des Marais, and S. Chang (1984) Carbon isotope composition of low
7 molecular weight hydrocarbons and monocarboxylic acids from Murchison meteorite.
8 *Nature* **307**, 252-254.

9
10
11
12

SUPPLEMENTARY MATERIAL

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15

EPA Method #556

The EPA Method #556 (Scheme 2), used previously for the analysis of the Murchison meteorite (Pizzarello and Holmes, 2009), is carried out as follows: carbonyl compounds are derivatized in water (pH = 4) by adding 1 mL of a 15 mg/mL *O*-(2,3,4,5,6-Pentafluorobenzyl) hydroxylamine hydrochloride (PFBHA) solution and heating the solution at 35°C for 2 hours. Following the derivatization step, 2-4 drops of concentrated H₂SO₄ are added to the sample to prevent extraction of excess PFBHA reagent. The PFBHA derivatives are extracted from solution with hexane and transferred to a vial containing 0.2 N H₂SO₄, as an acid-wash step. The hexane layer, containing the PFBHA derivatives, is then isolated, concentrated to a small volume via evaporation, and analyzed via GC-MS.

Table S1. A summary, adapted from Rodigast et al. (2015), of the experimental conditions used for aqueous phase derivatization and extraction of aldehydes and/or ketones.

PFBHA (mg/mL)	Derivatization time	Derivatization temperature	Extraction solvent	Extraction time (min)	Reference
0.1	40 min aldehydes, 24 h ketones	Room temp.	Ethyl acetate	–	Kobayashi et al. (1980)
0.1	2 h (longer for ketones)	Room temp.	Hexane	0.5	Glaze et al. (1989)
1	1 h aldehydes, 1 day ketones	Room temp.	Hexane	–	Yamada and Somiya (1989)
>10-fold excess	24 h	Room temp.	Hexane, MTBE	1	Le Lacheur et al. (1993)
0.75	2 h	35°C	Hexane	3	EPA Method 556 (1998)
0.8	20 s	900 W	Toluene	–	Strassnig et al. (2000)
0.5	24 h	Room temp.	Dichloromethane	–	Spaulding and Charles (2002)
0.06	4 h	60°C	–	–	Sugaya et al. (2004)
0.2	24-96 h	Room temp.	Hexane	–	Seaman et al. (2006)
0.06	2 h	Room temp.	–	–	Hudson et al. (2007)
–	4 h	Room temp.	–	–	Takeuchi et al. (2007)
0.75	10 min	Room temp.	–	–	Saison et al. (2009)
<i>Followed EPA Method 556</i>					Pizzarello and Holmes (2009)
0.05	<10 min	Room temp.	Chlorobenzene	2	Ye et al. (2011)
0.5	1 min	60°C	Hexane	1	Serrano et al. (2013)
–	45 min	–	Dichloromethane	–	de Marcellus et al. (2015)
0.4	24 h	–	Dichloromethane	30	Rodigast et al. (2015)

Dash line – parameter not reported




Figure S1. Abundances (GC-MS peak areas) of aldehyde and ketone derivatives (initial concentrations: 50 $\mu\text{g/mL}$) before and after storage at 4°C or room temperature. Dichloromethane was used as the extraction solvent for all samples. Solutions were evaporated to 1 mL volumes under a stream of nitrogen prior to the first GC-MS run (Day 0). Samples stored in solution were stored in their 1 mL volumes for 10 days. Samples stored dry were evaporated to dryness under a stream of nitrogen, immediately re-capped, stored dry for 10 days, and then re-constituted in 1 mL dichloromethane prior to the second GC-MS run (Day 10). All measurements were calibrated against an external standard (50 $\mu\text{g/mL}$ underivatized acetophenone) to adjust for instrumental variations. Each bar represents an average of 3 replicate samples, and error bars represent one standard deviation from the mean.




Figure S2. GC-MS chromatogram of aldehyde and ketone derivatives in dichloromethane after dry storage at 4°C for 10 days (initial concentrations: 50 µg/mL). The identities of the peaks are presented in Table 1. The GC-MS peak labeled with an asterisk is a decomposition product that was only observed in samples that had been stored dry. The mass spectrum displayed above the chromatogram represents the mass spectrum and compound identification provided by the NIST Mass Spectral Library for the decomposition product(*).

1

2

SUPPLEMENTARY REFERENCES

- 1
2
- 3 de Marcellus, P., C. Meinert, I. Myrgorodska, L. Nahon, T. Buhse, L. Le Sergeant D'Hendecourt,
4 and U. J. Meierhenrich (2015) Aldehydes and sugars from evolved precometary ice
5 analogs: Importance of ices in astrochemical and prebiotic evolution. *PNAS* **112**, 965-970.
- 6 EPA Method 556: Determination of carbonyl compounds in drinking water by pentafluorobenzyl-
7 hydroxylamine derivatization and capillary gas chromatography with electron capture
8 detection (1998) National exposure research laboratory office of research and
9 development. US Environmental Protection Agency, Cincinnati, Ohio, 45268.
- 10 Glaze, W. H., M. Koga and D. Cancilla (1989) Ozonation byproducts. 2. Improvement of an
11 aqueous-phase derivatization method for the detection of formaldehyde and other carbonyl
12 compounds formed by the ozonation of drinking water. *Environmental Science &*
13 *Technology* **23**, 838-847.
- 14 Hudson, E. D., K. Okuda, and P. A. Ariya (2007) Determination of acetone in seawater using
15 derivatization solid-phase microextraction. *Analytical and Bioanalytical Chemistry* **388**,
16 1275-1282.
- 17 Kobayashi, K., M. Tanaka, and S. Kawai (1980) Gas chromatographic determination of low-
18 molecular-weight carbonyl compounds in aqueous solution as their O-(2,3,4,5,6-
19 pentafluorobenzyl) oximes. *Journal of Chromatography* **187**, 413-417.
- 20 Le Lacheur, R. M., L. B. Sonnenberg, P. C. Singer, R. F. Christman, and M. J. Charles (1993)
21 Identification of carbonyl compounds in environmental samples. *Environmental Science*
22 *& Technology* **27**, 2745-2753.
- 23 Pizzarello, S. and W. Holmes (2009) Nitrogen-containing compounds in two CR2 meteorites: ¹⁵N
24 composition, molecular distribution and precursor molecules. *Geochimica et*
25 *Cosmochimica Acta* **73**, 2150-2162.
- 26 Rodigast, M., A. Mutzel, Y. Inuma, S. Haferkorn, and H. Herrmann (2015) Characterisation and
27 optimization of a sample preparation method for the detection and quantification of
28 atmospherically relevant carbonyl compounds in aqueous medium. *Atmospheric*
29 *Measurement Technique* **8**, 2409-2416.
- 30 Saison, D., D. P. De Schutter, F. DeIvaux, F. R. Delvaux (2009) Determination of carbonyl
31 compounds in beer by derivatization and headspace solid-phase microextraction in

1 combination with gas chromatography and mass spectrometry. *Journal of*
2 *Chromatography A* **1216**, 5061-5068. Seaman, V. Y., M. J. Charles, and T. M. Cahill
3 (2006) A sensitive method for the quantification of acrolein and other volatile carbonyls in
4 ambient air. *Analytical Chemistry* **78**, 2405-2412.

5 Serrano, M., M. Silva, and M. Gallego (2013) Development of an environment-friendly
6 microextraction method for the determination of aliphatic and aromatic aldehydes in water.
7 *Analytica Chimica Acta* **784**, 77-84.

8 Spaulding, R. S., and M. J. Charles (2002) Comparison of methods for extraction, storage, and
9 silylation of pentafluorobenzyl derivatives of carbonyl compounds and multi-functional
10 carbonyl compounds. *Analytical and Bioanalytical Chemistry* **372**, 808-816.

11 Strassnig, S., T. Wenzl, and E. P. Lankmayr (2000) Microwave-assisted derivatization of volatile
12 carbonyl compounds with *O*-(2,3,4,5,6-pentafluorobenzyl)hydroxylamine. *Journal of*
13 *Chromatography* **891**, 267-273.

14 Sugaya, N., K. Sakurai, T. Nakagawa, N. Onda, S. Onodera, M. Morita, and M. Tezuka (2004)
15 Development of a headspace GC-MS analysis of carbonyl compounds (aldehydes and
16 ketones) in household products after derivatization with *O*-(2,3,4,5,6-pentafluorobenzyl)-
17 hydroxylamine. *Analytical Sciences* **20**, 865-870.

18 Takeuchi, A., T. Takigawa, M. Abe, T. Kawai, Y. Endo, T. Yasugi, G. Endo, and K. Ogino (2007)
19 Determination of formaldehyde in urine by headspace gas chromatography. *Bulletin of*
20 *Environmental Contamination and Toxicology* **79**, 1-4.

21 Yabuta, H., G. D. Cody, and C. M. O'D. Alexander (2007) Chondritic organic matter as an
22 indicator of nebular and parent body processing: Extended pyrolysis studies for CM, CI,
23 CR, CO, CV, ordinary and Tagish Lake group meteorites. *38th Lunar and Planetary*
24 *Science Conference* contribution #2304.

25 Yamada, H., and I. Somiya (1989) The determination of carbonyl compounds in ozonated water
26 by the PFBOA method. *Ozone Science & Engineering* **11**, 127-141.

27 Ye, Q., D. Zheng, L. Liu, and L. Hong (2011) Rapid analysis of aldehydes by simultaneous
28 microextraction and derivatization followed by GC-MS. *Journal of Separation Science* **34**,
29 1607-1612.

30
31