

1 **The Paganica fault and surface co-seismic ruptures caused by the April 6, 2009 earthquake**
2 **(L'Aquila, central Italy)**

3

4 E. Falcucci (1,2), S. Gori (1,4), E. Peronace (3), G. Fubelli (4), M. Moro (5), M. Saroli (6), B.
5 Giaccio (7), P. Messina (7), G. Naso (3), G. Scardia (1), A. Sposato (7), M. Voltaggio (7), P. Galli
6 (3), F. Galadini (1).

7

8

9 1 Istituto Nazionale di Geofisica e Vulcanologia, sezione Milano-Pavia

10 2 Dipartimento di Scienze della Terra, Università degli Studi "La Sapienza", Roma

11 3 Dipartimento della Protezione Civile Nazionale

12 4 Dipartimento di Scienze Geologiche, Università degli Studi "Roma Tre", Roma

13 5 Istituto Nazionale di Geofisica e Vulcanologia, sezione CNT

14 6 Dipartimento di Meccanica, Strutture, Ambiente e Territorio, Università degli Studi di Cassino,
15 Cassino

16 7 Consiglio Nazionale delle Ricerche, Istituto di Geologia Ambientale e Geoingegneria

17

18

19

20 Corresponding author: Dr. Emanuela Falcucci

21 Istituto Nazionale di Geofisica e Vulcanologia, sezione Milano-Pavia

22 Via Bassini, 15

23 20133 Milan

24 Italy

25 e-mail: falcucci@mi.ingv.it

26 **1. Introduction**

27 On April 6, 2009, at 01:32 GMT, a Mw 6.3 seismic event (Istituto Nazionale di Geofisica e
28 Vulcanologia, 2009; MedNet,
29 http://mednet.rm.ingv.it/procedure/events/QRCMT/090406_013322/qrcmt.html) hit the central
30 Apennines, severely damaging the town of L'Aquila and dozens of neighbouring villages, resulting
31 in almost 300 casualties. This earthquake was the strongest in central Italy since the devastating
32 1915 Fucino event (Mw 7.0). The INGV national seismic network located the hypocentre 5 km SW
33 of L'Aquila, 8-9 km deep. Based on this information and on the seismotectonic framework of the
34 region, earthquake geologists travelled to the field to identify possible surface faulting (Emergeo,
35 Working Group, 2009; Rilievi geologici di terreno effettuati nell'area epicentrale della sequenza
36 sismica dell'Aquilano del 6 Aprile 2009; <http://www.ingv.it>; Emergeo, Working Group, 2009,
37 submitted to Terra Nova).

38 The most convincing evidence of primary surface rupture is along the Paganica fault, the geometry
39 of which is consistent with seismological, synthetic aperture radar (SAR) and GPS data.

40 Investigation of other known normal faults of the area, i.e. the Mt. Pettino, Mt. San Franco and Mt
41 Stabiata normal faults, (Emergeo, Working Group, 2009; Rilievi geologici di terreno effettuati
42 nell'area epicentrale della sequenza sismica dell'Aquilano del 6 Aprile 2009; <http://www.ingv.it>;
43 Emergeo, Working Group, 2009, submitted to Terra Nova) suggested that these structures were not
44 activated during the April 6 shock.

45 In this report, we first describe the seismotectonic framework of the area and then we present the
46 field information that supports the occurrence of surficial displacement on the Paganica fault.

47

48 **2. Seismotectonic framework**

49 The shock of April 6, 2009, occurred in a region that has been struck by past earthquakes (Working
50 Group CPTI, 2004; Fig. 1) and is in an Apennine sector which contains a complex array of active

51 normal faults (e.g. Barchi et al., 2000; Galadini and Galli, 2000; Boncio et al., 2004; Roberts and
52 Michetti, 2004).

53 In the central Apennines, active faulting and earthquakes result from the ongoing extensional
54 tectonics. The strongest earthquakes in the CPTI04 seismic catalogue (Working Group CPTI, 2004)
55 occurred on September 9, 1349 (Mw 6.5), and February 2, 1703 (Mw 6.7). While the origin of the
56 former is still debated with regard to its source, the latter event was associated with the Upper
57 Aterno Valley fault system (UAVFS; Fig. 1), an ensemble of 7-to-10-km-long normal-oblique fault
58 segments (Galadini and Galli, 2000; Galadini and Messina, 2001). The 1703 earthquake induced
59 heavy damage in L'Aquila (Is 9, MCS scale; Stucchi et al., 2007), and the association of this
60 earthquake with the UAVFS has been demonstrated by paleoseismological investigations (Moro et
61 al., 2002) and by the distribution of the damage along the Upper Aterno River valley. Moreover, the
62 recent activity of the UAVFS is demonstrated by displacement of Late Pleistocene-Holocene
63 colluvial and alluvial deposits along the different segments of the system (Galadini and Galli, 2000;
64 Moro et al., 2002).

65 Other damaging earthquakes that have affected the L'Aquila region in the past (i.e., 1461,
66 Mw 6.4; 1762, Mw 5.9; 1916, Mw 5.2; 1958, Mw 5.2; see also Rossi et al., 2005) are more
67 comparable to the April 6 damage distribution than to the damage from the 1349 and 1703 events.
68 In particular, the 1461 event caused damage in Onna, Sant'Eusanio Forconese, Poggio Picenze,
69 Castelnuovo and L'Aquila (10, 10, 10, 10 and 9, MCS scale, respectively; Stucchi et al., 2007),
70 which is comparable to that of the April 6 earthquake (Galli and Camassi, 2009). Moreover, the
71 magnitude calculated on the basis of 1461 earthquake macroseismic data point is evidently close to
72 that of the April 6 shock. The most evident characteristic of all of these events is that, as in the case
73 of the April 6 shock, most of the damage affected L'Aquila and villages south of L'Aquila, while
74 little damage was recorded in the villages north of L'Aquila.

75 The UAVFS aside, other faults that could have been responsible for Mw >6 earthquakes

76 have been reported in the literature, with slightly different interpretations regarding their geometry
77 (Fig. 1). In particular, the Ovindoli-Pezza and Campo Felice-Colle Cerasitto faults (OPF-CFCF;
78 Fig. 1) have normal or normal-oblique kinematics, and delineate a 28- to 30 km-long, NNW-SSE to
79 NW-SE trending fault system south of L'Aquila that has Late Pleistocene-Holocene displacements,
80 as indicated by paleoseismological investigations (Pantosti et al., 1996; Salvi et al., 2003). The 20-
81 km-long, NW-SE trending Laga Mts. fault (LMF; Fig. 1) north of L'Aquila shows evidence of
82 Holocene activity with normal kinematics, as indicated by displacement of 8ka deposits (Galadini
83 and Galli, 2003). NE of L'Aquila, Holocene displacements have also been detected on the Campo
84 Imperatore fault system (CIFS; Fig. 1) (Galli et al., 2002; Galadini et al., 2003) where trenches
85 across secondary normal faults exposed evidence of surface faulting that occurred since about 3,4
86 ka. The different segments have a total length on the order of 33 km. Also, an impressive fault
87 system exists between San Demetrio ne' Vestini and Molina Aterno, and further south, between
88 Castelvechio Subequo and Goriano Sicoli (Bosi and Bertini, 1970; Bagnaia et al., 1992; Galadini
89 and Galli, 2000; Boncio et al., 2004) that were partly described as the Middle Aterno Valley fault
90 system (MAVFS; Fig. 1) (Galadini and Galli, 2000). Paleoseismological investigations that were
91 ongoing when the earthquake of April 6 occurred have revealed late Holocene activity of the
92 segment between Castelvechio Subequo and Goriano Sicoli (SVF; Fig. 1) (Falcucci et al., in
93 preparation).

94 The location of the main shock, the aftershocks, and the distribution of the April 6 damage
95 (Istituto Nazionale di Geofisica e Vulcanologia, 2009; Analisi dati di sismicit , <http://www.ingv.it>),
96 however, are not consistent with the activation of the above-described faults (Fig. 2).

97 To identify the causative fault of this April 6 earthquake, we consider the structural
98 framework of the area E and SE of L'Aquila, for where different structural patterns have been
99 proposed in the past (Fig. 3). Indeed, since the 1970s, this sector has been repeatedly investigated
100 regarding its neotectonic and active tectonic frameworks (Bosi and Bertini, 1970; Bosi, 1989;

101 Bagnaia et al., 1992; Bertini and Bosi, 1993). In particular, the two most recent studies present
102 substantially different structural frameworks regarding the Quaternary activity. Bagnaia et al.
103 (1992) reported a normal fault in the zone of Paganica that connects it with another normal fault in
104 the area of San Demetrio ne' Vestini, providing a simple structural image of this area (Fig. 3a) that
105 is affected by a ca. 20-km-long fault that partly affects Quaternary deposits. In contrast, Bertini and
106 Bosi (1993) summarised a complicated structural framework of Quaternary faults in this area. The
107 scheme proposed excludes a link between the Paganica fault and that in the area of San Demetrio
108 ne' Vestini. The information on the Quaternary faults was summarised in a map by Vezzani and
109 Ghisetti (1998), where the Paganica fault is mapped as a 9-km-long segment independent of the San
110 Demetrio ne' Vestini fault (Fig. 3b).

111 Finally, these data have been included in different ways in seismotectonic studies at a
112 regional scale. When mapping the surficial expression of sources that could potentially be
113 responsible for $M \geq 6.5$ earthquakes, Galadini and Galli (2000) did not include the short Paganica
114 fault segment at the NW end of the MAVFS. In contrast, Boncio et al. (2004) linked one segment of
115 the UAVFS mapped by Galadini and Galli (2000), i.e. the southeastern-most one (the Mt. Pettino
116 fault), to the Paganica fault, defining a single 13-km-long fault segment in the area of L'Aquila that
117 is not linked to the San Demetrio ne' Vestini fault (Fig. 3c). Roberts and Michetti (2004) and
118 Papanikolaou et al. (2005) reported a segment of more than 30 km in this area, not including,
119 however, the Paganica fault.

120 The geological surveys carried out *ad-hoc* after the earthquake along the Paganica fault
121 (which is presently considered the surface expression of the seismogenic source of the earthquake
122 of April 6) provide a different picture from that in the published literature before this earthquake.

123

124 **3. The Paganica fault**

125 Our geological surveys and analyses of aerial photographs taken after the earthquake define the

126 surficial geometry of the long-term expression of this fault. We define three NW-SE trending
127 segments that have a dextral en-echelon relationship (Fig. 4). The northernmost segment (segment
128 1, 3 km long) is located between the area east of Collebrincioni and Vallone delle Grotte (NE of
129 Tempèra). The central segment is detected in the area of Paganica (segment 2; 3 km long) and
130 terminates towards the SE along the road that connects Paganica and Pescomaggiore. The
131 southernmost segment (segment 3; 2 km long) extends to the NW from the village of San Gregorio.

132 The surficial expression of the northern fault segment is marked by a prominent fault scarp
133 in carbonate bedrock, at the base of which the slickenside surface is exposed (Fig. 5a). In the central
134 part of the segment, next to Colle Cocurello and Colle Enzano, the fault splits into two splays (Fig.
135 4). Here, unconsolidated slope deposits (i.e., not older than the Last Glacial Maximum period; see
136 Castiglioni et al. 1979; Dramis, 1983; Coltorti and Dramis, 1988, for more on this topic) are
137 exposed at the base of the fault scarp and seems to be displaced along the eastern fault splay, where
138 the “Vigne Basse” viaduct of the A24 motorway crosses the fault zone (Fig. 5b). Moreover, a small
139 hand-made excavation (few tens of centimetres deep) performed along the fault scarp slightly to the
140 north exposed post-Last Glacial Maximum debris dragged along the fault scarp, due to the fault
141 movement. Hence, overall, these findings confirm the Late Pleistocene-Holocene activity of the
142 Paganica fault.

143 The scarp on the central segment of the Paganica fault is present at the base of the slopes
144 between Tempèra and Paganica. The fault scarp is formed both in the carbonate bedrock – in the
145 northernmost part of the branch – and in fluvial deposits (Fig. 5c) – in the central and southern
146 portions of the segment. Here, according to Sheet 359 “L’Aquila” of the Geologic Map of Italy at
147 1:50,000 (Foglio CARG 1:50,000, 2009. Cartografia geologica ufficiale Foglio CARG 1:50,000 N.
148 359, L’Aquila), the tectonic structure places Pliocene/Middle Pleistocene continental deposits in
149 contact with Late Pleistocene sediments.

150 Many outcrops in the fault zone show displaced middle Pleistocene-Holocene fluvial and

151 slope deposits (Fig. 6a) along several synthetic fault planes (about N140° trending).

152 The oldest fluvial sediments displaced along the fault contain partially-to-strongly reworked
153 and/or weathered ash layers. Locally, ash deposits occur as primary air-fall deposits, as centimetre-
154 thick, tephra layers (Fig. 6b). Chemical, lithological and isotope analyses on three of the thickest,
155 best-preserved ashes have allowed us to assign them to the eruptions of “Tufo Pisolitico di
156 Trigoria”, “Pozzolane Rosse” and “Tufo Rosso a Scorie Nere” (authors’ unpublished data) from the
157 Colli Albani and Sabatini volcanic districts (central Italy); these ashes have been dated to about 560,
158 456 and 450 ka, respectively (Marra et al., 2009, and references therein). These data provide the
159 oldest chronological point-control for the dating of the activity of the Paganica fault.

160 More recent activity on the central segment of the Paganica fault is documented by displaced
161 colluvial and fluvial deposits along at least three shear planes that are exposed in the walls of a deep
162 trench (up to 5 m high) that is perpendicular to the fault. The trench, which is in north of Paganica,
163 was dug out by a strong jet of water from a leak in an aqueduct that ruptured during the earthquake
164 (Fig.6c). In the uppermost part of the trench, one fault plane places carbonate breccia in contact
165 with fluvial gravel, colluvial and soil levels that are displaced by some metres (Fig. 6c). Other shear
166 planes exposed in the trench displace fluvial and slope deposits (Fig. 6d). Organic-rich material in
167 the colluvium yielded radiocarbon ages of 5,718 BC/5,467 B.C. to 5,403 BC/5,387 B.C. (calibrated,
168 2σ) (Fig. 6c) and 34,970 \pm 470 BP. The latter age is supported by U/Th dating of a calcareous tufa
169 that is embedded in the dated organic-rich deposit that gave an age of 33,000 \pm 4100 BP. These data
170 support the Late Pleistocene-Holocene activity of the Paganica fault.

171 The southern end of the central segment of the Paganica fault ends against a N-S trending
172 tectonic structure that dips about 80° to the E. This fault is visible along the road that connects
173 Paganica to Pescomaggiore and shows normal-to-transpressive kinematics (Fig. 6e). Here, alluvial-
174 fan and colluvial deposits are displaced on the Paganica fault and the N-S fault. The sediments
175 contain the above-described Middle Pleistocene tephra layers, thus indicating a post-Middle

176 Pleistocene activity of the N-S fault.
177 The central and the southern branches of the Paganica fault are separated by a transfer zone (e.g.
178 Peacock, 2002) that is here marked by a wide landsurface that gently dips towards the NW.
179 The southern segment of the Paganica fault is located SE of Paganica from the area E of the San
180 Vittorino to the village of San Gregorio. Its surficial expression is defined by a scarp in carbonate
181 bedrock, at the base of which the slickenside surface is often exposed (Fig. 7a, b). At its northern
182 end, the fault branch intersects a fault plane trending about N20° and dipping steeply eastward,
183 which affects the carbonate bedrock (Fig. 7c). This fault and the above-mentioned N-S trending
184 tectonic structure are probably part of to the same fault system.

185

186 **4. Coseismic surface deformation along the Paganica fault zone**

187 The field surveys performed in the days subsequent to the main shock allowed us to identify
188 coseismic surface deformation along the Paganica fault that produced sets of NW-SE trending
189 ground cracks aligned along the fault zone, and parallel the fault segments, and which comprise a
190 belt not wider than a few tens of metres. These fractures were up to a few centimetres wide and
191 several tens of metres long (with no solution of continuity), and they commonly had an en-echelon
192 (mainly dextral step-over) arrangement.

193 These ground cracks were almost continuous 1) through fields and urban areas, where roads and
194 buildings were affected by newly formed ruptures and damage; 2) along slopes and plain areas; 3)
195 across any morphological features of the landscape, such as small ridges, valleys and landsurfaces;
196 and 4) affecting both natural and man-made terrain.

197 Moreover, most cracks along the central fault segment had vertical and horizontal offset between
198 the sides (with the downslope side lowered) of a few centimetres; the displacement increased during
199 the days subsequent to the April 6 shock, reaching a maximum of about 15 cm. In contrast, cracks
200 along the northern and the southern branches did not have readily observable vertical offsets rather

201 the ground was gently flexed, with an amplitude of less than 10 to 15 cm.

202

203 4.1 Northern fault segment

204 Along the northern segment of the Paganica fault, discontinuous ground ruptures, no more than
205 about 20 m long and 1 cm wide, were observed at the base of the fault scarp (Fig. 8a, b, c, d). In
206 particular, in the central part of the segment, the cracks are located where the fault splits into two
207 splays (i.e. between Colle Cocurello and the Vigne Basse locality, see previous section), with
208 different sets that trend roughly 120° to 170°. We observed cracks on both splays. A crack was also
209 present where the above-described Late Pleistocene-Holocene slope debris was perhaps displaced
210 on the tectonic structure (Fig. 8e).

211 Here, the “Vigne Basse” viaduct of the A24 motorway that crosses the Paganica fault zone (see
212 previous section) suffered damage during the shock of April 6. This damage to the viaduct, mainly
213 consisting in the re-opening of pre-existing “injuries” due to aging (i.e. old cracks with traces of
214 water percolation), occurred where it crosses the Paganica fault. Hence, the damage might have
215 been mostly caused by the strong ground motion caused by the earthquake, combined
216 (subordinately) with the coseismic ground deformation (which is marked by cracks) at the fault
217 zone (where piers of the viaduct are positioned) .

218 Moreover, evidence of reactivation of the fault plane is visible along the northern part of the
219 segment where a fresh-looking free-face up to 8-10 cm high is present at the base of the slickenside
220 surface. This free-face extends continuously for several metres, and marks the contact between the
221 carbonate bedrock and the slope scree at the base of the scarp (Fig. 8f). In some places, the free-face
222 was marked by the remnants of soil that coats the base of the scarplet. Moreover, the slope scree
223 was separated from the bedrock scarp surface by an about 1-3-cm-wide fissure.

224

225 4.2 Central fault segment

226 We found ground cracks in the area between Tempèra and Paganica that parallel the fault branch
227 where synthetic shear planes affect the Late Pleistocene-Holocene deposits (Fig. 9). Cracks are
228 present continuously along the entire segment. In the village of Paganica village, these ground
229 ruptures intersected the trench at the aqueduct (Fig. 9c), so we can reasonably hypothesise that the
230 damage to the aqueduct can be attributed to the coseismic surface displacement along the tectonic
231 structure.

232 To the southeast, we found ground ruptures as far as the southern end of the central
233 segment. Along the transfer zone that separates the central segment and southern of the Paganica
234 fault, we found two intersecting sets of ground fractures, one set trending about roughly N20° and
235 the other trending about NW-SE (about 150°) (Fig. 10a). The former set was aligned with the
236 aforementioned N-S fault, while the latter has a direction similar to the Paganica fault trend. Some
237 of these intersecting ground cracks appeared in the days subsequent to the mainshock, probably
238 related to the occurrence of afterslip (Fig. 10b).

239 Rare and discontinuous E-W trending fractures extend for less than 1 km in length in the
240 area of Tempèra. These secondary features might result from local, surficial craking due to passive
241 movement caused by ground shaking along secondary E-W striking tectonic structures, which have
242 been mapped in this area (e.g. Bagnaia et al., 1992; Vezzani and Ghisetti, 1998; Foglio CARG
243 1:50,000, 2009. Cartografia geologica ufficiale Foglio CARG 1:50,000 N. 359, L'Aquila); the Late
244 Quaternary activity of these E-W tectonic structures has never been demonstrated by thorough
245 geological studies.

246

247 4.3 Southern fault segment

248 A few tens of metres south-eastwards, we found NW-SE trending ground cracks at the northern end
249 of the southern fault segment that extend towards the SE and parallel the carbonate bedrock fault
250 scarp (Fig. 10c). The width of the fractures increased in the days subsequent to the mainshock. This

251 confirmed the occurrence of a certain amount of afterslip.
252 Further south, we found several ground ruptures close to the southern end of the fault branch, in the
253 area of San Gregorio (Fig. 10d, e, f), with some cracks being tens of metres long. According to eye
254 witnesses, some of these fracture formed during the aftershock of April 7 (Mw 5.6; Istituto
255 Nazionale di Geofisica e Vulcanologia, 2009; MedNet,
256 http://mednet.rm.ingv.it/procedure/events/QRCMT/090406_013322/qrcmt.html).
257 Here, the cracks were arranged in two sets trending between N120° and N150°. We also found, a
258 fresh fracture with N350° trend that affected the carbonate bedrock (Fig. 10g). A fresh-looking
259 crack on the ground occurred where the fracture intersected the surface (Fig. 10g, inset).

260

261 **5. Investigation along the Mt. Bazzano antithetic normal fault**

262 We also examined a previously mapped normal fault (e.g. Vezzani and Ghisetti, 1998) on the
263 eastern slopes of Mt. Bazzano, and that is an antithetic fault to the Paganica fault. The surficial
264 expression of this structure is a prominent fault scarp at the base of which the fault plane is exposed.

265 We identify local evidence of fault-plane reactivation that consists of a free face of up to 15
266 cm high, at the base of the slickenside and which, in places, had remnants of the soil coating the
267 scree deposited at the base of the fault plane. This new free face was discontinuous as it was
268 typically observable for no more than a few metres in length (Fig. 10h). Many sectors of the fault,
269 indeed, showed no evidence of rejuvenation along the bedrock scarp.

270 Taken together with 1) the remarkable steepness of the slopes, and 2) the absence of evident ground
271 cracks along the fault, – comparably to the Paganica fault – this suggests that the formation of the
272 free face along the Bazzano normal faults was probably mainly due to non-tectonic gravitational
273 processes and/or compaction of the unconsolidated scree at the base of the fault planes.
274 Nevertheless, we cannot preclude a coseismic tectonic component to the new free face because, the
275 fault is antithetic to the Paganica fault.

276

277 **6. Concluding remarks**

278 Our field surveys and studies of aerial photographs in the L'Aquila area (central Apennines) after
279 the April 6 seismic event (Mw 6.3) permitted us to identify evidence of coseismic surface faulting
280 on the Paganica fault, a NW-SE striking and SW dipping Quaternary normal fault located E of
281 L'Aquila.

282 This structure had been mapped by previous studies (Bagnaia et al., 1992, Vezzani and
283 Ghisetti, 1998; Boncio et al., 2004; Foglio CARG 1:50,000, 2009) but none of the previous works
284 had documented its latest Pleistocene and Holocene activity.

285 Our data define three en-echelon segments of the Paganica fault, based on its surficial
286 expression, each one 3-4 km long. The total length of the structure is about 10 km.

287 Apart from the obvious association with the April 6 earthquake, recent activity of the
288 Paganica fault is defined by the displacement of ¹⁴C- and U-series-dated Late Pleistocene-Holocene
289 deposits along the tectonic structure. In particular, a trench eroded along a ruptured aqueduct
290 exposed several tens of centimetres of offset in Late Holocene deposits along the Paganica fault,
291 which suggests that the fault has probably generated stronger earthquakes in the past.

292 We identified evidence of surface faulting along the Paganica fault due to the April 6 shock
293 including set of continuous aligned ground cracks along the central segment of the fault; some
294 cracks had both vertical and horizontal offset up to about 15 cm. The offset of some cracks
295 increased as much as a few centimetres in the days subsequent to the mainshock. This is probably
296 evidence of a small amount of afterslip related to the aftershocks. The presence of ground cracks,
297 although discontinuous and with no vertical offset along the northern and the southern fault
298 segments suggest that also these branches of the Paganica fault were probably activated during the
299 April 6 shock.

300 Overall, our observations show to relatively little coseismic surficial movements on the Paganica

301 fault, which probably was less than a few centimetres. Further evidence that supports the occurrence
302 of surface faulting on the Paganica fault includes the rejuvenation of the fault plane, i.e. the free-
303 face observed at the base of the slickenside, along the northern and central segments of the tectonic
304 structure. It is worth noting that base on empirical relationships between moment magnitude and the
305 surface rupture length (Wells and Coppersmith, 1994), the length of the Paganica fault rupture is
306 consistent (taking into account the uncertainty range proposed by Wells and Coppersmith) with the
307 reported magnitude ($M_w=6.3$; Istituto Nazionale di Geofisica e Vulcanologica, 2009; MedNet,
308 http://mednet.rm.ingv.it/procedure/events/QRCMT/090406_013322/qrcmt.html).

309 Finally, our evidence of surface rupture combined with the location and the length of the Paganica
310 fault is consistent with it being the source of the April 6 earthquake. This conclusion is also
311 consistent with the results from studies of 1) interferometric SAR observations (Atzori et al., in
312 press); 2) the location and geometry of the fault modelled using the coseismic ground deformation
313 patterns from GPS data (Anzidei et al., in press); and 3) seismological data, that is the location and
314 focal mechanism of the main shock and the aftershock distribution (Istituto Nazionale di Geofisica e
315 Vulcanologia, 2009; Analisi dati di sismicit , <http://www.ingv.it>).

316

317 **Acknowledgements**

318 **The anonymous reviewer strongly improved the manuscript; we are grateful to Daniela**
319 **Pantosti and to David Schwartz for the useful discussion and suggestions. Davide Piccarreda**
320 **and Carlo Piccarreda helped us in the field work. We are grateful to all the people who live in**
321 **the area struck by the earthquake and that, often, helped us in the field investigations.**

322

323

324 **References**

325

326 Anzidei, M., Boschi, E., Cannelli, V., Devoti, R., Esposito, A., Galvani, A., Melini, D.,

327 Pietrantonio, G., Riguzzi, F., Sepe, V. and E. Serpelloni (in press). Coseismic deformation of the
328 destructive April 6, 2009 L'Aquila earthquake (central Italy) from GPS data. *Geophys. Res. Lett.*
329

330 Atzori, S., I. Hunstad, M. Chini, S. Salvi, C. Tolomei, C. Bignami, S. Stramondo, E. Trasatti, A.
331 Antonioli, and E. Boschi (2009), Finite fault inversion of DInSAR coseismic displacement of the
332 2009 L'Aquila earthquake (central Italy), *Geophys. Res. Lett.*, doi:10.1029/2009GL039293, in
333 press.
334

335 Bagnaia, R., D'Epifanio, A., and S. Sylos Labini (1992). Aquila and Subequan basins: an example
336 of Quaternary evolution in central Apennines, Italy. *Quaternaria Nova II*, 187-209.
337

338 Barchi, M., Galadini, F., Lavecchia, G., Messina, P., Michetti, A.M., Peruzza, L., Pizzi, A., Tondi,
339 E. and E. Vittori (editors) (2000). Sintesi delle conoscenze sulle faglie attive in Italia Centrale:
340 parametrizzazione ai fini della caratterizzazione della pericolosità sismica. CNR-Gruppo Nazionale
341 per la Difesa dai Terremoti - Roma, 2000, 62 p..
342

343 Bertini, T. and C. Bosi (1993). La tettonica quaternaria della conca di Fossa (L'Aquila). *Il*
344 *Quaternario* **6**, 293-314.
345

346 Boncio, P., Lavecchia, G. and B. Pace (2004). Defining a model of 3D seismogenic sources for
347 seismic hazard assessment applications: the case of central Apennines (Italy). *Journal of*
348 *Seismology* **8/3**, 407-425.
349

350 Bosi, C. (1989). Tentativo di correlazione fra le successioni plio-pleistoceniche. In: "Guida
351 all'escursione sul tema: Elementi di tettonica pliocenico-quaternaria e indizi di sismicità olocenica
352 nell'Appennino laziale-abruzzese". Tip. Esagrafica, Roma.
353

354 Bosi, C. and T. Bertini (1970). La geologia della media valle dell'Aterno. *Memorie della Società*

355 *Geologica Italiana* **9**, 719-777.

356

357 Castiglioni, G.B., Girardi, A., Sauro, U. and F. Tessari (1979). Grèzes litées e falde detritiche
358 stratificate di origine crionivale. *Geografia Fisica e Dinamica Quaternaria* **2**, 64-82.

359

360 Coltorti, M. and F. Dramis (1988). The significance of stratified slope-waste deposits in the
361 Quaternary of Umbria-Marche Apennines, central Italy. *Z Geomorph N F* **71** (Suppl. Bd), 59-70.

362

363 Dramis, F. (1983). Morfogenesi di versante nel Pleistocene superiore in Italia: i depositi detritici
364 stratificati. *Geografia Fisica e Dinamica Quaternaria* **6**, 180-182.

365

366 Emergeo Working Group (2009). Rilievi geologici nell'area epicentrale della sequenza sismica
367 dell'Aquilano del 6 aprile 2009. *Quaderni di Geofisica*, 70, ISSN 1590-2595,
368 <http://portale.ingv.it/produzione-scientifica/quaderni-di-geofisica/quaderni-di-geofisica-2009>

369

370 Emergeo Working Group (submitted to *Terra Nova*). Geological coseismic effects of the 2009
371 L'Aquila seismic sequence (central Italy).

372

373 Falcucci, E., Gori, S., Moro, M., Galadini, F., Ceccaroni, E., Pisani A.R. and P. Fredi (in
374 preparation). Paleoseismological investigations along the Middle Aterno-Subequana Valley fault
375 system (Abruzzi Apennines, central Italy): Implications for seismotectonics and seismic hazard in
376 the sector struck by the April 6, 2009 earthquake.

377

378

379 Foglio CARG 1:50,000, 2009. Cartografia geologica ufficiale Foglio CARG 1:50,000 N. 359,

380 L'Aquila

381

382 Galadini, F., and P. Galli (2000). Active tectonics in the central Apennines (Italy) - input data for
383 seismic hazard assessment. *Natural Hazards* **22**, 225-270.

384

385 Galadini, F. and P. Galli (2003). Paleoseismology of silent faults in the central Apennines (Italy):
386 the Mt. Vettore and Laga Mts. faults. *Annals of Geophysics* **46/5**, 815-836.

387

388 Galadini, F., Galli, P. and M. Moro (2003). Paleoseismology of silent faults in the central
389 Apennines (Italy): the Campo Imperatore fault (Gran Sasso Range fault system). *Annals of*
390 *Geophysics* **46**, 793-814.

391

392 Galadini, F. and P. Messina (2001). Plio-Quaternary changes of normal fault architecture in the
393 central Apennines (Italy). *Geodinamica Acta* **14**, 321-344.

394

395 Galli, P. and R. Camassi (2009). Rapporto sugli effetti del terremoto aquilano del 6 aprile 2009,
396 <http://www.mi.ingv.it/eq/090406/quest.html>

397

398 Galli, P., Galadini, F., Moro, M. and C. Giraudi (2002). New paleoseismological data from the Gran
399 Sasso d'Italia area (central Apennines). *Geophysical Research Letters* **29**,
400 doi:10.1029/2001GL013292.

401

402 Istituto Nazionale di Geofisica e Vulcanologia (2009). <http://www.ingv.it>

403

404 Marra, F., Karner, D.B., Freda, C., Gaeta, M. and P. Renne (2009). Large mafic eruptions at Alban

405 Hills Volcanic District (central Italy): Chronostratigraphy, petrography and eruptive behaviour.
406 Journal of Volcanology and Geothermal Research **179**, 217-232.
407
408 Moro, M., Bosi, V., Galadini, F., Galli, P., Giaccio, B., Messina, P. and A. Sposato (2002). Analisi
409 paleosismologiche lungo la faglia del M. Marine (alta valle dell'Aterno): risultati preliminari. Il
410 Quaternario (Italian Journal of Quaternary Sciences) **15**, 267-278.
411
412 Pantosti, D., D'Addezio, G. and F.R. Cinti (1996). Paleoseismicity of the Ovindoli-Pezza fault,
413 central Apennines, Italy: a history including a large previously unrecorded earthquake in the Middle
414 Ages (890-1300). Journal of Geophysical Research **101**, 5,937-5,959.
415
416 Papanikolaou, I.D., Roberts, G.P. and A.M. Michetti (2005). Fault scarps and deformation rates in
417 Lazio–Abruzzo, central Italy: comparison between geological fault slip-rate and GPS data.
418 Tectonophysics **408**, 147-176.
419
420 Peacock, D.C.P. (2002). Propagation, interaction and linkage in normal fault systems. Earth-Science
421 Reviews **58**, 121-142.
422
423 Roberts, G.P. and A.M. Michetti (2004). Spatial and temporal variations in growth rates along
424 active normal fault systems: an example from the Lazio-Abruzzo Apennines, central Italy. Journal
425 of Structural Geology **26**, 339-376.
426
427 Rossi, A., Tertulliani, A. and M. Vecchi (2005). Studio macrosismico del terremoto dell'Aquilano
428 del 24 giugno 1958. Il Quaternario (Italian Journal of Quaternary Sciences) **18**, 101-112.
429

430 Salvi, S., Cinti, F.R., Colini, L., D'Addezio, G., Doumaz, F. and E. Pettinelli (2003). Investigation
431 of the active Celano-L'Aquila fault system, Abruzzi (central Apennines, Italy), with combined
432 ground-penetrating radar and paleosismic trenching. *Geophysical Journal International* **155**, 805-
433 818.

434

435 Stucchi et al. (2007). DBMI04, il database delle osservazioni macrosismiche dei terremoti italiani
436 utilizzate per la compilazione del catalogo parametrico CPTI04.
437 <http://emidius.mi.ingv.it/DBMI04/Quaderni di Geofisica, V. 49, 38 p.>

438

439 Vezzani, L. and F. Ghisetti (1998). Carta geologica dell'Abruzzo, Scala 1:100.000, S.El.Ca.,
440 Firenze.

441

442 Wells, D.L. and K.J. Coppersmith (1994). New empirical relationships among magnitude, rupture
443 length, rupture width, rupture area, and surface displacement. *Bulletin of Seismological Society of*
444 *America* **84**, 974-1,002.

445

446 Working Group CPTI (2004), Catalogo Parametrico dei Terremoti Italiani, versione 2004 (CPTI04).
447 INGV, Bologna, Italy. <http://emidius.mi.ingv.it/CPTI/> (Last check of the availability: May 2009).

448

449 **Figure captions**

450 Figure 1 – Seismotectonic framework of the central Apennines (modified after Galadini and Galli,
451 2000). MVEF - Mt. Vettore fault; NFS – Norcia fault system; LMF - Laga Mts. fault; UAVFS -
452 Upper Aterno Valley fault system; CIFS – Campo Imperatore fault system; CFCF - Campo Felice-
453 Colle Cerasitto fault; OPF - Ovindoli-Pezza fault; MAVFS - Middle Aterno Valley fault system;
454 SVF – Subequana Valley fault; MMF - Mt. Morrone fault; FF - Fucino fault; MPFS – Maiella-

455 Porrara fault system; ACF - Aremogna-Cinquemiglia fault; USFS - Upper Sangro Valley fault
456 system.

457

458 Figure 2 – Digital terrain model of the study area. Black lines with bars on downthrown side mark
459 the Paganica fault segments as defined in the present study; the black dotted line encircles the area
460 that suffered intensities (MCS scale) \geq VIII from the April 6 shock; epicentres of the aftershocks
461 that occurred in the area are marked by white circles (for Mw up to 4) and white squares (i.e., Mw
462 comprised between 4 and 5), and the white stars indicate the epicentres of *i*) the Mw 6.3 April 6
463 shock (with the focal mechanism) and *ii*) the Mw 5.7 aftershock of April 7.

464

465 Figure 3 – Tectonic structures (white lines) affecting the area of Paganica, as reported in a) Bagnaia
466 et al. (1992), b) Vezzani and Ghisetti (1998) and c) Boncio et al. (2004).

467

468 Figure 4 – Satellite image on which the segments of the Paganica fault (as defined by the study) are
469 shown (white lines; white bars and points mark the possible southern prolongation of the coseismic
470 surface displacement), together with the approximately N-S trending fault plane (white dashed line)
471 detected between the central and the southern branches of the Paganica fault; the black and the
472 white squares mark sites where *i*) the Paganica fault and *ii*) the evidence of surface faulting/
473 rupturing (i.e. ground cracks and free-faces) have been observed, respectively (left panel). The
474 location of the figures showing the Paganica fault segments and the evidence of co-seismic ground
475 deformation along the tectonic structure is reported on the right panel.

476

477 Figure 5 – a) Fault scarp formed on the carbonate bedrock, along the northern segment of the
478 Paganica fault; b) slope deposits, the attitude of which is highlighted by the white dashed line,
479 probably placed in contact (marked by the black arrows) with the carbonate bedrock by the northern

480 fault branch (black dashed lines mark secondary shear planes); c) fault scarp on the northern fault
481 branch formed on alluvial deposits, near to the southern end of the segment.

482

483 Figure 6 – a) Fluvial gravels (the black dashed line marks the attitude) displaced along a synthetic
484 fault plane (indicated by the black arrows) related to the Paganica fault, characterised by rotated
485 clasts in the shear zone (inset); b) fluvial and slope deposits containing Middle Pleistocene tephra
486 layers displaced along a synthetic fault plane (indicated by the white arrows) of the Paganica fault;
487 c) synthetic fault plane, detected along the walls of the trench of the aqueduct, displacing alluvial,
488 pedogenic and colluvial deposits; d) synthetic shear plane, detected along the wall of the trench,
489 placing fluvial gravels in contact with colluvial deposits dated to the Late Holocene.

490

491 Figure 7 – a) Fault plane of the southern fault branch, exposed at the base of the carbonate bedrock
492 fault scarp; b) Paganica fault plane in the area of San Gregorio. c) Fault plane roughly N20°
493 trending, seen at the northern end of the southern branch of the Paganica fault;

494

495 Figure 8 – Ground cracks along the northern fault branch near the “Vigne Basse” viaduct of the
496 A24 motorway (a, b, c, d) and next to the slope deposits shown in Figure 5b (e); free-face exposed
497 at the base of the slickenside surface along the northern fault segment (f);

498

499 Figure 9 – Ground cracks (marked by black and white arrows) along the central segment of the
500 Paganica fault affected paved courtyards, walls, fields, and dirt roads (a, b, c, d, e, f, g); some
501 fractures have vertical and horizontal offsets between their sides (h, inset). Some ground cracks are
502 located close to the trench of the aqueduct (c).

503

504 Figure 10 – Intersecting sets of ground cracks (the white arrows indicate the 20°N trending set while

505 the black ones indicate the NW-SE (150°) trending set) in the transfer zone between the central and
506 southern segments of the Paganica fault (a, b); Ground fractures formed in the days subsequent to
507 the mainshock, in fields close to the northern end of the northern fault branch (c); ground cracks at
508 San Gregorio along paved and dirt roads, courtyards and building walls (d, e, f); fracture affecting
509 the carbonate bedrock near the southern termination of the southern segment of the Paganica fault
510 (g); free-face (white band at the base of the slickenside, indicated by the black arrows) exposed in
511 places at the base of the bedrock of the Bazzano fault (h).