

BIOTECHNOLOGY TECHNIQUES
Volume 9 No.8 (August 1995) pp.583-588
Received as revised 19th June

ADSORPTION EQUILIBRIUM AND DYNAMICS OF LACTASE/CM-SEPHADEX SYSTEM

Şebnem HARSA^{1*} Yekta GÖKSUNGUR² Ulgar GÜVENÇ²

1 Izmir Institute of Technology, Anafartalar cad.
No:904, Basmane 35220, Izmir-Turkey

2 Dept. of Food Eng., Faculty of Engineering,
Ege University 35100, Bornova, Izmir-Turkey

SUMMARY

Partitioning behaviour and adsorption isotherms of lactase/CM-Sephadex system at equilibrium were investigated together with the adsorption kinetics in this study. Maximum adsorption was obtained at the pH values between 5.5-6.0. Adsorption isotherm was a close fit to the Langmuir model.

INTRODUCTION

Lactase (β -galactosidase) enzyme preparations are widely used to hydrolyse lactose in milk, milk products and whey (by-product of dairy industry which cause environmental problems due to their high biological oxygen demand), giving a mixture of glucose and galactose for consumers who suffer from lactose intolerance. These monosaccharides are sweeter and more soluble than lactose; furthermore, glucose and galactose are more readily fermented than lactose.

Lactase can be produced by *Kluyveromyces marxianus* which is a potential new source of commercial enzymes. Our aim is to obtain data in order to design an appropriate separation and immobilization scheme for lactases from *K. marxianus* using low-cost ion-exchangers.

Within this context, we studied the equilibrium and dynamic behaviour of lactase/Sephadex system. Firstly, ionisation of adsorbate and adsorbent was determined from the titration curves. The effect of pH and gel concentration on the equilibrium partition coefficients was investigated. Finally, adsorption isotherms and kinetics were generated under the optimal conditions and suitable mathematical models were discussed. These will unveil information relevant to both adsorption based separation schemes and immobilized enzyme reaction systems. In both cases, knowledge of the equilibrium and dynamic behaviour of relevant species is key to process design and optimisation.

MATERIALS and METHOD

Carboxymethyl (CM)-Sephadex C-50 and lactase (Lactozym) were the products of Pharmacia and Novo Nordisk, respectively. This enzyme was the purified preparation of Novo and contained no other proteins (3000 Units/mL).

The binding capacity of CM-Sephadex and enzyme as a function of pH was obtained by pH titration with 0.1M HCL; 0.5 mL lactase enzyme (1500 Units) was used in 10 mL suitable buffer solutions to conduct the adsorption experiments. Enzyme solutions in

different pH (pH 2.5-10) buffers (citrate, acetate and phosphate buffers, 0.1M) were contacted with 0.5 g of hydrated and equilibrated gel. To determine the effect of gel:liquid ratio, different bulk volume of hydrated gel ($V_g/V = 0.02-0.5$) were used. Samples were placed to the incubator with shaking at 28°C for 24 hours and analysed for protein (using the Lowry method) when the solution approaches to equilibrium. Partition coefficients indicate the fraction of adsorbed protein at equilibrium, and estimated from the equation of ($f = q / p$), where q and p are the equilibrium concentrations of adsorbed protein and free protein.

Adsorption isotherm experiments were conducted for different enzyme concentrations at pH 6.0 at 28°C. Gels with different enzyme concentrations were placed into an incubator and stirred continuously. Equilibrium solution concentrations were determined approximately 24 hours later and the amount of enzyme adsorbed by 1 mL of hydrated was calculated from the difference of the initial and equilibrium solution concentrations.

A graph was plotted against the solution concentration at equilibrium (p) and adsorbed enzyme concentration inside the pellets (q). The calculations for isotherm equations were done and model line was also drawn on the graph.

To conduct the dynamic experiments, samples were taken at intervals and protein concentration at each free solution was determined. A graph p/p_0 against time was set up.

RESULTS and DISCUSSION

Ionisation of adsorbate and adsorbent. From the titration curves of lactase and CM-Sephadex, the ionisation of adsorbate and adsorbent was calculated and values are shown in Figure 1.




Figure 1: Ionisation of adsorbate and adsorbent

The gel is fully (negatively) charged at pH above 7.0; below that level, its net charge decreases sharply, and at pH 5.0 only half of its potential binding capacity is active. On the other hand, at the isoelectric point (here 6.5), the net charge of the enzyme is zero. At lower pH, the molecule becomes (positively) charged: at pH 5.0 about 50% of its total ionisation capacity is present, whereas at pH 3.0 this increases to 83%. The lactase is fully charged at very low pH (pH 2.0). At above the isoelectric point, the net charge of the enzyme is negative (ie. the same as the adsorbent) and no electrostatic binding will occur.

Effect of pH on partitioning. In Figure 2, the minimum value of partition coefficients were found at the pH values above 7.0 and below 4.0. Partitioning behaviour shows a rapid increase up to pH 6.0 and at pH values between 5.5-6.0 the optimum was observed. At this pH value the partition coefficient reaches its value of about 15.0. Below this pH partition coefficient started decreasing and at pH 9.0 and above nearly no adsorption was recorded.




Figure 2: Effect of pH on the partition coefficient

According to Coulomb's law the interaction force between unlike charges e_1 , e_2 is proportional to $e_1 \cdot e_2$. Thus, the net interaction force between the protein and matrix should be proportional to the product of the fractional degree of ionisation of each component (Harsa et al., 1990). The region of maximum partition coefficient corresponds to the region of maximum ionic interaction, as seen from Figure 2.

Effect of gel concentration. A mass balance on protein shows that:

$$V p_o = V p + V_g q \quad (1)$$

Where V and V_g are the bulk liquid and gel volume respectively, p_o is the initial concentration of protein in the liquid phase, and p and q are the final concentrations of unbound and bound protein respectively. It can be shown that:

$$f = \frac{(p_o / p) - 1}{V_g / V} \quad (2)$$

Equation (2) shows the relation between V_g/V and the partition coefficient. The partition coefficient can be expressed in terms of the Langmuir parameters as follows:

$$f = q / p = q_m / (K_p + p) \quad (3)$$

From equation the upper limit value for K is obtained when $p \rightarrow 0$, and is given by:

$$f = q_m / K_p \quad (4)$$

To conduct the experiments V_g/V was changed, the same

concentration of enzyme was used all the time. As V_g/V increased the partition coefficient decreased. The maximum value of partition coefficient was obtained for $V_g/V = 0.03-0.05$. At the value of 0.1 partition coefficient start to fall down and reaches its minimum at 0.3. In all adsorption experiments the V_g/V value of 0.05 was used.




Figure 4: Effect of gel concentration on equilibrium

Adsorption isotherms. Adsorption isotherm experiments were conducted at pH 6.0, temperature 28°C, since these values were found to be the optimal for the enzyme stability and maximum adsorption. Figure 4 shows the adsorption isotherm.




Figure 4: Adsorption isotherm of lactase on CM-Sephadex resin

Adsorption isotherms show the distribution of protein concentration between the liquid phase at equilibrium (p) and gel phase (q). There are several equations to describe the adsorption isotherms for ion exchange chromatography. These are linear, Freundlich and Langmuir isotherms. The most applicable and robust one is Langmuir model, which is shown below:

$$q = \frac{q_m p}{(K_p + p)} \quad (5)$$

Using this model it is possible to calculate the parameters which are quite important for the design and scale-up of chromatographic separation purposes. K_p is a dissociation constant for adsorbent-adsorbate complex, q_m is the maximum adsorption capacity of the gel. The value of q_m for the lactase/CM-Sephadex system is approximately 42 mg protein/ml gel. Dissociation constant K_p was calculated as 3.1 mg/ml solvent.

Adsorption kinetics. Determination of adsorption kinetics of an enzyme/ion-exchange system is important to understand how fast the equilibrium has been reached. To know the mechanism involved during adsorption has great importance in the design and scale-up of an adsorption system.

At equilibrium the rate of adsorption can be written:

$$- V \frac{d(p)}{d(t)} = k A (p - p^*) \quad (6)$$

where V is the bulk solvent volume and A is the bead interfacial surface area.

Equations (5), and (6) provide a simplified model for the kinetics of adsorption. Details of calculations were given elsewhere (Harsa et al., 1993).

$$Z = 1 - e^{-ka\alpha' t} / \alpha' \quad (7)$$

Equation (7) represents a simplified model for batch adsorption kinetics in a stirred vessel, in terms of a global mass transfer coefficient, ka , the Langmuir parameters q_m & K_p , and the gel and solvent volume ratio.




Figure 5: Adsorption kinetics of lactase/CM-Sephadex system

As seen in Figure 5, approximately 22% of adsorption occurred in about 4 hours time. The equilibrium has been attained after 24 hours with an adsorption yield of approximately 40-50%.

The dynamic model proposed for the protein/ion-exchanger systems described above closely fit the experimental data. Adsorption kinetics of this system is considered to be slow, this can be because of the size of the enzyme molecule (MW 200.000 Da, $D_m = 4.2 \cdot 10^{-11} \text{ m}^2/\text{sec}$).

This model assumes that kinetics can be described in terms of a lumped mass transfer coefficient involving internal and

external resistances. The global mass transfer coefficient k_a can be estimated from the initial rate of adsorption, assuming that at $t \approx 0$, $p \gg p^*$ in Equation (6), ie:

$$\text{at } t \approx 0 \quad - dp/dt = k a p$$

$$\text{Integration yields: } \ln(p/p_0) = -k a t$$

Therefore a plot of $\ln(p/p_0)$ versus t for experimental points at early stages of adsorption should yield a straight line with slope = $-k_a$.

Estimated apparent mass transfer coefficient (k) values are shown in Table 1 as a function of particle size.

Table 1: Mass transfer coefficients

Hydrated bead diameter of CM-Sephadex (m)	Experimental k (ms^{-1})	Theoretical k (ms^{-1})
$2.0 \cdot 10^{-4}$	$0.433 \cdot 10^{-7}$	$3.0 \cdot 10^{-7}$
$6.0 \cdot 10^{-4}$	$1.3 \cdot 10^{-7}$	$1.0 \cdot 10^{-7}$

It can be seen that the experimental value of k is of the same order of magnitude as that predicted (the differences can be accounted for by the assumption that $Sh=2$ which corresponds to very low values of Re , for a sphere in an unstirred liquid).

Experiments show that this single parameter model gives a reasonable description of the adsorption kinetics for lactase/CM-Sephadex system.

It can be concluded that data obtained from the experiments can be used for the designing of a lactase enzyme immobilization + separation and purification process using ion-exchange chromatography.

NOMENCLATURE

a	specific mass transfer area
D_m	molecular diffusion coefficient (m^2/sec)
e_1, e_2	charge of the protein and the gel
k	apparent mass transfer coefficient (s^{-1})
k_a	global mass transfer coefficient
f	partition coefficient
K_p	dissociation constant for adsorbent-adsorbate complex, (mg/mL solvent)
p	equilibrium concentration of free enzyme, (mg free enzyme/mL solution)
q	equilibrium concentration of adsorbed enzyme, (mg ads./mL gel)
q_m	maximum adsorption capacity, (mg ads./ml gel)
Re	particle Reynolds number
Sh	Sherwood number
V_g/V	gel volume (mL)/bulk solvent volume (mL)
Z	dimensionless extent of adsorption
α	K_p/p_0 , model parameter
α'	$(\alpha/\beta) + 1$, model parameter
β	$V_g q_m / V p_0$, model parameter

REFERENCES

- Harsa, S., Pyle, D.L. & Zaror, C.A. (1990). In: Separations for Biotechnology 2, (Ed. D.L. Pyle). Elsevier Applied Science pp 658.
- Harsa, S., Zaror, C.A. & Pyle, D.L. (1993). Enzyme and Microbial Techn. 15: 906-915.