

Association between red meat consumption and cancer risk

Marija Boskovic¹, Milan Z. Baltic¹

Abstract: Cancer is leading cause of mortality worldwide. It is assumed that cancer risk is mainly affected by environmental factors including diet habits. Among other food, it is considered that red meat consumption is linked to increased risk of cancer. Formation of mutagens, heterocyclic aromatic amines, and polycyclic aromatic hydrocarbons, mainly depend on cooking methods and is the highest in pan fried and barbecued meat. Fat, N-nitroso compounds and haeme iron which are also found in meat and meat products are considered to influence the occurrence of cancer. Colorectal cancer is the most often associated with red meat consumption, but studies show that meat eating may increase the risk for other types of cancer including breast cancer, gastric cancer, prostate cancer, pancreatic cancer, lung cancer, laryngeal cancer and bladder, kidney, and endometrial tumours.

Keywords: red meat, PAH, HAA, colorectal cancer, diet and lifestyle habits.

Introduction

Changes in dietary habits and lifestyle have resulted in civilisation diseases including coronary heart disease, obesity, hypertension, type 2 diabetes, cancers, autoimmune disease and osteoporosis, among which, cancer is leading cause of mortality worldwide (Carrera-Bastos *et al.*, 2011; Stewart and Wild, 2014). As reported by a WHO (World Health Organization) update from 2014, approximately 14 million new cases of cancer and 8.2 million cancer-related deaths were reported in 2012 (Stewart and Wild, 2014). According to the same report, it is expected in the next two decades that cancer cases per year will rise from 14 million in 2012 to 22 million (Stewart and Wild, 2014). The most frequent types of cancer which cause mortality are lung (1.59 million deaths), liver (745 000 deaths) and stomach (723 000 deaths) cancer followed by colorectal (694 000 deaths) breast (521 000 deaths) and oesophageal cancer (400 000 deaths). Serbia is a developing country where stress, pollution, irregular diet and bad eating habits, lack of physical activity and insufficient sleep are part of everyday life. All of this has led to increasing cancer prevalence in Serbia. WHO data shows that cancer mortality rate in Serbia was 160.1 cases per 100,000 population in 2000, and 170.8 cases per 100,000 population in 2012 (Stewart and Wild, 2014). In 2012, only

Zimbabwe, Hungary, Armenia and Mongolia had higher cancer mortality rates than Serbia. Cancer risk is influenced 5% by genetic factors, while the remaining impact is due to environmental factors, of which 30 to 35% represents diet (Baena Ruiz and Salinas Hernández, 2014). Meat, especially red meat, and meat products are the types of foods most often associated with some types of cancer, which highlights the need to discuss the role of meat in cancer aetiology. In October 2015, the Working Group from the International Agency for Research on Cancer (IARC) classified consumption of processed meat as “carcinogenic to humans” (Group 1) (International Agency for Research on Cancer, 2015). This classification was based on their being sufficient evidence to associate meat product (i.e. processed meat) consumption with colorectal cancer as well as with stomach cancer. Taking into account substantial epidemiological data showing a positive association between consumption of red meat and colorectal cancer, the IARC classified consumption of red meat as “probably carcinogenic to humans” (Group 2A).

Carcinogens from red meat

Fat, heterocyclic aromatic amines (HAAs), polycyclic aromatic hydrocarbons (PAHs), N-nitroso compounds (NOc) and haeme iron are substances

¹University of Belgrade, Faculty of Veterinary Medicine, Bulevar Oslobođenja 18, 11000 Belgrade, Republic of Serbia.

Corresponding author: Marija Boskovic, marijaboskovic116@gmail.com

isolated from unprocessed or processed red meat which are considered to influence the occurrence of cancer (Higgs, 2000; Williamson et al., 2005; Ferguson, 2010; Corpet, 2011; Baena Ruiz and Salinas Hernández, 2014; Boskovic et al., 2015). It is also suggested that intake of red meat, as a high energy food, could have an impact on obesity, which presents a major risk factor for cancer (*World Cancer Research Fund/American Institute for Cancer Research*, 2007; Ferguson, 2010). Although the exact mechanism by which meat influences the increased cancer risk is not completely known, there are a few proposed hypotheses. It is supposed that intake of fat, responsible for the high energy density of meat, has an important role in carcinogenesis because it increases insulin resistance, and impacts on higher production of secondary bile acids, which act as aggressive surfactants for the mucosa (Ferguson, 2010; Corpet, 2011; Baena Ruiz and Salinas Hernández, 2014).

NOc are produced by the reaction of nitrite and nitrogen oxides with secondary amines and N-alkylamides (Lijinsky, 1999; Ferguson, 2010). NOc can be formed endogenously after consumption of red and processed meat or could be found in some processed meats, although some authors suggested that N-nitrosamines produced from additives in processed meat have low carcinogenic potential because of small amounts in which they are found (Higgs, 2000; Demeyer et al., 2008; Santarelli et al., 2008; Ferguson, 2010). It was showed that haeme iron may catalyse the formation of NOc from natural precursors in the gastrointestinal tract (Santarelli et al., 2008). Haeme iron may play a critical role in understanding links between red meat and particularly colorectal cancer risk, since the higher haeme iron content is one of the major differences between red meats and white meats including poultry and fish, consumption of which is not associated with increased risk of cancer (Egeberg, 2013). Iron increases cell proliferation in the mucosa, through lipoperoxidation and/or cytotoxicity of faecal water (Sesink et al., 1999; Ferguson, 2010). Furthermore, haeme iron may act as a prooxidant, and induce lipid oxidation and also cause DNA damage (Tappel, 2007; Egeberg, 2013). While some studies found iron to be carcinogenic metal, other authors consider that it is more probable that iron has a co-carcinogenic effect (Huang, 2003; Ferguson, 2010).

The method of cooking red meat also influences the risk of cancer (Berjia et al., 2014). Although meat cooking, especially at high temperatures, reduces the risk of foodborne diseases, this practice causes formation of chemical compounds including carcinogens and mutagens (Navarro et al., 2004;

WHO/FAO, 2004; Jägerstad and Skog, 2005; Badry, 2010; Aaslyng et al., 2013). The most common are HAAs and PAHs (Shin et al., 2007; Kimura et al., 2007; Hogg, 2007; Larsson et al., 2009; Ferguson, 2010; Berjia et al., 2014; Boskovic et al., 2015). These genotoxic substances act directly on DNA, cause point mutations, deletions, insertions and initiate carcinogenesis process for a number of cancers (Baena Ruiz and Salinas Hernández, 2014; Baltic and Boskovic, 2015; Boskovic et al., 2015). Their formation depends on meat type, temperature, and method of cooking (barbecuing/grilling, frying/broiling, roasting/baking) (Jägerstad and Skog, 2005; Badry, 2010; Aaslyng et al., 2013; Berjia et al., 2014; Boskovic et al., 2015). High amounts of 2-amino-1-methyl-6-phenylimidazo [4,5-b] pyridine are found in very well cooked chicken and bacon, while 2-amino-1,7-dimethylimidazo [4,5-g] quinoxaline was found in very well done pan-fried beef and steak, and in beef gravy (Ferguson, 2010). A recent study conducted by Berjia et al. (2014) showed that barbecued meat is associated with the formation of the highest concentrations of HAAs and PAHs, followed by fried meat and roasted red meat. Time of cooking has a great role in the HAAs production. Thus meat cooked for extended period mainly at higher temperatures contains a higher concentration of HAAs, which is why very well cooked meat is associated with cancer risk (Knize et al., 1994; Navarro et al., 2004; Sugimura et al., 2004; Felton and Knize, 2006; Ni et al., 2008; Ferguson, 2010).

The type of marinade also affects the formation of HAAs in meat (Joshi et al., 2015). Results obtained from some studies showed that use of Asian-type marinades reduces HAA formation, while BBQ (barbecue) sauce marinade increased formation of these compounds (Nerurkar et al. 1999; Viegas et al., 2012).

Red meat and colorectal cancer

Colorectal cancer is the third most common cancer in men and the second most common cancer in women, but it is also the cancer type most often associated with red meat consumption (Ishibe et al., 2002; Le Marchand et al., 2002; Butler et al., 2003; Gunter et al., 2005; Wu et al., 2006; Cross et al., 2007; Wyness et al., 2011; Kim et al., 2013; Berjia et al., 2014).

In the meta-analysis conducted by Norat et al. (2002) based on case-control and cohort studies published between 1973 and 1999, it was concluded that high consumption of red and processed meat was associated with an increased risk of colorectal

cancer, while total meat intake, including poultry intake, was not associated with increased cancer risk. In a meta-analysis involving almost 8,000 cases from 19 prospective studies, *Larsson and Wolk (2006)* found consistent associations between high consumption of red and processed meat and an increased colorectal cancer risk. In this meta-analysis, they also found that high intake of red and processed meat is most associated with an increase of rectal cancer, followed by distal and proximal colon cancer, respectively. The relationship between red and processed meat consumption and increase the risk of colorectal cancer was also reported by *Norat et al. (2005)*. *Chan et al. (2011)* reported that consumption of red and processed meats, assessed separately, were associated with increased risk of colorectal and colon cancers, but no statistically significant association with rectal cancer was found. This finding was different than those observed when red and processed meats were combined into a single food item.

A meta-analysis of 14 prospective studies where 725,258 subjects and 7,743 cases of colorectal cancer were evaluated, during 5 to 20 years, found no association between red meat and colorectal cancer (*Cho and Smith-Warner, 2004; McNeill and Van Elswyk, 2012*). Also, findings from a prospective study conducted by *Oba et al. (2006)* in Japan showed no association between red meat consumption and colon cancer risk, while a high intake of processed meat was associated with increased risk. These studies indicate that fresh meat *per se* is not carcinogenic and that, as well as other types of cancer, the risk of colorectal cancer is mainly associated with the method of cooking and carcinogens produced during meat processing and preparation. Higher risks were observed for darkly browned surfaces produced by barbecuing or iron pan cooking at temperatures above 150°C, and this risk was explained due to the formation of HAAs, while there was no association found with roasted and boiled meat (*Navarro et al., 2004*). The incidence of meat consumption also has an impact on risk of colorectal cancer. *Norat et al. (2002)* found that consumption of 120 g per day of red meat (compared with no consumption) was associated with a 24% increase in risk, while consumption of 30 g per day of processed meat (compared with no consumption) increased the risk by 36% (*Williamson et al., 2005*). Results obtained from thirteen studies by *Sandhu et al. (2001)* showed that a daily increase of 100 g in total or red meat intake was associated with a significant 12 to 17% increase in colorectal cancer risk (*Williamson et al., 2005*). Results from meta-analysis conducted by *Smolińska and Paluszkiwicz (2010)* showed that the frequency of red meat consumption is a crucial

risk factor for colon and rectum carcinogenesis, even more important than the amount of consumed meat. In the same study, it was found that risk of rectal and colon cancer increased with consumption of red meat more than once daily and that red meat intake in amounts higher than 50 g day⁻¹ could lead to colon cancer, while the dependence between red meat consumption of over 50 g daily and rectal cancer was not conclusive (*Smolińska and Paluszkiwicz, 2010*). Moreover, some studies showed that exposure to meat carcinogens increase the risk of colorectal cancer only in genetically susceptible individuals (*Le Marchand et al., 2002*).

Apart from haeme iron, PAHs and HAAs discussed above, some findings suggest that changes in gut microbiota also influenced colorectal cancer risk. Bacterial fermentation induced by large amounts of undigested proteins and compounds formed from amino acids during bacterial metabolism caused changes in colon epithelial homeostasis which may be predisposing factors for colorectal cancer (*Kim et al., 2013*). Further, some studies showed that meat from different origins had differing impacts on colorectal cancer. *Egeberg et al. (2013)* did not find an association between all red meat consumption and risk of colon cancer, but they observed that intake of higher amounts of beef and lamb is related to elevated colorectal cancer risk. This could be explained due to the higher amount of haeme iron in lamb and beef compared to other types of meat.

Red meat and other types of cancer

Breast cancer (*Cho et al., 2006; Steck et al., 2007; Larsson et al., 2009; Berjia et al., 2014*), gastric cancer (*Zhu et al., 2013*) prostate cancer (*Cross et al., 2005; Koutros et al., 2008; John et al., 2011; Berjia et al., 2014*), pancreatic cancer (*Anderson et al., 2002; Li et al., 2007; Stolzenberg-Solomon et al., 2007; Berjia et al., 2014*), lung cancer (*Dosil-Diaz et al., 2007*), laryngeal cancer (*Oreggia et al., 2001; Bosetti et al., 2002*) as well as bladder, kidney, and endometrial tumours are also associated with meat intake (*Ferguson, 2010; Béjar et al., 2012; Baena Ruiz and Salinas Hernández, 2014*). Scientific data about the association between meat consumption and breast cancer are inconsistent. Breast cancer is associated with some meat components including meat-derived haeme iron, NOC, HAAs and PAHs (*Cho et al., 2006; Kabat and Rohan, 2007; Lauber and Gooderham, 2007; Kallianpur et al., 2008; Larsson et al., 2009*). It is important to mention that oestrogen receptor-positive breast tumours are etiologically different from receptor-negative tumours.

Red meat consumption could increase the risk, especially of hormone receptor-positive breast cancers. This was confirmed in Nurses' Health Study II which showed that a high intake of red meat by premenopausal women was associated with a significantly increased risk of ER+/PR+ breast cancer but not ER-/PR- breast cancer (Cho et al., 2006; Larsson et al., 2009). Larsson et al. (2009) conducted an epidemiologic study on 61,433 Swedish women to try to determine if there was any association of meat intake with incidence of breast cancer as defined by oestrogen receptor and progesterone receptor status. During 17.4 years, 2,952 incident cases of invasive breast cancer were determined, but statistical analyses showed that consumption of neither fresh red meat nor processed red meat had any significant association with risk of breast cancer. Missmer et al. (2002) also found no significant association between total meat, red or white meat consumption (all considered separately) and breast cancer risk. These results are in accordance with those of the NIH-AARP Diet and Health Study as well as other studies (Fraser, 1989; Ambrosone et al., 1998; Gertig, et al., 1999; Missmer et al., 2002; Larsson et al., 2009).

Along with smoking and drinking alcohol, consumption of red or processed meat is considered to be a predisposing factor for gastric cancer (González et al., 2013). High levels of salt are used in the production of some processed meats, and there are data that salt and salt-rich foods probably increase the risk of gastric cancer (D'Elia et al., 2012; González et al., 2013) which can explain the link between meat products and gastric cancer. Meta-analysis based on epidemiological studies showed that consumption of red and/or processed meat increased gastric cancer risk (Zhu et al., 2013). However, the same authors noted that further investigation should be conducted in order to confirm the association, especially for red meat. Other authors did not find association between red and processed meat consumption with increased risk of gastric cancer, but they reported increased risk for oesophageal squamous cell carcinoma (Keszei et al., 2012).

In a population-based case-control study Anderson et al. (2002) did not find a significant association between pancreas cancer and grilled/barbecued or fried red meat intake, but when fried meat was consumed, the risk of cancer was elevated. These results are consistent with those reported by Bosetti et al. (2013), while Polesel et al. (2010) found that eating red meat is associated with pancreatic cancer.

In studies carried by Brennan et al. (2000) and Alavanja et al. (2001), it was found that meat intake

influenced the incidence of lung cancer. Contrary to these results, Norat et al., (2005) found that consumption of pork meat has a positive effect on lung cancer. This result was not confined only to meat *per se*, but to the fact that this meat is often eaten with green leafy vegetables, and often tends to be consumed with red wine, which contains resveratrol and tannins and has a protective effect.

An association between red meat and laryngeal cancer was reported by Oreggia et al. (2001). They reported that red meat and total meat intake were associated with a strong increase in laryngeal cancer risk, but this effect disappeared after adjusting for total fat intake. In the same study, HAAs had an impact on laryngeal cancer risk, but no association with salted meat was reported, although some authors did find that connection and attributed it to the presence of nitrosamines (Oreggia et al., 2001). This association between red or processed meat and risk of laryngeal cancer was confirmed in the study conducted by Bosetti et al. (2002).

Moreover, a meta-analysis of 15 prospective studies found no positive association between consumption of red or processed meat and prostate cancer risk (Alexander et al., 2010).

Discussion and recommendations

Although consumption of red meat could be associated with some types of cancer, the fact that meat contains anticancer components which are also essential in human nutrition, such as selenium, zinc, omega-3 fatty acids, vitamins B₆, B₁₂, D and folic acid, indicates that meat should not be eliminated from the human diet, but meat intake should be reduced to the recommended level of up to 500g per week (cooked weight) and little, if any, processed meat (*World Cancer Research Fund/American Institute for Cancer Research*, 2007; Chan et al., 2011; Pérez-Cueto and Verbeke, 2012; *World Cancer Research Fund*, 2012; Baena Ruiz and Salinas Hernández, 2014; Baltic and Boskovic, 2015). Moreover, some peptides obtained from meat and fish possess anti-cancer properties, inhibit cell proliferation and have cytotoxic effects against tumour cells (Shahidi and Zhong, 2008; Ryan et al., 2011; Najafian and Babji, 2012; Udenigwe and Aluko, 2012; Baltic et al., 2014). Red meat also contains other compounds with beneficial effects on human health, including carnitine, coenzyme Q10 and creatinine (Schmid, 2009). Moreover, red meat, especially beef, has a better ratio of n6:n3 fatty acids and more vitamins (A, B₆, and B₁₂) and mineral substances (Fe and Zn) than does white meat

(Oostindjer *et al.*, 2014). Many authors agree that red meat is not carcinogenic *per se* and that risk of cancer is influenced by cooking practices and dietary patterns in which meat is consumed (Biesalski, 2005; Ferguson, 2010). Cooking methods such as barbecuing and frying should be avoided, as well as excessive external brownness or charring of meat during preparation, while baking, boiling, and stewing do not produce high levels of carcinogens (HAAs and PAHs) (Anderson *et al.*, 2002). Also, it was established that not only nitrate but also other chemical additives used in meat products to extend shelf life have adverse effects on human health and there is a need for more natural additives, such as natural plant extracts. Plant extracts exhibit antimicrobial and antioxidative properties, have positive effects on meat safety and quality and also prolong shelf life (Boskovic *et al.*, 2013). Phytochemicals reduce the formation of nitrosamines in the human body when consumed simultaneously with meat products (Chung *et al.*, 2013; Oostindjer *et al.*, 2014). It is suggested that addition of garlic or curcumin have cancer protective effects (Shu *et al.*, 2010; El-Bayoumy *et al.*, 2011; Oostindjer *et al.*, 2014), which is why it could be beneficial if they can be added to a marinade or in meat dishes.

Since in general, civilisation diseases are influenced by different factors, eating and lifestyle habits should be considered as the main predisposing factors for various diseases, including cancer. People's habits and diets have changed during recent decades. It has been observed in various studies that individuals who consume high amounts of red or processed meat can also commonly consume more energy rich food products, including sugar-rich drinks and condiments, eat less vegetables, drink more alcohol, take less vitamins and are less physically active (Alexander, 2013; Oostindjer *et al.*, 2014).

Conclusion

Although lately associated with number of diseases and conditions, because of the nutritional value of red meat, as well as the presence of minerals and vitamins, red meat still has a useful role in the human diet, and positively impacts on human health when it is consumed in moderate amounts after recommended cooking methods, and within the context of a well balanced diet.

Acknowledgment: This paper was supported by the Ministry of Education, Science and Technological Development, Republic of Serbia, through the funding of Project “Selected biological hazards to the safety/quality of food of animal origin and the control measures from farm to consumer” (TR 31034).

References

- Aaslyng, M. D., Duedahl-Olesen, L., Jensen, K. & Meinert, L. (2013). Content of heterocyclic amines and polycyclic aromatic hydrocarbons in pork, beef and chicken barbecued at home by Danish consumers. *Meat Science*, 93, 85–91.
- Alavanja, M. C., Field, R. W., Sinha, R., Brus, C. P., Shavers, V. L., Fisher, E. L. & Lynch, C. F. (2001). Lung cancer risk and red meat consumption among Iowa women. *Lung Cancer*, 34 (1), 37–46.
- Alexander, D. D., Mink, P. J., Cushing, C. A. & Scurman, B. (2010). A review and meta-analysis of prospective studies of red and processed meat intake and prostate cancer. *Nutrition Journal*, 9 (1), 50.
- Alexander, D. D. (2013). No association between meat intake and mortality in Asian countries. *The American journal of clinical nutrition*, 98 (4), 865–866.
- Ambrosone, C. B., Freudenheim, J. L., Sinha, R., Graham, S., Marshall, J. R., Vena, J. E. & Shields, P. G. (1998). Breast cancer risk, meat consumption and N-acetyltransferase (NAT2) genetic polymorphisms. *International Journal of Cancer*, 75 (6), 825–830.
- Anderson, K. E., Sinha, R., Kulldorff, M., Gross, M., Lang, N. P., Barber, C., Hanack, L., DiMaggio, E., Bliss, R. & Kadlubar, F. F. (2002). Meat intake and cooking techniques: associations with pancreatic cancer. *Mutation Research*, 506–507, 225–231.
- Badry, N. E. L. (2010). Effect of household cooking methods and some food additives on polyaromatic hydrocarbons (PAHs) formation in chicken meat. *World Applied Sciences Journal*, 9, 963–974.
- Baena Ruiz, R. & Salinas Hernández, P. (2014). Diet and cancer: Risk factors and epidemiological evidence. *Maturitas*, 77, 202–208.
- Baltic, M. Z. & Boskovic, M. (2015). When man met meat: meat in human nutrition from ancient times till today. *Procedia Food Science*, 5, 6–9.
- Baltic, M. Z., Boskovic, M., Ivanovic, J., Janjic, J., Dokmanovic, M., Markovic, R. & Baltic, T. (2014). Bioactive peptides from meat and their influence on human health. *Tehnologija Mesa*, 55 (1), 8–21.

- Béjar, L. M., Gili, M., Infantes, B. & Marcott, P. F. (2002). Incidence of colorectal cancer and influence of dietary habits in fifteen European countries from 1971 to 2002. *Gaceta Sanitaria*, 26 (1), 69–73.
- Berjia, F. L., Poulsen, M. & Nauta, M. (2014). Burden of diseases estimates associated to different red meat cooking practices. *Food and Chemical Toxicology*, 66, 237–244.
- Biesalski, H. K. (2005). Meat as a component of a healthy diet—are there any risks or benefits if meat is avoided in the diet? *Meat Science*, 70 (3), 509–524.
- Bosetti, C., Bravi, F., Turati, F., Edefonti, V., Polesel, J., Decarli, A. & Zeegers, M. P. (2013). Nutrient-based dietary patterns and pancreatic cancer risk. *Annals of Epidemiology*, 23 (3), 124–128.
- Bosetti, C., La Vecchia, C., Talamini, R., Negri, E., Levi, F., Dal Maso, L. & Franceschi, S. (2002). Food groups and laryngeal cancer risk: A case-control study from Italy and Switzerland. *International Journal of Cancer*, 100 (3), 355–360.
- Boskovic, M., Baltic, M. Z., Ivanovic, J., Djuric, J., Dokmanovic, M., Markovic, R., Sarcevic, D. & Baltic, T. (2015). The impact of pork meat and lard on human health. *Tehnologija Mesa*, 56 (1), 8–15.
- Boskovic, M., Baltic, M. Z., Ivanovic, J., Djuric, J., Loncina, J., Dokmanovic, M. Markovic, R. (2013). Use of essential oils in order to prevent foodborne illnesses caused by pathogens in meat. *Tehnologija Mesa*, 54 (1), 14–20.
- Brennan, P., Fortes, C., Butler, J., Agudo, A., Benhamou, S., Darby, S. & Boffetta, P. (2000). A multicenter case-control study of diet and lung cancer among non-smokers. *Cancer Causes and Control*, 11 (1), 49–58.
- Butler, L. M., Sinha, R., Millikan, R. C., Martin, C. F., Newman, B., Gammon, M. D., Ammerman, A. S. & Sandler, R. S. (2003). Heterocyclic amines, meat intake, and association with colon cancer in a population-based study. *The American Journal of Epidemiology*, 157, 434–445.
- Carrera-Bastos, P., Fontes-Villalba, M., O’Keefe, J. H., Lindberg, S. & Cordain, L. (2011). The western diet and lifestyle and diseases of civilization. *Research Reports in Clinical Cardiology*, 2, 15–35.
- Chan, D. S., Lau, R., Aune, D., Vieira, R., Greenwood, D. C., Kampman, E. & Norat, T. (2011). Red and processed meat and colorectal cancer incidence: meta-analysis of prospective studies. *PloS One*, 6 (6), 20456.
- Cho, E. & Smith-Warner, S. A. (2004). Meat and fat intake and colorectal cancer risk: a pooled analysis of 14 prospective studies. *Cancer Research*, 64 (7 Supplement), 113–113.
- Cho, E., Chen, W. Y., Hunter, D. J., Stampfer, M. J., Colditz, G. A., Hankinson, S. E. & Willett, W. C. (2006). Red meat intake and risk of breast cancer among premenopausal women. *Archives of Internal Medicine*, 166 (20), 2253–2259.
- Chung, M. Y., Lim, T. G. & Lee, K. W. (2013). Molecular mechanisms of chemopreventive phytochemicals against gastroenterological cancer development. *World Journal of Gastroenterology*, 19 (7), 984–993.
- Corpet, D. E. (2011). Red meat and colon cancer: should we become vegetarians, or can we make meat safer? *Meat Science*, 89 (3), 310–316.
- Cross, A. J., Leitzmann, M. F., Gail, M. H., Hollenbeck, A. R. & Schatzkin, A. (2007). A prospective study of red and processed meat intake in relation to cancer risk. *PLOS Medicine*, 4, 325.
- Cross, A. J., Peters, U., Kirsh, V. A., Andriole, G. L., Reding, D., Hayes, R. B. & Sinha, R. (2005). A prospective study of meat and meat mutagens and prostate cancer risk. *Cancer Research*, 65, 11779–11784.
- D’Elia, L., Rossi, G., Ippolito, R., Cappuccio, F. P. & Strazzullo, P. (2012). Habitual salt intake and risk of gastric cancer: a meta-analysis of prospective studies. *Clinical Nutrition*, 31 (4), 489–498.
- Demeyer, D., Honikel, K. & De Smet, S. (2008). The World Cancer Research Fund report 2007: A challenge for the meat processing industry. *Meat Science*, 80 (4), 953–959.
- Dosil-Diaz, O., Ruano-Ravina, A., Gestal-Otero, J. J. & Barros-Dios, J. M. (2007). Meat and fish consumption and risk of lung cancer: a case-control study in Galicia, Spain. *Cancer Letters*, 252 (1), 115–122.
- Egeberg R., Olsen, A., Christensen, J., Halkjær, J., Jacobsen, M. U., Overvad, K. & Tjønneland, A. (2013). Associations between red meat and risks for colon and rectal cancer depend on the type of red meat consumed. *The Journal of Nutrition*, 143 (4), 464–472.
- El-Bayoumy, K., Sinha, R., Cooper, A. J. & Pinto, J. T. (2011). The Role of alliums and their sulfur and selenium constituents in cancer prevention. In *Vegetables, Whole Grains, and Their Derivatives in Cancer Prevention* (pp. 91–121). Springer, Netherlands.
- Felton, J. S. & Knize, M. G. (2006). A meat and potato war: Implications for cancer etiology. *Carcinogenesis*, 27 (12), 2367–2370.
- Ferguson, L. R. (2010). Meat and cancer. *Meat Science*, 84 (2), 308–313.
- Fraser, G. E. (1989). Dietary habits and breast cancer incidence among Seventh-day Adventists. *Cancer*, 64, 582–590.
- Gertig, D. M., Hankinson, S. E., Hough, H., Spiegelman, D., Colditz, G. A., Willett, W. C. & Hunter, D. J. (1999). N-acetyl transferase 2 genotypes, meat intake and breast cancer risk. *International Journal of Cancer*, 80 (1), 13–17.
- González, C. A., Sala, N. Rokkas, T. (2013). Gastric cancer: epidemiologic aspects. *Helicobacter*, 18 (s1), 34–38.
- Gunter, M. J., Probst-Hensch, N. M., Cortessis, V. K., Kulldorff, M., Haile, R. W. & Sinha, R. (2005). Meat intake, cooking-related mutagens and risk of colorectal adenoma in a sigmoidoscopy-based case-control study. *Carcinogenesis*, 26, 637–642.
- Higgs, J. D. (2000). The changing nature of red meat: 20 years of improving nutritional quality. *Trends in Food Science & Technology*, 11 (3), 85–95.
- Hogg, N. (2007). Red meat and colon cancer: heme proteins and nitrite in the gut. A commentary on “diet-induced endogenous formation of nitroso compounds in the GI tract”. *Free Radical Biology and Medicine*, 43 (7), 1037–1039.
- Huang, X. (2003). Iron overload and its association with cancer risk in humans: evidence for iron as a carcinogenic metal. *Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis*, 533 (1), 153–171.
- <http://apps.who.int/gho/data/node.main.A864>
- <http://www.who.int/mediacentre/factsheets/fs297/en/>
- International Agency for Research on Cancer (2015). Working Monograph. Carcinogenicity of Consumption of Red and Processed Meat.
- Ishibe, N., Sinha, R., Hein, D. W., Kulldorff, M., Strickland, P., Fretland, A. J., Chow, W. H., Kadlubar, F. F., Lang, N. P. & Rothman, N. (2002). Genetic polymorphisms in heterocyclic amine metabolism and risk of colorectal adenomas. *Pharmacogenetics*, 12, 145–150.

- Jägerstad, M. & Skog, K. (2005). Genotoxicity of heat-processed foods. *Mutation Research*, 574, 156–172.
- John, E. M., Stern, M. C., Sinha, R. & Koo, J. (2011). Meat consumption, cooking practices, meat mutagens, and risk of prostate cancer. *Nutrition and Cancer*, 63, 525–537.
- Joshi, A. D., Kim, A., Lewinger, J. P., Ulrich, C. M., Potter, J. D., Cotterchio, M. & Stern, M. C. (2015). Meat intake, cooking methods, dietary carcinogens, and colorectal cancer risk: findings from the Colorectal Cancer Family Registry. *Cancer Medicine*, 4 (6), 936–952.
- Kabat, G. C. & Rohan, T. E. (2007). Does excess iron play a role in breast carcinogenesis? An unresolved hypothesis. *Cancer Causes and Control*, 18, 1047–1053.
- Kallianpur, A. R., Lee, S. A. & Gao, Y. T. (2008). Dietary animal-derived iron and fat intake and breast cancer risk in the Shanghai Breast Cancer Study. *Breast Cancer Research and Treatment*, 107, 123–132.
- Keszei, A. P., Schouten, L. J., Goldbohm, R. A. & Van Den Brandt, P. A. (2012). Red and processed meat consumption and the risk of esophageal and gastric cancer subtypes in The Netherlands Cohort Study. *Annals of Oncology*, 23, 2319–2326.
- Kim, E., Coelho, D. & Blachier, F. (2013). Review of the association between meat consumption and risk of colorectal cancer. *Nutrition Research*, 33 (12), 983–994.
- Kimura, Y., Kono, S., Toyomura, K., Nagano, J., Mizoue, T., Moore, M. A., Mibu, R., Tanaka, M., Kakeji, Y., Maehara, Y., Okamura, T., Ikejiri, K., Futami, K., Yasunami, Y., Maekawa, T., Takenaka, K., Ichimiya, H. & Imaizumi, N. (2007). Meat, fish and fat intake in relation to subsite-specific risk of colorectal cancer: the Fukuoka colorectal cancer study. *Cancer Science*, 98, 590–597.
- Knize, M. G., Dolbeare, F. A., Carroll, K. L., Moore, D. H. & Felton, J. S. (1994). Effect of cooking time and temperature on the heterocyclic amine content of fried beef patties. *Food and Chemical Toxicology*, 32, 532–595.
- Koutros, S., Cross, A. J., Sandler, D. P., Hoppin, J. A., Ma, X., Zheng, T., Alavanja, M. C. & Sinha, R. (2008). Meat and meat mutagens and risk of prostate cancer in the Agricultural Health Study. *Cancer Epidemiology Biomarkers and Prevention*, 17, 80–87.
- Larsson, S. C. & Wolk, A. (2006). Meat consumption and risk of colorectal cancer: a meta-analysis of prospective studies. *International Journal of Cancer*, 119 (11), 2657–2664.
- Larsson, S.C., Bergkvist, L. & Wolk, A. (2009). Long-term meat intake and risk of breast cancer by oestrogen and progesterone receptor status in a cohort of Swedish women. *European Journal of Cancer*, 45 (17), 3042–3046.
- Lauber, S. N. & Gooderham, N. J. (2007). The cooked meat derived genotoxic carcinogen 2-amino-3-methylimidazo [4, 5-b] pyridine has potent hormone-like activity: mechanistic support for a role in breast cancer. *Cancer Research*, 67, 9597–9602.
- Le Marchand, L., Hankin, J. H., Pierce, L. M., Sinha, R., Nerurkar, P. V., Franke, A. A. & Chang, W. (2002). Well-done red meat, metabolic phenotypes and colorectal cancer in Hawaii. *Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis*, 506, 205–214.
- Li, D., Day, R. S., Bondy, M. L., Sinha, R., Nguyen, N. T., Evans, D. B., Abbruzzese, J. L. & Hassan, M. M. (2007). Dietary mutagen exposure and risk of pancreatic cancer. *Cancer Epidemiology Biomarkers and Prevention*, 16, 655–661.
- Lijinsky, W. (1999). N-Nitroso compounds in the diet. *Mutation Research*, 443 (1–2), 129–138.
- McNeill, S. & Van Elswyk, M. E. (2012). Red meat in global nutrition. *Meat Science*, 92 (3), 166–173.
- Missmer, S. A., Smith-Warner, S. A., Spiegelman, D., Yaun, S. S., Adami, H. O., Beeson, W. L. & Hunter, D. J. (2002). Meat and dairy food consumption and breast cancer: a pooled analysis of cohort studies. *International Journal of Epidemiology*, 31 (1), 78–85.
- Najafian, L. & Babji, A. S. (2012). A review of fish-derived antioxidant and antimicrobial peptides: their production, assessment, and applications. *Peptides*, 33 (1), 178–185.
- Navarro, A., Muñoz, S. E., Lantieri, M. J., del Pilar Diaz, M., Cristaldo, P. E., de Fabro, S. P. & Eynard, A. R. (2004). Meat cooking habits and risk of colorectal cancer in Cordoba, Argentina. *Nutrition*, 20 (10), 873–877.
- Nerurkar, P. V., Marchand, L. L. & Cooney, R. V. (1999). Effects of marinating with Asian marinades or western barbecue sauce on PhIP and MeIQx formation in barbecued beef. *Nutrition and Cancer*, 34 (2), 147–152.
- Ni, W., McNaughton, L., LeMaster, D. M., Sinha, R. & Turesky, R. J. (2008). Quantization of 13 heterocyclic aromatic amines in cooked beef, pork, and chicken by liquid chromatography electro spray ionization/tandem mass spectrometry. *Journal of Agricultural and Food Chemistry*, 56 (1), 68–78.
- Norat, T., Bingham, S., Ferrari, P., Slimani, N., Jenab, M., Mazuir, M. & Riboli, E. (2005). Meat, fish, and colorectal cancer risk: the European Prospective Investigation into cancer and nutrition. *Journal of the National Cancer Institute*, 97 (12), 906–916.
- Norat, T., Lukanova, A. & Ferrari, P. (2002). Meat consumption and colorectal cancer risk: a dose–response meta-analysis of epidemiological studies. *International Journal of Cancer*, 98, 241–256.
- Oostindjer, M., Alexander, J., Amdam, G. V., Andersen, G., Bryan, N. S., Chen, D. & Karlsson, A. H. (2014). The role of red and processed meat in colorectal cancer development: a perspective. *Meat Science*, 97 (4), 583–596.
- Oreggia, F., De Stefani, E., Boffetta, P., Brennan, P., Deneo-Pellegrini, H. & Ronco, A. L. (2001). Meat, fat and risk of laryngeal cancer: a case-control study in Uruguay. *Oral Oncology*, 37 (2), 141–145.
- Pérez-Cueto, F. J. & Verbeke, W. (2012). Consumer implications of de WCRF’ permanent update on colorectal cancer. *Meat Science*, 90, 977–978.
- Polesel, J., Talamini, R., Negri, E., Bosetti, C., Boz, G. & Lucifora, E. (2010). Dietary habits and risk of pancreatic cancer: an Italian case-control study. *Cancer Causes and Control*, 21, 493–500.
- Ryan, J. T., Ross, R. P., Bolton, D., Fitzgerald, G. F. & Stanton, C. (2011). Bioactive peptides from muscle sources: meat and fish. *Nutrients*, 3 (9), 765–791.
- Sandhu, M. S., White, I. R. & McPherson, K. (2001). Systematic review of the prospective cohort studies on meat consumption and colorectal cancer risk: a meta-analytical approach. *Cancer Epidemiology Biomarkers and Prevention*, 10, 439–446.
- Santarelli, R. L., Pierre, F. & Corpet, D. E. (2008). Processed meat and colorectal cancer: A review of epidemiologic and experimental evidence. *Nutrition and Cancer*, 60 (2), 131–144.

- Sesink, A. L., Termont, D. S., Kleibeuker, J. H. & Van der Meer, R. (1999). Red meat and colon cancer: The cytotoxic and hyperproliferative effects of dietary heme. *Cancer Research*, 59 (22), 5704–5709.
- Shahidi, F. & Zhong, Y. (2008). Bioactive peptides. *Journal of AOAC International*, 91 (4), 914–931.
- Shin, A., Shrubsole, M. J., Ness, R. M., Wu, H., Sinha, R., Smalley, W. E. & Shyr, Y., Zheng, W. (2007). Meat and meat-mutagen intake, doneness preference and the risk of colorectal polyps: the Tennessee colorectal polyp study. *International Journal of Cancer*, 121 (1), 136–142.
- Shu, L., Cheung, K. L., Khor, T. O., Chen, C. & Kong, A. N. (2010). Phytochemicals: cancer chemoprevention and suppression of tumor onset and metastasis. *Cancer and Metastasis Reviews*, 29 (3), 483–502.
- Schmid, A. (2009). Bioactive substances in meat and meat products. *Fleisch Wirtschaft*, 89, 83–90.
- Smolinska, K. & Paluszkiwicz, P. (2010). Risk of colorectal cancer in relation to frequency and total amount of red meat consumption. Systematic review and meta-analysis. *Archives of Medical Science*, 6 (4), 605–610.
- Steck, S. E., Gaudet, M. M., Eng, S. M., Britton, J. A., Teitelbaum, S. L., Neugut, A. I., Santella, R. M. & Gammon, M. D. (2007). Cooked meat and risk of breast cancer—lifetime versus recent dietary intake. *Epidemiology*, 18, 373–382.
- Stewart, B. W. & Wild, C. P. (2014). World Cancer Report 2014. International Agency for Research on Cancer. *World Health Organization*, 505.
- Stolzenberg-Solomon, R. Z., Cross, A. J., Silverman, D. T., Schairer, C., Thompson, F. E., Kipnis, V., Subar, A. F., Hollenbeck, A., Schatzkin, A. & Sinha, R. (2007). Meat and meat-mutagen intake and pancreatic cancer risk in the NIH-AARP cohort. *Cancer Epidemiology Biomarkers and Prevention*, 16, 2664–2675.
- Sugimura, T., Wakabayashi, K., Nakagama, H. & Nagao, M. (2004). Heterocyclic amines: Mutagens/carcinogens produced during cooking of meat and fish. *Cancer Science*, 95 (4), 290–299.
- Tappel, A. (2007). Heme of consumed red meat can act as a catalyst of oxidative damage and could initiate colon, breast and prostate cancers, heart disease and other diseases. *Medical Hypotheses*, 68 (3), 562–564.
- Udenigwe, C. C. & Aluko, R. E. (2012). Food protein-derived bioactive peptides: production, processing, and potential health benefits. *Journal of Food Science*, 77 (1), 11–24.
- Viegas, O., Amaro, L. F., Ferreira, I. M. & Pinho, O. (2012). Inhibitory effect of antioxidant-rich marinades on the formation of heterocyclic aromatic amines in pan-fried beef. *Journal of Agricultural and Food Chemistry*, 60 (24), 6235–6240.
- WHO/FAO. (2004). Risk assessment of *Listeria monocytogenes* in ready-to-eat foods. Technical Report of Microbiological Risk Assessment Series 5, WHO, Geneva.
- Williamson, C. S., Foster, R. K., Stanner, S. A. & Buttriss, J. L. (2005). Red meat in the diet. *Nutrition Bulletin*, 30 (4), 323–355.
- World Cancer Research Fund. (2012). Red and processed meat: finding the balance for cancer prevention.
- World Cancer Research Fund/American Institute for Cancer Research (2007). Food, nutrition, physical activity, and the prevention of cancer: A global perspective. Washington, DC: American Institute for Cancer Research.
- Wu, K., Giovannucci, E., Byrne, C., Platz, E.A., Fuchs, C., Willett, W.C. & Sinha, R., (2006). Meat mutagens and risk of distal colon adenoma in a cohort of US men. *Cancer Epidemiology Biomarkers and Prevention*, 15, 1120–1125.
- Wyness, L., Weichselbaum, E., O'Connor, A., Williams, E. B., Benelam, B., Riley, H. & Stanner, S. (2011). Red meat in the diet: an update. *Nutrition Bulletin*, 36 (1), 34–77.
- Zhu, H., Yang, X., Zhang, C., Zhu, C., Tao, G., Zhao, L., Tang, S., Shu, Z., Cai, J., Dai, S. & Qin, Q. (2013). Red and processed meat intake is associated with higher gastric cancer risk: a meta-analysis of epidemiological observational studies. *PloS One*, 8 (8), 70955.

Paper received: 14.10.2016.

Paper corrected: 19.10.2016.

Paper accepted: 19.10.2016.