



POLITECNICO DI TORINO
Repository ISTITUZIONALE

SO₂ deactivation mechanism of NO oxidation and regeneration of the LaCoO₃ perovskite

Original

SO₂ deactivation mechanism of NO oxidation and regeneration of the LaCoO₃ perovskite / Martinovity, F.; Tran, Q.; Deorsola, F. A.; Bensaid, S.; Palkovits, R.; Paulus, W.; Bonelli, B.; Di Renzo, F.; Pirone, R.. - In: CATALYSIS SCIENCE & TECHNOLOGY. - ISSN 2044-4753. - 10:7(2020), pp. 2193-2202.

Availability:

This version is available at: 11583/2830736 since: 2020-05-28T22:09:06Z

Publisher:

Royal Society of Chemistry

Published

DOI:10.1039/c9cy02478f

Terms of use:

openAccess

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

default

No description

(Article begins on next page)

SO₂ deactivation mechanism of NO oxidation and regeneration of the LaCoO₃ perovskite

Ferenc Martinovic^{a,b}, Quang Nguyen Tran^{a,c}, Fabio Alessandro Deorsola^{*a}, Samir Bensaid^a, Regina Palkovits^b, Werner Paulus^c, Barbara Bonelli^a, Francesco di Renzo^c, Raffaele Pirone^a

^a DISAT, Politecnico di Torino, Corso Duca degli Abruzzi 24, 10129 Torino, Italy

^b Institut für Technische und Makromolekulare Chemie, RWTH Aachen University, Worringerweg 2, 52074 Aachen, Germany

^c Institut Charles Gerhardt, Université de Montpellier-CNRS-ENSCM, Place Eugène Bataillon, 34095 Montpellier, France

*Corresponding author: fabio.deorsola@polito.it




Fig. S1 SEM images of the a) LaCoO₃ b) LaCoO₃-S and c) LaCoO₃-R5 catalysts.




Fig. S2 XPS spectra of the sputtered $\text{LaCoO}_3\text{-S}$ after calcination at 700 °C.




Fig. S3 Deactivation of LaCoO₃ as pure and mixed with Ca(OH)₂. Reaction conditions: 200 mg catalyst, 100 mg Ca(OH)₂, 600 mL/min 500 ppm NO, 4 % O₂, in N₂ and 60 ppm SO₂ when added, T = 500 °C.